

# Global solvability of a networked integrate-and-fire model of McKean-Vlasov type

François Delarue, James Inglis, Sylvain Rubenthaler, Etienne Tanré

# ▶ To cite this version:

François Delarue, James Inglis, Sylvain Rubenthaler, Etienne Tanré. Global solvability of a networked integrate-and-fire model of McKean-Vlasov type. 2012. hal-00747565v1

# HAL Id: hal-00747565 https://inria.hal.science/hal-00747565v1

Submitted on 31 Oct 2012 (v1), last revised 2 Jun 2014 (v4)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# GLOBAL SOLVABILITY OF A NETWORKED INTEGRATE-AND-FIRE MODEL OF MCKEAN-VLASOV TYPE

#### F. DELARUE, J. INGLIS, S. RUBENTHALER, E. TANRÉ

ABSTRACT. We here investigate the well-posedness of a networked integrate-andfire model describing an infinite population of neurons which interact with one another through their common statistical distribution. The interaction is of the self-excitatory type as, at any time, the potential of a neuron increases when some of the others fire: precisely, the kick it receives is proportional to the instantaneous proportion of firing neurons at the same time. From a mathematical point of view, the coefficient of proportionality, denoted by  $\alpha$ , is of great importance as the resulting system is known to blow-up for large values of  $\alpha$ . In the current paper, we focus on the complementary regime and prove that existence and uniqueness hold for all time when  $\alpha$  is small enough. The critical value for existence and uniqueness is made explicit in some popular examples in neurosciences and is also compared with numerical experiments.

KEYWORDS: McKean nonlinear diffusion process; renewal process; density estimates; integrate-and-fire model; network of neurons; neuroscience.

### 1. INTRODUCTION

The stochastic integrate-and-fire model of the membrane potential V across a neuron in the brain has received a huge amount of attention since its introduction (see [18] for a comprehensive review). The central idea is to model V by threshold dynamics, in which the potential is described by a simple linear (stochastic) differential equation up until it reaches a fixed threshold value  $V_F$ , at which point the neuron emits a 'spike'. Experimentally, at this point an action potential is observed, whereby the potential increases very rapidly to a peak (hyperpolarization phase) before decreasing quickly to a reset value (depolarization phase). It then relatively slowly increases once more to the resting potential (refractory period).

Since spikes are stereotyped events, they are fully characterized by the times at which they occur. The integrate-and-fire model is part of a family of spiking neuron models which take advantage of this by modeling only the spiking times and disregarding the nature of the spike itself. Specifically, in the integrate-and-fire model we observe jumps in the action potential as the voltage is immediately reset to a value  $V_R$  whenever it reaches the threshold  $V_F$ , which is motivated by the fact that the time period during which the action potential is observed is very small (of the order of 2-3ms). Despite its simplicity, the integrate-and-fire model has been

This work has been supported by the Agence National de la Recherche through the ANR Project MANDy "Mathematical Analysis of Neuronal Dynamics", ANR-09-BLAN-0008-01.

able to predict the spiking times of a neuron with a reasonable degree of accuracy [10, 11], and as such provides the basis for a good first study.

Many extensions of the basic integrate-and-fire model have been studied, including ones in which attempts are made to include noise and to describe the situation when many integrate-and-fire neurons are placed in a network and interact with each other. Indeed, in [13] and [14] the following equation describing how the potential  $V_i$  of the *i*-th neuron in a network of N behaves in time is proposed:

$$\frac{d}{dt}V_i(t) = -\lambda V_i(t) + \frac{\alpha}{N} \sum_j \sum_{n_j = -\infty}^{\infty} \delta(t - t_{n_j}) + \frac{\beta}{N} \sum_{j \neq i} V_j(t) + I_i^{ext}(t) + \sigma \eta_i(t) \quad (1.1)$$

for  $V_i(t) < V_F$  and where  $V_i(t)$  is immediately reset to  $V_R$  when it reaches  $V_F$ . Here  $I_i^{ext}(t)$  represents the external input current to the neuron,  $\eta_i(t)$  is the noise (a white noise) which is importantly supposed to be independent from neuron to neuron, and the constants  $\lambda, \beta, \alpha$  and  $\sigma$  are chosen according to experimental data. Moreover, the interaction term is described in terms of  $t_{n_j}$ , which is the time of the *n*-th spike of neuron *j*, and the Dirac function  $\delta$ . Precisely, it says that when ever one of the other neurons in the network spikes, the potential across neuron *i* receives a 'kick' of size  $\alpha/N$ .

In the case of a large network i.e. when N is large, many authors approximate the interaction term in the above finite system by an instantaneous rate (the socalled mean-firing rate)  $\nu(t)$  (see for example [1, 2, 14, 16]), but in the neuroscience literature little attention is paid to how this convergence is achieved. Mathematically the mean-field limit as  $N \to \infty$  must be taken, and it is a not at all clear that one observes a propagation of chaos phenomenon. This is the subject of a forthcoming manuscript [5].

Just as important is the question of whether solutions exist for the resulting nonlinear equation when the mean-field limit is taken, and if so what type of solutions are admissible. Again this is not treated rigorously in the neuroscience literature, and it is in this diction that we proceed here. Since the mathematical difficulties lie within the jump interaction term, we in fact isolate it by making two simplifications. Indeed, we suppose that there is no external input current ( $I_i^{ext}(t) \equiv 0$ ), and that the interaction term is composed solely of the jump or reset part ( $\beta = 0$ ). Although this is a big simplification from a neuroscience perspective, it still captures all the mathematical complexity of the resulting mean-field equation, and standard approaches (see for example [20]) would then allow extensions to more realistic models.

Without loss of generality, we also take the firing threshold  $V_F = 1$  and the reset value  $V_R = 0$  for notational simplicity. We finally choose to describe the problem in the language of stochastic processes, so that the precise nonlinear stochastic differential equation under study here becomes

$$X_t = X_0 + \int_0^t b(X_s)ds + \alpha \mathbb{E}(M_t) + \sigma W_t - M_t, \qquad t \ge 0, \tag{1.2}$$

where  $X_0 < 1$  almost surely,  $\alpha \in \mathbb{R}$ ,  $\sigma > 0$ ,  $W_t$  is a standard Brownian motion in  $\mathbb{R}$  and  $b : \mathbb{R} \to \mathbb{R}$  is Lipschitz continuous. The jumps, or resets, of the system are

described by

$$M_t = \sum_{k \ge 1} \mathbf{1}_{[0,t]}(\tau_k)$$

where  $(\tau_k)_{k \ge 1}$  stands for the sequence of hitting times of 1 by the process  $(X_t)_{t \ge 0}$ . That is  $(M_t)_{t \ge 0}$  counts the number of times  $X_t$  hits the threshold before time t, so that  $\mathbb{E}(M_t)$  denotes the expected number of times the threshold is reached before t. This equation corresponds to what we would expect when the limit  $N \to \infty$  is taken in (1.1) when the above simplifications are made, where the resets are now included in the equation by the presence of the counting process  $(M_t)_{t \ge 0}$ . It is indeed the case that the associated finite system of N neurons converges to this nonlinear limit equation (again this is shown in our forthcoming manuscript [5]).

In fact a less general version of equation (1.2) has been rigorously studied from the PDE viewpoint before, though not completely satisfactorily. If one applies the generalized Itô formula for jump processes to  $f(X_t)$  for a suitable function f, then in the standard way one arrives at the Fokker-Planck equation for the density  $p(t, y)dy = \mathbb{P}(X_t \in dy)$ :

$$\partial_t p(t,y) + \partial_y \left[ (b(y) + \alpha e'(t)) \, p(t,y) \right] - \frac{1}{2} \partial_{yy}^2 p(t,x) = \delta_0 (y - V_R) e'(t), \qquad y < 1,$$

where  $e(t) = \mathbb{E}(M_t)$ , subject to p(t, 1) = 0,  $p(t, -\infty) = 0$ ,  $p(0, y)dy = \mathbb{P}(X_0 \in dy)$ and  $\delta_0$  is the Dirac measure at 0. Moreover, the condition that p(t, y) must remain a probability density translates into the fact that we must have

$$e'(t) = \frac{d}{dt} \mathbb{E}(M_t) = -\frac{1}{2} \partial_y p(1, t), \qquad \forall t > 0.$$

Now, in the case when b(x) = -x, this nonlinear Fokker-Planck equation is exactly the one studied in [3] and [4]. Therein, the authors define a solution to be one in which e'(t) remains bounded, since the explosion of e'(t) corresponds to the situation where a large proportion of the neurons in the network all spike at the same time. Given this definition, the equation is shown to be uniquely solvable in the cases  $\alpha = 0$  and  $\alpha < 0$  (see Theorem 1.1 in [4]), the latter one being referred to as "self-inhibitory" in neuroscience. In the so-called "self-excitatory" framework, i.e. for  $\alpha > 0$ , existence of a solution for all time has been left open. Instead, very interestingly, a negative result was established in [3], stating that, for any  $\alpha > 0$ , it is possible to find an initial probability distribution  $\mathbb{P}(X_0 \in dy)$  such that there is a blow-up, i.e. such that  $e'(t) = \infty$  in finite time. As a remarkable point, it says that solvability in the long run may fail for possibly small values of  $\alpha$ . On the contrary, it does not come as a surprise that solutions may explode in a finite time for the prescribed values of  $V_R$  and  $V_F$  when  $\alpha \ge 1$ : indeed, if Np particles in (1.1) reach the threshold  $V_F = 1$  between some times  $t_1$  and  $t_2$  for some proportion  $p \in (0, 1]$ , then all the particles perform a jump of size greater than  $\alpha p(N-1)/N$  within the same period, which reads as a size greater than  $\alpha p$  in the ideal case  $N = \infty$  corresponding to (1.2); in particular, when  $\alpha p \ge 1$  and the reset is fixed as 0, particles keep on crossing the threshold an infinite number of times between  $t_1$  and  $t_2$ . Such a picture

indicates that the system enters a sort of "super-self-excitatory" regime when  $\alpha$  is above  $V_F - V_R$ , which matches 1 with our convention.

Here we thus investigate the case  $\alpha \in (0, 1)$ . We then contribute the associated positive and until now unknown result, by proving that, given a starting point  $X_0 = x_0$ , we can find an explicit  $\alpha$  small enough so that there is a unique solution to (1.2) (and hence to the associated Fokker-Planck equation) for all time, such that  $e'(t) = [d/dt]\mathbb{E}(M_t)$  remains bounded. The surprising difficulty of this result is reflected in the rather involved nature of our proofs.

Our result is for a general Lipschitz function b, but there are two important specific cases that we study further: the Brownian case when  $b \equiv 0$  and the Ornstein-Uhlenbeck case when  $b(x) = -\lambda x$ ,  $\lambda \ge 0$ . The Ornstein-Uhlenbeck case is most relevant to neuroscience since it is often used in the literature, but surprising difficulties remain in the purely Brownian case. In both these cases we are able to give an explicit  $\alpha_0$  depending on the starting point  $x_0$  such that (1.2) has a solution for all  $\alpha < \alpha_0$ . However, it is interesting to note that our explicit values do not appear to be optimal, in the sense that from our numerical simulations there appears to be a real 'gap' between our positive result and the negative result of [3]. Precisely, our simulations show that for a given  $x_0$  there exist classical solutions for all time for  $\alpha$  bigger than our explicit  $\alpha_0$ , while there exist solutions that blow-up that do not satisfy the conditions of [3]. Thus a very interesting, and perhaps very difficult question is to determine the true critical value  $\alpha_c$  for a given initial starting point. We discuss this further at the end of the paper.

Equation (1.2) can be thought as of McKean-Vlasov-type, since the process  $X_t$  depends on the distribution of the solution itself. However, it is highly non-standard, since it actually depends on the distribution of the *hitting times* of the threshold by the solution. This renders the traditional approaches to McKean-Vlasov equations and propagation of chaos, such as those presented by Sznitman in [20], inapplicable, because we have no *a priori* smoothness. Thus our results are also new in this context.

The general structure of the proof is quite typical of the methods used to investigate well-posedness of nonlinear systems, and especially well-posedness of Markovian stochastic differential equations (or equivalently of partial differential equations of second order) involving some non-trivial nonlinearity. Precisely, the first point is to tackle unique solvability in small time: when the parameter  $\alpha$  is (strictly) less than 1 and the mass of the initial condition decays linearly at the threshold, it is proven that the system induces a natural contraction in a well-chosen space provided the time duration is small enough. In this framework, the specific notion of solution at hand plays a crucial role as it defines the right space for the contraction. Below, solutions are sought in such a way that the mapping  $t \mapsto \mathbb{E}(M_t)$  is continuously differentiable. The proof is at the intersection between probability and PDEs. The second point is then to extend existence and uniqueness from small to long times. Exactly as for PDEs of quasilinear type, the point is to prove that some key quantity is preserved as time goes by. Here, the point is to prove that the system cannot accumulate too much mass in the right vicinity of 1. Equivalently, this amounts to showing that the Lipschitz constant of the mapping  $t \mapsto \mathbb{E}(M_t)$  cannot blow-up in a finite time. This is where the condition  $\alpha$  small enough comes in: when  $\alpha$  is small enough, we manage to give some estimates for the density of  $X_t$  in the neighbourhood of 1, the critical value of  $\alpha$  explicitly depending upon the available bound of the density. Generally speaking, we make use of standard Gaussian estimates of Aronson type of the density; in the cases  $b \equiv 0$  and  $b(x) = -\lambda x$ ,  $x \in \mathbb{R}$ , for some  $\lambda > 0$ , the bounds are explicit enough to derive a numerical  $\alpha_0$  up to which unique solvability holds. Unfortunately, the estimates we use are rather poor as they mostly neglect the right behavior of the density of  $X_t$  at the boundary, thus yielding a nonoptimal value. Anyhow, they serve as a starting point for proving a refined estimate of the gradient of the density at the boundary: this is the required ingredient for proving that, at any time t, the mass of  $X_t$  decays linearly in the neighbourhood of 1, uniformly in t in compact sets, and thus to apply the existence and uniqueness argument in small time. Such a refined estimate is proven on any interval where a solution does exist: below, it is thus referred to as an *a priori* estimate. Coupled with the result of existence and uniqueness in small time, it permits to prove by induction that existence and uniqueness hold on any finite interval and thus on the whole  $[0,\infty)$ .

The layout of the paper is as follows. In the first section below we make the necessary definitions and describe the setting in which we will be working. Section 3 is devoted to proving the existence and uniqueness of a solution to (1.2) in small time. Long-time *a priori* estimates are proven in Section 4. This section contains the crux of our argument, since we here show that, for  $\alpha$  small enough, if there exists a solution, it does not exhibit the blow-up property highlighted in [3]. In Section 5 the final step is made, whereby we show that under the *a priori* Lipschitz bound, a unique solution exists to (1.2) that does not blow-up at any time. We conclude the paper in Section 6 with a discussion of the critical value  $\alpha_c$  such that for  $\alpha < \alpha_c$  (1.2) has a solution, in both the Brownian and the Ornstein-Uhlenbeck cases, by calculating an explicit lower bound  $\alpha_0$  for  $\alpha_c$ , for fixed  $x_0$ . However, we show numerically that  $\alpha_0 < \alpha_c$ .

#### 2. Setting

As stated in the introduction, we are interested in solutions to the nonlinear McKean-Vlasov-type SDE

$$X_{t} = X_{0} + \int_{0}^{t} b(X_{s})ds + \alpha \mathbb{E}(M_{t}) + W_{t} - M_{t}, \qquad t \ge 0,$$
(2.1)

where  $X_0 < 1$  almost surely,  $\alpha \in (0, 1)$  and  $W_t$  is a standard Brownian motion with respect to a filtration  $(\mathcal{F}_t)_{t \ge 0}$  satisfying the usual conditions, with values in  $\mathbb{R}$  and starting from 0. The jumps, or resets, of the system are described by

$$M_t = \sum_{k \ge 1} \mathbf{1}_{[0,t]}(\tau_k), \qquad (2.2)$$

where the sequence of stopping times  $(\tau_k)_{k \ge 0}$  are defined by  $\tau_0 = 0$  and

$$\tau_k = \inf\{t > \tau_{k-1} : X_{t-} \ge 1\}, \quad k \ge 1.$$
(2.3)

We assume that  $b: (-\infty, 1] \to \mathbb{R}$  is Lipschitz continuous such that

$$|b(x)| \leq \Lambda(|x|+1), \qquad |b(x) - b(y)| \leq K|x-y|, \qquad \forall x, y \in (-\infty, 1].$$

**Remark 2.1.** Although we take the intensity of the noise in (2.1) to be 1 rather than a general  $\sigma > 0$  as stated in the introduction, this results in no loss of generality. Indeed, by setting  $u = t/\sigma^2$  we can rewrite (2.1) as

$$X_{\sigma^{2}u} = X_{0} + \int_{0}^{\sigma^{2}u} b(X_{s})ds + \alpha \mathbb{E}(M_{\sigma^{2}u}^{X}) + W_{\sigma^{2}u} - M_{\sigma^{2}u}^{X}, \qquad u \ge 0,$$

for an arbitrary  $\sigma > 0$ , where we have emphasized the dependence of the process Mon X by writing  $M = M^X$ . Now, by Brownian scaling we have that  $W_{\sigma^2 u} = \sigma \widetilde{W}_u$ where  $\widetilde{W}$  is another Brownian motion starting from 0. Set  $\widetilde{X}_u = X_{\sigma^2 u}$ . Then, since the number of times the process X hits the threshold before time  $\sigma^2 u$  is equal to the number of time the process  $\widetilde{X}$  hits the threshold before time u, we have that  $M_{\sigma^2 u}^X = M_u^{\widetilde{X}}$ . Hence it follows that

$$\widetilde{X}_u = X_0 + \int_0^{\sigma^2 u} b(X_s) ds + \alpha \mathbb{E}(M_u^{\widetilde{X}}) + \sigma \widetilde{W}_u - M_u^{\widetilde{X}}, \qquad u \ge 0.$$

By a change of variable we then have

$$\widetilde{X}_u = X_0 + \int_0^u \sigma^2 b(\widetilde{X}_s) ds + \alpha \mathbb{E}(M_u^{\widetilde{X}}) + \sigma \widetilde{W}_u - M_u^{\widetilde{X}}, \qquad u \ge 0.$$

Thus, if we modify the function b accordingly,  $\widetilde{X}$  satisfies exactly the equation of interest, but with now an arbitrary noise intensity.

As discussed in the introduction, the key point is to look for a solution for which  $t \mapsto \mathbb{E}(M_t)$  is continuously differentiable, which would correspond to a solution whose derivative does not exhibit a finite time blow-up. To this end we introduce the following objects:

• The space  $\mathcal{L}$ : For  $T, A \ge 0$ , let

$$\mathcal{L}(T,A) = \left\{ e \in \mathcal{C}^1[0,T] : e(0) = 0, \ e(s) \leqslant e(t) \ \forall s \leqslant t, \sup_{0 \leqslant t \leqslant T} |e'(t)| \leqslant A \right\}$$

where as usual  $\mathcal{C}^1[0,T]$  denotes the space of continuously differentiable functions on [0,T].

• The map  $\Gamma$ : For  $e \in \mathcal{L}(T, A)$  define

$$\Gamma(e)(t) := \mathbb{E}(M_t^e) \tag{2.4}$$

where

$$M_t^e = \sum_{k \ge 1} \mathbf{1}_{[0,t]}(\tau_k^e) \tag{2.5}$$

and  $\tau_k^e = \inf\{t > \tau_{k-1}^e : X_{t-}^e \ge 1\}, \tau_0 = 0$  and  $(X_t^e, M_t^e)$  is a solution to

$$X_t^e = X_0 + \int_0^t b(X_s^e) ds + \alpha e(t) + W_t - M_t^e, \qquad t \ge 0, \qquad X_0 < 1.$$
(2.6)

Note that the solution is well defined as b is Lipschitz continuous. The map  $\Gamma$  is thus defined as a map from  $\mathcal{L}(T, A)$  into the set of non-decreasing functions on [0, T]. It thus depends on A as its domain of definition depends on A; for this reason, it should be denoted by  $\Gamma^A$ . Anyhow, since the family  $(\Gamma^A)_{A \ge 0}$  is consistent in the sense that, for any  $A' \le A$ , the restriction of  $\Gamma^A$  to  $\mathcal{L}(T, A')$  coincides with  $\Gamma^{A'}$ , we can get rid of the superscript A in  $\Gamma^A$  and then use the simpler notation  $\Gamma$ .

The idea is to look for a fixed point of  $\Gamma$  in a certain space, which means that we are to find a space that is kept invariant by  $\Gamma$ . With this in mind, the following *a* priori stability result for  $\Gamma$  will be useful, the proof of which we leave until the end of the section.

**Proposition 2.2.** Given T > 0,  $e \in \mathcal{L}(T, A)$  and some real  $a \ge \mathbb{E}[(X_0)_+]$  (where  $(x)_+$  indicates the positive part of  $x \in \mathbb{R}$ ), it holds that

$$\left(\forall t \in [0, T], \ e(t) \leqslant g(t)\right) \Rightarrow \left(\forall t \in [0, T], \ \Gamma(e)(t) \leqslant g(t)\right), \tag{2.7}$$

with

$$g(t) := \frac{a + (4 + \Lambda T^{1/2})t^{1/2}}{1 - \alpha} \exp\left(\frac{2\Lambda t}{1 - \alpha}\right).$$

With thus also introduce the space  $\mathcal{H}$ , and our definition of a solution to (2.1):

• The space  $\mathcal{H}$ : For an initial value  $X_0 < 1$ , let

$$\mathcal{H}(T, A, X_0) = \left\{ e \in \mathcal{L}(T, A) : e(t) \leqslant C_T(\sqrt{t} + \mathbb{E}|X_0|)e^{\theta t} \right\},\$$

where  $C_T$  and  $\theta$  are chosen according to Proposition 2.2, that is

$$C_T := \frac{4 + \Lambda T^{1/2}}{1 - \alpha}, \quad \theta := \frac{2\Lambda}{1 - \alpha}$$

(The parameter *a* in Proposition 2.2 being chosen as  $\mathbb{E}[X_0]$ .) Equip  $\mathcal{H}(T, A, X_0)$  with the norm  $||e||_{\mathcal{H}(T,A,X_0)} = ||e||_{\infty,T} + ||e'||_{\infty,T}$ . Then  $\mathcal{H}(T,A,X_0)$  is a complete metric space, since it is a closed subspace of  $\mathcal{C}^1[0,T]$ . We set:

**Definition 2.3** (Solution to (2.1)). The coupled process  $(X_t, M_t)_{t \in [0,T]}$  will be said to be a solution to (2.1) up until time T if  $(M_t)_{t \in [0,T]}$  satisfies (2.2) subject to (2.3),  $([0,T] \ni t \mapsto \mathbb{E}(M_t)) \in \mathcal{H}(A,T,X_0)$  for some  $A \ge 0$ , and  $(X_t)_{t \in [0,T]}$  is a strong solution of (2.1) up until time T.

With the above definitions, it is clear that, for any  $A \ge 0$ , any fixed point of  $\Gamma$  that belongs to  $\mathcal{H}(A, T, X_0)$  provides a solution to (2.1), and conversely that, for any solution  $(X_t, M_t)_{t \in [0,T]}$  of (2.1),  $([0,T] \ni t \mapsto \mathbb{E}(M_t))$  must be a fixed point for  $\Gamma$  in  $\mathcal{H}(A, T, X_0)$ .

### 2.1. Proof of Proposition 2.2:

*Proof of Proposition 2.2:* Fix T > 0. We first note that we may write

$$M_t^e = \sup_{s \leqslant t} \lfloor (Z_s^e)_+ \rfloor, \quad Z_t^e = X_t^e + M_t^e, \quad t \in [0, T],$$
(2.8)

Indeed, one can see that for  $t \in [\tau_k^e, \tau_{k+1}^e), k \ge 0$ 

$$\sup_{s \leqslant t} \lfloor (Z_s^e)_+ \rfloor = \max_{0 \leqslant j \leqslant k-1} \left\{ \sup_{s \in [\tau_j^e, \tau_{j+1}^e)} \lfloor (X_s^e + j)_+ \rfloor, \sup_{s \in [\tau_k^e, t)} \lfloor (X_s^e + k)_+ \rfloor \right\}$$
$$= \max_{0 \leqslant j \leqslant k-1} \left\{ j+1, k \right\} = M_t^e,$$

using the fact that  $X_t^e < 1$  for all  $t \ge 0$ .

Then, given  $t \in [0, T]$  such that  $Z_t^e \ge 0$ , we denote by  $\rho^e := \sup\{s \in [0, t] : Z_s^e < 0\}$ (sup  $\emptyset = 0$ ). Pay attention that  $\rho^e$  is not a stopping time. Then, for  $s \in [\rho^e, t]$ ,

$$|b(X_s^e)| \leqslant \Lambda(1+|X_s^e|) \leqslant \Lambda(1+|Z_s^e|+M_s^e) = \Lambda(1+(Z_s^e)_++M_s^e).$$
(2.9)

By (2.8), we know that  $M_s^e \leq \sup_{0 \leq r \leq s} (Z_r^e)_+$ . Therefore,

$$|b(X_s^e)| \leqslant \Lambda \left(1 + 2 \sup_{0 \leqslant r \leqslant s} (Z_r^e)_+\right).$$

By (2.6), we obtain:

$$Z_t^e \leqslant Z_{\rho^e}^e + \Lambda \int_{\rho^e}^t \left( 1 + 2 \sup_{0 \leqslant r \leqslant s} (Z_r^e)_+ \right) ds + \alpha e(t) + W_t - W_{\rho^e}.$$
(2.10)

If  $\rho^e > 0$ , then  $Z^e_{\rho^e} = 0$  as, obviously,  $(Z^e_s)_{0 \leq s \leq T}$  is a continuous process. If  $\rho^e = 0$ , then  $Z^e_{\rho^e} = X_0 \ge 0$ . Therefore,

$$(Z_t^e)_+ \leqslant (X_0)_+ + \Lambda \int_0^t \left(1 + 2\sup_{0 \leqslant r \leqslant s} (Z_r^e)_+\right) ds + \alpha e(t) + 2\sup_{0 \leqslant s \leqslant t} |W_s|.$$
(2.11)

Obviously, the above inequality still holds if  $Z_t^e \leq 0$ . Therefore, taking the supremum in the left-hand side, taking the expectation and applying Doob's maximal inequality for martingales,

$$\mathbb{E}\Big[\sup_{0 \leqslant s \leqslant t} (Z_s^e)_+\Big] \leqslant \mathbb{E}[(X_0)_+] + \Lambda \int_0^t \left(1 + 2\mathbb{E}\Big[\sup_{0 \leqslant r \leqslant s} (Z_r^e)_+\Big]\right) ds + \alpha e(t) + 4t^{1/2}, \quad (2.12)$$

which shows that  $\mathbb{E}[\sup_{0 \leq t \leq T} (Z_t^e)_+]$  is finite. In particular, if

$$\forall t \in [0,T], e(t) \leq g(t), \text{ with } g(t) := \frac{\mathbb{E}[(X_0)_+] + (4 + \Lambda T^{1/2})t^{1/2}}{1 - \alpha} \exp\left(\frac{2\Lambda t}{1 - \alpha}\right),$$

and  $R^e$  stands for the deterministic hitting time:

$$R^e := \inf\left\{t \in [0,T] : \mathbb{E}\left[\sup_{0 \leqslant s \leqslant t} (Z_s^e)_+\right] > g(t)\right\} \quad (\inf \emptyset = +\infty),$$

then, for any  $t \in [0, R^e \wedge T]$ 

$$\begin{split} & \mathbb{E}\Big[\sup_{0 \leqslant s \leqslant t} (Z_s^e)_+\Big] \\ & \leqslant \mathbb{E}[(X_0)_+] + \Lambda \int_0^t (1+2g(s))ds + \alpha g(t) + 4t^{1/2} \\ & < \left(\mathbb{E}[(X_0)_+] + (4+\Lambda T^{1/2})t^{1/2}\right) \left[1 + \int_0^t \frac{2\Lambda}{1-\alpha} \exp\left(\frac{2\Lambda s}{1-\alpha}\right)ds\right] + \alpha g(t) \\ & = (1-\alpha)g(t) + \alpha g(t) = g(t). \end{split}$$

Now, by continuity of the paths of  $Z^e$  and by finiteness of  $\mathbb{E}[\sup_{0 \leq t \leq T} (Z_t^e)_+]$ , we deduce that  $\mathbb{E}[\sup_{0 \leq s \leq t} (Z_s^e)_+]$  is continuous in t. Therefore, if  $R^e \leq T$ , then  $\mathbb{E}[\sup_{0 \leq s \leq R^e} (Z_s^e)_+]$  must be equal to  $g(R^e)$ , but, by the above inequalities, this sounds as a contradiction. By (2.8), this proves the announced bound when a in the statement exactly matches  $\mathbb{E}[(X_0)_+]$ . The proof is completed in the same way when  $a > \mathbb{E}[(X_0)_+]$ .

**Remark 2.4.** In the case when  $X_0 = 0$ , we have that  $\Gamma$  exhibits the same stability on functions e such that  $e(t) \leq C_T \sqrt{t} e^{\theta t}$ .

#### 3. EXISTENCE AND UNIQUENESS IN SMALL TIME

The main result of this section is the following:

**Theorem 3.1.** Suppose there exist  $\beta, \epsilon > 0$  such that  $\mathbb{P}(X_0 \in dx) \leq \beta(1-x)dx$  for any  $x \in (1-\epsilon, 1]$ . Then there exist constants  $A_1 \geq 1$  and  $T_1 \in (0, 1]$ , depending upon  $X_0$  through  $\beta, \epsilon$  and  $\mathbb{E}|X_0|$  only, such that

$$\Gamma(\mathcal{H}(A_1, T_1, X_0)) \subset \mathcal{H}(A_1, T_1, X_0).$$

Moreover, for all  $e_1, e_2 \in \mathcal{H}(A_1, T_1, X_0)$ 

$$\|\Gamma(e_1) - \Gamma(e_2)\|_{\mathcal{H}(A_1, T_1, X_0)} \leq \frac{1}{2} \|e_1 - e_2\|_{\mathcal{H}(A_1, T_1, X_0)}$$

Hence there exists a unique fixed point of the restriction of  $\Gamma$  to  $\mathcal{H}(A_1, T_1, X_0)$ , which provides a solution to (2.1) according to Definition 2.3 up until time  $T_1$ .

3.1. **PDEs.** Given  $e \in \mathcal{L}(T, A)$ , we have

$$\Gamma(e)(t) = \mathbb{E}(M_t^e) = \sum_{k \ge 1} \int_0^t \mathbb{P}\big(\tau_{k+1}^e \in (s,t] \, | \tau_k^e = s\big) \mathbb{P}(\tau_k^e \in ds) + \mathbb{P}(\tau_1^e \leqslant t),$$

where  $\mathbb{P}(\tau_k^e \in ds)$  is a convenient abuse of notation for denoting the law of  $\tau_k^e$  and  $\mathcal{B}(\mathbb{R}) \ni A \mapsto \mathbb{P}(\tau_{k+1}^e \in A | \tau_k^e = s)$  stands for the conditional law of  $\tau_{k+1}^e$  given  $\tau_k^e = s$ . Here  $\mathcal{B}(\mathbb{R})$  is the Borel  $\sigma$ -algebra on  $\mathbb{R}$ . Observing that the solution  $X^e$  to (2.6) is a Markov process, we may write

$$\Gamma(e)(t) = \mathbb{E}(M_t^e) = \sum_{k \ge 1} \int_0^t \mathbb{P}\big(\tau_1^{e^{\sharp s}} \leqslant t - s | X_0 = 0\big) \mathbb{P}(\tau_k^e \in ds) + \mathbb{P}(\tau_1^e \leqslant t), \quad (3.1)$$

where  $e^{\sharp_s}$  stands for the mapping  $([0, T - s] \ni t \mapsto e(t + s) - e(s)) \in \mathcal{L}(T - s, A)$ . Thus in order to understand the behaviour of  $\Gamma$ , one must study the density of the killed process

$$\frac{d}{dy}\mathbb{P}(X_t^e \in dy, t < \tau_1^e).$$

Note here we have used the notation of the Radon-Nikodym derivative to denote the density of the killed process with respect to the Lebesgue measure, which makes sense thanks to the following Lemma: **Lemma 3.2.** Given an initial condition  $X_0$  satisfying the same assumption as in the statement of Theorem 3.1 together with a continuously differentiable deterministic mapping  $[0,T] \ni t \mapsto e(t)$  satisfying e(0) = 0, consider the solution  $(\chi_t)_{0 \leq t \leq T}$  of the SDE

 $d\chi_t = b(\chi_t)dt + \alpha e'(t)dt + dW_t, \quad t \in [0, T] ; \quad \chi_0 = X_0, \tag{3.2}$ 

together with the stopping time

$$\tau_1 = \inf\{t \in [0, T] : \chi_t \ge 1\}.$$

Then, for any  $t \in (0, T]$ , the measure

$$\mathcal{B}((-\infty,1]) \ni A \mapsto \mathbb{P}(\chi_t \in A, t < \tau_1)$$

has a density with respect to the Lebesgue measure. Here,  $\mathcal{B}((-\infty, 1])$  denotes the Borel subsets of  $(-\infty, 1]$ . Pay also attention that the density is not normalized. Denoting the density by

$$p(t,y) = \frac{d}{dy} \mathbb{P}(\chi_t \in dy, t < \tau_1), \quad t \in [0,T], \ y \leqslant 1,$$
(3.3)

it is continuous and continuously differentiable in y on any compact subset of  $([0,T] \times (-\infty,1]) \setminus (\{0\} \times (-\infty,1-\epsilon])$  and admits bounded Sobolev derivatives of order 1 in t and of order 2 in y on any compact subset of  $(0,T] \times (-\infty,1)$ . Moreover, it satisfies the Kolmogorov equation:

$$\partial_t p(t,y) + \partial_y \left[ \left( b(y) + \alpha e'(t) \right) p(t,y) \right] - \frac{1}{2} \partial_{yy}^2 p(t,y) = 0, \quad t \in (0,T], \ y < 1, \quad (3.4)$$

with the Dirichlet boundary condition p(t,1) = 0 and the measure-valued initial condition  $p(0,y)dy = \mathbb{P}(X_0 \in dy)$ , both p(t,y) and  $\partial_y p(t,y)$  decaying to 0 as  $y \to -\infty$ . Finally, the distribution of  $\tau_1$  has a density on [0,T], given by

$$\frac{d}{dt}\mathbb{P}(\tau_1 \leqslant t) = -\frac{1}{2}\partial_y p(t,1), \quad t \in (0,T],$$
(3.5)

the supremum norm of which can be bounded in terms of  $\alpha$ , b, T and the supremum norms of e and e' only.

Lemma 3.2 is quite standard. Anyhow, it is difficult to find a complete proof of it under our assumption. For this reason, we provide a complete proof in appendix, see Section 7. Here is an application to (3.1):

**Proposition 3.3.** Given  $e \in \mathcal{L}(T, A)$ , the mapping  $[0, T] \ni t \mapsto \Gamma(e)(t)$  is continuously differentiable. Moreover,

$$\frac{d}{dt} \big[ \Gamma(e) \big](t) = -\int_0^t \frac{1}{2} \partial_y p_e^{(0,s)}(t-s,1) \frac{d}{ds} \big[ \Gamma(e) \big](s) ds - \frac{1}{2} \partial_y p_e(t,1), \quad t \in [0,T], \quad (3.6)$$

where  $p_e$  represents the density of the process  $X^e$  killed at 1 and  $p_e^{(0,s)}$  represents the density of the process  $X^{e^{\sharp s}}$  killed at 1 with  $X_0^{e^{\sharp s}} = 0$ .

*Proof.* We first check that  $\Gamma(e)$  is Lipschitz continuous on [0, T]. Considering a finite difference in (3.1), we get, for  $t, t + h \in [0, T]$ ,

$$\Gamma(e)(t+h) - \Gamma(e)(t) = \sum_{k \ge 1} \int_{t}^{t+h} \mathbb{P}\left(\tau_{1}^{e^{\sharp s}} \le t+h-s\right) \mathbb{P}\left(\tau_{k}^{e} \in ds\right)$$
$$-\frac{1}{2} \sum_{k \ge 1} \int_{0}^{t} \int_{t-s}^{t+h-s} \partial_{y} p_{e}^{(0,s)}(r,1) dr \mathbb{P}\left(\tau_{k}^{e} \in ds\right) \qquad (3.7)$$
$$-\frac{1}{2} \int_{t}^{t+h} \partial_{y} p_{e}(s,1) ds.$$

By Lemma 3.2, we can handle the two last terms in the above to find a constant C > 0 such that

$$\Gamma(e)(t+h) - \Gamma(e)(t) \leqslant \sum_{k \ge 1} \int_{t}^{t+h} \mathbb{P}\big(\tau_1^{e^{\sharp s}} \leqslant t+h-s\big) \mathbb{P}(\tau_k^e \in ds) + Ch\big(1 + \Gamma(e)(T)\big),$$

the last term in the right-hand side being finite thanks to Proposition 2.2. Moreover, from the continuous differentiability of e, we deduce that

$$\lim_{h \searrow 0} \sup_{0 \leqslant s \leqslant T-h} \mathbb{P}\left(\tau_1^{e^{\sharp s}} \leqslant h | X_0^{e^{\sharp s}} = 0\right)$$
  
= 
$$\lim_{h \searrow 0} \sup_{0 \leqslant s \leqslant T-h} \mathbb{P}\left(\sup_{0 \leqslant r \leqslant h} Z_r^{e^{\sharp s}} \geqslant 1 | X_0^{e^{\sharp s}} = 0\right) = 0,$$
(3.8)

where  $Z^{e^{\sharp s}}$  is given by (2.8), the second line following from (2.11). Therefore, there exists a mapping  $\eta : \mathbb{R}_+ \to \mathbb{R}_+$  matching 0 at 0 and continuous at 0 such that

$$\Gamma(e)(t+h) - \Gamma(e)(t) \leqslant \eta(h) \big[ \Gamma(e)(t+h) - \Gamma(e)(t) \big] + Ch \big( 1 + \Gamma(e)(T) \big).$$

Choosing h small enough, Lipschitz continuity easily follows.

As a consequence, we can divide both sides of (3.7) by h and then let h tend to 0. By (3.8), we have for a given  $t \in [0, T)$ ,

$$\lim_{h \searrow 0} h^{-1} \sum_{k \ge 1} \int_{t}^{t+h} \mathbb{P}\left(\tau_{1}^{e^{\sharp s}} \leqslant t+h-s | X_{0}^{e^{\sharp s}} = 0\right) \mathbb{P}\left(\tau_{k}^{e} \in ds\right)$$
$$\leqslant \lim_{h \searrow 0} \left[ \sup_{0 \leqslant s \leqslant T-h} \mathbb{P}\left(\tau_{1}^{e^{\sharp s}} \leqslant h | X_{0}^{e^{\sharp s}} = 0\right) \frac{\Gamma(e)(t+h) - \Gamma(e)(t)}{h} \right] = 0.$$

Handling the second term in (3.7) by Lemma 3.2 and using the Lebesgue Dominated Convergence Theorem, we deduce that

$$\frac{d}{dt}\Gamma(e)(t) = \sum_{k \ge 1} \int_0^t \frac{d}{dt} \mathbb{P}\left(\tau_1^{e^{\sharp s}} \leqslant t - s | X_0^{e^{\sharp s}} = 0\right) \mathbb{P}\left(\tau_k^e \in ds\right) + \frac{d}{dt} \mathbb{P}\left(\tau_1^e \leqslant t\right)$$
$$= -\sum_{k \ge 1} \int_0^t \frac{1}{2} \partial_y p_e^{(0,s)}(t - s, 1) \mathbb{P}\left(\tau_k^e \in ds\right) - \frac{1}{2} \partial_y p_e(t, 1).$$

Formula (3.6) follows from the relationship

$$\Gamma(e)(t) = \sum_{k \ge 1} \int_0^t \mathbb{P}(\tau_k^e \in ds), \quad t \in [0, T].$$

$$(3.9)$$

3.2. Density estimates in short time. Suppose that  $X_0 = x_0 < 1$ . Following the parametrix method in [8, Chapter 1], one can show that the solution to (3.4) (then denoted by  $p_e^{x_0}$ ) is given by

$$p_e^{x_0}(t,y) = q(t,x_0,y) - \int_0^t \int_{-\infty}^1 \left(\alpha e'(s) + b(1)\right) \partial_z p_e^{x_0}(s,z) q(t-s,z,y) dz ds, \quad (3.10)$$

for  $t \in (0, T]$  and y < 1, where q(t, x, y) is a solution to the PDE

$$\begin{cases} \partial_t q(t, x, y) &= (1/2) \partial_{yy}^2 q(t, x, y) - \partial_y \left[ (b(y) - b(1)) q(t, x, y) \right] \\ q(t, x, 1) &= 0 \\ q(0, x, y) &= \delta_0(x - y) \end{cases}$$
(3.11)

on  $[0,T] \times (-\infty,1] \times (-\infty,1]$ . We will need the following bound on the behaviour of q.

**Proposition 3.4.** For any T > 0 there exists a constant  $B_T > 0$  such that

$$\left|\partial_{y}q(t,x,y)\right| \leqslant \frac{B_{T}}{t} \exp\left(-\frac{|\xi_{t}^{x}-y|^{2}}{B_{T}t}\right),\tag{3.12}$$

and

$$\left|\partial_{y}q(t,x,y)\right| \leqslant \frac{B_{T}}{t} \exp\left(-\frac{|x-\xi_{-t}^{y}|^{2}}{B_{T}t}\right),\tag{3.13}$$

for all  $t \in [0,T]$ ,  $x, y \leq 1$ , where

$$\xi_t^x = x + \int_0^t \widetilde{b}(\xi_r^x) dr, \quad t \in \mathbb{R},$$
(3.14)

and  $\tilde{b}(v) = b(v) - b(1)$  if v < 1 while  $\tilde{b}(v) = -\tilde{b}(2-v)$  if  $v \ge 1$ .

*Proof.* Let  $(\tilde{X}_t)_{t \ge 0}$  denote the solution of the SDE:

$$dX_t = b(X_t)dt + dW_t. (3.15)$$

We can then see that q(t, x, y) is the transition density of  $\widetilde{X}_t$  killed at 1 (equation (3.11) is the Fokker-Planck equation for the killed process).

Note that  $\widetilde{b}(1+v) = -\widetilde{b}(1-v)$  for  $v \in \mathbb{R}$ , so that  $\widetilde{b}$  is odd with respect to 1. Therefore, when initialised at  $\widetilde{X}_0 = 1$  the process  $(\widetilde{X}_t - 1)_{0 \leq t \leq T}$  satisfies

$$\widetilde{X}_t - 1 = \int_0^t \widetilde{b}(\widetilde{X}_s - 1 + 1)ds + W_t, \qquad t \in [0, T]$$

and the process  $(1 - \widetilde{X}_t)_{0 \leq t \leq T}$  satisfies

$$1 - \widetilde{X}_t = -\int_0^t \widetilde{b}(\widetilde{X}_s)ds - W_t = \int_0^t \widetilde{b}(2 - \widetilde{X}_s)ds - W_t$$
$$= \int_0^t \widetilde{b}(1 - \widetilde{X}_s + 1)ds - W_t,$$

which is the same equation. Hence  $(1 - \widetilde{X}_t)_{0 \leq t \leq T}$  and  $(\widetilde{X}_t - 1)_{0 \leq t \leq T}$  have the same distribution when  $\widetilde{X}_0 = 1$ . Therefore the reflection principle (see [17, Section I.13]) applies and

$$q(t,x,y) = \widetilde{q}(t,x,y) - \widetilde{q}(t,x,2-y), \qquad t \in [0,T], \quad x,y \le 1,$$
(3.16)

where  $\tilde{q}(t, x, y)$  is the transition density of the non-killed process.

We thus need to estimate the transition density  $\tilde{q}$  of the non-killed process. By Theorem 1 of [6] it follows that

$$\widetilde{q}(t,x,y) \leqslant \frac{B_T}{\sqrt{t}} \exp\left(-\frac{|y-\xi_t^x|^2}{B_T t}\right), \quad t > 0, \ x,y \in \mathbb{R},$$
(3.17)

where

$$\xi_t^x = x + \int_0^t \widetilde{b}(\xi_s^x) ds, \qquad t \ge 0.$$

We claim that a similar bound holds for  $|\partial_y \tilde{q}(t, x, y)|$ , i.e.

$$\left|\partial_{y}\widetilde{q}(t,x,y)\right| \leqslant \frac{B_{T}}{t} \exp\left(-\frac{|y-\xi_{t}^{x}|^{2}}{B_{T}t}\right), \quad t > 0, \ x,y \in \mathbb{R}.$$
(3.18)

Before we prove (3.18), we show that it implies (3.12) and (3.13). By (3.16), we deduce that

$$|\partial_y q(t, x, y)| \leqslant \frac{B_T}{t} \left[ \exp\left(-\frac{|y - \xi_t^x|^2}{B_T t}\right) + \exp\left(-\frac{|2 - y - \xi_t^x|^2}{B_T t}\right) \right], \quad t > 0, \ x, y \leqslant 1.$$

Since b(1) = 0, we have  $\xi_t^1 = 1$  for any  $t \ge 0$ . Therefore, by the comparison principle for ODEs, it must hold  $\xi_t^x \le \xi_t^1 = 1$  for any  $t \ge 0$  and  $x \le 1$ . As a consequence, for t > 0 and  $x, y \le 1$ ,

$$\begin{aligned} |y - \xi_t^x| &= |(y - 1) - (\xi_t^x - 1)| \leqslant |y - 1| + |\xi_t^x - 1| \\ &= 1 - y + 1 - \xi_t^x = 2 - y - \xi_t^x = |2 - y - \xi_t^x|. \end{aligned}$$

Therefore, the largest exponential term in the above bound for  $|\partial_y \tilde{q}(t, x, y)|$  is the first one. Equation (3.12) follows directly (adjusting  $B_T$  as necessary). Now, by Gronwall's Lemma and thanks to the Lipschitz property of b, we have:

$$e^{-Ks}|\xi_t^x - y| \leqslant |\xi_{t-s}^x - \xi_{-s}^y| \leqslant e^{Ks}|\xi_t^x - y|$$
(3.19)

for all  $s \in [0, t]$ , so that, choosing s = t, we get (3.13).

We now turn to the proof of (3.18). Without any loss of generality, we can assume  $\tilde{b}$  to be a twice continuously differentiable function with a bounded secondorder derivative. Indeed, if we can prove that, in such a case, (3.18) holds with respect to a constant  $B_T$  that does not refer to the second-order differentiability of  $\tilde{b}$ , then (3.18) holds in the original setting as well by a standard mollification argument. Since  $\tilde{q}$  is the transition probability of the solution to (3.15), it satisfies the Fokker-Planck equation

$$\partial_t \widetilde{q}(t, x, y) = \frac{1}{2} \partial_{yy}^2 \widetilde{q}(t, x, y) - \widetilde{b}(y) \partial_y \widetilde{q}(t, x, y) - \widetilde{b}'(y) \widetilde{q}(t, x, y),$$

for  $t \in [0,T]$  and  $x, y \in \mathbb{R}$ . Define  $(\widetilde{Y}_t^y)_{t \ge 0}$  to be the solution of the SDE

$$\begin{cases} d\widetilde{Y}_t^y &= -\widetilde{b}(\widetilde{Y}_t^y)dt + dW_t, \\ \widetilde{Y}_0^y &= y. \end{cases}$$
(3.20)

By Theorem 39 [15, p. 312], we know that, a.s., the mapping  $[0, +\infty) \times \mathbb{R} \ni (t, y) \mapsto Y_t^y$  is continuously differentiable with respect to y and that

$$\partial_y \widetilde{Y}_s^y = \exp\left(-\int_0^s \widetilde{b}'(\widetilde{Y}_u^y) du\right). \tag{3.21}$$

Moreover, by Lemma 3.2, we know that  $\tilde{q}$  is regular enough in order to apply the Itô-Krylov formula (see [12, Section II.10]) to  $[0, t/2] \ni s \mapsto \tilde{q}\left(t-s, x, \tilde{Y}_s^y\right)$ :

$$\widetilde{q}(t,x,y) = \mathbb{E}\left[\widetilde{q}\left(t/2,x,\widetilde{Y}_{t/2}^y\right)\right] - \int_0^{t/2} \mathbb{E}\left[\partial_y \widetilde{b}(\widetilde{Y}_s^y) \widetilde{q}\left(t-s,x,\widetilde{Y}_s^y\right)\right] ds.$$

By the Malliavin-Bismut-Elworthy formula (see, for example, Theorem 2.1 of [7]),

$$\partial_{y}\widetilde{q}(t,x,y) = \frac{2}{t} \mathbb{E} \left[ \widetilde{q}(t/2,x,\widetilde{Y}_{t/2}^{y}) \int_{0}^{t/2} \partial_{y}\widetilde{Y}_{s}^{y} dW_{s} \right] \\ - \int_{0}^{t/2} \frac{1}{s} \mathbb{E} \left[ \partial_{y}\widetilde{b}(\widetilde{Y}_{s}^{y})\widetilde{q}\left(t-s,x,\widetilde{Y}_{s}^{y}\right) \int_{0}^{s} \partial_{y}\widetilde{Y}_{r}^{y} dW_{r} \right] ds.$$
(3.22)

By (3.21),  $|\partial_y \widetilde{Y}_s^y| \leq \exp(Ks)$ . Thus using (3.17) we can compute for any  $0 < s \leq t/2$  (where the constant  $B_T$  changes from line to line below)

$$\begin{split} I(s,t) &:= \left| \frac{1}{s} \mathbb{E} \left[ \widetilde{q}(t-s,x,\widetilde{Y}_s^y) \int_0^s \partial_y \widetilde{Y}_r^y dW_r \right] \right| \\ &\leqslant \frac{1}{s} \left[ \mathbb{E} \left( \widetilde{q}^2(t-s,x,\widetilde{Y}_s^y) \right) \right]^{1/2} \left[ \mathbb{E} \left( \int_0^s \partial_y \widetilde{Y}_r^y dW_r \right)^2 \right]^{1/2} \\ &\leqslant \frac{B_T}{\sqrt{s}} \left[ \mathbb{E} \left( \widetilde{q}^2(t-s,x,\widetilde{Y}_s^y) \right) \right]^{1/2} \\ &\leqslant \frac{B_T}{\sqrt{s}} \left[ \frac{1}{t-s} \mathbb{E} \left( \exp \left( -\frac{|\widetilde{Y}_s^y - \xi_{t-s}^x|^2}{B_T(t-s)} \right) \right) \right]^{1/2} \\ &= \frac{B_T}{\sqrt{s}} \left[ \frac{1}{t-s} \int_{\mathbb{R}} \exp \left( -\frac{|z - \xi_{t-s}^x|^2}{B_T(t-s)} \right) \mathbb{P}(\widetilde{Y}_s^y \in dz) \right]^{1/2}. \end{split}$$

By an estimate similar to (3.17) but for the density of  $\widetilde{Y}$ , we deduce

$$\begin{split} I(s,t) &\leqslant \frac{B_T}{\sqrt{s}} \left[ \frac{1}{(t-s)\sqrt{s}} \int_{\mathbb{R}} \exp\left(-\frac{|z-\xi_{t-s}^x|^2}{B_T(t-s)}\right) \exp\left(-\frac{|z-\xi_{-s}^y|^2}{B_Ts}\right) dz \right]^{1/2} \\ &\leqslant \frac{B_T}{\sqrt{s}} \left[ \frac{1}{\sqrt{t-s}\sqrt{t}} \exp\left(-\frac{|\xi_{t-s}^x-\xi_{-s}^y|^2}{B_Tt}\right) \right]^{1/2}, \end{split}$$

where, to write the first line, we have used the fact the flow associated with the ODE driven by  $-\tilde{b}$  is nothing but the backward flow associated with the ODE driven by  $\tilde{b}$ , i.e.  $(\xi_{-t}^x)_{t \ge 0}$  satisfies

$$\xi_{-t}^x = x - \int_0^t \widetilde{b}(\xi_{-s}^x) ds, \quad t \ge 0,$$

and, to pass from the first to the second line, we have recognised a Gaussian convolution. By (3.19), we deduce that

$$I(s,t) \leqslant \frac{B_T}{\sqrt{s}} \left[ \frac{1}{\sqrt{t-s}\sqrt{t}} \exp\left(-\frac{|\xi_t^x - y|^2}{B_T t}\right) \right]^{1/2}$$

from which (3.22) yields

$$\begin{aligned} |\partial_y \widetilde{q}(t, x, y)| &\leqslant \frac{B_T}{t} \exp\left(-\frac{|\xi_t^x - y|^2}{B_T t}\right) \\ &+ B_T \exp\left(-\frac{|\xi_t^x - y|^2}{B_T t}\right) \int_0^{t/2} \frac{1}{\sqrt{s}} \left[\frac{1}{\sqrt{t - s}\sqrt{t}}\right]^{1/2} ds \\ &\leqslant \frac{B_T}{t} \exp\left(-\frac{|\xi_t^x - y|^2}{B_T t}\right) + \frac{B_T}{\sqrt{t}} \exp\left(-\frac{|\xi_t^x - y|^2}{B_T t}\right) \int_0^{t/2} \frac{1}{\sqrt{s}} ds. \end{aligned}$$
proves (3.18).

This proves (3.18).

**Lemma 3.5.** Let T > 0. Suppose  $X_0 = x_0 < 1$  and  $e \in \mathcal{L}(T, A)$ . Then there exists a constant  $\kappa_1(T)$  (independent of A and  $x_0$ ) which increases with T such that

$$\sup_{0 \leq s \leq t} \left[ \sqrt{s} \left\| \partial_y p_e^{x_0}(s, .) - \partial_y q(s, x_0, .) \right\|_{\infty} \right] \leq (\alpha A + |b(1)|) \kappa_1(T),$$

for all non-negative  $t \leq \min \{ [2(\alpha A + |b(1)|)\kappa_1(T)]^{-2}, T \}.$ 

*Proof.* For notational sake, define

$$F_e^{x_0}(t,y) := \partial_y p_e^{x_0}(t,y) - \partial_y q(t,x_0,y)$$

By (3.10), we see that

$$|F_{e}^{x_{0}}(t,y)| \leq (\alpha A + |b(1)|) \int_{0}^{t} \int_{-\infty}^{1} |\partial_{z}q(s,x_{0},z)\partial_{y}q(t-s,z,y)|dzds + (\alpha A + |b(1)|) \int_{0}^{t} \int_{-\infty}^{1} |\partial_{z}p_{e}^{x_{0}}(s,z) - \partial_{z}q(s,x_{0},z)| |\partial_{y}q(t-s,z,y)| dzds.$$
(3.23)

Thus defining  $\widetilde{A} := \alpha A + |b(1)|$  and allowing  $B_T$  to increase as necessary from line to line below, by Proposition 3.4 we see that for  $t \leq T$ 

$$\begin{aligned} |F_e^{x_0}(t,y)| \\ &\leqslant \widetilde{A}B_T \int_0^t \int_{-\infty}^1 s^{-1} (t-s)^{-1} \exp\left(-\frac{|z-\xi_s^{x_0}|^2}{B_T s}\right) \exp\left(-\frac{|z-\xi_{-(t-s)}^y|^2}{B_T (t-s)}\right) dz ds \\ &+ \widetilde{A}B_T \int_0^t (t-s)^{-1} \|F_e^{x_0}(s,\cdot)\|_{\infty} \int_{-\infty}^1 \exp\left(-\frac{|z-\xi_{-(t-s)}^y|^2}{B_T (t-s)}\right) dz ds. \end{aligned}$$

We can recognise the first term in the above as a Gaussian convolution. Thus

$$|F_{e}^{x_{0}}(t,y)| \leq \frac{\widetilde{A}B_{T}}{\sqrt{t}} \int_{0}^{t} \frac{1}{\sqrt{s}\sqrt{t-s}} \exp\left(-\frac{|\xi_{-(t-s)}^{y} - \xi_{s}^{x_{0}}|^{2}}{B_{T}t}\right) ds + \widetilde{A}B_{T} \int_{0}^{t} \frac{\sqrt{s}}{\sqrt{t-s}\sqrt{s}} \|F_{e}^{x_{0}}(s,\cdot)\|_{\infty} ds$$
(3.24)
$$\leq \widetilde{A}B_{T} \left(\frac{1}{\sqrt{t}} \exp\left(-\frac{|\xi_{-t}^{y} - x_{0}|^{2}}{B_{T}t}\right) + \sup_{0 \leq s \leq t} \left[\sqrt{s} \|F_{e}^{x_{0}}(s,\cdot)\|_{\infty}\right]\right),$$

where again we have used (3.19). Therefore, by multiplying both sides of the inequality by  $\sqrt{t}$ , we deduce

$$\sup_{0 \leqslant s \leqslant t} \left[ \sqrt{s} \left\| F_e^{x_0}(s, \cdot) \right\|_{\infty} \right] \leqslant \widetilde{A}B_T + \sqrt{t}\widetilde{A}B_T \sup_{0 \leqslant s \leqslant t} \left[ \sqrt{s} \left\| F_e^{x_0}(s, \cdot) \right\|_{\infty} \right],$$

We can conclude that, for  $t \in [0,T]$  such that  $t \leq [2\widetilde{A}B_T]^{-2} = [2(\alpha A + |b(1)|)B_T]^{-2}$ , we have that

$$\sup_{0 \leqslant s \leqslant t} \left[ \sqrt{s} \left\| \partial_y p_e^{x_0}(s, .) - \partial_y q(s, x_0, .) \right\|_{\infty} \right] \leqslant 2\widetilde{A}B_T.$$

$$(3.25)$$

**Proposition 3.6.** Let T > 0. Suppose  $X_0 = x_0 < 1$ , and that  $e \in \mathcal{L}(T, A)$ . Then there exists a constant  $\kappa_2(T)$  (independent of A and  $x_0$ ) which increases with T such that

$$|\partial_y p_e^{x_0}(t,y) - \partial_y q(t,x_0,y)| \leq \kappa_2(T)(\alpha A + |b(1)|) \frac{1}{\sqrt{t}} \exp\left(-\frac{|\xi_t^{x_0} - y|^2}{\kappa_2(T)t}\right),$$

for all  $t \leq \min \{ [(\alpha A + |b(1)|)\kappa_2(T)]^{-2}, T \}$  and  $y \leq 1$ , where  $(\xi_t^{x_0})_{t \geq 0}$  is as in Proposition 3.4.

*Proof.* Throughout the proof, we will use the fact the mapping

$$\varphi : \mathbb{R} \ni x \mapsto \frac{\exp(x) - 1}{x} \quad (\varphi(0) = 1),$$
(3.26)

is non-decreasing. We will also use the same notation as in the proof of Lemma 3.5:

$$F_e^{x_0}(t,y) := \partial_y p_e^{x_0}(t,y) - \partial_y q(t,x_0,y)$$

First Step. As noted in (3.23), we have for  $t \in [0, T]$ 

$$|F_{e}^{x_{0}}(t,y)| \leqslant \widetilde{A} \int_{0}^{t} \int_{-\infty}^{1} |\partial_{z}q(s,x_{0},z)\partial_{y}q(t-s,z,y)|dzds + \widetilde{A} \int_{0}^{t} \int_{-\infty}^{1} |F_{e}^{x_{0}}(t_{1},y_{1})| |\partial_{y}q(t-t_{1},y_{1},y)| dy_{1}dt_{1}$$
(3.27)

where we have denoted  $\widetilde{A} = \alpha A + |b(1)|$ . The fact that  $\varphi$  given by (3.26) is nondecreasing says that, for  $t \in [0, T]$ ,

$$\frac{1 - \exp(-2KT)}{2KT} \leqslant \frac{1 - \exp(-2Kt)}{2Kt} \leqslant 1 \leqslant \frac{\exp(2Kt) - 1}{2Kt} \leqslant \frac{\exp(2KT) - 1}{2KT},$$

from which, together with Proposition 3.4, we deduce there exists  $B_T > 0$  such that

$$|\partial_y q(t, x, y)| \leqslant \frac{B_T}{1 - e^{-2Kt}} \exp\left(-\frac{|x - \xi_{-t}^y|^2}{2B_T(1 - e^{-2Kt})}\right)$$
(3.28)

$$|\partial_y q(t, x, y)| \leq \frac{B_T}{e^{2Kt} - 1} \exp\left(-\frac{|\xi_t^x - y|^2}{2B_T(e^{2Kt} - 1)}\right).$$
(3.29)

Unlike in the previous lemma,  $B_T$  is now fixed. We then introduce the kernel

$$\mathcal{K}(t,x,y) = \begin{cases} \frac{1}{\sqrt{B_T(e^{2Kt}-1)}} \exp\left(-\frac{|\xi_t^x - y|^2}{2B_T(e^{2Kt}-1)}\right) & \text{if } t > 0, \\ \frac{1}{\sqrt{B_T(1-e^{2Kt})}} \exp\left(-\frac{|x - \xi_t^y|^2}{2B_T(1-e^{2Kt})}\right) & \text{if } t < 0. \end{cases}$$
(3.30)

For  $0 < s < t \leq T$ ,  $\mathcal{K}$  satisfies the Gaussian convolution property:

$$\begin{split} \mathcal{K}(s, x, \cdot) &\otimes \mathcal{K}\big(-(t-s), \cdot, y\big) \\ &:= \int_{\mathbb{R}} \mathcal{K}(s, x, z) \mathcal{K}\big(-(t-s), z, y\big) dz \\ &= \frac{\sqrt{2\pi}}{\sqrt{B_T(e^{2Ks} - e^{-2K(t-s)})}} \exp\left(-\frac{|\xi_s^x - \xi_{-(t-s)}^y|^2}{2B_T(e^{2Ks} - e^{-2K(t-s)})}\right). \end{split}$$

By (3.19), we know that  $|\xi_t^x - y| \leq e^{K(t-s)} |\xi_s^x - \xi_{-(t-s)}^y|$  for all 0 < s < t. Therefore,

$$\mathcal{K}(s,x,\cdot) \otimes \mathcal{K}\left(-(t-s),\cdot,y\right) \leqslant \frac{\sqrt{2\pi}}{\sqrt{B_T(1-e^{-2Kt})}} \exp\left(-\frac{|\xi_t^x - y|^2}{2B_T(e^{2Kt} - 1)}\right) \qquad (3.31)$$
$$\leqslant \sqrt{2\pi} e^{KT} \mathcal{K}(t,x,y).$$

By (3.27), (3.28) and (3.29), we deduce that, for  $t \in [0, T]$ ,

$$\begin{aligned} |F_e^{x_0}(t,y)| &\leqslant \widetilde{A}B_T^3 e^{KT} \mathcal{K}(t,x_0,y) \int_0^t \frac{\sqrt{2\pi}}{\sqrt{e^{2Ks} - 1}\sqrt{1 - e^{-2K(t-s)}}} ds \\ &+ \widetilde{A}B_T^{3/2} \int_0^t \int_{-\infty}^1 \frac{|F_e^{x_0}(t_1,y_1)|}{\sqrt{1 - e^{-2K(t-t_1)}}} \mathcal{K}(-(t-t_1),y_1,y) dy_1 dt_1. \end{aligned}$$
(3.32)

Using (3.26), we notice that

$$\int_{0}^{t} \frac{1}{\sqrt{e^{2Ks} - 1}\sqrt{1 - e^{-2K(t-s)}}} ds \leqslant \frac{\sqrt{2KT}}{\sqrt{1 - \exp(-2KT)}} \int_{0}^{t} \frac{1}{\sqrt{2Ks}\sqrt{2K(t-s)}} ds$$
$$= \frac{\sqrt{T}}{\sqrt{2K(1 - \exp(-2KT))}} \int_{0}^{t} \frac{1}{\sqrt{s}\sqrt{t-s}} ds$$
$$\leqslant \frac{4\sqrt{T}}{\sqrt{2K(1 - \exp(-2KT))}} =: C_{T}.$$
(3.33)

Finally, (3.32) yields

$$\begin{aligned} |F_e^{x_0}(t,y)| &\leqslant \widetilde{A}\widetilde{C}_T B_T^3 \mathcal{K}(t,x_0,y) \\ &+ \widetilde{A} B_T^{3/2} \int_0^t \int_{-\infty}^1 \frac{|F_e^{x_0}(t_1,y_1)|}{\sqrt{1-e^{-2K(t-t_1)}}} \mathcal{K}(-(t-t_1),y_1,y) dy_1 dt_1, \end{aligned}$$
(3.34)

with  $\widetilde{C}_T := \sqrt{2\pi} C_T \exp(KT)$ .

Second Step. We now prove by induction, that for any  $N \ge 0$ ,

$$|F_e^{x_0}(t,y)| \leqslant \widetilde{A}\widetilde{C}_T B_T^3 \mathcal{K}(t,x_0,y) \sum_{i=0}^N \left(\sqrt{K}\widetilde{A}\widetilde{C}_T B_T^{3/2} \sqrt{t}\right)^i + R_{N+1}(t,y), \qquad (3.35)$$

where

$$R_N(t,y) = (\widetilde{A}B_T^{3/2})^N \int_{0 \leq t_N \leq \dots \leq t_0} dt_N \dots dt_1 \prod_{i=0}^{N-1} \frac{1}{\sqrt{1 - e^{-2K(t_i - t_{i+1})}}} \int_{(-\infty,1]^N} dy_N \dots dy_1 |F_e^{x_0}(t_N, y_N)| \prod_{i=0}^{N-1} \mathcal{K}\big(-(t_i - t_{i+1}), y_{i+1}, y_i\big),$$

with the convention  $t_0 = t$  and  $y_0 = y$ . In the first step, we established (3.35) when N = 0. Assume now that, for some  $N \ge 0$ , (3.35) holds for any  $t \in [0, T]$  and  $y \le 1$ . Then, plugging the induction assumption at rank N into (3.34), we get:

$$\begin{aligned} |F_e^{x_0}(t,y)| \\ &\leqslant \widetilde{A}\widetilde{C}_T B_T^3 \mathcal{K}(t,x_0,y) \\ &+ \widetilde{A}\widetilde{C}_T B_T^3 \left( \sum_{i=0}^N \left( \sqrt{K}\widetilde{A}\widetilde{C}_T B_T^{3/2} \sqrt{t} \right)^i \right) \widetilde{A} B_T^{3/2} \int_0^t \frac{\mathcal{K}(t_1,x_0,\cdot) \otimes \mathcal{K}(-(t-t_1),\cdot,y)}{\sqrt{1-e^{-2K(t-t_1)}}} dt_1 \\ &+ R_{N+2}(t,y). \end{aligned}$$

By (3.31), we deduce that

$$\begin{aligned} |F_e^{x_0}(t,y)| \\ &\leqslant \widetilde{A}\widetilde{C}_T B_T^3 \mathcal{K}(t,x_0,y) \\ &+ \widetilde{A}\widetilde{C}_T B_T^3 \bigg( \sum_{i=0}^N \bigg( \sqrt{K}\widetilde{A}\widetilde{C}_T B_T^{3/2} \sqrt{t} \bigg)^i \bigg) \widetilde{A} B_T^{3/2} \sqrt{2\pi} e^{KT} \mathcal{K}(t,x_0,y) \int_0^t \frac{1}{\sqrt{1-e^{-2K(t-t_1)}}} dt_1 \\ &+ R_{N+2}(t,y). \end{aligned}$$

Following (3.33), we have:

$$\int_0^t \frac{ds}{\sqrt{1 - e^{-2K(t-s)}}} \leqslant \frac{\sqrt{2KT}}{\sqrt{1 - \exp(-2KT)}} \int_0^t \frac{1}{\sqrt{2K(t-s)}} ds$$
$$= \frac{\sqrt{T}}{\sqrt{(1 - \exp(-2KT))}} \int_0^t \frac{1}{\sqrt{t-s}} ds$$
$$\leqslant \frac{2\sqrt{t}\sqrt{T}}{\sqrt{(1 - \exp(-2KT))}} \leqslant \sqrt{t}\sqrt{KC_T}.$$

Since  $\widetilde{C}_T = \sqrt{2\pi} \exp(KT) C_T$ , we deduce that (3.35) holds at rank N + 1.

Third Step. Suppose that  $\sqrt{t} \leq [\sqrt{K}\widetilde{A}\widetilde{C}_T B_T^{3/2}]^{-1}/2$ . Then, the series in (3.35) is convergent (and bounded above by 2). The point is thus to prove that  $R_N(t, y) \to 0$  as  $N \to \infty$  for all  $y \leq 1$ . By Lemma 3.5, we deduce that, for  $\sqrt{t} \leq [2\widetilde{A}\kappa_1(T)]^{-1}$ ,

$$R_{N}(t,y) \leq (\sqrt{2\pi}\widetilde{A}B_{T}^{3/2})^{N}\widetilde{A}\kappa_{1}(T) \int_{0 \leq t_{N} \leq t_{N-1} \leq \dots \leq t_{0}} \frac{1}{\sqrt{t_{N}}} \prod_{i=0}^{N-1} \frac{1}{\sqrt{1-e^{-2K(t_{i}-t_{i+1})}}} dt_{1} \dots dt_{N}.$$

Using (3.26), we obtain

$$R_N(t,y) \leqslant \left(\sqrt{2\pi}\widetilde{A}B_T^{3/2}\frac{\sqrt{T}}{\sqrt{1-e^{-2KT}}}\right)^N \widetilde{A}\kappa_1(T)$$

$$\times \int_{0 \leqslant t_N \leqslant t_{N-1} \leqslant \dots \leqslant t_0} \frac{1}{\sqrt{t_N}} \prod_{i=0}^{N-1} \frac{1}{\sqrt{t_i - t_{i+1}}} dt_1 \dots dt_N.$$
(3.36)

To compute the integral in the right-hand side of the above, define

$$J_n(t) = \int_0^t u^{(n-1)/2} (t-u)^{-1/2} du, \quad t \ge 0, \ n \in \mathbb{N}.$$

By the change of variable u = ts, we have  $J_n(t) = t^{n/2}J_n(1)$ . Note that  $J_0(1) \leq 4$ . Moreover, for  $n \geq 1$ ,

$$J_n(1) \leqslant n^{1/2} \int_0^{1-1/n} u^{(n-1)/2} du + \int_{1-1/n}^1 (1-u)^{-1/2} du$$
  
$$\leqslant \frac{2n^{1/2}}{n+1} + 2n^{-1/2} \leqslant 4n^{-1/2}.$$
(3.37)

With this notation

$$I_{N,0}(t) := \int_{0 \leqslant t_N \leqslant t_{N-1} \leqslant \dots \leqslant t_0} \frac{1}{\sqrt{t_N}} \prod_{i=0}^{N-1} \frac{1}{\sqrt{t_i - t_{i+1}}} dt_1 \dots dt_N$$
$$= \int_{0 \leqslant t_{N-1} \leqslant \dots \leqslant t_0} J_0(t_{N-1}) \prod_{i=0}^{N-2} \frac{1}{\sqrt{t_i - t_{i+1}}} dt_1 \dots dt_{N-1},$$

where the second equality holds for  $N \ge 2$ . More generally, setting for any  $n \le N-2$ 

$$I_{N,n}(t) := \int_{0 \leq t_{N-(n+1)} \leq \dots \leq t_0} J_n(t_{N-(n+1)}) \prod_{i=0}^{N-(n+2)} \frac{1}{\sqrt{t_i - t_{i+1}}} dt_1 \dots dt_{N-(n+1)},$$

we have, for any  $0 \leq n \leq N-3$ ,

$$\begin{split} I_{N,n}(t) &= J_n(1) \int_{0 \,\leq \, t_{N-(n+1)} \,\leq \, \dots \,\leq \, t_0} t_{N-(n+1)}^{n/2} \prod_{i=0}^{N-(n+2)} \frac{1}{\sqrt{t_i - t_{i+1}}} dt_1 \dots dt_{N-(n+1)} \\ &= J_n(1) \int_{0 \,\leq \, t_{N-(n+2)} \,\leq \, \dots \,\leq \, t_0} \left[ \int_0^{t_{N-(n+2)}} \frac{t_{N-(n+1)}^{n/2}}{\sqrt{t_{N-(n+2)} - t_{N-(n+1)}}} dt_{N-(n+1)} \right] \\ &\times \prod_{i=0}^{N-(n+3)} \frac{1}{\sqrt{t_i - t_{i+1}}} dt_1 \dots dt_{N-(n+2)} \\ &= J_n(1) I_{N,n+1}(t), \end{split}$$

so that, by (3.37),

$$I_{N,0}(t) = \left[\prod_{n=0}^{N-3} J_n(1)\right] I_{N,N-2}(t) = \left[\prod_{n=0}^{N-2} J_n(1)\right] J_{N-1}(t)$$
$$= \left[\prod_{n=0}^{N-1} J_n(1)\right] t^{(N-1)/2} \leq [4t^{1/2}]^{N-1} [(N-1)!]^{-1/2},$$

using the fact that  $I_{N,N-2}(t) = J_{N-2}(1)J_{N-1}(t)$ . Going back to (3.36), we deduce that  $R_N(t,y) \to 0$  as  $N \to \infty$ . Hence taking the limit as  $N \to \infty$  in (3.35) yields (with  $\sqrt{t} \leq [\sqrt{K}\widetilde{A}\widetilde{C}_T B_T^{3/2}]^{-1}/2$ )

$$|F_e^{x_0}(t,y)| \leq 2\widetilde{A}\widetilde{C}_T B_T^3 \mathcal{K}(t,x_0,y),$$

from which the result follows.

Complementary to the above two results, we have the following two, which estimate the difference between the densities of two processes driven by two different drifts.

**Lemma 3.7.** Let T > 0. Suppose  $X_0 = x_0 < 1$ , and that  $e_1, e_2 \in \mathcal{L}(T, A)$  where  $(w.l.o.g.) \quad \alpha A + |b(1)| \ge 1$ . Then there exists a constant  $\kappa_3(T)$  (independent of A and  $x_0$ ) which increases with T such that

$$\sup_{0 \leqslant s \leqslant t} \left[ \sqrt{s} \left\| \partial_y p_{e_1}^{x_0}(s, .) - \partial_y p_{e_2}^{x_0}(s, .) \right\|_{\infty} \right] \leqslant (\alpha A + |b(1)|) \kappa_3(T) \|e_1' - e_2'\|_{\infty, t},$$

for all  $t \leq \min\{[(\alpha A + |b(1)|)\kappa_3(T)]^{-2}, T\}.$ 

*Proof.* Let  $y \leq 1$ . Again by (3.10), we see that

$$\begin{aligned} \left| \partial_{y} p_{e_{1}}^{x_{0}}(t,y) - \partial_{y} p_{e_{2}}^{x_{0}}(t,y) \right| \\ &\leqslant \left( \alpha A + |b(1)| \right) \int_{0}^{t} \int_{-\infty}^{1} \left| \partial_{z} p_{e_{1}}^{x_{0}}(s,z) - \partial_{z} p_{e_{2}}^{x_{0}}(s,z) \right| \left| \partial_{y} q(t-s,z,y) \right| dz ds \\ &+ \alpha \left\| e_{1}^{\prime} - e_{2}^{\prime} \right\|_{\infty,t} \int_{0}^{t} \int_{-\infty}^{1} \left| \partial_{z} p_{e_{2}}^{x_{0}}(s,z) - \partial_{z} q(s,x_{0},z) \right| \left| \partial_{y} q(t-s,z,y) \right| dz ds \\ &+ \alpha \left\| e_{1}^{\prime} - e_{2}^{\prime} \right\|_{\infty,t} \int_{0}^{t} \int_{-\infty}^{1} \left| \partial_{z} q(s,x_{0},z) \right| \left| \partial_{y} q(t-s,z,y) \right| dz ds. \end{aligned}$$
(3.38)

Using Proposition 3.4 once again and the notation (3.30) (see also (3.28) and (3.29)) together with (3.26), we see that

$$\begin{split} \left| \partial_{y} p_{e_{1}}^{x_{0}}(t,y) - \partial_{y} p_{e_{2}}^{x_{0}}(t,y) \right| \\ &\leqslant \widetilde{A} \kappa_{3}(T) \sup_{0 \leqslant s \leqslant t} \left[ \sqrt{s} \left\| \partial_{z} p_{e_{1}}^{x_{0}}(s,.) - \partial_{z} p_{e_{2}}^{x_{0}}(s,.) \right\|_{\infty} \right] \\ &+ \kappa_{3}(T) \left\| e_{1}' - e_{2}' \right\|_{\infty,t} \int_{0}^{t} \int_{-\infty}^{1} \frac{\left| \partial_{z} p_{e_{2}}^{x_{0}}(s,z) - \partial_{z} q(s,x_{0},z) \right|}{\sqrt{t-s}} \mathcal{K}(-(t-s),z,y) dz ds \\ &+ \kappa_{3}(T) \left\| e_{1}' - e_{2}' \right\|_{\infty,t} \int_{0}^{t} \int_{-\infty}^{1} \frac{1}{\sqrt{s}} \frac{1}{\sqrt{t-s}} \mathcal{K}(s,x_{0},z) \mathcal{K}(-(t-s),z,y) dz ds, \end{split}$$

for some constant  $\kappa_3(T) > 0$ , where we have denoted  $\widetilde{A} := \alpha A + |b(1)|$ . Using Proposition 3.6 and (3.31), and allowing the constant  $\kappa_3(T)$  to increase as necessary from line to line below, it follows that

$$\begin{aligned} \left| \partial_{y} p_{e_{1}}^{x_{0}}(t,y) - \partial_{y} p_{e_{2}}^{x_{0}}(t,y) \right| \\ &\leqslant \widetilde{A} \kappa_{3}(T) \sup_{0 \leqslant s \leqslant t} \left[ \sqrt{s} \left\| \partial_{y} p_{e_{1}}^{x_{0}}(s,.) - \partial_{y} p_{e_{2}}^{x_{0}}(s,.) \right\|_{\infty} \right] \\ &+ \kappa_{3}(T) \widetilde{A} \left\| e_{1}' - e_{2}' \right\|_{\infty,t} \int_{0}^{t} \int_{-\infty}^{1} \frac{1}{\sqrt{t-s}} \mathcal{K}(s,x_{0},z) \mathcal{K}(-(t-s),z,y) dz ds \\ &+ \kappa_{3}(T) \left\| e_{1}' - e_{2}' \right\|_{\infty,t} \mathcal{K}(t,x_{0},y), \end{aligned}$$

for all  $t \leq \min\{[\widetilde{A}\kappa_2(T)]^{-2}, T\}$ . Using (3.31) again, together with the assumption  $\widetilde{A} \geq 1$ , we deduce:

$$\begin{aligned} \left| \partial_{y} p_{e_{1}}^{x_{0}}(t,y) - \partial_{y} p_{e_{2}}^{x_{0}}(t,y) \right| &\leq \widetilde{A} \kappa_{3}(T) \sup_{0 \leq s \leq t} \left[ \sqrt{s} \left\| \partial_{y} p_{e_{1}}^{x_{0}}(s,.) - \partial_{y} p_{e_{2}}^{x_{0}}(s,.) \right\|_{\infty} \right] \\ &+ \kappa_{3}(T) \widetilde{A} \left\| e_{1}' - e_{2}' \right\|_{\infty,t} \mathcal{K}(t,x_{0},y), \end{aligned}$$
(3.39)

Assuming that  $\kappa_3(T) \ge \kappa_2(T)$  and multiplying the above inequality by  $\sqrt{t}$ , we deduce that, for  $t \le \min\{[2\widetilde{A}\kappa_3(T)]^{-2}, T\}$  we have

$$\sup_{0 \leq s \leq t} \left[ \sqrt{s} \left\| \partial_{y} p_{e_{1}}^{x_{0}}(s, .) - \partial_{y} p_{e_{2}}^{x_{0}}(s, .) \right\|_{\infty} \right]$$
  
$$\leq 2\kappa_{3}(T) \sup_{0 < t \leq T} \left[ \frac{\sqrt{t}}{\sqrt{B_{T}(1 - e^{-2Kt})}} \right] \widetilde{A} \left\| e_{1}' - e_{2}' \right\|_{\infty, t}.$$

**Proposition 3.8.** Let T > 0. Suppose  $X_0 = x_0 < 1$ , and that  $e_1, e_2 \in \mathcal{L}(T, A)$ , where (w.l.o.g.)  $\alpha A + |b(1)| \ge 1$ . Then there exists a constant  $\kappa_4(T)$  (independent of A and  $x_0$ ) which increases with T such that

$$\left|\partial_{y} p_{e_{1}}^{x_{0}}(t,y) - \partial_{y} p_{e_{2}}^{x_{0}}(t,y)\right| \leq (\alpha A + |b(1)|)\kappa_{4}(T) \frac{1}{\sqrt{t}} \exp\left(-\frac{|\xi_{t}^{x_{0}} - y|^{2}}{\kappa_{4}(T)t}\right) \|e_{1}' - e_{2}'\|_{\infty,t},$$

for all  $t \leq \min\{[(\alpha A + |b(1)|)\kappa_4(T)]^{-2}, T\}$  and  $y \leq 1$ , where  $(\xi_t^{x_0})_{t \geq 0}$  is as in Proposition 3.4.

*Proof.* We follow the strategy of the proof of Proposition 3.6. We thus define

$$G_{e_1,e_2}^{x_0}(t,y) = \partial_y p_{e_1}^{x_0}(t,y) - \partial_y p_{e_2}^{x_0}(t,y).$$

Going back to (3.38) we can proceed in the same way as Lemma 3.7 in order to bound the second two terms of this expression. We then get that, for some constant  $\kappa_4(T) > 0$ ,

$$\begin{split} \left| G_{e_1,e_2}^{x_0}(t,y) \right| &\leqslant \widetilde{A}\kappa_4(T) \left\| e_1' - e_2' \right\|_{\infty,t} \mathcal{K}(t,x_0,y) \\ &+ \widetilde{A}\kappa_4(T) \int_0^t \int_{-\infty}^t \frac{\left| G_{e_1,e_2}^{x_0}(t_1,y_1) \right|}{\sqrt{1 - e^{-2K(t-t_1)}}} \mathcal{K}(-(t-t_1),y_1,y) dy_1 dt_1. \end{split}$$

We can iterate this inequality in exactly the same way as in Proposition 3.6 (see (3.35)), and the convergence follows from Lemma 3.7. This yields the result.  $\Box$ 

# 3.3. Conclusion.

**Proposition 3.9.** Let T > 0 and suppose  $\mathbb{P}(X_0 \in dx) \leq \beta(1-x)dx$ , for  $x \in (1-\epsilon, 1]$ , with (w.l.o.g.)  $\beta \geq 1$ . Suppose  $e_1, e_2 \in \mathcal{H}(T, A, X_0)$  with (w.l.o.g)  $\widetilde{A} := \alpha A + |b(1)| \geq 1$ . Then there exist a constant  $\kappa(T)$ , independent of A,  $\beta$  and  $\epsilon$ , and increasing in T, and a constant  $\widetilde{\kappa}(T, \beta, \epsilon)$ , independent of A and increasing in T, such that

$$\sup_{0 \leqslant s \leqslant t} \left| \frac{d}{ds} \mathbb{E}(M_s^{e_1}) - \frac{d}{ds} \mathbb{E}(M_s^{e_2}) \right| \leqslant \widetilde{A} \widetilde{\kappa}(T, \beta, \epsilon) \sqrt{t} \left( 1 + \mathbb{E}|X_0| \right) \|e_1' - e_2'\|_{\infty, t},$$

for  $t \leq \min\{[(\alpha A + |b(1)|)\kappa(T)]^{-2}, T\}$ . Proof. We have by (2.5) and (3.6)

$$\frac{d}{dt} \mathbb{E}(M_t^{e_1}) - \frac{d}{dt} \mathbb{E}(M_t^{e_2}) \left| \\ \leq \frac{1}{2} \int_{-\infty}^1 \left| \partial_y p_{e_1}^x(t,1) - \partial_y p_{e_2}^x(t,1) \right| \mathbb{P}(X_0 \in dx) \\ + \frac{1}{2} \sum_{k \ge 1} \int_0^t \left| \partial_y p_{e_1}^{(0,s)}(t-s,1) - \partial_y p_{e_2}^{(0,s)}(t-s,1) \right| \mathbb{P}(\tau_k^{e_1} \in ds) \qquad (3.40) \\ + \frac{1}{2} \int_0^t \left| \partial_y p_{e_2}^{(0,s)}(t-s,1) \right| \left| \frac{d}{ds} \mathbb{E}(M_s^{e_1}) - \frac{d}{ds} \mathbb{E}(M_s^{e_2}) \right| ds \\ := \frac{1}{2} (L_1 + L_2 + L_3).$$

Suppose  $t \leq T$  and  $\sqrt{t} \leq [(\alpha A + |b(1)|)\kappa_4(T)]^{-1}$ , where  $\kappa_4(T)$  is as in Proposition 3.8. Considering the first term only, we can use this result to see that

$$L_1 \leqslant \widetilde{A}\beta\kappa_4(T) \left( \int_{1-\epsilon}^1 \frac{1}{\sqrt{t}} \exp\left(-\frac{(1-x)^2}{\kappa_4(T)t}\right) (1-x)dx \right) \|e_1' - e_2'\|_{\infty,t} + \widetilde{A}\kappa_4(T) \left( \int_{-\infty}^{1-\epsilon} \frac{1}{\sqrt{t}} \exp\left(-\frac{(1-x)^2}{\kappa_4(T)t}\right) \mathbb{P}(X_0 \in dx) \right) \|e_1' - e_2'\|_{\infty,t}$$

where we have used the bi-Lipschitz property (3.19) in the statement of Proposition 3.8, together with the fact that  $\xi_t^1 = 1$  for all t and  $\tilde{A} = \alpha A + |b(1)|$  as usual. Indeed, by definition (see the statement of Proposition 3.4), we have

$$\xi_t^1 = 1 + \int_0^t \widetilde{b}(\xi_s^1) ds, \quad \widetilde{b}(1) = 0.$$

We deduce that there exists a constant  $\kappa_5(T, \beta, \epsilon) > 0$ , which is independent of A and which is allowed to increase as necessary from line to line below, such that (the value of  $\kappa_4(T)$  being also allowed to increase from line to line)

$$L_{1} \leqslant \widetilde{A}\beta\kappa_{4}(T)\sqrt{t} \left(\int_{0}^{\infty} z \exp\left(-\frac{z^{2}}{\kappa_{4}(T)}\right) dz\right) \|e_{1}' - e_{2}'\|_{\infty,t}$$

$$+ \widetilde{A}\kappa_{4}(T)\frac{1}{\sqrt{t}} \exp\left(-\frac{\epsilon^{2}}{\kappa_{4}(T)t}\right) \|e_{1}' - e_{2}'\|_{\infty,t}.$$

$$\leqslant \widetilde{A}\kappa_{5}(T,\beta,\epsilon)\sqrt{t}\|e_{1}' - e_{2}'\|_{\infty,t}.$$
(3.41)

We can then use Proposition 3.8 again to see that

$$L_2 \leqslant \widetilde{A}\kappa_4(T) \sup_{0 < s \leqslant t} \left[ s^{-1/2} \exp\left(-\frac{1}{\kappa_4(T)s}\right) \right] \mathbb{E}(M_t^{e_1}) \|e_1' - e_2'\|_{\infty,t}.$$

By Proposition 2.2, we deduce that

$$L_2 \leqslant \widetilde{A}\kappa_4(T)\sqrt{t}\left(\sqrt{t} + \mathbb{E}|X_0|\right) \|e_1' - e_2'\|_{\infty,t}, \qquad (3.42)$$

where we use the elementary inequality  $e^{-\frac{1}{v}} \leq v$  for all  $v \geq 0$ . We finally turn to  $L_3$  in (3.40). By Proposition 3.6, Proposition 3.4 and (3.19), we also have that

$$\begin{aligned} \left|\partial_{y} p_{e_{2}}^{(0,s)}(t-s,1)\right| &\leqslant \kappa_{4}(T) \widetilde{A} \frac{1}{\sqrt{t-s}} \exp\left(-\frac{1}{\kappa_{4}(T)(t-s)}\right) + \left|\partial_{y} q(t-s,0,1)\right| \\ &\leqslant \kappa_{4}(T) \exp\left(-\frac{1}{\kappa_{4}(T)(t-s)}\right) \left[\frac{\widetilde{A}}{\sqrt{t-s}} + \frac{1}{t-s}\right] \\ &\leqslant \kappa_{4}(T) \widetilde{A}. \end{aligned}$$

$$(3.43)$$

Thus, from (3.40), (3.41), (3.42) and (3.43), we deduce

$$\left|\frac{d}{dt}\mathbb{E}(M_t^{e_1}) - \frac{d}{dt}\mathbb{E}(M_t^{e_2})\right| \leqslant \widetilde{A}\kappa_5(T,\beta,\epsilon)\sqrt{t}(1+\mathbb{E}|X_0|)\|e_1' - e_2'\|_{\infty,t} + \widetilde{A}\kappa_4(T)\int_0^t \left|\frac{d}{ds}\mathbb{E}(M_s^{e_1}) - \frac{d}{ds}\mathbb{E}(M_s^{e_2})\right| ds.$$

By taking the supremum over all  $s \leq t$  in the above, we have, for  $t \leq (2\kappa_4(T)\widetilde{A})^{-1}$ ,

$$\sup_{0 \leqslant s \leqslant t} \left| \frac{d}{ds} \mathbb{E}(M_s^{e_1}) - \frac{d}{ds} \mathbb{E}(M_s^{e_2}) \right| \leqslant 2\widetilde{A}\kappa_5(T,\beta,\epsilon)\sqrt{t} \left(1 + \mathbb{E}|X_0|\right) \|e_1' - e_2'\|_{\infty,t}.$$

Proof. [Proof of Theorem 3.1] Choose

$$A_1 = 2 \sup_{0 \leqslant t \leqslant 1} \left| \frac{d}{dt} \mathbb{E}(M_t^0) \right| + 1.$$

Note that  $A_1$  depends on  $\beta$ . Then choose  $T_1 \leq \min\{[(\alpha A_1 + |b(1)|)\kappa(1)]^{-2}, 1\}$  such that

$$\sqrt{T_1}\widetilde{\kappa}(1,\beta,\epsilon)(\alpha A_1 + |b(1)|)(1 + \mathbb{E}|X_0|) \leqslant \frac{1}{4}, \qquad (3.44)$$

where  $\kappa(1)$  and  $\tilde{\kappa}(1,\beta,\epsilon)$  are as in the Proposition 3.9. By that result, if  $e \in \mathcal{H}(A_1,T_1,X_0)$  then

$$\left|\frac{d}{dt}\mathbb{E}(M_t^e)\right| \leq \sqrt{t}\widetilde{\kappa}(T_1,\beta,\epsilon)(\alpha A_1 + |b(1)|)(1 + \mathbb{E}|X_0|)A_1 + \left|\frac{d}{dt}\mathbb{E}(M_t^0)\right|$$

for all  $t \leq \min\{[(\alpha A_1 + |b(1)|)\kappa(T_1)]^{-2}, 1\}$ . By definition, we have  $T_1 \leq 1$  so that  $\kappa(T_1) \leq \kappa(1)$  and  $\tilde{\kappa}(T_1, \beta, \epsilon) \leq \tilde{\kappa}(1, \beta, \epsilon)$ . Therefore

$$\left|\frac{d}{dt}\mathbb{E}(M_t^e)\right| \leqslant \sqrt{t}\widetilde{\kappa}(1,\beta,\epsilon)(\alpha A_1 + |b(1)|)(1 + \mathbb{E}|X_0|)A_1 + \left|\frac{d}{dt}\mathbb{E}(M_t^0)\right|$$

for all 
$$t \leq \min\{[(\alpha A_1 + |b(1)|)\kappa(1)]^{-2}, 1\}$$
. Hence for all  $t < T_1$   
$$\left|\frac{d}{dt}\mathbb{E}(M_t^e)\right| \leq \sqrt{t}\widetilde{\kappa}(1, \beta, \epsilon)(\alpha A_1 + |b(1)|)(1 + \mathbb{E}|X_0|)A_1 + \left|\frac{d}{dt}\mathbb{E}(M_t^0)\right|$$
$$\leq \frac{A_1}{2} + \sup_{0 \leq t \leq 1} \left|\frac{d}{dt}\mathbb{E}(M_t^0)\right| \leq A_1$$

by (3.44), so that  $\Gamma(e) \in \mathcal{H}(A_1, T_1)$ .

To prove that  $\Gamma$  is a contraction on  $\mathcal{H}(A_1, T_1, X_0)$ , first note that for  $e \in \mathcal{H}(A_1, T_1, X_0)$ 

$$||e'||_{\infty,T_1} \leq ||e||_{\mathcal{H}(A,T_1,X_0)} \leq 2||e'||_{\infty,T_1}$$

by the mean value theorem, since e(0) = 0 and  $T_1 \leq 1$ . Thus for any  $e_1, e_2 \in \mathcal{H}(A_1, T_1, X_0)$ 

$$\begin{split} \|\Gamma(e_{1}) - \Gamma(e_{2})\|_{\mathcal{H}(A_{1},T_{1},X_{0})} &\leq 2 \|\Gamma(e_{1})' - \Gamma(e_{2})'\|_{\infty,T_{1}} \\ &\leq 2 \sup_{t \leq T_{1}} \left| \frac{d}{dt} \mathbb{E}(M_{t}^{e_{1}}) - \frac{d}{dt} \mathbb{E}(M_{t}^{e_{2}}) \right| \\ &\leq 2\sqrt{T_{1}} \widetilde{\kappa}(T_{1},\beta,\epsilon) \left(1 + \mathbb{E}|X_{0}|\right) \left(\alpha A + |b(1)|\right) \|e_{1}' - e_{2}'\|_{\infty,T_{1}} \\ &\leq \frac{1}{2} \|e_{1} - e_{2}\|_{\mathcal{H}(A_{1},T_{1},X_{0})}, \end{split}$$

by our choice of  $T_1$  and using Proposition 3.9 once more. Since  $\mathcal{H}(A_1, T_1, X_0)$  is a closed subspace of  $\mathcal{C}^1[0, T]$  (a complete metric space), the existence of a fixed point for  $\Gamma$  follows from the Banach Fixed Point Theorem.

#### 4. Long-time estimates

In order to extend the existence and uniqueness from small time to any arbitrarily prescribed interval, we need an a priori bound for the Lipschitz constant of  $t \mapsto \mathbb{E}(M_t)$  on any finite interval [0, T]. In the whole section, for a given initial condition  $X_0 = x_0 \in (-\infty, 1)$ , we thus assume that there exists a solution to (2.1) according to Definition 2.3 i.e. such that  $e : [0, T] \ni t \mapsto \mathbb{E}(M_t)$  is continuously differentiable. The aim is to then prove a universal bound for the Lipschitz constant of  $[0, T] \ni t \mapsto \mathbb{E}(M_t)$ :

**Theorem 4.1.** For a given  $\epsilon \in (0,1)$ , there exists a positive constant  $\alpha_0 \in (0,1]$ , only depending upon  $\epsilon$ , the Lipschitz constant K and the linear growth constant  $\Lambda$ of b, such that, for any  $\alpha \in (0, \alpha_0)$  and any positive time T > 0, there exist a constant  $\mathcal{M}_T$ , only depending on T,  $\epsilon$ , K and  $\Lambda$ , such that, for any initial condition  $X_0 = x_0 \leq 1 - \epsilon$ , any solution to (2.1) according to Definition 2.3 satisfies

$$\forall t \in [0,T], \quad e'(t) = \frac{d}{dt} \mathbb{E}(M_t) \leqslant \mathcal{M}_T.$$

**Remark 4.2.** In the proof below, we provide an expression for  $\alpha_0$  (or put it differently a lower bound for the best possible choice of  $\alpha_0$ ): we refer to the statement of Proposition 4.5, where the constant  $\alpha_0$  therein is the same as the one in the statement of Theorem 4.1. Without any further assumptions on the coefficients, the expression

in Proposition 4.5 is not completely explicit as it depends upon a non-explicit constant. However, it becomes tractable when  $b(x) = -\lambda x$ , which is the common case treated in the neuroscience literature. We thus refer the reader to Subsection 6.1 for a detailed discussion about the critical threshold  $\alpha_0$  in this framework.

4.1. Reformulation of the Equation and *a priori* Bounds for the Solution. In the whole proof, we shall use a reformulated version of the nonlinear equation (2.1). Indeed, given a solution  $(X_t, M_t)_{0 \leq t \leq T}$  to (2.1) on some interval [0, T] according to Definition 2.3, we set

$$Z_t = X_t + M_t, \quad t \in [0, T].$$
 (4.1)

Then, the dynamics of  $(Z_t)_{0 \leq t \leq T}$  have the form:

$$Z_t = X_0 + \int_0^t b(X_s) ds + \alpha \mathbb{E}(M_t) + W_t, \quad t \in [0, T],$$
(4.2)

so that, by continuity of the mapping  $[0,T] \ni t \mapsto e(t) = \mathbb{E}(M_t)$ , the process  $(Z_t)_{0 \leq t \leq T}$  has continuous paths. Moreover, as noted above in Proposition 2.2,  $M_t$  is completely defined in terms of  $Z_t$ , since it holds that

$$M_t = \lfloor \left( \sup_{0 \leqslant s \leqslant t} Z_s \right)_+ \rfloor = \sup_{0 \leqslant s \leqslant t} \lfloor (Z_s)_+ \rfloor.$$

$$(4.3)$$

Following the proof of Proposition 2.2, we claim:

**Lemma 4.3.** There exists a constant  $B(T, \alpha, b)$ , only depending upon T,  $\alpha$ , b and non-decreasing in  $\alpha$ , such that

$$\sup_{0 \leqslant t \leqslant T} e(t) = e(T) \leqslant \mathbb{E} \Big[ \sup_{0 \leqslant t \leqslant T} (Z_t)_+ \Big] \leqslant B(T, \alpha, b).$$
(4.4)

A possible choice for B is

$$B(T, \alpha, b) = \frac{\mathbb{E}[(X_0)_+] + 4T^{1/2} + \Lambda T}{1 - \alpha} \exp\left(\frac{2\Lambda T}{1 - \alpha}\right).$$

*Proof.* The proof is a straightforward consequence of (2.12) in the proof of Proposition 2.2.

4.2. Local Hölder Bound of the Solution. We now turn to the critical point of the proof. Indeed, in the next subsection, we shall prove that, for  $\alpha$  small enough, the function  $t \mapsto e(t) = \mathbb{E}(M_t)$  generated by some solution to (2.1) according to Definition 2.3 (so that *e* is continuously differentiable) satisfies an a priori 1/2-Hölder bound, with an explicit Hölder constant. This acts as the keystone of the argument to extend the local existence and uniqueness result into a global one. As a first step, the proof consists in establishing a local Hölder bound for *e* in the case when the probability that the process X lies in the neighbourhood of 1 is not too large:

**Lemma 4.4.** Keep the same notation as in the previous subsection and consider in particular a solution  $(X_t)_{0 \leq t \leq T}$  to (2.1) on some interval [0,T], with T > 0 and initial condition  $X_0 = x_0$ . Assume in addition that there exist some time  $t_0 \in [0,T]$ 

and two constants  $\epsilon \in (0,1)$  and  $c \in (0,1/\alpha)$  such that  $x_0 \leq 1-\epsilon$ , and, for any Borel subset  $A \subset [1-\epsilon, 1]$ ,

$$\mathbb{P}(X_{t_0} \in A) \leqslant c|A|, \tag{4.5}$$

where |A| stands for the Lebesgue measure of A. Then, with

$$\mathcal{B}_{0} = \frac{\exp(2\Lambda)[(8+5c+8\epsilon^{-1})\Lambda + 4(1+c+\epsilon^{-1})]}{1-c\alpha}.$$

it holds, for any  $h \in (0, 1)$ ,

$$\mathcal{B}_0 \exp(2\Lambda h) h^{1/2} \leqslant \epsilon/2 \\ t_0 + h \leqslant T$$
  $\} \Rightarrow e(t_0 + h) - e(t_0) \leqslant \mathcal{B}_0 h^{1/2}$ 

*Proof.* Without any loss of generality, we can assume  $t_0 = 0$ , with T being understood as  $T - t_0$ . Indeed, setting

$$X_t^{\sharp_{t_0}} := X_{t_0+t}, \quad t \in [0, T - t_0], \tag{4.6}$$

we observe that, for  $t \in [0, T - t_0]$ ,

$$X_t^{\sharp_{t_0}} = X_{t_0} + \int_0^t b(X_r^{\sharp_{t_0}}) dr + \alpha \mathbb{E}(M_{t+t_0} - M_{t_0}) + W_{t+t_0} - W_{t_0} - (M_{t+t_0} - M_{t_0}).$$
(4.7)

Here  $M_{t+t_0} - M_{t_0}$  represents the number of times the process X reaches 1 within the interval  $(t_0, t+t_0]$ . Therefore, this also matches the number of times the process  $X^{\sharp t_0}$  hits 1 within the interval (0, t], so that  $X^{\sharp t_0}$  indeed satisfies the nonlinear equation (2.1) on  $[0, T - t_0]$ , with  $X_0^{\sharp t_0} = X_{t_0}$  as initial condition and with respect to the shifted Brownian motion  $(W_t^{\sharp t_0} := W_{t_0+t} - W_{t_0})_{0 \leq t \leq T-t_0}$ . In what follows,  $t_0$  is thus assumed to be zero, the new T standing for the previous  $T - t_0$  and the new  $X_0$  matching the previous  $X_{t_0}$  and thus satisfying (4.5).

For a given  $h \in (0, 1)$ , such that  $h \leq T$ , and a given  $\mathcal{B}_0 > 0$  (the value of which will be fixed later), we then define the deterministic hitting time:

$$R = \inf \left\{ t \in [0,h] : \mathbb{E}(M_t) = e(t) \ge \mathcal{B}_0 h^{1/2} \right\}.$$

Following the proof of (2.11) (see more specifically (2.9)), we have, for any  $t \in [0, h \wedge R]$ ,

$$M_{t} \leq \sup_{0 \leq s \leq t} (Z_{s})_{+} \leq (X_{0})_{+} + \Lambda \int_{0}^{t} (1 + (Z_{s})_{+} + M_{s}) ds + \alpha e(t) + 2 \sup_{0 \leq s \leq t} |W_{s}|$$
$$\leq (X_{0})_{+} + 2\Lambda \int_{0}^{t} (1 + M_{s}) ds + \alpha \mathcal{B}_{0} h^{1/2} + 2 \sup_{0 \leq s \leq t} |W_{s}|$$
$$\leq (X_{0})_{+} + 2\Lambda h + 2\Lambda \int_{0}^{t} M_{s} ds + \alpha \mathcal{B}_{0} h^{1/2} + 2 \sup_{0 \leq s \leq t} |W_{s}|,$$

where we have used (4.3) to pass from the first to the second line. By Gronwall's Lemma, we obtain

$$M_t \leqslant \exp(2\Lambda h) \big[ (X_0)_+ + 2\Lambda h + \alpha \mathcal{B}_0 h^{1/2} + 2 \sup_{0 \leqslant s \leqslant h} |W_s| \big],$$
  
$$\leqslant (X_0)_+ + \exp(2\Lambda h) \big[ 4\Lambda h + \alpha \mathcal{B}_0 h^{1/2} + 2 \sup_{0 \leqslant s \leqslant h} |W_s| \big],$$
(4.8)

as  $\exp(2\Lambda h) \leq 1 + 2\Lambda h \exp(2\Lambda h)$  and  $(X_0)_+ \leq 1$ .

Assume that  $\mathcal{B}_0 \exp(2\Lambda h) h^{1/2} \leq \epsilon/2 \leq 1/2$ . Then, by Doob's maximal inequality for martingales,

$$\sum_{k \ge 2} \mathbb{P}(M_t \ge k) \leqslant \sum_{k \ge 2} \mathbb{P}\left(\exp(2\Lambda h)\left[4\Lambda h + 2\sup_{0 \le s \le h} |W_s|\right] \ge k - 3/2\right)$$
  
$$\leqslant 2\exp(2\Lambda h)\mathbb{E}\left[4\Lambda h + 2|W_h|\right] \leqslant \exp(2\Lambda h)\left[8\Lambda h + 4h^{1/2}\right].$$
(4.9)

Moreover,

$$\mathbb{P}(M_t \ge 1)$$

$$\leq \mathbb{P}((X_0)_+ + \exp(2\Lambda h) \left[ 4\Lambda h + \alpha \mathcal{B}_0 h^{1/2} + 2 \sup_{0 \le s \le h} |W_s| \right] \ge 1)$$

$$\leq \mathbb{P}(X_0 \in [1 - \epsilon, 1], X_0 + \exp(2\Lambda h) \left[ 4\Lambda h + \alpha \mathcal{B}_0 h^{1/2} + 2 \sup_{0 \le s \le h} |W_s| \right] \ge 1)$$

$$+ \mathbb{P}(\exp(2\Lambda h) \left[ 4\Lambda h + 2 \sup_{0 \le s \le h} |W_s| \right] \ge \epsilon/2)$$

$$:= I_1 + I_2,$$

where we used  $\mathcal{B}_0 \exp(2\Lambda h) h^{1/2} \leq \epsilon/2$  in the third line.

By Doob's maximal inequality again, we deduce that

$$I_2 \leqslant 2\exp(2\Lambda h)\epsilon^{-1}\mathbb{E}\left[4\Lambda h + 2|W_h|\right] \leqslant \exp(2\Lambda h)\epsilon^{-1}\left[8\Lambda h + 4h^{1/2}\right].$$
(4.10)

We now switch to  $I_1$ . By independence of  $X_0$  and  $(W_s)_{0 \leq s \leq T}$  and by (4.5),

$$I_{1} \leq c \int_{0}^{\epsilon} \mathbb{P}\left\{\exp(2\Lambda h)\left[4\Lambda h + \alpha \mathcal{B}_{0}h^{1/2} + 2\sup_{0 \leq s \leq h}|W_{s}|\right] \geq x\right\} dx$$
$$\leq c \int_{0}^{+\infty} \mathbb{P}\left\{\exp(2\Lambda h)\left[4\Lambda h + \alpha \mathcal{B}_{0}h^{1/2} + 2\sup_{0 \leq s \leq h}|W_{s}|\right] \geq x\right\} dx$$
$$= c \exp(2\Lambda h)\mathbb{E}\left[4\Lambda h + \alpha \mathcal{B}_{0}h^{1/2} + 2\sup_{0 \leq s \leq h}|W_{s}|\right].$$

By Doob's  $L^2$  inequality,

$$I_1 \leqslant c \exp(2\Lambda h) \left[ 4\Lambda h + \alpha \mathcal{B}_0 h^{1/2} + 4h^{1/2} \right].$$

Together with (4.10), we deduce that

$$\mathbb{P}\{M_t \ge 1\} \leqslant \exp(2\Lambda h) \left[4(c+2\epsilon^{-1})\Lambda h + 4(c+\epsilon^{-1})h^{1/2} + c\alpha \mathcal{B}_0 h^{1/2}\right].$$
  
From (4.9), we finally obtain, for  $t \leqslant R \wedge h$ ,

$$\mathbb{E}(M_t) = \sum_{k \ge 1} \mathbb{P}(M_t \ge k)$$
  

$$\leqslant \exp(2\Lambda h) \left[ 4(2+c+2\epsilon^{-1})\Lambda h + 4(1+c+\epsilon^{-1})h^{1/2} + c\alpha \mathcal{B}_0 h^{1/2} \right]$$
  

$$\leqslant \exp(2\Lambda h) \left[ (8+5c+8\epsilon^{-1})\Lambda h + 4(1+c+\epsilon^{-1})h^{1/2} \right] + c\alpha \mathcal{B}_0 h^{1/2},$$

provided  $\mathcal{B}_0 \exp(2\Lambda h) h^{1/2} \leq \epsilon/2 \leq 1/2$ , which implies

$$c\alpha \mathcal{B}_0 \exp(2\Lambda h)h^{1/2} \leq c\alpha \mathcal{B}_0 h^{1/2} + c\Lambda h$$

(using the fact that  $e^{2\Lambda h} \leq 1 + 2\Lambda h e^{2\Lambda h}$ ). Therefore, if  $R \leq h$ , then we can choose t = R in the left-hand side above. By continuity of e on [0, T], it then holds  $e(R) = \mathcal{B}_0 h^{1/2}$ , so that

$$(1 - c\alpha)\mathcal{B}_0 h^{1/2} \leq \exp(2\Lambda h) \left[ (8 + 5c + 8\epsilon^{-1})\Lambda h + 4(1 + c + \epsilon^{-1})h^{1/2} \right] < \exp(2\Lambda) \left[ (8 + 5c + 8\epsilon^{-1})\Lambda + 4(1 + c + \epsilon^{-1}) \right] h^{1/2},$$

which is not possible when

$$\mathcal{B}_{0} = \frac{\exp(2\Lambda)[(8+5c+8\epsilon^{-1})\Lambda + 4(1+c+\epsilon^{-1})]}{1-c\alpha}$$

Precisely, with  $\mathcal{B}_0$  as above and  $\mathcal{B}_0 \exp(2\Lambda h)h^{1/2} \leqslant \epsilon/2$  it cannot hold  $R \leqslant h$ .

## 4.3. Global Hölder Bound. In this subsection, we shall prove

**Proposition 4.5.** For a given  $\epsilon \in (0, 1)$ , there exists a positive constant  $\alpha_0 \in (0, 1]$ , only depending upon  $\epsilon$ , K and  $\Lambda$ , such that, for any  $\alpha < \alpha_0$ , there exists a constant  $\mathcal{B}$ , only depending on  $\alpha$ ,  $\epsilon$ , K and  $\Lambda$ , such that, for any positive time T > 0 and any initial condition  $X_0 = x_0 \leq 1 - \epsilon$ , any solution to (2.1) according to Definition 2.3, so that the mapping  $e : [0,T] \ni t \mapsto \mathbb{E}(M_t)$  is continuously differentiable, satisfies, for any  $h \in (0,1)$  and  $t_0 \in [0,T]$ ,

$$\begin{array}{l} \mathcal{B}h^{1/2} \leqslant \epsilon/2 \\ t_0 + h \leqslant T \end{array} \right\} \Rightarrow e(t_0 + h) - e(t_0) \leqslant \mathcal{B}h^{1/2}.$$

Note that  $\mathcal{B}$  above may differ from  $\mathcal{B}_0$  in the statement of Lemma 4.4. The constant  $\alpha_0$  can be described as follows. Defining  $T_0$  as the largest time less than 1 such that

$$(1-\epsilon)\exp(\Lambda T_0) \leq 1-7\epsilon/8, \quad \Lambda T_0\exp(\Lambda T_0) \leq \epsilon/8,$$

then  $\alpha_0$  can be chosen as the largest (positive) real satisfying (with  $B(T_0, \alpha_0, b)$  as in Lemma 4.3)

$$\begin{aligned} \alpha_0 B(T_0, \alpha_0, b) &\leqslant \epsilon/4, \\ \alpha_0 2^{3/2} (c')^{3/2} \exp\left(-\frac{1}{2}\right) \left[\epsilon^{-1} + B(T_0, \alpha_0, b)\right] &\leqslant 1, \\ \alpha_0 \left[c' T_0^{-1/2} + 2^{3/2} (c')^{3/2} \exp\left(-\frac{1}{2}\right) B(T_0, \alpha_0, b)\right] &\leqslant 1 \end{aligned}$$

where the constant c' is defined by the following property: c' > 0 depending on K only is such that for any diffusion process  $(U_t)_{0 \leq t \leq 1}$  satisfying

$$dU_t = F(t, U_t)dt + dW_t, \quad t \in [0, 1],$$

where  $U_0 = 0$  and  $F : [0, T] \times \mathbb{R} \to \mathbb{R}$  is K-Lipschitz in x and satisfies F(t, 0) = 0for any  $t \in [0, 1]$ , it holds that

$$\frac{d}{dx}\mathbb{P}(U_t \in dx) \leqslant \frac{c'}{\sqrt{t}} \exp\left(-\frac{x^2}{c't}\right), \quad x \in \mathbb{R}, \ t \in (0,1].$$

The proof relies on the following

**Lemma 4.6.** Given an initial condition  $X_0 = x_0 \leq 1 - \epsilon$ , with  $\epsilon \in (0, 1)$ , and a solution  $(X_t)_{0 \leq t \leq T}$  to (2.1) on some interval [0, T], the random variable  $X_t$  has a density on  $(-\infty, 1]$ , for any  $t \in (0, T]$ . Moreover, defining  $T_0$  as in the statement of Proposition 4.5 and choosing  $\alpha \leq \alpha_1$  satisfying

$$\alpha_1 B(T_0, \alpha_1, b) \leqslant \epsilon/4,$$

*then, for*  $x \in [1 - \epsilon/4, 1)$ *,* 

$$\frac{d}{dx}\mathbb{P}(X_t \in dx) \leqslant 2^{3/2} (c')^{3/2} \exp\left(-\frac{1}{2}\right) \left[\epsilon^{-1} + B(T_0, \alpha, b)\right] \quad \text{if } t \leqslant T_0$$
$$\frac{d}{dx}\mathbb{P}(X_t \in dx) \leqslant c' T_0^{-1/2} + 2^{3/2} (c')^{3/2} \exp\left(-\frac{1}{2}\right) B(T_0, \alpha, b) \quad \text{if } t > T_0,$$

where the constant c' is as in the statement of Proposition 4.5.

Before we prove Lemma 4.6, we introduce some materials. As usual, we set  $e(t) = \mathbb{E}(M_t)$ , for  $t \in [0, T]$ , the mapping e being assumed to be continuously differentiable on [0, T]. Moreover, with  $(X_t)_{t \in [0,T]}$ , we associate the sequence of hitting times  $(\tau_k)_{k \geq 0}$  given by (2.3). We then investigate the marginal distributions of  $(X_t)_{t \in [0,T]}$ . Given a Borel subset  $A \subset (-\infty, 1]$ , we write in the same way as in the proof of (3.1):

$$\mathbb{P}(X_t \in A) = \mathbb{P}(X_t \in A, \tau_1 > t) + \sum_{k \ge 1} \int_0^t \mathbb{P}(X_t \in A, \tau_{k+1} > t | \tau_k = s) \mathbb{P}(\tau_k \in ds),$$

$$(4.11)$$

where the notation  $\mathbb{P}(\cdot | \tau_k = s)$  stands for the conditional law given  $\tau_k = s$ . Following (4.6) and (4.7), we can shift the system by length  $s \in [0, T]$ . Precisely, we know that  $(X_r^{\sharp_s} := X_{s+r})_{r \in [0, T-s]}$  satisfies

$$X_{r}^{\sharp_{s}} = X_{s} + \int_{0}^{r} b(X_{u}^{\sharp_{s}}) du + \alpha e^{\sharp_{s}}(r) + W_{s+r} - W_{s} - M_{r}^{\sharp_{s}}, \qquad (4.12)$$

with

$$e^{\sharp_{s}}(r) := e(s+r) - e(s) \quad ; \quad M_{r}^{\sharp_{s}} := \sum_{k \ge 1} \mathbf{1}_{(0,r]}(\tau_{k}^{\sharp_{s}}) \quad ;$$

$$\tau_{k}^{\sharp_{s}} := \inf\{u > \tau_{k-1}^{\sharp_{s}} : X_{s+u-} \ge 1\}, \quad k \ge 1 \quad ; \quad \tau_{0}^{\sharp_{s}} := 0.$$

$$(4.13)$$

Conditionally on  $\tau_k = s$ , the law of  $(X_r^{\sharp_s})_{0 \leq r \leq T-s}$  until  $\tau_1^{\sharp_s}$  coincides with the law of  $(\hat{Z}_r^{\sharp_s,0})_{0 \leq r \leq T-s}$  until the first time it reaches 1, where, for a given  $\mathcal{F}_0$ -measurable initial condition  $\zeta$  with values in  $(-\infty, 1)$ ,  $(\hat{Z}_r^{\sharp_s,\zeta})_{0 \leq r \leq T-s}$  stands for the solution of the SDE:

$$\hat{Z}_{r}^{\sharp_{s},\zeta} = \zeta + \int_{0}^{r} b(\hat{Z}_{u}^{\sharp_{s},\zeta}) du + \alpha e^{\sharp_{s}}(r) + W_{r}, \quad t \in [0, T-s].$$
(4.14)

Below, we will write  $\hat{Z}_r^{\zeta}$  for  $\hat{Z}_r^{\sharp_0,\zeta}$ . In the end, we thus have (by (4.11)):

$$\mathbb{P}(X_t \in A) \leqslant \mathbb{P}(\hat{Z}_t^{X_0} \in A) + \sum_{k \geqslant 1} \int_0^t \mathbb{P}(\hat{Z}_{t-s}^{\sharp_s,0} \in A) \mathbb{P}(\tau_k \in ds)$$

$$= \mathbb{P}(\hat{Z}_t^{X_0} \in A) + \int_0^t \mathbb{P}(\hat{Z}_{t-s}^{\sharp_s,0} \in A) e'(s) ds,$$
(4.15)

for any Borel set  $A \subset (-\infty, 1]$ , the passage from the first to the second line following from (3.9).

We can now turn to:

Proof of Lemma 4.6. Given an initial condition  $x_0 \in (-\infty, 1-\epsilon]$  for  $\varepsilon \in (0,1)$ , we know from Delarue and Menozzi [6] that  $\hat{Z}_t^{x_0}$  has a density for any  $t \in (0,T]$  (and thus  $\hat{Z}_{t-s}^{\sharp_{s,0}}$  as well for  $0 \leq s < t$ ). From (4.15), we deduce that the law of  $X_t$  has a density on  $(-\infty, 1]$  since  $\mathbb{P}(X_t \in A) = 0$  when |A| = 0 where |A| stands for the Lebesgue measure of A. Moreover, there exists a constant  $c' \geq 1$ , depending on K only, such that, for any  $t \in [0, T \land 1]$ :

$$\frac{d}{dx}\mathbb{P}(\hat{Z}_t^{x_0} \in dx) \leqslant \frac{c'}{\sqrt{t}} \exp\left(-\frac{[x - \vartheta_t^{x_0}]^2}{c't}\right),\tag{4.16}$$

where  $\vartheta_t^{x_0}$  is the solution of the ODE:

$$\frac{d}{dt}\vartheta_t = b(\vartheta_t) + \alpha e'(t), \quad t \in [0, T],$$
(4.17)

with  $\vartheta_0^{x_0} = x_0$ . Above, the function  $[0,T] \ni t \mapsto e(t)$  represents  $[0,T] \ni t \mapsto \mathbb{E}(M_t)$ given  $X_0 = x_0$ , which means that the initial condition  $x_0$  of  $X_0$  upon which *e* depends is fixed once for all, independently of the initial condition of  $\vartheta$ . In particular, as the initial condition of  $\vartheta$  varies, the function *e* does not. We emphasize that *c'* is independent of *e* and can be taken to be that defined in Proposition 4.5. Indeed, we can write  $\mathbb{P}(\hat{Z}_t^{x_0} \in dx)$  as  $\mathbb{P}(\hat{Z}_t^{x_0} - \vartheta_t^{x_0} \in d(x - \vartheta_t^{x_0}))$ , with

$$d(\hat{Z}_t^{x_0} - \vartheta_t^{x_0}) = F(t, \hat{Z}_t^{x_0} - \vartheta_t^{x_0})dt + dW_t, \quad t \in [0, T] ; \quad \hat{Z}_0^{x_0} - \vartheta_0^{x_0} = 0 ;$$
  
$$F(t, x) = b(x + \vartheta_t^{x_0}) - b(\vartheta_t^{x_0}), \quad t \in [0, T], \quad x \in \mathbb{R}.$$

We then notice that  $F(t, \cdot)$  is K-Lipschitz continuous (since b is) and satisfies F(t, 0) = 0, so that, referring to [6], all the parameters involved in the definition of the constant c' are independent of e. The fact that c' is independent of e is crucial. As a consequence, we can bound  $(d/dx)\mathbb{P}(\hat{Z}_{t-s}^{\sharp_{s},0} \in dx)$  in a similar way, that is, with the same constant c' as in (4.16): for any  $0 \leq s < t \leq T$ , with  $t - s \leq 1$ ,

$$\frac{d}{dx}\mathbb{P}(\hat{Z}_{t-s}^{\sharp_{s},0} \in dx) \leqslant \frac{c'}{\sqrt{t-s}} \exp\left(-\frac{[x-\vartheta_{t-s}^{\sharp_{s},0}]^2}{c'(t-s)}\right),\tag{4.18}$$

where  $\vartheta^{\sharp_s,0}$  is the solution of the ODE:

$$\frac{d}{dt}\vartheta_t^{\sharp_s} = b\big(\vartheta_t^{\sharp_s}\big) + \alpha \frac{d}{dt}e^{\sharp_s}(t), \quad t \in [0, T-s],$$

with  $\vartheta_0^{\sharp_{s},0} = 0$  as initial condition.

Bound of the density in small time. Keep in mind that  $X_0 = x_0 \leq 1-\epsilon$ . Therefore, by the comparison principle for ODEs,  $\vartheta_t^{x_0} \leq \vartheta_t^{1-\epsilon}$  for any  $t \in [0, T]$ , so that by Gronwall's Lemma

$$\vartheta_t^{x_0} \leqslant \vartheta_t^{1-\epsilon} \leqslant (1-\epsilon + \Lambda T + \alpha e(T)) \exp(\Lambda T).$$

By Lemma 4.3, we know that  $e(T) \leq B(T, \alpha, b)$ , so that

$$\vartheta_t^{x_0} \leqslant \left(1 - \epsilon + \Lambda T + \alpha B(T, \alpha, b)\right) \exp(\Lambda T).$$
(4.19)

Now choose  $T_0$  as in Proposition 4.5, i.e.  $T_0 \leq 1$  such that

$$(1-\epsilon)\exp(\Lambda T_0) \leqslant 1-7\epsilon/8, \quad \Lambda T_0\exp(\Lambda T_0) \leqslant \epsilon/8,$$

and then take  $\alpha_1 \in (0, 1)$  such that

$$\alpha_1 B(T_0, \alpha_1, b) \exp(\Lambda T_0) \leqslant \epsilon/4.$$

Then, whenever  $\alpha \leq \alpha_1$ , it holds that

$$\vartheta_t^{x_0} \leqslant 1 - \epsilon/2, \quad t \in [0, T_0 \wedge T]$$

Therefore, for  $x \ge 1 - \epsilon/4$ ,

$$\exp\left(-\frac{[x-\vartheta_t^{x_0}]^2}{c't}\right) \leqslant \exp\left(-\frac{\epsilon^2}{16c't}\right), \quad t \in [0, T_0 \wedge T].$$
(4.20)

Similarly,

$$\vartheta_{t-s}^{\sharp_s,0} \leqslant 3\epsilon/8 \leqslant 3/8, \quad 0 \leqslant s \leqslant t \leqslant T_0 \wedge T.$$

Indeed,  $e^{\sharp_s}(T-s) \leq e(T)$  for  $s \in [0,T]$ , so that (4.19) applies to  $\vartheta_{t-s}^{\sharp_s,0}$  with  $1-\epsilon$  therein being replaced by 0. Therefore, for  $x \geq 1-\epsilon/4$ , it holds  $x - \vartheta_{t-s}^{\sharp_s,0} \geq 3/4 - 3/8 = 3/8 \geq 1/4$ , so that

$$\exp\left(-\frac{[x-\vartheta_{t-s}^{\sharp_s,0}]^2}{c'(t-s)}\right) \leqslant \exp\left(-\frac{1}{16c'(t-s)}\right), \quad 0 \leqslant s < t \leqslant T_0 \wedge T.$$
(4.21)

In the end, for  $x \in (1 - \epsilon/4, 1)$  and  $t \leq T_0 \wedge T$ , we deduce from (4.15), (4.16), (4.18), (4.20), (4.21) and Lemma 4.4 again, that

$$\frac{d}{dx}\mathbb{P}(X_t \in dx) \leqslant c'\varpi_0 \big[\epsilon^{-1} + e(T \wedge T_0)\big] \leqslant c'\varpi_0 \big[\epsilon^{-1} + B(T_0, \alpha, b)\big],$$

where

$$\varpi_0 = \sup_{t>0} \left[ t^{-1/2} \exp\left(-\frac{1}{16c't}\right) \right] = 4\sqrt{c'} \sup_{u>0} \left[ u \exp(-u^2) \right] = 2^{3/2} \sqrt{c'} \exp\left(-\frac{1}{2}\right).$$

Bound of the density in long time. We now discuss what happens for  $T > T_0$  and  $t \in [T_0, T]$ . Then,

$$\frac{d}{dx}\mathbb{P}(X_t \in dx) \leqslant \frac{d}{dx}\mathbb{P}\left(X_t \in dx, \tau_1^{\sharp_{t-T_0}} \leqslant T_0\right) + \frac{d}{dx}\mathbb{P}\left(X_t \in dx, \tau_1^{\sharp_{t-T_0}} > T_0\right)$$
(4.22)  
=  $\pi_1 + \pi_2$ ,

with  $\tau_1^{\sharp_{t-T_0}} = \inf\{u > 0 : X_{t-T_0+u-} \ge 1\} = \inf\{u > 0 : X_{u-}^{\sharp_{t-T_0}} \ge 1\}$ . The above expression says that we split the event  $(X_t \text{ is in the neighbourhood of } x)$  into two disjoint parts according to the fact that X reaches the threshold or not within the

time window  $[t - T_0, t]$ . We have chosen this interval to be of length  $T_0$  in order to apply the results in small time.

We first investigate  $\pi_2$ . The point is that, on the event that  $\tau_1^{\sharp_{t-T_0}} > T_0$  and within the time window  $[t - T_0, t]$ , X behaves as a standard diffusion process without any jumps, namely as a process with the same dynamics as  $\hat{Z}^{\sharp_{t-T_0}, X_{t-T_0}}$ . Following (4.16), we then have

$$\pi_{2} = \frac{d}{dx} \mathbb{P} \left( \hat{Z}_{T_{0}}^{\sharp_{t-T_{0}}, X_{t-T_{0}}} \in dx, \tau_{1}^{\sharp_{t-T_{0}}} \leqslant T_{0} \right)$$
$$\leqslant \frac{d}{dx} \mathbb{P} \left( \hat{Z}_{T_{0}}^{\sharp_{t-T_{0}}, X_{t-T_{0}}} \in dx \right) \leqslant \sup_{z \leqslant 1} \frac{d}{dx} \mathbb{P} \left( \hat{Z}_{T_{0}}^{\sharp_{t-T_{0}}, z} \in dx \right) \leqslant c' T_{0}^{-1/2}.$$
(4.23)

We now turn to  $\pi_1$ . Here we write that

$$\begin{aligned} \pi_1 &= \mathbb{P}\left(X_t \in dx, \tau_1^{\sharp_{t-T_0}} \leqslant T_0\right) = \sum_{k \ge 1} \mathbb{P}\left(X_t \in dx, \tau_k^{\sharp_{t-T_0}} \leqslant T_0 < \tau_{k+1}^{\sharp_{t-T_0}}\right) \\ &= \sum_{k \ge 1} \int_0^{T_0} \mathbb{P}\left(X_t \in dx, \tau_k^{\sharp_{t-T_0}} \leqslant T_0 < \tau_{k+1}^{\sharp_{t-T_0}} | \tau_k^{\sharp_{t-T_0}} = s\right) \mathbb{P}(\tau_k^{\sharp_{t-T_0}} \in ds) \\ &= \sum_{k \ge 1} \int_0^{T_0} \mathbb{P}\left(\hat{Z}_{T_0-s}^{\sharp_{s+t-T_0},0} \in dx, T_0 < \tau_{k+1}^{\sharp_{t-T_0}}\right) \mathbb{P}(\tau_k^{\sharp_{t-T_0}} \in ds), \end{aligned}$$

since on the event that  $\tau_k^{\sharp_{t-T_0}} \leq T_0 < \tau_{k+1}^{\sharp_{t-T_0}}$  and given that the k-th (and last) jump of X in the interval  $[t - T_0, t]$  occurs at time  $t - T_0 + s$  with  $s \in [0, T_0]$ , we have that the process  $X_r$  for  $r \in [t - T_0 + s, t]$  coincides with the process  $\hat{Z}_u^{\sharp_{s+t-T_0}, 0}$  for  $u \in [0, T_0 - s]$ . Thus

$$\pi_{1} \leqslant \sum_{k \geqslant 1} \int_{0}^{T_{0}} \frac{d}{dx} \mathbb{P}\left(\hat{Z}_{T_{0}-s}^{\sharp_{s+t-T_{0}},0} \in dx\right) \mathbb{P}(\tau_{k}^{\sharp_{t-T_{0}}} \in ds)$$

$$= \int_{0}^{T_{0}} \frac{d}{dx} \mathbb{P}\left(\hat{Z}_{T_{0}-s}^{\sharp_{s+t-T_{0}},0} \in dx\right) e'(s+t-T_{0}) ds.$$
(4.24)

By (4.18), we have

$$\int_{0}^{T_{0}} \frac{d}{dx} \mathbb{P}\left(\hat{Z}_{T_{0}-s}^{\sharp_{s+t-T_{0}},0} \in dx\right) e'(s+t-T_{0})ds$$
$$\leqslant \int_{0}^{T_{0}} \frac{c'}{\sqrt{T_{0}-s}} \exp\left(-\frac{[x-\vartheta_{T_{0}-s}^{\sharp_{s+t-T_{0}},0}]^{2}}{c'(T_{0}-s)}\right) e'(s+t-T_{0})ds$$

Following the shift in (4.12), it is well seen that the mapping  $[0, T_0] \ni s \mapsto e^{\sharp_{t-T_0}}(s)$  satisfies Lemma 4.3, that is

$$\sup_{0 \leq s \leq T_0} e^{\sharp_{t-T_0}}(s) = \sup_{0 \leq s \leq T_0} \left[ e(s+t-T_0) - e(t-T_0) \right] = e(t) - e(t-T_0) \leq B(T_0, \alpha, b).$$

Therefore, we can follow the same strategy as in short time. Indeed, for  $\alpha \leq \alpha_1$ , by the choice of  $T_0$  as before, it holds that

$$\pi_1 \leqslant c' \varpi_0 B(T_0, \alpha, b),$$

for  $x \in [1-\epsilon/4, 1)$ . Using (4.23) and the above bound, we deduce that, for  $t \in [T_0, T]$ ,

$$\frac{d}{dx}\mathbb{P}\left(X_t \in dx\right) \leqslant c' \big[T_0^{-1/2} + \varpi_0 B(T_0, \alpha, b)\big].$$

Now, we can complete the proof of Proposition 4.5.

Proof of Proposition 4.5. Proposition 4.5 follows from the combination of Lemmas 4.4 and 4.6. Indeed, given  $T_0$  and  $\alpha_0$  as defined in Proposition 4.5, then by Lemma 4.6 it follows that  $\mathbb{P}(X_t \in A) < (1/\alpha)|A|$  for any Borel subset  $A \subset [1 - \epsilon/4, 1]$ , any  $\alpha < \alpha_0$  and any  $t \in [0, T]$ . The result follows by Lemma 4.4, with  $\mathcal{B}$  being given by  $\mathcal{B}_0 \exp(2\Lambda)$  with  $\epsilon$  in  $\mathcal{B}_0$  replaced by  $\epsilon/4$ .

4.4. Estimate of the Density of the Killed Process. In light of the previous subsection, for a solution  $(X_t)_{0 \leq t \leq T}$  to (2.1) such that the mapping  $[0,T] \ni t \mapsto e(t) = \mathbb{E}(M_t)$  is continuously differentiable, we here investigate:

$$\frac{d}{dx}\mathbb{P}\left(X_t \in dx, t < \tau_1\right), \quad t \in [0, T], \ x \leqslant 1,$$

where  $\tau_1 = \inf\{t > 0 : X_{t-} \ge 1\}$  as usual, which is the density of the killed process  $(X_{t \land \tau_1})_{t \in [0,T]}$  and which makes sense because of Lemma 4.6.

Here is the main result of this subsection:

**Lemma 4.7.** For any  $\epsilon \in (0, 1)$ , any T > 0 and any  $\mathcal{B} > 0$ , there exist two constants  $\mu_T$  and  $\eta_T$ , only depending upon T,  $\mathcal{B}$ ,  $\epsilon$ , K and  $\Lambda$ , such that, for any initial condition  $x_0 \leq 1 - \epsilon$  and any continuously differentiable non-decreasing deterministic mapping  $[0, T] \ni t \mapsto e(t)$  satisfying

$$e(0) = 0$$
 ;  $|e(t) - e(s)| \leq \mathcal{B}(t-s)^{1/2}$ ,  $(s,t) \in [0,T]^2$ ,

if  $(\chi_t)_{0 \leq t \leq T}$  denotes the solution of the SDE

$$d\chi_t = b(\chi_t)dt + \alpha e'(t) + dW_t, \quad t \in [0, T] ; \quad \chi_0 = x_0$$

then, with p(t, y) denoting the density of  $\chi_t$  killed at 1 as in (3.2), we have

$$p(t,y) \leq \mu_T (1-y)^{\eta_T}, \quad t \in [0,T], \ y \in [1-\epsilon/4,1].$$
 (4.25)

*Proof. First Step.* The first step is to provide a probabilistic representation for p. For a given  $(T, x) \in (0, +\infty) \times (-\infty, 1)$ , we consider the solution to the SDE:

$$dY_t = -[b(Y_t) + \alpha e'(T-t)]dt + dW_t, \quad Y_0 = y,$$
(4.26)

together with some stopping time  $\rho \leq \rho_0 \wedge T$ , where  $\rho_0 = \inf\{t \in [0, T] : Y_t \geq 1\}$ (with  $\inf \emptyset = +\infty$ ). Then, by Lemma 3.2 and by Itô-Krylov formula (see [12, Chapter II, Section 10]),

$$\begin{aligned} d\big(p(T-t,Y_t)\big) \\ &= -\partial_t p(T-t,Y_t) dt - \big[b(Y_t) + \alpha e'(T-t)\big] \partial_y p(T-t,Y_t) dt + \frac{1}{2} \partial_{yy}^2 p(T-t,Y_t) dt \\ &+ \partial_y p(T-t,Y_t) dW_t \\ &= b'(Y_t) p(T-t,Y_t) dt + \partial_y p(T-t,Y_t) dW_t, \end{aligned}$$

for  $0 \leq t \leq \rho$ . Therefore, the Feynman-Kac formula yields

$$p(T,y) = \mathbb{E}\left[p(T-\rho,Y_{\rho})\mathbf{1}_{\{Y_{\rho}\neq 1\}}\exp\left(-\int_{0}^{\rho}b'(Y_{s})ds\right)\right],\tag{4.27}$$

the indicator function following from the Dirichlet boundary condition satisfied by  $p(\cdot, 1)$ .

Second Step. We now specify the choice of  $\rho$ . Given some free parameters  $L \ge 1$ and  $\delta \in (0, \epsilon/4)$  such that  $L\delta \le \epsilon/4$ , we assume that the initial condition y in (4.26) is in  $(1 - \delta, 1)$  and then consider the stopping time:

$$\rho = \inf\{t \in [0, T] : Y_t \notin (1 - L\delta, 1)\} \land 4\delta^2.$$
(4.28)

Assume that  $4\delta^2 \leq T$ . By (4.27), we deduce that

$$p(T,y) \leqslant \exp(4K\delta^2)(1 - \mathbb{P}(Y_\rho = 1)) \sup_{(t,z)\in\mathcal{Q}(\delta,L)} p(t,z), \qquad (4.29)$$

with

$$\mathcal{Q}(\delta,L) = \left\{ (t,z) \in [T-4\delta^2,T] \times [1-L\delta,1] \right\}.$$

The point is then to give a lower bound for  $\mathbb{P}(Y_{\rho} = 1)$ . By assumption, we know that e is  $\mathcal{B}$  Hölder continuous on [0, T]. Therefore, since  $Y_0 = y \in (1 - \delta, 1)$ , we have, for any  $t \in [0, \delta^2 \wedge \rho]$ ,

$$W_t \ge 1 - \delta - m\delta^2 - \alpha \mathcal{B}\delta + W_t,$$

with

$$m = \sup_{0 \leqslant z \leqslant 1} |b(z)|. \tag{4.30}$$

Therefore, for  $m\delta \leq 1$ ,

$$W_t \ge 1 - 2\delta - \alpha \mathcal{B}\delta + W_t, \quad t \in [0, \delta^2 \wedge \rho],$$

so that

$$\left\{Y_{\rho}=1\right\}\supset\left\{\sup_{0\leqslant t\leqslant\delta^{2}}W_{t}>(2+\alpha\mathcal{B})\delta\right\}\cap\left\{\inf_{0\leqslant t\leqslant\delta^{2}}W_{t}>(2+\alpha\mathcal{B}-L)\delta\right\}.$$
 (4.31)

Choosing  $L = 3 + \alpha \mathcal{B}$  and applying a scaling argument, we deduce that

$$\mathbb{P}\left\{\sup_{\substack{0 \leq t \leq \delta^{2}}} W_{t} > (2 + \alpha \mathcal{B})\delta\right\} \cap \left\{\inf_{\substack{0 \leq t \leq \delta^{2}}} W_{t} > (2 + \alpha \mathcal{B} - L)\delta\right\} \\
= \mathbb{P}\left\{\left\{\sup_{\substack{0 \leq t \leq 1}} W_{t} > (2 + \alpha \mathcal{B})\right\} \cap \left\{\inf_{\substack{0 \leq t \leq 1}} W_{t} > -1\right\}\right\} =: c'' \in (0, 1).$$
(4.32)

We note that the above quantity c'' is independent of  $\delta$  and T. Moreover, we deduce from (4.31) that  $\mathbb{P}(Y_{\rho} = 1) \ge c''$  and therefore, from (4.29), that

$$p(T,y) \leqslant (1-c'') \exp(4K\delta^2) \sup_{z \in \mathcal{I}(L\delta)} \sup_{t \in [0,T]} p(t,z),$$

with  $\mathcal{I}(r) = [1 - r, 1]$ , for r > 0. Choosing  $\delta$  small enough such that  $(1 - c'') \exp(4K\delta^2) \leq (1 - c''/2)$ , we obtain

$$p(T,y) \leqslant \left(1 - \frac{c'}{2}\right) \sup_{z \in \mathcal{I}(L\delta)} \sup_{t \in [0,T]} p(t,z), \quad y \in \mathcal{I}(\delta).$$

Modifying c'' if necessary (c'' being chosen as small as needed), we can summarize the above inequality as follows: for  $\delta \leq c''$ ,

$$p(T,y) \leqslant (1-c'') \sup_{z \in \mathcal{I}(L\delta)} \sup_{t \in [0,T]} p(t,z), \quad y \in \mathcal{I}(\delta).$$

$$(4.33)$$

We now look at what happens when  $T \leq 4\delta^2$  in (4.29). In such a case  $\rho$  in the previous argument must be replaced by  $\rho \wedge T$ . Observing that  $p(T - \rho \wedge T, Y_{\rho \wedge T}) = 0$  on the event  $\{\rho \geq T\} \cup \{Y_{\rho \wedge T} = 1\}$  and following (4.29), we obtain, for  $y \in \mathcal{I}(\delta)$ ,

$$p(T,y) \leqslant \exp(4K\delta^2) \left[ 1 - \mathbb{P}\left( \{ Y_{\rho \wedge T} = 1 \} \cup \{ \rho \ge T \} \right) \right] \sup_{(t,z) \in \mathcal{Q}'(\delta,L)} p(t,z), \quad (4.34)$$

with  $\mathcal{Q}'(\delta, L) = \{(t, z) \in [0, T] \times [1 - L\delta, 1]\}$ . Now, the right-hand side of (4.31) is included in the event  $\{Y_{\rho \wedge T} = 1\} \cup \{\rho \ge T\}$  so that (4.32) yields a lower bound for  $\mathbb{P}(\{Y_{\rho \wedge T} = 1\} \cup \{\rho \ge T\})$ . Therefore, we can repeat the previous arguments in order to prove that (4.33) also holds when  $T \le 4\delta^2$ , which means that (4.33) holds true in both cases.

Therefore, by replacing T by t in the left-hand side in (4.33) and by letting t vary within [0, T], we have in any case,

$$\sup_{y \in \mathcal{I}(\delta)} \sup_{t \in [0,T]} p(t,y) \leq (1-c'') \sup_{z \in \mathcal{I}(L\delta)} \sup_{t \in [0,T]} p(t,z).$$

By induction, for any integer  $n \ge 1$  such that  $L^n \delta \le r_0$ , with  $r_0 = c'' \land (\epsilon/4)$ ,

$$\sup_{y \in \mathcal{I}(\delta)} \sup_{t \in [0,T]} p(t,y) \leqslant (1-c'')^n \sup_{z \in \mathcal{I}(L^n\delta)} \sup_{t \in [0,T]} p(t,z).$$

Given  $\delta \in (0, r_0/L)$ , the maximal value for n is  $n = \lfloor \ln[r_0/\delta]/\ln L \rfloor$ . We deduce that, for any  $\delta \in (0, r_0/L)$ ,

$$\sup_{y \in \mathcal{I}(\delta)} \sup_{t \in [0,T]} p(t,y) \leqslant (1 - c'')^{(\ln[r_0/\delta]/\ln L) - 1} \sup_{z \in \mathcal{I}(\epsilon/4)} \sup_{t \in [0,T]} p(t,z).$$
(4.35)

Following (4.16), we know that

$$\sup_{z \in \mathcal{I}(\epsilon/4)} \sup_{t \in [0,T]} p(t,z) \leq \sup_{z \in \mathcal{I}(\epsilon/4)} \sup_{t \in [0,T]} \left[ \frac{c_T}{\sqrt{t}} \exp\left(-\frac{[z - \vartheta_t^{x_0}]^2}{c_T t}\right) \right],$$
(4.36)

for some constant  $c_T$  only depending upon T and K and where  $(\vartheta_t^{x_0})_{0 \leq t \leq T}$  stands for the solution of the ODE

$$\frac{d\vartheta}{dt} = b(\vartheta_t) + \alpha e'(t), \quad t \in [0,T] ; \quad \vartheta_0 = x_0.$$

Pay attention that we here use the same notation as in (4.17) for the solution of the above ODE whereas e(t) is not given as some  $\mathbb{E}(M_t)$ . Actually, we feel that there is no possible confusion here. Notice also that e is fixed and does not depend upon the initial condition  $x_0$ .

By the comparison principle for ODEs and then by Gronwall's Lemma, we deduce from the fact that e is  $\mathcal{B}$ -Hölder continuous of exponent 1/2 that

$$\vartheta_t^{x_0} \leqslant \vartheta_t^{1-\epsilon} \leqslant \left[1-\epsilon + \Lambda t + \mathcal{B}t^{1/2}\right] \exp(\Lambda t), \quad t \in [0,T].$$

Using the above inequality, we can bound the right-hand side in (4.36): precisely, the above inequality says that the exponential term in the supremum decays exponentially fast as t tends to 0 so that the term inside the supremum can be bounded when t is small; when t is bounded away from 0, the term inside the supremum is bounded by  $1/\sqrt{t}$ . It is plain to deduce that

$$\sup_{z \in \mathcal{I}(\epsilon/4)} \sup_{t \in [0,T]} p(t,z) \leqslant c_T, \tag{4.37}$$

for a new value of  $c_T$ , possibly depending on  $\epsilon$  as well. Therefore, for  $\delta \in (0, r_0/L)$ , (4.35) yields

$$\sup_{y \in \mathcal{I}(\delta)} \sup_{t \in [0,T]} p(t,y) \leqslant \frac{c_T}{(1-c'')} \left(\frac{\delta}{r_0}\right)^{\eta},$$

with  $\eta = -\ln(1-c'')/\ln L$ . This proves (4.25) for  $y \in (1-r_0/L, 1)$ . Using (4.37), we can easily extend the bound to any  $y \in (1-\epsilon/4, 1)$  by modifying if necessary the parameters  $\mu_T$  and  $\eta_T$  therein. This completes the proof.

4.5. **Bound for the gradient.** Here is the final step to complete the proof of Theorem 4.1:

**Proposition 4.8.** For any  $\epsilon \in (0,1)$ , any  $\mathcal{B} > 0$  and any T > 0, there exists a constant  $\mathcal{M}_T > 0$ , only depending upon T,  $\mathcal{B}$ ,  $\epsilon$ , K and  $\Lambda$ , such that, for any initial condition  $x_0 < 1 - \epsilon$ , any continuously differentiable non-decreasing deterministic mapping  $[0,T] \ni t \mapsto e(t)$  satisfying

$$e(0) = 0$$
 ;  $|e(t) - e(s)| \leq \mathcal{B}(t-s)^{1/2}$ ,  $(s,t) \in [0,T]^2$ ,

if  $(\chi_t)_{0 \leq t \leq T}$  denotes the solution of the SDE

$$d\chi_t = b(\chi_t)dt + \alpha e'(t) + dW_t, \quad t \in [0, T] ; \quad \chi_0 = x_0,$$

then, for any integer n such that  $n \ge \lfloor 4/\epsilon \rfloor$ ,

$$|\partial_y p(t,1)| \leq \frac{\mathcal{M}_T n^{-\eta_T}}{1 - \exp[-\mathcal{M}_T^{-1}(1 + \alpha C_T)n^{-1}]} (1 + \alpha C_T), \quad t \in [0,T],$$

where p(t, y) is the density of  $\chi_t$  killed at 1 as in (3.2),  $\eta_T$  is as in Lemma 4.7, and

$$C_T = \sup_{0 \leqslant t \leqslant T} e'(t).$$

*Proof.* We consider the barrier function:

$$q(t,y) = \Theta \exp(Kt) \left[ 1 - \exp(\gamma(y-1)) \right], \quad t \ge 0, \ y \in \mathbb{R},$$

$$(4.38)$$

where  $\gamma$  and  $\Theta$  are free nonnegative parameters. Then, for t > 0 and y < 1,

$$\partial_t q(t,y) + (b(y) + \alpha e'(t)) \partial_y q(t,y) - \frac{1}{2} \partial_{yy}^2 q(t,y)$$
  
=  $\Theta \exp(Kt) \exp(\gamma(y-1)) \left( -(b(y) + \alpha e'(t))\gamma + \frac{1}{2}\gamma^2 \right) + Kq(t,y).$ 

Keeping in mind that

$$\sup_{0 \leqslant t \leqslant T} e'(t) = C_T$$

and choosing

$$\gamma = 2(\max(m, 1) + \alpha C_T), \qquad (4.39)$$

where  $m = \sup_{0 \leqslant z \leqslant 1} |b(z)|$  as before, we obtain, for  $t \in [0, T]$  and  $y \in (0, 1)$ ,

$$-(b(y) + \alpha e'(t))\gamma + \frac{1}{2}\gamma^2 \ge -2(\max(m, 1) + \alpha C_T)^2 + 2(\max(m, 1) + \alpha C_T)^2 = 0.$$

We obtain, for  $t \in [0, T]$  and  $y \in (0, 1)$ ,

$$\partial_t q(t,y) + \left(b(y) + \alpha e'(t)\right) \partial_y q(t,y) - \frac{1}{2} \partial_{yy}^2 q(t,y) \ge Kq(t,y) \ge -b'(y)q(t,y),$$

which reads

$$\partial_t q(t,y) + \partial_y \left[ \left( b(y) + \alpha e'(t) \right) q(t,y) \right] - \frac{1}{2} \partial_{yy}^2 q(t,y) \ge 0.$$
(4.40)

For a given integer  $n \ge \lfloor 4/\epsilon \rfloor$ , we choose  $\Theta$  as the solution of

$$\Theta\left[1 - \exp\left(-\frac{2(\max(m, 1) + \alpha C_T)}{n}\right)\right] = \mu_T n^{-\eta_T}, \qquad (4.41)$$

with  $\mu_T$  and  $\eta_T$  as in the statement of Lemma 4.7. Pay attention that the factor in the left-hand side cannot be 0 as  $\max(m, 1) > 0$ . Notice also q thus depends upon n. By Lemma 4.7, we deduce that

$$q\left(t,1-\frac{1}{n}\right) \ge p\left(t,1-\frac{1}{n}\right), \quad 0 \le t \le T.$$

Now, we can apply the comparison principle for PDEs. Indeed, we also observe that  $q(0,y) \ge p(0,y) = 0$  for  $y \in [1 - 1/n, 1]$  and q(t,1) = p(t,1) = 0 for  $t \in [0,T]$ . Therefore, by (4.40), we have

$$p(t,y) \leq q(t,y), \quad t \in [0,T], \ y \in \left[1 - \frac{1}{n}, 1\right].$$
 (4.42)

Since p(t, 1) = 0 = q(t, 1), we deduce

$$|\partial_y p(t,1)| \leq |\partial_y q(t,1)| = \frac{2\mu_T(\max(m,1) + \alpha C_T)n^{-\eta_T}}{1 - \exp[-2(\max(m,1) + \alpha C_T)/n]} \exp(Kt).$$
(4.43)

We now complete the proof of Theorem 4.1. We make use of Proposition 3.3. Recall (3.6)

$$e'(t) = -\int_0^t \frac{1}{2} \partial_y p^{(0,s)}(t-s,1) e'(s) ds - \frac{1}{2} \partial_y p(t,1), \quad t \in [0,T],$$

where p represents the density of the process X killed at 1 and  $p^{(0,s)}$  represents the density of the process  $X^{\sharp_s}$  driven by  $e^{\sharp_s} = e(\cdot + s) - e(s)$  (see (4.12)) killed at 1 with  $X_0^{\sharp_s} = 0$  as initial condition.

By Proposition 4.5, we know that, for a given  $s \in [0, T)$  and for the prescribed values of  $\alpha$ , the mapping  $[0, T-s] \ni r \mapsto e^{\sharp_s}(r)$  is 1/2-Hölder continuous, the Hölder

constant only depending upon  $\alpha$ ,  $\epsilon$ , K and  $\Lambda$ . Therefore, by Proposition 4.8, we know that

$$|\partial_y p^{(0,s)}(t-s,1)| \leqslant \frac{\mathcal{M}_T n^{-\eta_T}}{1 - \exp[-\mathcal{M}_T^{-1}(1+\alpha C_T)n^{-1}]} (1+\alpha C_T), \quad t \in [s,T], \quad (4.44)$$

for  $n \ge \lceil 4/\epsilon \rceil$  and for some constant  $\mathcal{M}_T$  only depending upon T,  $\alpha$ ,  $\epsilon$ , K and  $\Lambda$ . The same bound also holds true for  $\partial_y p(t, 1)$ .

We deduce that, for any  $t \in [0, T]$  and any n such that  $n \ge \lfloor 4/\epsilon \rfloor$ ,

$$e'(t) \leq \frac{\mathcal{M}_T n^{-\eta_T}}{1 - \exp[-\mathcal{M}_T^{-1}(1 + \alpha C_T)n^{-1}]} (1 + \alpha C_T) \frac{e(T) + 1}{2}.$$

By Lemma 4.3, we have a bound for  $e(T) = \mathbb{E}(M_T)$ , which means that we can bound (e(T)+1)/2 in the right-hand side above by modifying the constant  $\mathcal{M}_T$ . Recalling

$$C_T = \sup_{0 \leqslant t \leqslant T} e'(t),$$

we deduce that

$$C_T \left( 1 - \exp\left[-\mathcal{M}_T^{-1}(1 + \alpha C_T)n^{-1}\right] \right) \leqslant \mathcal{M}_T (1 + \alpha C_T)n^{-\eta_T}.$$

$$(4.45)$$

Choosing n large enough such that the right hand side is less than  $(1 + \alpha C_T)/2$  (so that n depends on T) and multiplying by  $\alpha$ , we get (since  $\alpha \in (0, 1)$ ):

$$\frac{\alpha C_T}{2} \leqslant \frac{1}{2} + \alpha C_T \exp[-\mathcal{M}_T^{-1}(1+\alpha C_T)n^{-1}].$$

This shows that  $\alpha C_T$  must be bounded in terms of  $\mathcal{M}_T$  and n. Precisely, we have

$$\alpha C_T \leq 1 + 2 \sup_{r \geq 0} \left[ x \exp[-\mathcal{M}_T^{-1}(1+r)n^{-1}] \right] := R < +\infty.$$

By (4.45), we deduce that

$$C_T \leqslant \sup_{0 \leqslant r \leqslant R} \left[ \frac{\mathcal{M}_T(1+r)n^{-\eta_T}}{1 - \exp[-\mathcal{M}_T^{-1}(1+r)n^{-1}]} \right],$$

which is independent of  $\alpha$  (for  $\alpha \in (0, \alpha_0]$ ), as required.

#### 5. EXISTENCE AND UNIQUENESS FOR ALL TIME

We now put everything together to arrive at our goal, which is Theorem 5.2 below.

**Lemma 5.1.** For any T > 0, initial condition  $X_0 = x_0 < 1$ , and  $\alpha < \alpha_0$ , where  $\alpha_0$  is as in Theorem 4.1, there exists a constant  $C_{den}(T)$  depending only on T,  $x_0$ , K and  $\Lambda$  such that any solution to (2.1) on [0,T] according to Definition 2.3 satisfies

$$\frac{d}{dy}\mathbb{P}\left(X_t \in dy\right) \leqslant C_{\mathrm{den}}(T)(1-y),$$

for all  $y \in (1 - \epsilon/8, 1)$  and  $t \in [0, T]$ , with  $\epsilon = \min(1, 1 - x_0)$ .

*Proof.* We assume that  $(X_t)_{0 \leq t \leq T}$  is the unique solution to (2.1) with  $X_0 = x_0$  up until time T, and set  $e(t) = \mathbb{E}(M_t)$ . Following the notation of Section 3 (see also the last part of the proof of Theorem 4.1), for  $y \leq 1$  and  $t \leq T$ , let

$$p(t,y) := \frac{d}{dy} \mathbb{P} \left( X_t \in dy, t \leqslant \tau_1 \right),$$
$$p^{(0,s)}(t,y) := \frac{d}{dy} \mathbb{P} \left( X_t^{\sharp_s} \in dy, t \leqslant \tau_1^{\sharp_s} | X_0^{\sharp_s} = 0 \right).$$

By Theorem 4.1, we know that e is  $\mathcal{M}_T$ -Lipschitz continuous, so that by (4.42),

$$p(t,y) \leqslant q(t,y), \quad t \in [0,T], \ y \in \left[1 - \frac{1}{n}, 1\right],$$

where n stands for  $\lceil 4/\epsilon \rceil$  and q is given by (4.38), with  $\gamma$  and  $\Theta$  being fixed by (4.39) and (4.41), with  $C_T = \mathcal{M}_T$ . By the specific form of q, this says that there exists a constant  $C'_T$ , depending only on T,  $x_0$ , K and  $\Lambda$ , such that

$$p(t,y) \leq C'_T(1-y), \quad t \in [0,T], \ y \in \left[1 - \frac{\epsilon}{8}, 1\right]$$

using the elementary inequality  $1 - e^{-x} \leq x$  for  $x \in \mathbb{R}$ . Clearly, the same argument applies to  $p^{(0,s)}(t-s,y)$ , i.e.

$$p^{(0,s)}(t-s,y) \leq C'_T(1-y), \quad 0 \leq s < t \leq T, \ y \in \left[1 - \frac{\epsilon}{8}, 1\right].$$

Now, following the proof of (4.11), we get for  $t \in [0, T]$  and  $y \in [1 - \epsilon/8, 1]$ ,

$$\frac{d}{dy}\mathbb{P}(X_t \in dy) = p(t,y) + \int_0^t p^{(0,s)}(t-s,y)e'(s)ds$$
$$\leqslant C'_T(1+e(T))(1-y),$$

where we use Lemma 3.2 for justifying the passage to the density in (4.11). By Lemma 4.3, this completes the proof.

**Theorem 5.2.** For any T > 0, initial condition  $X_0 = x_0 < 1$ , and  $\alpha < \alpha_0$ , where  $\alpha_0$  as in Theorem 4.1, there exists a unique solution to the nonlinear equation (2.1) on [0,T] according to Definition 2.3.

*Proof.* We would like a solution up until fixed time T > 0. The idea is to iterate Theorem 3.1, which is possible thanks to Lemma 5.1. Indeed, by Theorem 3.1, we have that there exists a solution to (2.1) with  $X_0 = x_0$  up until some small time  $T_1 > 0$ . By Lemma 5.1, we therefore have that

$$\mathbb{P}(X_{T_1} \in dy) \leqslant C_{\text{den}}(T_1)(1-y), \quad y \in \left[1 - \frac{\epsilon}{8}, 1\right],$$

where  $\epsilon = \min(1 - x_0, 1)$ . If  $T_1 \ge T$  we are done. If not, since we have the above density bound and since we have an explicit bound for  $\mathbb{E}|X_{T_1}|$  deriving from Lemma 4.3 and from Gronwall's Lemma, we can apply Theorem 3.1 again to see that there exists a solution to (2.1) on some interval  $[T_1, T_1 + T_2]$  starting from  $X_{T_1}$ . As  $T_2$  depends upon  $X_{T_1}$  through  $\epsilon$ ,  $C_{den}(T_1)$  and  $\mathbb{E}|X_{T_1}|$  only and as these quantities can be bounded in terms of T,  $\epsilon$ , K,  $\Lambda$  only, we then see that

$$T_2 \geqslant \phi(T)$$

for some constant  $\phi(T)$  that refers to T,  $\epsilon$ , K,  $\Lambda$  only. Now we know that there exists a solution to (2.1) with  $X_0 = x_0$  on  $[0, T_1 + T_2]$ . If  $T_1 + T_2 > T$  we are done. If not, by Lemma 5.1 once again,

$$\mathbb{P}(X_{T_1+T_2} \in dy) \leqslant C_{\mathrm{den}}(T_1+T_2)(1-y), \quad y \in \left[1-\frac{\epsilon}{8}, 1\right],$$

and we can then repeat the argument n times to get a solution up until time  $T_1 + \cdots + T_n$ , where all  $T_k \ge \phi(T)$  for  $k \ge 2$  i.e. each time step is of size at least  $\phi(T)$ . It is then clear that there exists  $n \ge 1$  such that  $T_1 + \cdots + T_n \ge T$ , and so we are done for the existence of a solution.

Uniqueness of the solution proceeds in the same way. Given another solution  $(X'_t, M'_t)_{t \in [0,T]}$  on the interval [0, T] in the sense of Definition 2.3, it must satisfy the *a priori* estimates in the statements of Theorem 4.1 and Lemmas 4.3 and 5.1. In particular, dividing the interval [0, T] into subintervals of length  $\phi(T)$  (except for the last interval the length of which might be less than  $\phi(T)$ ), with the same  $\phi(T)$  as above, we can apply the contraction property in Theorem 3.1 on each subinterval iteratively. Precisely, choosing  $A_1$  accordingly in Theorem 3.1, we prove by induction that the two solutions coincide on  $[0, \phi(T)]$ ,  $[0, 2\phi(T)]$ , and so on.

**Remark 5.3** (A closed formula for  $e(t) = \mathbb{E}(M_t)$ ). This section is motivated by the proof of finite time blow-up in [3] (Theorem 2.2). In fact, thanks to the knowledge about the existence of a solution for  $\alpha$  sufficiently small provided by Theorem 5.2, we can extract some interesting formulae by modifying the methods in [3].

Indeed, fix  $x_0 < 1$  and suppose that we are in the Ornstein-Uhlenbeck case, so that  $b(x) = -\lambda x$  with  $\lambda \ge 0$ . Let  $\alpha_0$  be as in Theorem 5.2. Then for  $\alpha < \alpha_0$  a solution to (2.1) with  $X_0 = x_0 < 1$  exists for all time in the sense of Definition 2.3. Set  $e(t) = \mathbb{E}(M_t)$ .

For  $\mu > 0$ , let  $\varphi(x) = \exp(\mu x)$ . Then since e(t) is continuously differentiable, by the generalized Itô formula we see that

$$\mathbb{E}(\varphi(X_t)) = \mathbb{E}(\varphi(X_0)) + \int_0^t (\alpha \mu e'(s) \mathbb{E}(\varphi(X_s)) - \lambda \mu \mathbb{E}(X_s \varphi(X_s))) \, ds \\ + \frac{\mu^2}{2} \int_0^t \mathbb{E}(\varphi(X_s)) \, ds + (\varphi(0) - \varphi(1)) \mathbb{E}(M_t).$$

Differentiating with respect to t and writing  $w(t) = \mathbb{E}(\varphi(X_t))$  yields

$$\frac{d}{dt}w(t) = \alpha \mu e'(t)w(t) - \lambda \mu \mathbb{E}(X_t\varphi(X_t)) + \frac{\mu^2}{2}w(t) + (\varphi(0) - \varphi(1))e'(t)$$

for all t > 0, with initial condition  $w(0) = \exp(\mu x_0)$ . Setting

$$\upsilon := \frac{\varphi(1) - \varphi(0)}{\alpha \mu} = \frac{\exp(\mu) - 1}{\alpha \mu},$$

and using the fact that  $X_t < 1$  we then see that

$$\frac{d}{dt}w(t) \geqslant \left[\alpha\mu e'(t) - \lambda\mu + \frac{\mu^2}{2}\right]w(t) - \nu\alpha\mu e'(t).$$

In other words w(t) satisfies the differential inequality

$$\begin{cases} \frac{d}{dt}w(t) \geqslant \eta(t)w(t) + \nu(t), \quad t > 0\\ w(0) = \exp(\mu x_0), \end{cases}$$

with  $\eta(t) = \alpha \mu e'(t) - \lambda \mu + \mu^2/2$  and  $\nu(t) = -\nu \alpha \mu e'(t)$ . The associated differential equation is

$$\begin{cases} \frac{d}{dt}z(t) &= \eta(t)z(t) + \nu(t), \quad t > 0\\ z(0) &= \exp(\mu x_0), \end{cases}$$

which can be solved by the standard method to give

$$z(t) = \exp\left(\int_0^t \eta(s)ds\right) \left[z(0) + \int_0^t \nu(s)\exp\left(-\int_0^s \eta(u)du\right)ds\right].$$

It follows that  $w(t) \ge z(t)$  for all  $t \ge 0$ .

We now claim that  $z(t) \ge 0$  for all  $t \ge 0$ . Consider the process  $Y_t := X_0 + \alpha e(t) - \lambda t + W_t - M_t$  where  $(X_t, M_t)_{t \ge 0}$  is the solution to (2.1). Then by Itô's formula once more we arrive at

$$\frac{d}{dt}\mathbb{E}\left(\varphi(Y_t)\right) = \left[\mu\alpha e'(t) - \mu\lambda + \frac{\mu^2}{2}\right]\mathbb{E}\left(\varphi(Y_t)\right) - \nu\alpha\mu e'(t),$$

and  $\mathbb{E}(\varphi(Y_0)) = \mathbb{E}(\varphi(X_0))$ . Therefore (by uniqueness of solutions to the first-order linear ODE) we have that  $z(t) = \mathbb{E}(\varphi(Y_t)) = \mathbb{E}[\exp(\mu Y_t)] \ge 0$ .

Hence we have that  $w(t) \ge z(t) \ge 0$ . Setting  $\gamma(t) := \exp(\alpha \mu e(t) + (\mu^2/2 - \lambda \mu)t)$ , we see that

$$\begin{aligned} z(t) &= \exp\left(\int_0^t \eta(s)ds\right) \left[z(0) + \int_0^t \nu(s) \exp\left(-\int_0^s \eta(u)du\right)ds\right] \\ &= \gamma(t) \left[z(0) - \upsilon \int_0^t \alpha \mu e'(s) \exp\left(-\alpha \mu e(s) - \left(\frac{\mu^2}{2} - \lambda \mu\right)s\right)ds\right] \\ &= \gamma(t) \left[z(0) + \upsilon \int_0^t \frac{d}{ds} \left[\exp(-\alpha \mu e(s))\right] \exp\left(-\left(\frac{\mu^2}{2} - \lambda \mu\right)s\right)ds\right] \\ &= \gamma(t) \left[z(0) + \upsilon \left(\exp\left(-\alpha \mu e(t) - \left(\frac{\mu^2}{2} - \lambda \mu\right)t\right) - 1\right)\right] \\ &+ \gamma(t) \left(\frac{\mu^2}{2} - \lambda \mu\right)\upsilon \int_0^t \exp\left(-\alpha \mu e(s) - \left(\frac{\mu^2}{2} - \lambda \mu\right)s\right)ds \end{aligned}$$

by integration by parts. Simplifying

$$w(t) \ge z(t) = \gamma(t) \left[ z(0) + \upsilon \left( \gamma(t)^{-1} - 1 \right) \right] + \gamma(t) \left( \frac{\mu^2}{2} - \lambda \mu \right) \upsilon \int_0^t \gamma(s)^{-1} ds$$
$$= \upsilon + \gamma(t) \left[ z(0) - \upsilon + \left( \frac{\mu^2}{2} - \lambda \mu \right) \upsilon \int_0^t \gamma(s)^{-1} ds \right] \ge 0.$$
(5.1)

It is worth noting here that in the Brownian case  $(\lambda = 0)$ , we actually have that w(t) = z(t), so that we have an explicit formula for the exponential moments  $\mathbb{E}(e^{\mu X_t})$ .

Coming back to the general Ornstein-Uhlenbeck case, note that (as is done in [3]) that

$$w(t) = \mathbb{E}\left[\exp(\mu X_t)\right] \leqslant e^{\mu},$$

for all t > 0, i.e. w(t) is bounded. Moreover, for  $\mu \ge 2\lambda$  it is clear that  $\gamma(t) \to \infty$ as  $t \to \infty$ . Thus by (5.1) we must have that

$$w(0) - \upsilon + \left(\frac{\mu^2}{2} - \lambda\mu\right)\upsilon \int_0^t \gamma(s)^{-1} ds \to 0$$

as  $t \to \infty$ . Indeed, if the limit was positive then  $w(t) \to \infty$  as  $t \to \infty$ , which is a contradiction since we have just noted that it is bounded. On the other hand if the limit was negative, then  $z(t) \to -\infty$  as  $t \to \infty$ , which is also a contradiction since  $z(t) \ge 0$  for all t.

We have thus arrived at the following surprising closed formula, which characterizes e:

$$\left(\frac{\mu^2}{2} - \lambda\mu\right) \int_0^\infty \exp\left(-\alpha\mu e(s) - \left(\frac{\mu^2}{2} - \lambda\mu\right)s\right) ds = 1 - \frac{\alpha\mu \exp(\mu x_0)}{\exp(\mu) - 1}$$
(5.2)

for all  $\mu > 2\lambda$  and  $\alpha < \alpha_0$ .

#### 6. Explicit Bounds for the Criticallity

Theorem 5.2 says that, for a given initial condition  $x_0$ , existence and uniqueness of a solution to (2.1) holds when the self-excitation rate  $\alpha$  lies in  $(0, \alpha_0(x_0))$ , for some  $\alpha_0(x_0) \in (0, 1]$ , the value of which is given in the statements of both Theorem 4.1 and Proposition 4.5. This raises several questions:

- (1) Does the model indeed exhibit several regimes for global solvability according to the values of  $\alpha$ ?
- (2) If, so, are there two phases exactly (using an analogy with statistical mechanics)? To put it differently, does there exist a critical threshold  $\alpha_c$  such that global solvability holds for  $\alpha < \alpha_c$  and fails for  $\alpha > \alpha_c$ ? Or are there any more complex cases when global solvability fails for some  $\alpha^+$  and then holds for some  $\alpha^- > \alpha^+$ ?

(3) Conversely, are there any cases when global solvability holds for any  $\alpha < 1$ ? Although Theorem 5.2 does not provide any answer to these questions, it says that, for a given initial condition  $x_0 < 1$ , the critical rate

$$\alpha_c^-(x_0) = \sup \left\{ \alpha^- \in (0,1] : \mathcal{G}(\alpha, x_0) \text{ holds for any } \alpha \in (0,\alpha^-) \right\}$$

makes senses as it is (strictly) positive, since it is larger than  $\alpha_0(x_0)$ , where  $\mathcal{G}(\alpha, x_0)$  stands for the property of global unique solvability of Eq. (2.1) when driven by  $\alpha$  and initialized with  $x_0$ .

In this framework, the lower bound  $\alpha_c^-(x_0) \ge \alpha_0(x_0)$ , with  $\alpha_0(x_0)$  given by Proposition 4.5 and Theorem 4.1, is complementary to the previous result by Caceres, Carrillo and Perthame [3]: in [3], it is indeed proven that existence of a solution fails in the long run for some values of  $\alpha < 1$  and some initial conditions, including Dirac masses located right below the barrier 1, in the case when  $b(x) + \lambda x \ge 0$ , with  $\lambda > 0$ . This shows that, in some cases,  $\alpha_c^-(x_0)$  is indeed strictly less than 1. Actually, the result in [3] says more: for any value of  $\lambda > 0$ , there exists a critical threshold  $x_c < 1$  such that, for any  $x_0 \in (x_c, 1)$ , there exists some  $\alpha'_0(x_0) < 1$  such that global existence fails (in the sense that e' blows up in finite time) for any  $\alpha \in (\alpha'_0(x_0), 1)$ . In other words, the critical rate

$$\alpha_c^+(x_0) = \inf \left\{ \alpha^+ \in (0,1] : \mathcal{G}(\alpha, x_0) \text{ fails for any } \alpha \in (\alpha^+, 1) \right\}$$

is well-defined and strictly less than 1 for  $x_0$  close to 1. Obviously,

$$\alpha_c^-(x_0) \leqslant \alpha_c^+(x_0),$$

but, as written above, the following questions remain open:

- (1) Are there any cases when  $\alpha_c^-(x_0) \neq \alpha_c^+(x_0)$ ?
- (2) Are there any cases when  $\alpha_c^-(x_0) = 1$ ?

More generally, finding the optimal values of  $\alpha_c^-(x_0)$  or/and  $\alpha_c^+(x_0)$  turns out to be a really challenging problem.

In this section, we address two problems: (i) the first is to provide an explicit lower bound for  $\alpha_c^-(0)$  in the case when  $b(x) = -\lambda x$ ,  $x \in \mathbb{R}$ , for some  $\lambda > 0$ ; (ii) the second is to provide a lower bound for  $\alpha_c^-(x_0)$  and an upper bound for  $\alpha_c^+(x_0)$ in the case when b = 0 but  $x_0$  is chosen in such a way that  $\alpha_c^+(x_0)$  is known to be less than 1 and then to discuss the bounds in light of some numerical experiments.

6.1. **Ornstein-Uhlenbeck Case.** In this paragraph, we investigate further the case when  $b(x) = -\lambda x$  (referred to as the Ornstein-Uhlenbeck case, since the dynamics of X up until the first jump time coincide with those of the Ornstein-Uhlenbeck process), which is choice most often made in neuroscience. As above, we assume that the diffusion coefficient is 1. We then exhibit a lower bound for the optimal value of  $\alpha_c^-$  according to the methodology described in the previous subsection. As the method involves rather tedious computations, we will focus on the case  $X_0 = 0$  only: this makes things a bit simpler.

We thus consider

$$X_t = -\lambda \int_0^t X_s ds + \alpha e(t) + W_t - M_t, \quad t \ge 0, \tag{6.1}$$

where  $e(t) = \mathbb{E}(M_t)$ . When  $X_0 = 0$ , as is the case here, numerical experiments indicate that existence and uniqueness of a global solution may hold for any  $\alpha \in (0, 1)$ . Equation (6.1) is then solved by using a particle method: we refer to the forthcoming paper [5] for a complete description and analysis of the numerical scheme. For instance, when  $\lambda = 0.01$ , we get as a plot of the mapping  $t \mapsto e(t)$  associated to the approximate solution that given in Fig 1.



FIGURE 1. Plot of  $t \mapsto e(t)$  for  $X_0 = 0$ ,  $\lambda = 0.01$  and  $t \in [0, 10]$ .

Numerically speaking, existence and uniqueness is then suspected to hold since the graph of the mapping  $t \mapsto e(t)$  is smooth and, in particular, has no jump. We refer to Subsection 6.2 for a counter-example when a jump occurs for some  $x_0 \neq 0$ , indicating that existence of a solution, as defined above, cannot hold. Below, we are not able to prove that  $\alpha_c^-(0) = 1$ , showing that our methodology is not optimal. Anyhow, we manage to give an explicit lower bound for the critical value of  $\alpha_c^-(0)$ and to prove that it converges toward 1 when  $\lambda$  tends to  $+\infty$ .

The strategy is to seek  $\alpha_0$  such that, for any  $t \ge 0$  and y in a neighbourhood of 1,  $\mathbb{P}(X_t \in dy) < 1/\alpha$  for all  $\alpha < \alpha_0$  so that the condition (4.5) in Lemma 4.4 is satisfied.

First Step. The first step is to provide some a priori estimates for the solution on any interval [0, T] where existence holds. Precisely, the point is to refine the *a* priori  $L^{\infty}$  bound for the function  $(e(t) = \mathbb{E}(M_t))_{0 \leq t \leq T}$  given in the statement of Lemma 4.3, by taking advantage of the mean-reverting property of the Ornstein-Uhlenbeck process. (As in Section 4, in all the *a priori* estimates below, we assume that  $(X_t, M_t)_{t \in [0,T]}$  stands for a solution to (6.1) in the sense of Definition 2.3.)

Here is the first claim:

**Lemma 6.1.** For any  $t \in [0, T]$ ,

$$\max\left(\mathbb{E}\left[\left[(X_t)_{-}\right]^2\right]^{1/2}, \ \mathbb{E}\left[(X_t)_{+}\right], \ (1-\alpha)\exp(-\lambda t)\int_0^t \exp(\lambda s)e'(s)ds\right) \leqslant f_1(\lambda, t),$$
  
with  $f_1(\lambda, t) = \left[\frac{1-\exp(-2\lambda t)}{2\lambda}\right]^{1/2},$ 

where  $(x)_{-} = \max(-x, 0)$  for  $x \in \mathbb{R}$ .

*Proof.* We first investigate how far the process  $(X_t)_{0 \leq t \leq T}$  can go in the negative direction. The point is to apply Itô's formula to  $([(X_t)_{-}]^2)_{0 \leq t \leq T}$ :

$$\begin{split} &[(X_t)_{-}]^2 \\ &= 2\lambda \int_0^t (X_{s-})_{-} X_s ds - 2\alpha \int_0^t (X_{s-})_{-} e'(s) ds + \int_0^t \mathbf{1}_{\{X_{s-} < 0\}} ds - 2 \int_0^t (X_{s-})_{-} dW_s \\ &= -2\lambda \int_0^t (X_s)_{-}^2 ds + 2\alpha \int_0^t X_s \mathbf{1}_{\{X_s < 0\}} e'(s) ds + \int_0^t \mathbf{1}_{\{X_s < 0\}} ds - 2 \int_0^t (X_{s-})_{-} dW_s, \end{split}$$

for  $t \in [0, T]$ . Here,  $(X_{s-})_{-}$  stands for the negative part of the left-limit of X at time s. To pass from the second to the third line, notice that  $X_{s-} = X_s$  when  $X_{s-} \leq 0$ . Multiplying by  $\exp(2\lambda t)$ , reapplying Itô's formula and then taking the expectation, we deduce:

$$\mathbb{E}\left[\exp(2\lambda t)(X_t)_{-}^2\right] \leqslant \int_0^t \exp(2\lambda s) ds = \frac{\exp(2\lambda t) - 1}{2\lambda} = \exp(2\lambda t) f_1^2(\lambda, t).$$
(6.2)

Now, multiplying in the same way (6.1) by  $\exp(\lambda t)$  and applying Itô's formula, we get:

$$\exp(\lambda t)X_t = \alpha \int_0^t \exp(\lambda s)e'(s)ds + \int_0^t \exp(\lambda s)dW_s - \int_0^t \exp(\lambda s)dM_s.$$

Taking the expectation, we deduce:

$$(1-\alpha)\int_0^t \exp(\lambda s)e'(s)ds = -\mathbb{E}\big[\exp(\lambda t)X_t\big],$$

so that, writing  $X_t = (X_t)_+ - (X_t)_-$ , with  $(x)_+ = \max(x, 0)$ ,

$$\mathbb{E}\left[\exp(\lambda t)(X_t)_+\right] + (1-\alpha) \int_0^t \exp(\lambda s) e'(s) ds = \mathbb{E}\left[\exp(\lambda t)(X_t)_-\right] \\ \leqslant \exp(\lambda t) f_1(\lambda, t),$$
(6.3)

where the last inequality follows from (6.2) and Hölder's inequality.

We deduce:

**Lemma 6.2.** For any  $t, t' \in [0, T], t \leq t'$ 

$$e(t') - e(t) \leq \frac{f_2(\lambda, t' - t, T)}{1 - \alpha}$$

with

$$f_2(\lambda, r, T) = \min\left[\exp(1)f_1(\lambda, T)(1 + \lambda r), 1 + (\lambda/2)^{1/2}r + (2r/\pi)^{1/2}\right].$$

*Proof.* We prove first that the bound holds with respect to the first argument in the minimum defining  $f_2$ . By Lemma 6.1, we know that, for  $0 \le t \le t + h \le T$ ,

$$[e(t+h) - e(t)] \exp(-\lambda h) \leqslant \exp[-\lambda(t+h)] \int_0^{t+h} \exp(\lambda s) e'(s) ds \leqslant \frac{f_1(\lambda, T)}{1-\alpha}.$$

Choosing  $h = 1/\lambda$ , we deduce that for  $0 \leq t \leq s \leq T \land (t + 1/\lambda)$ 

$$e(s) - e(t) \leq \exp(1) \frac{f_1(\lambda, T)}{1 - \alpha}$$

Summing over a subdivision of [t, t'] made of small intervals of lengths less than  $1/\lambda$ , we complete the proof for the first argument in the minimum.

We now turn to the second argument in the minimum. As in the proof of Lemma 4.4, we can assume without loss of generality that t = 0 by shifting the system (the shift being of length t), provided the new initial condition  $X_0$  is chosen as an arbitrary random variable. Precisely, the new initial condition  $X_0$  is then understood as the previous  $X_t$ . Similarly, the new variables  $(X_s)_{0 \le s \le t'-t}$  are understood as the previous  $(X_s)_{t \le s \le t'}$  so that, by Lemma 6.1, we can assume the new  $(X_s)_{0 \le s \le t'-t}$  satisfy

$$\mathbb{E}\left[(X_s)_+\right] \leqslant f_1(\lambda, T) \leqslant \left[\frac{1}{2\lambda}\right]^{1/2}$$

Following the proof of (2.10)–(2.11), we then get for any  $s \in [0, t' - t]$ :

$$e(s) = \mathbb{E}(M_s) \leqslant \mathbb{E}\left[\sup_{0 \leqslant r \leqslant s} (Z_r)_+\right] \leqslant 1 + \lambda \int_0^s \mathbb{E}\left[(X_s)_+\right] ds + \alpha e(s) + \mathbb{E}\left[\sup_{0 \leqslant r \leqslant s} W_r\right]$$
$$\leqslant 1 + \left[\frac{\lambda}{2}\right]^{1/2} (t'-t) + \alpha e(s) + \mathbb{E}\left[\sup_{0 \leqslant r \leqslant s} W_r\right],$$

so that, for any  $s \in [0, t' - t]$ ,

$$e(s) \leqslant \frac{1 + (\lambda/2)^{1/2}(t'-t) + \sqrt{2(t'-t)/\pi}}{1-\alpha}$$

the last term in the right-hand side following from

$$\mathbb{E}\left(\sup_{0 \leqslant s \leqslant t} W_s\right) = \sqrt{\frac{2t}{\pi}}.$$
(6.4)

Indeed, by the reflection principle,

$$\mathbb{P}\left(\sup_{s \leqslant t} W_s \leqslant a\right) = 2\mathbb{P}\left(W_t \leqslant a\right) - 1 = \sqrt{\frac{2}{t\pi}} \int_0^a e^{-z^2/2t} dz$$

so that

$$\mathbb{E}\left(\sup_{s \leqslant t} W_s\right) = \sqrt{\frac{2}{t\pi}} \int_0^\infty z e^{-z^2/2t} dz = \sqrt{\frac{2t}{\pi}}.$$

Second Step. According to the program we fixed in the proof of Lemma 4.6, we now investigate the bound of the density in small time. In the Ornstein-Uhlenbeck

case, (4.16) (with  $x_0 = 0$ ) has the form:

$$\frac{d}{dx} \mathbb{P}\left(\hat{Z}_t^0 \in dx\right) = \frac{1}{(2\pi)^{1/2} f_1(\lambda, t)} \times \exp\left[-\frac{1}{2f_1^2(\lambda, t)} \left(x - \alpha \exp(-\lambda t) \int_0^t \exp(\lambda r) e'(r) dr\right)^2\right],$$
(6.5)

for  $t \leq T$ , with  $f_1(\lambda, t)$  as in the statement of Lemma 6.1 and  $\hat{Z}_t^0$  given by (4.14). More generally, (4.18) reads:

$$\frac{d}{dx} \mathbb{P}\left(\hat{Z}_{t-s}^{\sharp_{s},0} \in dx\right) = \frac{1}{(2\pi)^{1/2} f_{1}(\lambda, t-s)}$$

$$\times \exp\left[-\frac{1}{2f_{1}^{2}(\lambda, t-s)} \left(x - \alpha \exp\left[-\lambda(t-s)\right] \int_{0}^{t-s} \exp(\lambda r) e'(s+r) dr\right)^{2}\right],$$

for  $s < t \leq T$ . By Lemma 6.1, we notice that the "mean-term" in the exponential is bounded by

$$\alpha \exp\left[-\lambda(t-s)\right] \int_{0}^{t-s} \exp(\lambda r) e'(s+r) dr = \alpha \exp(-\lambda t) \int_{s}^{t} \exp(\lambda r) e'(r) dr$$

$$\leqslant \frac{\alpha f_{1}(\lambda, t)}{1-\alpha}.$$
(6.6)

By Lemma 6.2, it is also bounded by

$$\alpha \exp\left[-\lambda(t-s)\right] \int_0^{t-s} \exp(\lambda r) e'(s+r) dr \leqslant \alpha(e(t)-e(s)) \leqslant \frac{\alpha f_2(\lambda,t-s,t)}{1-\alpha}.$$
 (6.7)

Therefore, since  $r \mapsto f_2(\lambda, r, t)$  is non-decreasing, if, for some  $\varepsilon \in (0, 1)$ ,

$$\frac{\alpha}{1-\alpha}\min[f_1(\lambda,t),f_2(\lambda,t,t)] \leqslant 1-\varepsilon,$$

then, for  $x \in (1 - \varepsilon \delta, 1)$ , with  $\delta \in (0, 1)$ , we have for any  $s \in [0, t)$  (pay attention that this covers (6.5) as well):

$$\begin{split} \frac{d}{dx} \mathbb{P}\left(\hat{Z}_{t-s}^{\sharp_{s},0} \in dx\right) &\leqslant \frac{1}{(2\pi)^{1/2} f_{1}(\lambda,t-s)} \exp\left[-\frac{(1-\delta)^{2} \varepsilon^{2}}{2 f_{1}^{2}(\lambda,t-s)}\right] \\ &\leqslant \frac{1}{(1-\delta)\varepsilon} g\left(\frac{(1-\delta)\varepsilon}{f_{1}(\lambda,t)}\right), \end{split}$$

where

$$g(u) = \sup_{v>u} \left[ \frac{v}{(2\pi)^{1/2}} \exp\left(-\frac{v^2}{2}\right) \right] = \begin{cases} \frac{u}{(2\pi)^{1/2}} \exp\left(-\frac{u^2}{2}\right) & \text{if } u \ge 1, \\ \frac{1}{(2\pi)^{1/2}} \exp\left(-\frac{1}{2}\right) & \text{if } u < 1, \end{cases}$$

using the fact that  $u \mapsto g(u)$  is non-increasing and  $r \mapsto f_1(\lambda, r)$  is non-decreasing. Therefore, by (4.15), for  $x \in (1 - \varepsilon \delta, 1)$ 

$$\frac{d}{dx}\mathbb{P}\left(X_t \in dx\right) \leqslant \frac{1+e(t)}{(1-\delta)\varepsilon}g\left(\frac{(1-\delta)\varepsilon}{f_1(\lambda,t)}\right).$$

Define  $T_0 = 1 \wedge \lambda^{-1}$ . Then, by Lemma 6.2, for  $t \leq T \wedge T_0$ ,

$$\frac{d}{dx}\mathbb{P}\left(X_t \in dx\right) \leqslant \left(1 + \frac{f_2(\lambda, T_0, T_0)}{1 - \alpha}\right) \frac{1}{(1 - \delta)\varepsilon} g\left(\frac{(1 - \delta)\varepsilon}{f_1(\lambda, T_0)}\right)$$
(6.8)

again since g is non-increasing, provided

$$\frac{\alpha}{1-\alpha}\min[f_1(\lambda, T_0), f_2(\lambda, T_0, T_0)] \leqslant 1-\varepsilon$$
(6.9)

holds true.

Third Step. We now investigate the bound of the density for  $t \ge T_0$ . We first discuss  $\pi_1$  defined by (4.22). By (4.24) (plugging inside the current value of  $T_0$ ), and the same arguments as in the second step,

$$\pi_1 \leqslant \frac{1}{(1-\delta)\varepsilon} g\left(\frac{(1-\delta)\varepsilon}{f_1(\lambda,T_0)}\right) (e(t) - e(t-T_0)),$$

provided

$$\frac{\alpha}{1-\alpha} \min \left[ f_1(\lambda, \infty), f_2(\lambda, T_0, \infty) \right] \leqslant 1 - \varepsilon.$$
(6.10)

Pay attention that, in  $f_1(\lambda, \cdot)$  and  $f_2(\lambda, T_0, \cdot)$ ,  $\cdot$  is understood as  $T_0$  in (6.9) whereas it is understood as  $\infty$  in (6.10). The difference between both comes from the fact that, above, t may be large. By Lemma 6.2,

$$\pi_1 \leqslant \frac{1}{(1-\delta)\varepsilon} g\left(\frac{(1-\delta)\varepsilon}{f_1(\lambda,T_0)}\right) \frac{f_2(\lambda,T_0,\infty)}{1-\alpha}.$$
(6.11)

It now remains to investigate  $\pi_2$  also defined in (4.22). We start with the case  $\lambda \ge 1$ , so that  $T_0 = \lambda^{-1}$ . Following (4.23) and (6.5) (keeping in mind that  $T_0\lambda = 1$ ), we have

$$\pi_2 \leqslant \sup_{z \leqslant 1} \pi_2(z), \tag{6.12}$$

with

$$\pi_2(z) = \frac{1}{(2\pi)^{1/2} f_1(\lambda, 1/\lambda)} \times \exp\left[-\frac{1}{2f_1^2(\lambda, 1/\lambda)} \left(x - \exp(-1)\left[z + \alpha \int_0^{1/\lambda} \exp(\lambda r)e'(t - T_0 + r)dr\right]\right)^2\right].$$

Actually, the bound can be refined for  $\lambda$  large by taking into account the fact that z is understood as a realization of  $X_{t-T_0}$  and that the probability that  $X_{t-T_0}$  is away

from 0 is very small when  $\lambda$  is large because of the mean-reverting property of the Ornstein-Uhlenbeck process. Precisely, we claim:

$$\pi_{2} \leqslant \max \Big[ \sup_{z \leqslant 1/2} \pi_{2}(z), \sup_{z \leqslant 1} \pi_{2}(z) \mathbb{P} \{ X_{t-T_{0}} \ge 1/2 \} \Big] \\ \leqslant \max \Big[ \sup_{z \leqslant 1/2} \pi_{2}(z), (2/\lambda)^{1/2} \sup_{z \leqslant 1} \pi_{2}(z) \Big],$$
(6.13)

the last line following from Lemma 6.1 and Markov's inequality. As above, we know that the mean term in the exponential in  $\pi_2(z)$  is bounded by

$$\exp(-1)z + \alpha \exp(-1) \int_0^{1/\lambda} \exp(\lambda r) e'(t - T_0 + r) dr$$
  
$$= \exp(-1)z + \alpha \exp(-\lambda T_0) \int_0^{T_0} \exp(\lambda r) e'(t - T_0 + r) dr$$
  
$$= \exp(-1)z + \alpha \exp(-\lambda t) \int_{t-T_0}^t \exp(\lambda r) e'(r) dr$$
  
$$\leqslant \exp(-1)z + \frac{\alpha}{1-\alpha} \min[f_1(\lambda, \infty), f_2(\lambda, T_0, \infty)].$$

Therefore, if (6.10) holds for  $\varepsilon \in (\exp(-1), 1)$ , then, for  $x \in (1 - (\varepsilon - \exp(-1))\delta, 1)$ , we get from (6.12) and (6.13):

$$\pi_2 \leqslant \min \left[ f_4(\delta, \lambda), f_5(\delta, \lambda) \right],$$

with

$$f_4(\delta,\lambda) = \frac{1}{(2\pi)^{1/2} f_1(\lambda, 1/\lambda)} \exp\left[-\frac{(1-\delta)^2 (\varepsilon - \exp(-1))^2}{2f_1^2(\lambda, 1/\lambda)}\right],$$
  
$$f_5(\delta,\lambda) = \max\left\{\frac{1}{(2\pi)^{1/2} f_1(\lambda, 1/\lambda)} \exp\left[-\frac{(1-\delta)^2 (\varepsilon - \exp(-1)/2)^2}{2f_1^2(\lambda, 1/\lambda)}\right], \frac{2^{1/2} f_4(\delta,\lambda)}{\lambda^{1/2}}\right\}.$$

Finally, we have

$$\frac{d}{dx} \mathbb{P}\left\{X_t \in dx\right\} \leqslant \frac{1}{(1-\delta)\varepsilon} g\left(\frac{(1-\delta)\varepsilon}{f_1(\lambda,T_0)}\right) \frac{f_2(\lambda,T_0,\infty)}{1-\alpha} + \min\left[f_4(\delta,\lambda), f_5(\delta,\lambda)\right].$$
(6.14)

provided (6.10) holds true, since  $[d/dx]\mathbb{P}\{X_t \in dx\} \leq \pi_1 + \pi_2$  by (4.23).

We now turn to the case  $\lambda < 1$ . Then,  $T_0 = 1$  and

$$\pi_2 \leqslant \frac{1}{(2\pi)^{1/2} f_1(\lambda, 1)}.$$

Therefore,

$$\frac{d}{dx}\mathbb{P}\left\{X_t \in dx\right\} \leqslant \frac{1}{(1-\delta)\varepsilon}g\left(\frac{(1-\delta)\varepsilon}{f_1(\lambda,T_0)}\right)\frac{f_2(\lambda,T_0,\infty)}{1-\alpha} + \frac{1}{(2\pi)^{1/2}f_1(\lambda,1)}.$$
 (6.15)

Conclusion. For our purpose, it is worth noting that  $\delta$  can be chosen as small as desired, so that the critical value  $\alpha_c^-(0)$  can be obtained by putting  $\delta = 0$  in all

the previous. The reason is quite simple: the strategy for proving Proposition 4.5 applies provided the estimate of the density holds true on a small interval of the form  $(1 - \delta, 1)$ ,  $\delta$  being possibly really small. Obviously, when  $\delta$  tends to 0, the resulting Hölder constant in Proposition 4.5 blows up, but this does not affect the determination of  $\alpha_c^-(0)$ .

Therefore, when  $\lambda \ge 1$ . From (6.8), (6.10) (which implies (6.9)) and (6.14), we get as critical conditions for  $\alpha$  in order to satisfy (4.5):

$$\alpha \left[ 1 + \frac{\min[f_1(\lambda, \infty), f_2(\lambda, T_0, \infty)]}{1 - \varepsilon} \right] < 1,$$

$$\frac{1}{\varepsilon} g \left( \frac{\varepsilon}{f_1(\lambda, T_0)} \right) \alpha + \frac{1}{\varepsilon} f_2(\lambda, T_0, T_0) g \left( \frac{\varepsilon}{f_1(\lambda, T_0)} \right) \frac{\alpha}{1 - \alpha} < 1,$$

$$\min[f_4(\delta, \lambda), f_5(\delta, \lambda)] \alpha + \frac{1}{\varepsilon} f_2(\lambda, T_0, \infty) g \left( \frac{\varepsilon}{f_1(\lambda, T_0)} \right) \frac{\alpha}{1 - \alpha} < 1.$$
(6.16)

When  $\lambda < 1$ , we get similar conditions as in (6.16), but the third line in (6.16) is replaced by

$$\frac{1}{(2\pi)^{1/2}f_1(\lambda, T_0)}\alpha + \frac{1}{\varepsilon}f_2(\lambda, T_0, \infty)g\left(\frac{\varepsilon}{f_1(\lambda, T_0)}\right)\frac{\alpha}{1-\alpha} < 1.$$
(6.17)

The first inequality in (6.16) is explicit. The two last (and (6.17) as well) are of the form

$$c_1 \alpha + c_2 \frac{\alpha}{1 - \alpha} < 1 \Leftrightarrow c_1 \alpha^2 - (1 + c_1 + c_2)\alpha + 1 > 0$$
$$\Leftrightarrow \alpha < \frac{1 + c_1 + c_2 - [(1 + c_1 + c_2)^2 - 4c_1]^{1/2}}{2c_1},$$

where we used  $\alpha \in (0,1)$  in the two equivalences. Therefore, a lower bound for the critical value of  $\alpha$  can be computed numerically: we denote it by  $\alpha_0^{\varepsilon}$  as the coefficients in (6.16) depend upon  $\varepsilon \in (\exp(-1), 1)$ . Optimizing  $\alpha_0^{\varepsilon}$  over  $\varepsilon$ , we get a new bound, which must be read as the  $\alpha_0$  (or  $\alpha_0(0)$  if indicating the initial condition) in Proposition 4.5 and in Theorem 4.1. See Figure 2 for a plot of  $\alpha_0(0)$  in terms of  $\lambda$ (the optimization of  $\alpha_0^{\varepsilon}$  with respect to  $\varepsilon$  being performed numerically over a mesh of step size 0.01 for the parameter  $\varepsilon$ ) and Figure 3 for a zoom on the initial values.

The minimum is  $\alpha_0^{\min}(0) \approx 0.301$ . It is reached at  $\lambda = 0.23$ , which is the point at which Figure 3 has a singular behaviour. Actually, such a singularity is expected to be of computational essence and certainly indicates that the computations we have performed are far from being optimal, especially for  $\lambda$  small. Anyhow, for  $\lambda$  large, Figure 2 shows that the critical value  $\alpha_c^-(0)$  is (at worse) close to 1, which sounds as an interesting result from a practical point of view. Indeed, by scaling, the case  $\lambda \gg 1$  and  $\sigma = 1$  (with  $\sigma$  as in (1.2)) is equivalent to the case  $\lambda = 1$  and  $\sigma \ll 1$ : for a macroscopic value of  $\lambda$ , the system is thus uniquely solvable for most of the values of  $\alpha$  in (0, 1) when the noise is small. Actually, the convergence of  $\alpha_0(0)$  towards 1 as  $\lambda \to \infty$  can be proven rigorously, as stated in Proposition 6.3 below.



FIGURE 2. Plot of  $\alpha_0(0)$  in terms of  $\lambda \in [0; 80]$ .



FIGURE 3. Plot of  $\alpha_0(0)$  in terms of  $\lambda \in [0; 10]$ .

**Proposition 6.3.** In the case when  $X_0 = 0$  and  $b(x) = -\lambda x$  (with  $\sigma = 1$ ), the critical value  $\alpha_0$  in Theorem 4.1 and Proposition 4.5 can be chosen such that, for  $\lambda$  large,

$$\alpha_0 = \left[1 + \frac{1}{(1 - \exp(-1))(2\lambda)^{1/2}}\right]^{-1} \sim_{\lambda \to +\infty} 1 - \frac{1}{(1 - \exp(-1))(2\lambda)^{1/2}}.$$
  
$$\approx 1 - 1.12\lambda^{-1/2}.$$

In particular,  $\alpha_0 \to 1$  as  $\lambda \to +\infty$ .

*Proof.* The proof is quite straightforward and follows from the fact that, in (6.16), when  $\varepsilon$  is given, the coefficients of  $\alpha$  and  $\alpha/(1-\alpha)$  in the second and third lines tend to 0 as  $\lambda$  tends to the infinity, so that the conditions are automatically satisfied for  $\lambda$  large. In the first line, the condition is of the form

$$\alpha \Big[ 1 + \frac{1}{(1-\varepsilon)(2\lambda)^{1/2}} \Big] < 1.$$

Letting  $\varepsilon$  tend to the minimal authorized value  $\exp(-1)$ , we complete the proof.  $\Box$ 

The idea of the proof of Proposition 6.3 is thus very simple: the optimal strategy when  $\lambda$  is large consists in choosing  $\varepsilon$  close to the minimal authorized value  $\exp(-1)$ . Such a strategy is well-checked from a numerical point of view. Indeed, here is a plot of the optimal choice for  $\varepsilon$  in terms of  $\lambda$ , when optimizing  $\alpha_0^{\varepsilon}$  with respect to  $\varepsilon$ ( $\varepsilon$  being chosen amongst a mesh of step size 0.01):



FIGURE 4. Plot of the optimal value of  $\varepsilon$  in terms of  $\lambda \in [0; 40]$ .

6.2. A Case with Blow-Up. We now investigate the Brownian case (so that  $b \equiv 0$ ). We have already proven that, for any initial condition  $X_0 = x_0$ ,  $\alpha_c^-(x_0) > 0$ . On the other hand, by adapting the strategy in [3], we can prove that  $\alpha_c^+(x_0) < 1$  for some  $x_0$  (note that the result in [3] is written for a non-zero  $\lambda$ ).

Indeed, from Remark 5.3 (see in particular (5.1)), we know that, for any solution  $(X_t, M_t)_{t \ge 0}$  to (2.1), the Laplace transform of  $X_t$  at point  $\mu > 0$  (w(t) = z(t) in the notation of Remark 5.3) blows up as  $t \to \infty$  whenever

$$\exp(\mu x_0) > \frac{\exp(\mu) - 1}{\alpha \mu} \tag{6.18}$$

(i.e. whenever z(0) > v in the notation of Remark 5.3), which sounds as a contradiction since the Laplace transform at positive points keeps bounded as  $t \to \infty$ , since X remains bounded above by 1. In such cases, existence of a solution must fail.

Minimizing over  $\mu$  in (6.18), we deduce that existence fails for  $\alpha > \inf_{\mu>0}[[\exp(\mu) - 1]/[\mu \exp(\mu x_0)]]$ , so that  $\alpha_c^+(x_0) \leq \inf_{\mu>0}[[\exp(\mu) - 1]/[\mu \exp(\mu x_0)]]$  (if less than 1, which might not be the case). As in the previous subsection, this value must be compared with numerical experiments. Indeed, by investigating numerically the graphs of  $e(t) = \mathbb{E}(M_t)$  for different values of  $\alpha$ , we can detect the emergence of some discontinuity and thus give a numerical bound for  $\alpha_c^+(x_0)$ .

We choose  $x_0 = 0.8$ . Numerically, we find  $\inf_{\mu>0}[[\exp(\mu) - 1]/[\mu \exp(0.8\mu)]] \sim 0.539$ , so that  $\alpha_c^+(0.8) \leq 0.54$ . In Figure 5, we numerically observe that the graph of e is regular for  $\alpha = 0.38$  but has a jump for  $\alpha = 0.39$ . We also observe that the associated particle system keeps stable for  $\alpha = 0.38$  but blows up  $\alpha = 0.39$ . From the observations we have for other values of  $\alpha$ , it seems that, in this framework,

global solvability fails for  $\alpha \ge 0.39$  and holds for  $\alpha \le 0.38$ . We thus expect that, in this framework,  $\alpha_c^+(0.8)$  and  $\alpha_c^-(0.8)$  coincide, with a common value between 0.38 and 0.39. Figure 6 below summarizes these facts (when  $b(x) \equiv 0$  and  $x_0 = 0.8$ ). For  $\alpha$  in region **D**, by the above argument, we know that global uniqueness fails. For  $\alpha$  in region **C**, by numerical experiments it seems that global uniqueness also fails, but we are unable to prove it. Similarly, we have proven in this article that for  $\alpha$  in region **A**, for some  $\alpha_0(0.8) > 0$ , a unique solution for all time exists, while numerically it seems that global solutions also exist for  $\alpha$  in region **B**. We thus suspect that  $\alpha_c^-(0.8) = \alpha_c^+(0.8)$  lies in the region in between **B** and **C**.



FIGURE 5. Plot of  $t \mapsto e(t)$  for  $x_0 = 0.8$ ,  $b(x) \equiv 0$ ,  $\alpha = 0.38$  and  $\alpha = 0.39$ .



FIGURE 6. Critical regions of  $\alpha \in (0, 1)$ , for  $x_0 = 0.8$  and  $b(x) \equiv 0$ .

To complete Figure 6 we now explicitly find such an  $\alpha_0(0.8)$  in the case  $b \equiv 0$ under consideration, which can be used in Theorem 4.1 and Proposition 4.5 to prove global existence and uniqueness according to Theorem 5.2. Precisely, we prove that  $\alpha_0(0.8)$  can be taken as 0.104.

In fact we carry out the calculations for a general  $x_0 \in (0, 1)$ . As in the previous section, it suffices to find an  $\alpha_0(x_0) > 0$  such that in a neighbourhood of 1

$$\frac{d}{dx}\mathbb{P}(X_t \in dx) < \frac{1}{\alpha},\tag{6.19}$$

for all  $\alpha < \alpha_0(x_0)$  and all t > 0, where  $X_t$  is a solution to (2.1) with  $b \equiv 0$ , i.e.

$$X_t = x_0 + \alpha e(t) + W_t - M_t, \qquad e(t) = \mathbb{E}(M_t).$$

Suppose that  $\alpha \leq 1 - \eta$  for  $\eta \in (0, 1)$  with  $\eta$  to be chosen later and  $x_0 < 1 - \alpha$ . Then consider  $x \in [x_0/(1-\alpha) + \varepsilon, 1]$  with  $\varepsilon = (1/2) [1 - x_0/(1-\alpha)]$ . The following simple Gaussian estimate will be our starting point:

$$\frac{d}{dx}\mathbb{P}(X_t \in dx) = \frac{d}{dx}\mathbb{P}(Z_t - M_t \in dx) = \sum_{k=0}^{\infty} \frac{d}{dx}\mathbb{P}(Z_t - k \in dx, M_t = k)$$
$$\leqslant \frac{1}{\sqrt{2\pi t}} \sum_{k=0}^{\infty} \exp\left(-\frac{(x + k - x_0 - \alpha e(t))^2}{2t}\right), \tag{6.20}$$

where as usual  $Z_t = X_t + M_t$ .

We will need an *a priori* estimate of e(t), similar to the one given in Proposition 2.2. However, in this simple case when  $b \equiv 0$  we can do better. Indeed, using the fact that  $M_t = \lfloor \sup_{s \leq t} (Z_s)_+ \rfloor$  (see (4.3)) and following (2.10)–(2.11), we see that

$$e(t) = \mathbb{E}(M_t) = \mathbb{E}\left(\left|\sup_{s \leqslant t} (Z_s)_+\right|\right)$$
  
$$\leqslant x_0 + \alpha e(t) + \mathbb{E}\left(\sup_{s \leqslant t} W_s\right) \leqslant x_0 + \alpha e(t) + \left[\frac{2t}{\pi}\right]^{1/2}, \qquad (6.21)$$

the last inequality following from (6.4). By (6.21), this yields

$$e(t) \leqslant \frac{1}{1-\alpha} (x_0 + C\sqrt{t}) \tag{6.22}$$

where  $C = \sqrt{2/\pi}$ . As in the previous section we look at what happens first in small time t.

Small t: By (6.22), we have that, for any  $x \in [x_0/(1-\alpha) + \varepsilon, 1]$ ,

$$x - x_0 - \alpha e(t) \ge x - \frac{x_0}{1 - \alpha} - \frac{C\alpha}{1 - \alpha}\sqrt{t} \ge \varepsilon - \frac{C\alpha}{1 - \alpha}\sqrt{t}$$
$$\ge \frac{\varepsilon}{2} \tag{6.23}$$

when  $\sqrt{t} \leq \varepsilon(1-\alpha)/(2C\alpha)$ . Let  $t_0 = [\varepsilon\eta/(2C(1-\eta))]^2$ . Then since  $\alpha \leq 1-\eta$  we have that

$$t \leq t_0 \Rightarrow \sqrt{t} \leq \frac{\varepsilon \eta}{2C(1-\eta)} \leq \frac{\varepsilon(1-\alpha)}{2C\alpha}$$

so that (6.23) holds for all  $t \leq t_0$ . At this point we define  $\beta := C(1-\eta)/\eta$ , so that  $\sqrt{t_0} = \varepsilon/(2\beta)$ . From (6.23), we thus have that, for all  $t \in [0, t_0]$ ,

$$\exp\left(-\frac{(x-x_0-\alpha e(t))^2}{2t}\right) \leqslant \exp\left(-\frac{\varepsilon^2}{8t}\right)$$

and

$$\exp\left(-\frac{(x+k-x_0-\alpha e(t))^2}{2t}\right) \leqslant \exp\left(-\frac{k^2}{2t}\right)$$

for  $k \ge 1$ . Hence by (6.20)

$$\frac{d}{dx}\mathbb{P}(X_t \in dx) \leqslant \frac{1}{\sqrt{2\pi t}} \exp\left(-\frac{\varepsilon^2}{8t}\right) + \frac{1}{\sqrt{2\pi t}} \sum_{k=1}^{\infty} \exp\left(-\frac{k^2}{2t}\right), \qquad t \in [0, t_0].$$

Note that

$$\frac{1}{\sqrt{2\pi t}}\sum_{k=1}^{\infty}\exp\left(-\frac{k^2}{2t}\right) \leqslant \frac{1}{\sqrt{2\pi t}}\sum_{k=1}^{\infty}\int_{k-1}^{k}\exp\left(-\frac{x^2}{2t}\right)dx = \frac{1}{2}.$$

Moreover

$$\sup_{t \ge 0} \left[ \frac{1}{\sqrt{2\pi t}} \exp\left(-\frac{\varepsilon^2}{8t}\right) \right] = \sqrt{\frac{2}{\pi}} \exp\left(-\frac{1}{2}\right) \frac{1}{\varepsilon},$$
  
is attained when  $t = \varepsilon^2/4$ 

and the supremum is attained when  $t = \varepsilon^2/4$ .

We thus arrive at

$$\frac{d}{dx}\mathbb{P}(X_t \in dx) \leqslant \sqrt{\frac{2}{\pi}} \exp\left(-\frac{1}{2}\right)\frac{1}{\varepsilon} + \frac{1}{2}, \qquad t \in [0, t_0].$$
(6.24)

Large t (i.e.  $t \ge t_0$ ): For fixed  $x \in [x_0/(1-\alpha) + \varepsilon, 1]$  and t > 0 define

$$N(t, x) = \max \left\{ \lfloor \alpha e(t) + x_0 - x \rfloor, 0 \right\}.$$

Then, by (6.22) again, we have that

$$N(t,x) \leq \left(\alpha e(t) + x_0 - x\right)_+ \leq \left(\frac{C\alpha}{1-\alpha}\sqrt{t} + \frac{1}{1-\alpha}x_0 - x\right)_+ \\ \leq \left(-\varepsilon + \frac{C\alpha}{1-\alpha}\sqrt{t}\right)_+ \leq \frac{C\alpha}{1-\alpha}\sqrt{t}$$
(6.25)

for all t > 0. Moreover, for  $t \ge t_0$ ,

$$\frac{1}{\sqrt{2\pi t}} \sum_{k=0}^{\infty} \exp\left(-\frac{(x+k-x_0-\alpha e(t))^2}{2t}\right)$$

$$\leq \frac{1}{\sqrt{2\pi t}} \sum_{\substack{k \ge 0 \\ k:x+k-x_0-\alpha e(t) \le 1}} \exp\left(-\frac{(x+k-x_0-\alpha e(t))^2}{2t}\right) + \frac{1}{\sqrt{2\pi t}} \sum_{\substack{k \ge 1}}^{\infty} \exp\left(-\frac{k^2}{2t}\right)$$

$$\leq \frac{1}{\sqrt{2\pi t}} \left[\sum_{\substack{k=0 \\ k=0}}^{N(x,t)} \exp\left(-\frac{k^2}{2t}\right) + 1\right] + \frac{1}{2}$$

$$\leq \frac{1}{\sqrt{2\pi t}} \left[\sum_{\substack{k=0 \\ k=0}}^{\lfloor C\sqrt{t}\alpha/(1-\alpha)\rfloor} \exp\left(-\frac{k^2}{2t}\right) + 1\right] + \frac{1}{2}$$

$$\leq \frac{2}{\sqrt{2\pi t_0}} + \frac{C\alpha}{(1-\alpha)\sqrt{2\pi}} + \frac{1}{2} \leq 2\sqrt{\frac{2}{\pi}}\beta\frac{1}{\varepsilon} + \frac{\beta}{\sqrt{2\pi}} + \frac{1}{2},$$
(6.26)

using the fact that  $\sqrt{t_0} = \varepsilon/2\beta$ ,  $\beta = C(1-\eta)/\eta \ge C\alpha/(1-\alpha)$ .

We now consider two cases.

**Case 1:** when the RHS of (6.24) is bigger than the RHS of (6.26). This occurs if

$$2\sqrt{\frac{2}{\pi}}\beta\frac{1}{\varepsilon} + \frac{\beta}{\sqrt{2\pi}} \leqslant \sqrt{\frac{2}{\pi}} \exp\left(-\frac{1}{2}\right)\frac{1}{\varepsilon} \Leftrightarrow \qquad \frac{1}{2}(4+\varepsilon)\beta \leqslant \exp\left(-\frac{1}{2}\right)$$
$$\Leftrightarrow \qquad 1-\eta \leqslant \frac{2}{2+C\exp(1/2)(4+\varepsilon)}.$$

We therefore fix  $\eta$  such that  $1 - \eta = 2/(2 + C \exp(1/2)(4 + \varepsilon))$ . Thus we have a set of constraints on  $\alpha$  in order to achieve (6.19), which are

$$\alpha < \left(\sqrt{\frac{2}{\pi}}\exp\left(-\frac{1}{2}\right)\frac{1}{\varepsilon} + \frac{1}{2}\right)^{-1}$$
$$\alpha \leqslant 1 - \eta = \frac{2}{2 + C\exp(1/2)(4 + \varepsilon)},$$

subject to  $\alpha < 1 - x_0$ . Substituting in the value of  $\varepsilon$ , the constraints become

$$\alpha < \left(\frac{2C\exp(-1/2)(1-\alpha)}{1-\alpha-x_0} + \frac{1}{2}\right)^{-1} = f_1^{x_0}(\alpha) \tag{6.27}$$

$$\alpha \leqslant \frac{2}{2 + (4 + [1 - \alpha - x_0]/[2(1 - \alpha)]) C \exp(1/2)} = f_2^{x_0}(\alpha), \tag{6.28}$$

still subject to  $\alpha < 1 - x_0$ , with  $C = \sqrt{2/\pi}$ . The first thing to note is that  $f_1^{x_0}(\alpha)$  decreases to 0 on the interval  $\alpha \in [0, 1 - x_0)$ . By plotting the curves of  $f_1^{x_0}(\alpha)$  and  $f_2^{x_0}(\alpha)$  ( $\alpha \in [0, 1 - x_0)$ ) for different values of  $x_0$ , we see that (6.27) is more stringent than (6.28) for large values of  $x_0$ , while the reverse holds for small values of  $x_0$ . Indeed the critical value of  $x_0$  is approximately 0.51.

Hence for  $x_0 \gtrsim 0.51$ , if (6.27) is satisfied subject to  $\alpha < 1 - x_0$ , then we achieve (6.19). Thus for  $x_0 \gtrsim 0.51$ , we choose  $\alpha_0(x_0) = \min\{1 - x_0, \tilde{\alpha}_0(x_0)\}$  where  $\tilde{\alpha}_0(x_0)$  is such that

$$1 = \tilde{\alpha}_0(x_0) \left( \frac{2C \exp(-1/2)(1 - \tilde{\alpha}_0(x_0))}{1 - \tilde{\alpha}_0(x_0) - x_0} + \frac{1}{2} \right),$$

which solves as a second-order equation. For instance, this yields  $\alpha_0(0.8) \approx 0.104$ , which completes Figure 6 above.

Now, if  $x_0 \leq 0.51$  then (6.28) is the more stringent. In this case we must solve  $\alpha_0(x_0) = f_2^{x_0}(\alpha_0(x_0))$ . For example,  $\alpha_0(x_0) \approx 0.264$  when  $x_0 = 0.4$ . We note here, however, that as  $x_0 \downarrow 0$ ,  $f_2^{x_0}(\alpha) \to 4/[4 + 9C \exp(1/2)] \approx 0.2525$  for all  $\alpha \in (0, 1)$ . We can in fact hope to do better when  $x_0$  is small.

**Case 2:** when the RHS of (6.26) is bigger than the RHS of (6.24). In order to try and do better for  $x_0$  small, we now look at this case, which occurs when  $(4 + \varepsilon)\beta \ge 2\exp(-1/2)$ : here  $\varepsilon$  depends upon  $x_0$  but  $\beta$  does not so that  $\beta$  acts as a free parameter that can be tuned accordingly to the value of  $x_0$ . Such a strategy turns

out to be efficient when  $x_0$  is small. Indeed, our constraints now become

$$\alpha < \left(\sqrt{\frac{2}{\pi}}\frac{2\beta}{\varepsilon} + \frac{\beta}{\sqrt{2\pi}} + \frac{1}{2}\right)^{-1} = \left(\left(\frac{4}{\varepsilon} + 1\right)\frac{\beta}{\sqrt{2\pi}} + \frac{1}{2}\right)^{-1} \tag{6.29}$$

$$\alpha \leqslant 1 - \eta = 1 - \frac{C}{\beta + C} = \frac{\beta}{\beta + C},\tag{6.30}$$

again subject to  $\alpha < 1 - x_0$ , where we may vary  $\beta$ , as long as  $(4 + \varepsilon)\beta \ge 2 \exp(-1/2)$ . Now, optimizing (6.29) and (6.30) over  $\beta$  by equalizing the two right-hand sides and thus by solving

$$\beta\left(\left(\frac{4}{\varepsilon}+1\right)\frac{C\beta}{2}+\frac{1}{2}\right) = \beta + C \Leftrightarrow \left(\frac{4}{\varepsilon}+1\right)C\beta^2 - \beta - 2C = 0,$$

we would like to take  $\beta$  such that

$$2\left(\frac{4}{\varepsilon}+1\right)C\beta = 1 + \sqrt{1+4\left(\frac{4}{\varepsilon}+1\right)2C^2}.$$

This can be done provided it remains true that  $(4 + \varepsilon)\beta \ge 2\exp(-1/2)$ , i.e. that

$$(4+\varepsilon)\left(1+\sqrt{1+4\left(\frac{4}{\varepsilon}+1\right)2C^2}\right) \ge 4C\exp\left(-\frac{1}{2}\right)\left(\frac{4}{\varepsilon}+1\right),$$

which we can see numerically holds for  $\varepsilon > 0.15$  (which basically says that  $x_0$  must be small, as announced).

If we suppose that  $\alpha \leq 0.42$  (we will see below that we do not in fact lose anything by doing this), then since  $\varepsilon = (1 - x_0 - \alpha)/2(1 - \alpha)$ , it is certainly true that  $\varepsilon > 0.15$ when  $x_0$  is small. Indeed it is true when  $x_0 < 0.28$ . In this case, take  $\alpha_0(x_0)$  such that

$$\alpha_0(x_0) = 4 \left( 3 + \sqrt{1 + 4 \left(\frac{4}{\varepsilon} + 1\right) 2C^2} \right)^{-1}$$
$$= 4 \left( 3 + \sqrt{1 + 4 \left(\frac{8(1 - \alpha_0(x_0))}{1 - \alpha_0(x_0) - x_0} + 1\right) 2C^2} \right)^{-1}$$

Then it follows that whenever  $x_0 < 0.28$ , (6.19) holds for all  $\alpha < \alpha_0(x_0)$ . Taking  $x_0 = 0$  we calculate that

$$\alpha_0(0) = 4 \left(3 + \sqrt{1 + 72C^2}\right)^{-1} \approx 0.41,$$

which is much better than that achieved above for small  $x_0$ .

#### 7. Appendix: Proof of Lemma 3.2

*First Step.* We first discuss the solvability of (3.4). We start with the following case: we assume that  $\chi_0 = x_0 \in (-\infty, 1)$ , that b is smooth and bounded, with bounded derivatives of any order, and that e is smooth as well. Then, by [9, Th

1.10, Chap. VI], we know that the generator of  $\chi$ , namely the family of second-order differential operators

$$\left(\mathcal{L}_{s,x} := \left(b(\cdot) + \alpha e'(s)\right)\partial_x + \frac{1}{2}a\partial_{xx}^2\right)_{s \in [0,T]}$$

admits a Green function  $G : [0,T]^2 \times (-\infty,1)^2 \ni (s,t,x,y) \mapsto G(s,x,t,y)$ . For a given  $(t,y) \in [0,T] \times (-\infty,1]$ , the function  $[0,t) \times (-\infty,1] \ni (s,x) \mapsto G(s,x,t,y)$  is a classical solution of the PDE

$$\partial_s G(s, x, t, y) + \mathcal{L}_{s,x} G(s, x, t, y) = 0,$$

with G(s, 1, t, y) = 0, for  $s \in [0, t)$  and  $G(s, x, t, y) \to \delta_y(x)$  as  $s \nearrow t$ , where  $\delta_y$  is the Dirac mass at point y. Following [8, Th. 5, Sec. 5, Chap. 9], for a given  $(s, x) \in [0, T) \times (-\infty, 1)$ , the function  $(s, T] \times (-\infty, 1] \ni (t, y) \mapsto G(s, x, t, y)$  is also known to be the Green function of the adjoint operator

$$\partial_t + \partial_y \left[ \left( b(y) + \alpha e'(s) \right) \partial_y \right] - \frac{1}{2} \partial_{y,y}^2$$

In particular, it satisfies the Dirichlet boundary condition G(s, x, t, 1) = 0 together with  $G(s, x, t, y) \to \delta_x(y)$  as  $t \searrow s$ . When  $\chi_0 = x_0$ , we then set

$$p(t,y) = G(0, x_0, t, y), \quad t \in (0,T], \ y \in (-\infty, 1].$$

By [9, Th. 1.10, Chap. VI] (applied to the adjoint operator), we know that p(t, y) decays exponentially fast as t tends to 0 and y stays away from  $x_0$ . This proves that p is continuous on any compact subset of  $([0, T] \times (-\infty, 1]) \setminus \{(0, x_0)\}$ .

Second Step. Still in the smooth framework, we now make the connection with the diffusion process  $\chi$  when  $X_0 = x_0$ . Precisely, the point is to prove (3.3) when  $\chi_0 = x_0$ . Given a smooth function  $\phi : [0, T] \times (-\infty, 1] \to \mathbb{R}$ , with a compact support, the analysis of the Green function in [9, Th. 1.10, Chap. VI] says that the PDE

$$\partial_s u(t,x) + \left(b(x) + \alpha e'(s)\right) \partial_x u(s,x) + \frac{1}{2} \partial_{xx}^2 u(s,x) = \phi(s,x), \quad (s,x) \in (0,T] \times (-\infty,1),$$

with u(T, x) = 0,  $x \in (-\infty, 1)$ , as initial condition and u(s, 1) = 0,  $s \in [0, T]$ , as Dirichlet boundary condition, admits a (unique) classical solution:

$$u(s,x) = \int_{s}^{T} \int_{-\infty}^{1} G(s,x,t,y)\phi(t,y)dtdy, \quad s \in [0,T), \ x < 1.$$
(7.1)

Precisely, u is bounded and continuous on  $[0, T] \times (-\infty, 1]$  and is once continuously differentiable in time and twice differentiable in space on  $[0, T) \times (-\infty, 1)$ . Therefore, we can expand  $(u(t \wedge \tau_1, \chi_{t \wedge \tau_1}))_{0 \leq t \leq T}$  by Itô's formula, with  $\tau_1 = \inf\{t \geq 0 : \chi_t \geq 1\}$ . We then have the well-known representation formula:

$$u(0, x_0) = \mathbb{E} \int_0^{T \wedge \tau_1} \phi(t, X_t) dt$$
(7.2)

By equalizing (7.1) and (7.2), we deduce that

$$\mathbb{E}\int_0^{T\wedge\tau_1}\phi(t,X_t)dt = \int_0^T\int_{-\infty}^1 p(t,y)\phi(t,y)dtdy.$$
(7.3)

Writing

$$\mathbb{E} \int_0^{T \wedge \tau_1} \phi(t, X_t) dt = \int_0^T \mathbb{E} \left[ \phi(t, X_t) \mathbf{1}_{\{t < \tau_1\}} \right] dt$$
$$= \int_0^T \int_{-\infty}^1 \phi(t, y) \mathbb{P} \left( \chi_t \in dy, t < \tau_1 \right) dt$$

we deduce that

$$\int_{0}^{T} \int_{-\infty}^{1} \phi(t, y) \mathbb{P}(\chi_{t} \in dy, t < \tau_{1}) dt = \int_{0}^{T} \int_{-\infty}^{1} p(t, y) \phi(t, y) dt dy,$$
(7.4)

so that (3.3) holds in the smooth setting.

In the same framework, we then prove (3.5). The cumulative distribution function of  $\tau_1$  is given by

$$\mathbb{P}(\tau_1 \leqslant t) = 1 - \mathbb{P}(\tau_1 > t) = 1 - \int_{-\infty}^1 p(t, y) dy.$$

By [9, Th. 1.10, Chap. VI], we can differentiate the above expression with respect to t and exchange the derivative and the integral. From (3.4), we deduce:

$$\begin{aligned} \frac{d}{dt} \mathbb{P}(\tau_1 \leqslant t) &= -\int_{-\infty}^1 \partial_t p(t, y) dy \\ &= \int_{-\infty}^1 \partial_y \left( [\alpha e'(t) + b(y)] p(t, y) \right) dy - \frac{1}{2} \int_{-\infty}^1 \partial_{yy}^2 p(t, y) dy, \quad t \ge 0. \end{aligned}$$

Again by [9, Th. 1.10, Chap. VI], we know that both p(t, y) and  $\partial_y p(t, y)$  tend to 0 exponentially fast as  $y \to -\infty$ . So, using the boundary condition p(t, 1) = 0, we obtain (3.5).

Third Step. We now aim at proving the same results, still with  $\chi_0 = x_0$  but under the original assumptions on b and e. The strategy is to use a mollification argument. In order to so, we must prove that, in the smooth setting, p and  $\partial_y p$  can be bounded by constants that only depend upon T, A, K and  $\Lambda$ .

We thus go back to the case when b and e are smooth and bounded with bounded derivatives of any order. By Proposition 3.6, for  $t \in [0, T]$  small enough, we already have a Gaussian bound for  $\partial_y p(t, y)$  in terms of T, A, K and  $\Lambda$  only (notice that the argument applies since we know that p indeed satisfies PDE (3.4) in the smooth setting). With the same notation as in the statement of Proposition 3.6 (using in addition Proposition 3.4), the bound is of the form

$$\left|\partial_y p(t,y)\right| \leqslant \frac{C}{t} \exp\left(-\frac{|\xi_t^{x_0} - y|^2}{Ct}\right),\tag{7.5}$$

for  $t \in (0, \delta]$ , where  $(\xi_t^{x_0})_{t \ge 0}$  is given by (3.14) and C and  $\delta$  are constants that depend on T, A, K and  $\Lambda$  only. Plugging this bound into (3.10) and repeating the Gaussian convolution argument used in (3.31), we deduce that, for  $t \in (0, \delta]$ ,

$$p(t,y) \leqslant \frac{C}{\sqrt{t}} \exp\left(-\frac{|\xi_t^{x_0} - y|^2}{Ct}\right),\tag{7.6}$$

up to a new value of C.

Actually, (7.5) and (7.6) can be seen as bounds for the Green function G and its derivative in small time since  $p(t, y) = G(0, x_0, t, y)$ . In the same way, we could prove similar bounds for G(s, x, t, y) and  $\partial_y G(s, x, t, y)$  when  $t - s \in (0, \delta]$ . By a standard chaining argument, we then deduce that (7.5) and (7.6) are valid on the entire [0, T], for a possibly new value of C. Indeed, for a given  $t \in (0, T]$ , we can consider a sequence  $0 = t_1 < t_2 < \cdots < t_N = t$  such that  $t_{i+1} - t_i \leq \delta$ . Then, by using the Markov structure of  $\chi$ , we have:

$$p(t,y) = \int_0^t \prod_{i=1}^N G(t_{i-1}, t_i, z_{i-1}, z_i) dz_1 \dots dz_{N-1}$$

with the convention  $z_0 = 0$  and  $z_{N-1} = y$ . Noticing that N can be assumed to be bounded from above and using again a Gaussian convolution argument, we deduce that (7.5) and (7.6) can be extended to the entire (0, T].

Fourth Step. We still assume that the coefficients are smooth. By the third step, we are then able to reduce the PDE (3.4) to a heat PDE with a non-trivial source term:

$$\partial_t p(t,y) - \frac{1}{2} \partial_{yy}^2 p(t,y) = -\partial_y \big[ \big( b(y) + \alpha e'(t) \big) p(t,y) \big], \quad t \in (0,T], \ y < 1.$$

For any compact subset  $\mathcal{K} \subset (0,T] \times (-\infty,1]$ , we can consider a smooth cut-off function  $\eta : [0,T] \times \mathbb{R} \to \mathbb{R}_+$  matching 1 on  $\mathcal{K}$  and vanishing outside another compact subset  $\mathcal{K}' \subset (0,T] \times \mathbb{R}$ ,  $\mathcal{K} \subset \mathcal{K}'$ . Then, the function  $(0,T] \times (-\infty,1] \ni$  $(t,y) \mapsto [\eta p](t,y)$  satisfies the heat equation:

$$\left(\partial_t - \frac{1}{2}\partial_{yy}^2\right)\left[\eta p\right](t,y) = f(t,y),\tag{7.7}$$

with  $[\eta p](0, \cdot) = 0$  as initial condition and with  $[\eta p](t, 1) = 0$  and  $[\eta p](t, y) = 0$  for  $t \in [0, T]$  and |y| large enough as boundary conditions, where f is a smooth function which can be bounded in terms of A,  $\eta$ , K,  $\Lambda$  and T only. By [8, Th. 4, Sec. 2, Chap. 7], we deduce that  $\eta p$  and  $\partial_y [\eta p]$  are (1/4, 1/2)-Hölder continuous in (t, x), the Hölder constant depending on T, A,  $\eta$ , K and  $\Lambda$  only. Therefore, on any compact subset  $\mathcal{K}$  of  $(0, T] \times (-\infty, 1]$ , p and  $\partial_y p$  are (1/4, 1/2)-Hölder continuous in (t, x), the Hölder constant depending on T, A,  $\mathcal{K}$ , K and  $\Lambda$  only. By the same argument, on any compact subset  $\mathcal{K}$  of  $[0, T] \times \mathcal{I}$ ,  $\mathcal{I}$  being a finite interval included in  $(-\infty, 1]$  not containing  $x_0$ , p and  $\partial_y p$  are also (1/4, 1/2)-Hölder continuous in (t, x), the Hölder constant depending on T, A,  $\mathcal{K}$ , K and  $\Lambda$  only. Indeed, in such a case, p(t, y) and  $\partial_y p(t, y)$  tend to 0 as t tends to 0 and y stays in  $\mathcal{I}$ , so that the compact support  $\mathcal{K}'$  of  $\eta$  can be assumed to be included in  $[0, T] \times \mathbb{R}$  (and not necessarily in  $(0, T] \times \mathbb{R}$ ). In the end, we deduce that, on any compact subset  $\mathcal{K}$  of  $([0, T] \times (-\infty, 1]) \setminus \{(0, x_0)\}$ , p and  $\partial_y p$  are (1/4, 1/2)-Hölder continuous in (t, x), the Hölder constant depending on T,  $A, \mathcal{K}, \mathcal{K}$  and  $\Lambda$  only. Indeed, in such a case, p(t, y) and  $\partial_y p(t, y)$  tend to 0 as t tends to 0 and y stays in  $\mathcal{I}$ , so that the compact support  $\mathcal{K}'$  of  $\eta$  can be assumed to be included in  $[0, T] \times \mathbb{R}$  (and not necessarily in  $(0, T] \times \mathbb{R}$ ). In the end, we deduce that, on any compact subset  $\mathcal{K}$  of  $([0, T] \times (-\infty, 1]) \setminus \{(0, x_0)\}$ , p and  $\partial_y p$  are (1/4, 1/2)-Hölder continuous in (t, x), the Hölder constant depending on T,  $A, \mathcal{K}, K$  and  $\Lambda$  only.

In order to tackle the second-order derivatives in space, we assume that  $\mathcal{K}' \subset (0,T] \times (-\infty,1), \mathcal{K}'$  being the support of  $\eta$ . Then, the function  $(0,T] \times \mathbb{R} \ni (t,y) \mapsto [\eta p](t,y)$  satisfies (7.7) on the whole  $(0,T] \times \mathbb{R}$ , so that it can be represented as a

standard Gaussian convolution. Then, by Calderon and Zygmund estimates, see [19, Eq. (0.4), App. A], for any  $q \ge 1$ , the  $L^q([0,T] \times \mathbb{R}, dt \otimes dy)$ -norms of  $\partial_t[\eta p]$  and  $\partial_{yy}[\eta p]$  are bounded in terms of  $A, \eta, K, \Lambda$  and T only. Therefore, on any compact subset  $\mathcal{K}$  of  $(0,T] \times (-\infty,1)$ , for any  $q \ge 1$ , the  $L^q(\mathcal{K}, dt \otimes dy)$ -norms of  $\partial_t p$  and  $\partial_{yy}p$  are bounded in terms of  $T, A, \mathcal{K}, K$  and  $\Lambda$  only. For example, when  $\mathcal{K}$  is a cylinder of the form  $[\delta,T] \times [y-1,y+1]$ , (7.5) and (7.6) say that

$$\int_{\mathcal{K}} \left( |\partial_t p(t,z)|^q + |\partial_{yy}^2 p(t,z)|^q \right) dt dz \leqslant C_{q,\delta} \int_{\delta}^T \exp\left(-\frac{|\xi_s^{x_0} - y|^2}{C_{q,\delta}}\right) ds, \tag{7.8}$$

for a constant  $C_{q,\delta}$  that is independent of  $x_0$  and y.

Fifth Step. We now have all the required ingredients to go back to the original framework. The point is to approximate b and e by two sequences  $(b^n)_{n \ge 1}$  and  $(e^n)_{n \ge 1}$  that satisfy the previous smoothness conditions. The associated solutions to the PDE (3.4) are denoted by  $(p^n)_{n \ge 1}$ . On any compact subset  $\mathcal{K} \subset ([0,T] \times (-\infty,1]) \setminus \{(0,x_0)\}$ , the sequences  $(p^n)_{n \ge 1}$  and  $(\partial_y p^n)_{n \ge 1}$  are uniformly bounded and (1/4, 1/2)-Hölder continuous in (t, x). Therefore, there exists a subsequence  $(\varphi(n))_{n \ge 1}$  such that  $(p^{\varphi(n)})_{n \ge 1}$  and  $(\partial_y [p^{\varphi(n)}])_{n \ge 1}$  are uniformly convergent on compact subsets of  $([0,T] \times (-\infty,1]) \setminus \{(0,x_0)\}$ . Similarly, we can assume that the sequences  $(\partial_t [p^{\varphi(n)}])_{n \ge 1}$  and  $(\partial_{yy}^2 [p^{\varphi(n)}])_{n \ge 1}$  are weakly convergent for the  $L^q(\mathcal{K}, dt \otimes dy)$  topology, for any  $q \ge 1$  and any compact subset  $\mathcal{K} \subset (0,T] \times (-\infty,1)$ . The limit function of the sequence  $(p^{\varphi(n)})_{n \ge 1}$  is denoted by p: clearly, it is a solution of (3.4) (in the Sobolev sense), with p(t, 1) = 0, for t > 0, as Dirichlet boundary condition. By (7.6), it tends to 0 as t tends to 0 and y stays away from  $x_0$ . Moreover,  $\partial_y p$  exists and is continuous on  $([0,T] \times (-\infty,1]) \setminus \{(0,x_0)\}$ .

Then, we can prove (3.3). Indeed, we know that  $(p^{\varphi(n)})_{n \ge 1}$  converges toward p uniformly on compact subsets of  $([0,T] \times (-\infty,1]) \setminus \{(0,x_0)\}$ . Moreover, it is standard to prove that the solution  $\chi^n$  to the SDE (3.2), but driven by  $b^n$  and  $e^n$  instead of b and e, converges in law (as  $n \to \infty$ ) toward  $\chi$  on the space  $\mathcal{C}([0,T],\mathbb{R})$  of continuous functions on [0,T] endowed with the uniform topology. Denoting by  $\tau_1^n := \inf\{t \ge 0 : \chi_t^n \ge 1\}$ , we claim that the pair  $(\tau_1^n \wedge T, \chi^n)$  converges in law toward  $(\tau_1 \wedge T, \chi)$  on the product space  $\mathbb{R} \times \mathcal{C}([0,T],\mathbb{R})$  endowed with the product topology. Indeed, the mapping

$$\mathcal{C}([0,T],\mathbb{R}) \ni x \mapsto (\inf\{t \ge 0 : x(t) \ge 1\}) \land T$$

is continuous at any path x satisfying the "crossing property":

$$\left(\exists t \in [0,T) : x(t) \ge 1\right) \Rightarrow \left(\forall p \ge 1, \ \exists t_p \in (t,t+1/p) : x(t_p) > 1\right),\tag{7.9}$$

and (7.9) holds for a.e. trajectory of  $\chi$  because of the Brownian part in the dynamics of  $\chi$ , see (3.2). Recalling that (7.3) holds true in the smooth setting, that is with p and  $\tau_1$  therein replaced by  $p^n$  and  $\tau_1^n$ ,  $n \ge 1$ , we can pass to the limit along the subsequence  $(\varphi(n))_{n \ge 1}$ . We deduce that (7.3) holds in the general setting as well, from which we deduce that (7.4) also holds in the general setting, provided  $\chi_0 = x_0$ .

By the same approximation argument (using Scheffe's Lemma), we can prove that (3.5) holds in the general setting as well, provided  $\chi_0 = x_0$ .

Below, we shall denote  $p^{x_0}$  for p to indicate the dependence upon  $x_0$ . By (3.3),  $p^{x_0}$  depends upon  $x_0$  in a measurable way.

Sixth Step. To complete the proof, it remains to discuss what happens when  $\chi_0$  does not reduce to a Dirac mass. The point is then prove that

$$p(t,y) = \int_{-\infty}^{1} p^{x}(t,y) \mathbb{P}(\chi_{0} \in dx), \quad t \ge 0, \ y \le -1,$$
(7.10)

is the right candidate for solving the Fokker-Planck equation and for making the connection with  $\chi$ .

Actually, (3.3) is direct consequence of the Markov property for  $\chi$ . The point is thus to check that p satisfies (3.4) and (3.5). To this end, we must justify that we can differentiate under the integral in (7.10): this is a direct consequence of (7.5) and (7.8). Indeed, by (3.19), we have that, for any constant C > 0, there exists a constant C', only depending on T, C and K, such that

$$\begin{split} &\int_{-\infty}^{1} \exp\left(-\frac{|\xi_{t}^{x}-y|^{2}}{Ct}\right) \mathbb{P}\left(\chi_{0} \in dx\right) \\ &\leqslant \int_{-\infty}^{1} \exp\left(-\frac{|x-\xi_{-t}^{y}|^{2}}{C't}\right) \mathbb{P}\left(\chi_{0} \in dx\right) \\ &= \exp\left(-\frac{|1-\xi_{-t}^{y}|^{2}}{C't}\right) + \int_{-\infty}^{1} \frac{2(x-\xi_{-t}^{y})}{C't} \exp\left(-\frac{|x-\xi_{-t}^{y}|^{2}}{C't}\right) \mathbb{P}\left(\chi_{0} < x\right) dx \\ &\leqslant \exp\left(-\frac{|1-\xi_{-t}^{y}|^{2}}{C't}\right) + \int_{-\infty}^{1} \frac{2|x-\xi_{-t}^{y}| \mathbb{E}(|2-\chi_{0}|)}{C't(2-x)} \exp\left(-\frac{|x-\xi_{-t}^{y}|^{2}}{C't}\right) < +\infty, \end{split}$$

the third line following from an integration by parts and the last one form Markov's inequality. This also proves that both p(t, y) and  $\partial_y p(t, y)$  tend to 0 as y tends to  $-\infty$ .

#### References

- N. BRUNEL, Dynamics of sparsely connected networks of excitatory and inhibitory spiking neurons., J. Comput. Neurosci., 8 (2000), pp. 183–208.
- [2] N. BRUNEL AND V. HAKIM, Fast global oscillations in networks of integrate-and-fire neurons with low firing rates, Neural Comput., 11 (1999), pp. 1621–1671.
- [3] M. J. CÁCERES, J. A. CARRILLO, AND B. PERTHAME, Analysis of nonlinear noisy integrate & fire neuron models: blow-up and steady states, J. Math. Neurosci., 1 (2011), pp. Art. 7, 33.
- [4] J. A. CARRILLO, M. D. M. GONZALEZ, M. P. GUALDANI, AND M. E. SCHONBEK, Classical Solutions for a nonlinear Fokker-Planck equation arising in Computational Neuroscience. Preprint, 2011.
- [5] F. DELARUE, J. INGLIS, S. RUBENTHALER, AND E. TANRÉ, A particle method for the LIF model of McKean-Vlasov-type. Preprint, 2012.
- [6] F. DELARUE AND S. MENOZZI, Density estimates for a random noise propagating through a chain of differential equations, J. Funct. Anal., 259 (2010), pp. 1577–1630.
- [7] K. D. ELWORTHY AND X.-M. LI, Formulae for the derivatives of heat semigroups, J. Funct. Anal., 125 (1994), pp. 252–286.
- [8] A. FRIEDMAN, Partial differential equations of parabolic type, Prentice-Hall, Englewood Cliffs, N.J., first ed., 1964.

- [9] M. G. GARRONI AND J. L. MENALDI, Green functions for second order parabolic integrodifferential problems, Longman Scientific & Technical, Harlow, UK, 1992.
- [10] R. JOLIVET, T. J. LEWIS, AND W. GERSTNER, Generalized integrate-and-fire models of neuronal activity approximate spike trains of a detailed model to a high degree of accuracy, J. Neurophysiol., 92 (2004), pp. 959–976.
- [11] W. KISTLER, W. GERSTNER, AND J. L. VAN HEMMEN, Reduction of the hodgkin-huxley equations to a single-variable threshold model., Neural Comput., 5 (1997), pp. 1015–1045.
- [12] N. V. KRYLOV, Controlled diffusion processes, vol. 14 of Applications of Mathematics, Springer-Verlag, New York, 1980. Translated from the Russian by A. B. Aries.
- [13] T. J. LEWIS AND J. RINZEL, Dynamics of spiking neurons connected by both inhibitory and electrical coupling, J. Comput. Neurosci, 14 (2003), pp. 283–309.
- [14] S. OSTOJIC, N. BRUNEL, AND V. HAKIM, Synchronization properties of networks of electrically coupled neurons in the presence of noise and heterogeneities, J. Comput. Neurosci., 26 (2009), pp. 369–392.
- [15] P. E. PROTTER, Stochastic integration and differential equations, vol. 21 of Stochastic Modelling and Applied Probability, Springer-Verlag, Berlin, 2005. Second edition. Version 2.1, Corrected third printing.
- [16] A. RENART, N. BRUNEL, AND X.-J. WANG, Mean-field theory of irregularly spiking neuronal populations and working memory in recurrent cortical networks, in Computational neuroscience, Chapman & Hall/CRC Math. Biol. Med. Ser., Chapman & Hall/CRC, Boca Raton, FL, 2004, pp. 431–490.
- [17] L. C. G. ROGERS AND D. WILLIAMS, Diffusions, Markov Processes and Martingales. Volume 1 Foundations, John Wiley & Sons, Ltd, Chichester, second ed., 1994.
- [18] L. SACERDOTE AND M. T. GIRAUDO, Stochastic integrate and fire models: a review on mathematical methods and their applications, in Stochastic Biomathematical Models with Applications to Neuronal Modeling, vol. 2058 of Lecture Notes in Math., Springer, Berlin, 2013.
- [19] D. W. STROOCK AND S. R. S. VARADHAN, Multidimensional Diffusion Processes, vol. 233 of Grundlehren der mathematischen Wissenschaften, Springer-Verlag, Berlin, 1979.
- [20] A.-S. SZNITMAN, Topics in propagation of chaos, in École d'Été de Probabilités de Saint-Flour XIX—1989, vol. 1464 of Lecture Notes in Math., Springer, Berlin, 1991, pp. 165–251.

François Delarue, Sylvain Rubenthaler, Laboratoire J.-A. Dieudonné, Université de Nice Sophia-Antipolis, Parc Valrose, 06108 Nice Cedex 02, France.

*E-mail address*: delarue@unice.fr,rubentha@unice.fr

JAMES INGLIS, ETIENNE TANRÉ, EQUIPE TOSCA, INRIA SOPHIA-ANTIPOLIS MÉDITER-RANÉE, 2004 ROUTE DES LUCIOLES, BP 93, 06902 SOPHIA ANTIPOLIS CEDEX, FRANCE *E-mail address*: James.Inglis@inria.fr,Etienne.Tanre@inria.fr