



HAL
open science

The remarkable effect of network topology on calcium wave propagation in astrocyte networks

Jules Lallouette, Mati Goldberg, Maurizio de Pittà, Eshel Ben-Jacob, Hugues Berry

► **To cite this version:**

Jules Lallouette, Mati Goldberg, Maurizio de Pittà, Eshel Ben-Jacob, Hugues Berry. The remarkable effect of network topology on calcium wave propagation in astrocyte networks. Multidisciplinary approaches to quantify astrocyte neuron signaling, Jul 2012, Barcelona, Spain. 1 p. hal-00759232

HAL Id: hal-00759232

<https://inria.hal.science/hal-00759232>

Submitted on 1 Feb 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The remarkable effect of network topology on calcium wave propagation in astrocyte networks

Jules Lallouette¹, Mati Goldberg², Maurizio De Pittà², Eshel Ben-Jacob^{2,3} and Hugues Berry^{1,*}



1. BEAGLE Team, INRIA Rhône-Alpes, 69603 Villeurbanne, France
 2. School of Physics and Astronomy, Tel Aviv University, 69978 Ramat Aviv, Israel
 3. Center for Theoretical Biological Physics, UCSD, La Jolla, CA 92093-0319, USA





Calcium waves and network topologies


In astrocyte networks, the reported speed and extent of propagation of intercellular calcium signals can largely vary. Of course, this variability in the propagation patterns may reflect different intracellular properties (biochemical, signaling). But experimental evidence also suggests that the way astrocytes connect to each other in the network (topology) varies depending on the brain region. Such different topologies may already bring forth, by themselves, different modes of intercellular calcium propagation.

Different types of waves

Wave Type [exper. ref]	Activated cells	Speed ($\mu\text{m}\cdot\text{s}^{-1}$)
Locally synchronized [1]	~ 10	N.A.
Spatially restricted [2]	~ 40	~ 15
Regenerative [3]	∞	~ 35
Glissandi [4]	∞	~ 60



Different topologies



source : Giaume et al. (2010)


Can we account for the different wave properties by the different topologies ?

Calcium Dynamics model

$$\frac{dC^i}{dt} = J_{chan}(C^i, h^i, IP_3^i) + J_{leak}(C^i) - J_{pump}(C^i)$$


$$\frac{dh^i}{dt} = (h_{\infty}(C^i, IP_3^i) - h^i) / \tau_h(C^i, IP_3^i)$$

$$\frac{dIP_3^i}{dt} = P_{PLC\delta}(C^i, IP_3^i) - D_{3K}(C^i, IP_3^i) - D_{5P}(IP_3^i) + J_{net}^i$$





$$J_{j \rightarrow i} = \frac{F}{2} \left(1 + \tanh \left(\frac{|\Delta_{ji} IP_3| - IP_3^{thr}}{IP_3^{scale}} \right) \right) \frac{\Delta_{ji} IP_3}{|\Delta_{ji} IP_3|}$$

Simulation



3D Network topologies



- Stimulation of the center cell with IP₃ during the whole simulation.
- Several classes of spatial networks tested.



Influence of topology

Low mean degrees increases propagation

High mean-shortest path increases propagation




- The major classes of observed propagations can be emulated by a mere variation of the connection topology.


- Calcium wave propagation is favored when the connections between astrocytes are few and mainly restricted to small inter-cell distances.

- Propagation is improved when the mean-shortest path of the network is large.

Schematically :




Local rule of propagation



$$\chi = \frac{nb_{stim}^{crit}}{nb_{sink} + 1}$$


- Activated cells leak IP₃ to their inactive neighbors which act as sinks
- Unactivated cells receive IP₃ from all their active neighbors and get activated if the IP₃ total influx χ crosses a threshold

Effect of link parameters



$$\chi = A(IP_3^{thr})k + B(IP_3^{thr})$$

Simplified model



References

- [1] Sasaki et al. Locally synchronized astrocytes. Cereb Cortex, Jan 2011.
- [2] Charles. Intercellular calcium waves in glia. Glia, 24(1):39-49, 1998.
- [3] Scemes et al. Astrocyte calcium waves: What they are and what they do. Glia, 54(7):716-725, 2006.
- [4] Kuga et al. Large-scale calcium waves traveling through astrocytic networks in vivo. J Neurosci, 31(7):2607-2614, Feb 2011.
- [5] Giaume et al. Astroglial wiring is adding complexity to neuroglial networking. Front Neuroenergetics, 2, 2010.

Contact

jules.lallouette@inria.fr
 hugues.berry@inria.fr
 eshelbj@gmail.com