



HAL
open science

Using Salomé to reproduce the structure and to observe the diffusion of water molecules in biological tissue

Dang Van Nguyen, Denis S Grebenkov, Jing-Rebecca Li

► **To cite this version:**

Dang Van Nguyen, Denis S Grebenkov, Jing-Rebecca Li. Using Salomé to reproduce the structure and to observe the diffusion of water molecules in biological tissue. 2012. hal-00764328

HAL Id: hal-00764328

<https://inria.hal.science/hal-00764328>

Preprint submitted on 12 Dec 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Using Salomé to reproduce the structure and to observe the diffusion of water molecules in biological tissue

*Dang Van Nguyen¹, Denis Grebenkov², and Jing-Rebecca Li³


¹Neurospin, CEA Saclay, Gif-sur-Yvette, France, ²LPMC, Ecole Polytechnique, Palaiseau, France, ³Equipe DEFI, INRIA Saclay, Palaiseau, France

Introduction

Diffusion magnetic resonance imaging (DMRI) can give useful information on cellular structure and its structural changes [1]. Salomé [2] is used to reproduce some complicated shapes in d -dimensions ($d=2,3$) that are used to represent the natural structures of various biological tissue. The meshes representing these shapes are used as inputs to a finite element code that we built upon FENICS C++ [3]. Results were obtained for a model of glioblastoma (cerebral tumor) as a Voronoi diagram which was used to observe the convergence of the apparent diffusion tensor in long-time limit to the effective diffusion tensor computed by homogenization theory [4].


Problem formulation

- (1) The black points in the glioblastoma image are extracted to create a 2D Voronoi diagram by Matlab.
- (2) The common segments of two adjacent Voronoi cells are thickened to create extra-cellular space (Fig.1) in which the interior compartment contains cells, and the exterior compartment is between the cells.
- (3) Simulation




(3.a) Generate the triangular mesh from step 2 using Salomé and perform simulation for 2D using our finite element code.

(3.b) Make a thickness in the z-direction, create one layer, repeat this layer and put a thickness between two layers to create multi-layer structure, generate the tetrahedral mesh and do simulation for 3D.



The DMRI signals from which we define the apparent diffusion tensors are obtained by solving Bloch-Torrey equation for a Stejskal - Tanner PGSE sequence (Fig.2) and the effective infinite time diffusion tensors are obtained by solving Laplace equation on the same domain Ω which contains a representative sample of the cellular structure $\Omega = \prod_{k=1}^d [a_k, b_k]$ ($a_k < b_k$).



Bloch-Torrey PDE and the apparent diffusion tensor D^A

$$\frac{\partial M(\vec{x}, t)}{\partial t} = -i\gamma f(t)(\vec{g} \cdot \vec{x}) M(\vec{x}, t) + \nabla \cdot (D \nabla M(\vec{x}, t))$$

with initial conditions are the uniform distribution

$$D \nabla M^{in}(\vec{x}, t) \cdot \vec{n}^{in}(\vec{x}) = -D \nabla M^{ex}(\vec{x}, t) \cdot \vec{n}^{ex}(\vec{x}) = \kappa (M^{ex}(\vec{x}, t) - M^{in}(\vec{x}, t)), \vec{x} \in \Gamma,$$

$$[M(\vec{x}, t)]_{\vec{x}_i=a_i} = [M(\vec{x}, t)]_{\vec{x}_i=b_i} \exp(i(\vec{g} \cdot \vec{e}_i) \gamma l_i \int_0^t f(s) ds), \quad \left[\frac{\partial M(\vec{x}, t)}{\partial x_k} \right]_{\vec{x}_i=a_i} = \left[\frac{\partial M(\vec{x}, t)}{\partial x_k} \right]_{\vec{x}_i=b_i} \exp(i\gamma(\vec{g} \cdot \vec{e}_i) l_i \int_0^t f(s) ds)$$

$$\vec{g} = \vec{q} / \gamma \quad \Psi(\vec{q}, t) = \int_{\Omega} M(\vec{x}, t) d\vec{x} \quad \ln \Psi(\vec{q}, t) = \vec{q}^T D^A \vec{q} \int_0^t f(s) ds + O(\|\vec{q}\|^4)$$

Laplace's equation and the effective diffusion tensor D^{eff}

$$\nabla \cdot (D \nabla v_j(\vec{x})) = 0$$

$$D \nabla v_j^in(\vec{x}) \cdot \vec{n}^{in}(\vec{x}) = -D \nabla v_j^ex(\vec{x}) \cdot \vec{n}^{ex}(\vec{x}) = \kappa (v_j^ex(\vec{x}) - v_j^in(\vec{x})), \vec{x} \in \Gamma$$

$$[v_j(\vec{x})]_{\vec{x}_i=a_i} = [v_j(\vec{x})]_{\vec{x}_i=b_i} - (\vec{g} \cdot \vec{e}_i) l_i$$

$$\left[\frac{\partial v_j(\vec{x})}{\partial x_k} \right]_{\vec{x}_i=a_i} = \left[\frac{\partial v_j(\vec{x})}{\partial x_k} \right]_{\vec{x}_i=b_i} \quad D_{jk}^{eff} = [D_{jk}^{eff}]_{j=1, d, k=1, d}$$

$$D_{jk}^{eff} = D \int_{\Omega} \nabla v_j \cdot \vec{e}_k d\vec{x}$$

\vec{e}_k is the unit vector in the k^{th} direction

Results

We simulated on 2D and 3D voronoi cells in the box $[-25 \mu m; 25 \mu m]^d$ (Fig.2 and Fig.3) by taking the same intrinsic diffusion coefficient $D=3 \cdot 10^{-3} \mu m^2 / \mu s$ for interior and exterior compartments. The permeability of the inter-compartment interface was $\kappa = 10^{-5} \mu m / \mu s$. Seven b-values were taken from 0 to 4000 to get a curve of signal attenuation from which we obtain the tensor D^A .




Fig.2: 2D sample with 202 cells




Fig.3: 3D sample with 202x5 cells




Fig.4: The signal attenuation

The signals in both 2D and 3D were obtained at $\delta = \Delta = 10ms$ in Fig.4. The computation for 7 b-values costs about 11 minutes for 2D with the mesh-size of 8565 vertices and 1.5 hours for 3D with the mesh-size of 49035 vertices.

Conclusion and future works

A Voronoi diagram was created to observe the diffusion in glioblastoma in both 2D and 3D. Preliminary results showed that Salomé gives good meshes for this application. More realistic shapes are currently under development such as random cylinders to approach the structure of the human cerebral grey matter.




Fig.6: Human visual cortex (Le Bihan et al. PNAS 2006)

$$b\text{-value} = \gamma^2 \vec{g}^T \delta^2 (\Delta - \delta/3)$$

Fig.5 shows the convergence of the apparent diffusion coefficient D^A to the effective diffusion coefficient D^{eff} in 2D for $\delta = 20ms$ and $\Delta = 20 \cdot 180ms$.

The computation for each D^A computed from 7 b-values costs between 20 minutes to 50 minutes.




Fig.5: The convergence of D^A to D^{eff} for 2D

References

- [1] LeBihan (2007) Phys Med Bio 52.
- [2] <http://www.salome-platform.org/>
- [3] A. Logg, K.-A. Mardal, G. N. Wells et al. (2012). Automated Solution of Differential Equations by the Finite Element Method, Springer.
- [4] Bensoussan et al. (1978) Asymptotic analysis for periodic structures, North-Holland, Amsterdam.