



HAL
open science

Modeling a Case of Herding Behavior in a Multi-Player Game

Guy Fayolle, Jean-Marc Lasgouttes

► **To cite this version:**

Guy Fayolle, Jean-Marc Lasgouttes. Modeling a Case of Herding Behavior in a Multi-Player Game. [Research Report] RR-8190, 2012, pp.27. hal-00768324v1

HAL Id: hal-00768324

<https://inria.hal.science/hal-00768324v1>

Submitted on 21 Dec 2012 (v1), last revised 8 Jul 2013 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Modeling a Case of Herding Behavior in a Multi-Player Game

Guy Fayolle, Jean-Marc Lasgouttes

**RESEARCH
REPORT**

N° 8190

December 2012

Project-Team Imara



Modeling a Case of Herding Behavior in a Multi-Player Game*

Guy Fayolle[†], Jean-Marc Lasgouttes[†]

Project-Team Imara

Research Report n° 8190 — December 2012 — 24 pages

Abstract: The system mentioned in the title belongs to the family of the so-called massively multi-player online social games (MMOSG). It features a scoring system for the elements of the game that is prone to herding effects. We analyze in detail its stationary regime in the thermodynamic limit, when the number of players tends to infinity. In particular, for some classes of input sequences and selection policies, we provide necessary and sufficient conditions for the existence of a complete meanfield-like measure, showing off an interesting *condensation* phenomenon.

Key-words: Thermodynamical limit, condensation, ergodicity, non-linear differential system.

* This work was supported by grant 109-2167/R from the *Région Île-de-France*

[†] INRIA Paris–Rocquencourt — Domaine de Voluceau B.P. 105, 78153 Le Chesnay cedex (France)

**RESEARCH CENTRE
PARIS – ROCQUENCOURT**

Domaine de Voluceau, - Rocquencourt
B.P. 105 - 78153 Le Chesnay Cedex

Un modèle de comportement grégaire dans un jeu multi-joueurs

Résumé : Le système mentionné dans le titre fait partie de la famille des jeux sociaux en ligne dits « massivement multi joueurs ». Il repose sur un système de scores pour les éléments du jeu pouvant générer des comportements grégaires. Nous analysons en détail son régime stationnaire en limite thermodynamique, lorsque le nombre de joueurs tend vers l'infini. Pour certaines classes de paramètres, nous donnons des conditions nécessaires et suffisantes d'existence d'une mesure limite. Des phénomènes de condensation surprenants sont mis en évidence.

Mots-clés : Limite thermodynamique, condensation, ergodicité, système différentiel non linéaire.

The system mentioned in the title belongs to the family of the so-called massively multi-player online social games (MMOSG). It features a scoring system for the elements of the game that is prone to herding effects. We analyze in detail its stationary regime in the thermodynamic limit, when the number of players tends to infinity. In particular, for some classes of input sequences and selection policies, we provide necessary and sufficient conditions for the existence of a complete meanfield-like measure, showing off an interesting *condensation* phenomenon.

1 Description of the basic model

Ma Micro Planète [6] is a geolocalized massively multi-player online social game which entices players to use sustainable means of transport. At the heart of the game is a community-driven creation of a set of *points of interest* (POIs): interesting places, events, or even traffic incidents. The popularity of the POIs is assessed by scores. At his turn, the player is presented with a number of POIs to visit in his neighborhood, which have been selected through a mechanism that favors already popular ones. Then the selected POI has its score increased by a random value. It is not difficult to see that there is a risk of *herding* behavior, that would concentrate all the attention on a few places only. The aim of this paper is to present some results about the POIs dynamics by means of a probabilistic model, and in particular to determine when a stationary solution exists.

Assume there are N players in the game and let $n_i(t; N)$ be the number of POIs having an integer score i at time t . The state of the system at time t can be described by the infinite sequence

$$\vec{R}(t; N) = \frac{1}{N}[n_1(t; N), \dots, n_i(t; N), \dots].$$

The dynamics of the model is

- POIs are created according to a Poisson process of rate λN . Each new POI is granted an initial score k with probability $\varphi_k, k \geq 1$, where $\{\varphi_k, k \geq 1\}$ form a proper probability distribution.
- For any $i \geq 1$, each POI with score i decreases its score by 1 after a random time exponentially distributed at rate μ , where μ is a strictly positive constant.
- The N players are all independent and visit POIs at instants forming a global Poisson process with rate αN . Upon being visited, a POI sees its score increased by $j \geq 1$ with probability θ_j , independently of any other event, where the sequence $\{\theta_j, j \geq 1\}$ forms a proper probability distribution. We shall implicitly take $\theta_0 = 0$, which is in no way a restriction, as it is simply tantamount to modifying the parameter α .

- When he decides to play, at the instant of a Poisson process of parameter αN , a player selects a POI having score $i \geq 1$ with probability $\pi_i(\vec{R}(t; N))$, written as a functional of the state of the system, with the normalizing condition

$$\sum_{i \geq 1} \pi_i(\vec{R}(t; N)) = 1.$$

In accordance with the description of the game, we shall essentially consider selection policies such that the probability π of selecting a POI, given its score i , is a *non-decreasing* function of i .

The following typical scoring rules are used in *Ma Micro Planète*.

- Each player starts a journey at rate α and selects a POI to visit with a probability proportional to its current score.
 - with probability p , the player adds new information to the POI, the score of which is increased by 15 points;
 - otherwise, the POI is merely visited and its score gains 5 points; this happens with probability $q \stackrel{\text{def}}{=} 1 - p$.
- Each player creates POIs at rate λ and their initial score is set to 50.
- The POIs decrease their scores by 1 at a rate $\mu = 3$ per day.
- A POI disappears as soon as its score becomes zero or negative.
- The values of α , p and λ can only be measured empirically from concrete game statistics.

A way of describing this model might be in terms of queues. Consider a possibly infinite array of queues $M/M/1$ queues with service rate μ . New queues are created at rate λ and batches of customers join an existing queue with score $i \geq 1$ at rate $\alpha \pi_i(\vec{R}(t; N))$. Such systems have already been described, for example when the customers join the shortest of two queues [9]. The situation described in this paper is unusual due to the bias towards selecting the *longest* queue. This only makes sense because there is no cost associated to visiting a highly popular POI, whereas joining a long queue is expensive in terms of waiting time. A similar situation [8] occurs when customers, unsure of the quality of two restaurants, choose the busier one.

The model can have ramifications with other areas of interest (e.g., dynamics of populations, social networks, chemical or energy networks in physics).

In this work, we examine the system in the so-called *thermodynamic limit*, as $N \rightarrow \infty$, the existence of which is proved in Section 2, under general conditions, where the probability of increasing the score of a visited POI is

a function of the state of the system. Section 3 is devoted to the stationary regime, for which complete (but surprising!) answers are given for various input sequences and selection policies, and the existence of possible phase transition phenomena is enlightened. Section 4 presents miscellaneous simple limit cases. The final Section 5 outlines a possible method of solution for general input sequences, and also discusses a brief list of unsolved questions.

2 The thermodynamical limit

To capture information on the behavior of the system under various game policies, we proceed by scaling and analyze the *thermodynamical limit*, when the number N of players becomes large. In mathematical terms, we consider the system when $N \rightarrow \infty$, for any fixed t .

2.1 Notation

It will be convenient to gather here most of the notational material concerning input sequences and mathematical symbols.

- $\mathcal{E} \stackrel{\text{def}}{=} \{\vec{y} = [y_1, \dots, y_i, \dots], y_i \in \mathbb{Q}\}$, where \mathbb{Q}_+ is the set of positive rational numbers. \mathbb{R}_+^∞ being the set of non-negative real numbers, \vec{e}_j will stand for the j -th unit vector of \mathbb{R}_+^∞ . We shall also denote by $\mathbb{L} \in \mathbb{R}_+^\infty$ the subset of bounded sequences $\|\vec{y}\| \stackrel{\text{def}}{=} \sup_i |y_i| < M < \infty$.
- E being an arbitrary metric space, $\mathcal{B}(E)$ is the space of bounded real-valued Borel measurable functions on E , $\mathcal{C}(E) \subset \mathcal{B}(E)$ is the Banach subspace of bounded continuous functions, and $\mathcal{C}_1(E)$ is the subspace of functions with uniformly bounded continuous derivatives of first order.
- We shall need the two following subspaces

$$\mathcal{C}_2 \stackrel{\text{def}}{=} \left\{ f \in \mathcal{C}_1(\mathbb{L}) : \sum_{j \geq 1} \left| \frac{\partial f}{\partial y_j} + \frac{\partial^2 f}{\partial y_j^2} \right| \leq C \right\},$$

$$\mathcal{H}_2 \stackrel{\text{def}}{=} \left\{ f \in \mathcal{C}_1(\mathbb{L}) : \sup_{j,k \geq 0} \left| \frac{\partial^2 f}{\partial y_j^2} + \frac{\partial^2 f}{\partial y_k^2} - 2 \frac{\partial^2 f}{\partial y_j \partial y_k} \right| \leq H \right\},$$

C, H being arbitrary positive constants.

- For any complete separable metric space S , let $D_S[0, \infty]$ denote the space of right continuous functions $f : [0, \infty] \rightarrow S$ with left limits, endowed with the Skorokhod topology.

2.2 The main limit theorem

Clearly $\vec{R}(t; N)$ is a continuous time Markov chain, and its generator $G^{(N)}$, for $f \in \mathcal{B}(\mathbb{L})$, is the operator given by,

$$\begin{aligned} G^{(N)} f(\vec{y}) &= \lambda N \sum_{j \geq 1} \varphi_j \left[f \left(\vec{y} + \frac{\vec{e}_j}{N} \right) - f(\vec{y}) \right] \\ &\quad + \mu N \sum_{j \geq 1} y_j \left[f \left(\vec{y} + \frac{\vec{e}_{j-1}}{N} - \frac{\vec{e}_j}{N} \right) - f(\vec{y}) \right] \\ &\quad + \alpha N \sum_{j \geq 1} \pi_j(\vec{y}) \sum_{i \geq 1} \theta_i \left[f \left(\vec{y} - \frac{\vec{e}_j}{N} + \frac{\vec{e}_{i+j}}{N} \right) - f(\vec{y}) \right]. \end{aligned} \quad (2.1)$$

Throughout this study, the system will be subject to the following weak assumption.

Assumption (L) *The probability distribution $\{\pi_i(\vec{r}), i \geq 1\}$ is a functional supposed to be twice differentiable with respect to \vec{r} , for all $\vec{r} \in \mathbb{L}$.*

The above assumption (L) holds in particular for the forthcoming policy denoted by $\mathbf{P}_{\vec{a}}$, which will be analyzed in Section 3,

$$\pi_i(\vec{r}) = \frac{a_i r_i}{\sum_{j \geq 1} a_j r_j},$$

where the a_i form a sequence of positive numbers.

Then, with the notation $\theta_0 = \pi_0(\cdot) = 0$, the following proposition holds.

Theorem 2.1. *Suppose that, as $N \rightarrow \infty$, $\vec{R}(0; N)$ converges in distribution to $\vec{R}(0) \in \mathbb{L}$. Then the Markov process $\vec{R}(t; N)$ converges also weakly in $D[0, \infty]$ to a deterministic dynamical system $\vec{r}(\cdot) \in \mathbb{R}_+^\infty$, which evolves according to the following infinite system of nonlinear differential equations*

$$\frac{dr_i(t)}{dt} = \lambda \varphi_i + \mu [r_{i+1}(t) - r_i(t)] + \alpha \left[\sum_{j=0}^{j=i} \theta_j \pi_{i-j}(\vec{r}(t)) - \pi_i(\vec{r}(t)) \right], \quad (2.2)$$

where $\pi_i(\vec{r}(t))$ denotes the probability that the currently visited POI has score i . Moreover, under Assumption (L) and for any given finite initial condition $\vec{r}(0) \in \mathbb{L}$, the system (2.2) admits a unique solution, for all $t \geq 0$.

Proof. The method to solve this weak convergence problem is somehow classical (see for example the footsteps of [1]). First, a second order Taylor's expansion in (2.1) yields immediately, for any $f \in \mathcal{C}_2(\mathbb{L})$,

$$G^{(N)} f(\vec{y}) = Bf(\vec{y}) + \mathcal{O} \left(\frac{1}{N} \right),$$

where B is the operator with domain $\mathcal{C}_1(\mathbb{L})$ satisfying

$$\begin{aligned} Bf(\vec{y}) &= \lambda \sum_{j \geq 1} \varphi_j \frac{\partial f}{\partial y_j} + \mu \sum_{j \geq 1} y_j \left(\frac{\partial f}{\partial y_{j-1}} - \frac{\partial f}{\partial y_j} \right) \\ &+ \alpha \sum_{j \geq 1} \pi_j(\vec{y}) \sum_{i \geq 1} \theta_i \left(\frac{\partial f}{\partial y_{i+j}} - \frac{\partial f}{\partial y_j} \right). \end{aligned} \quad (2.3)$$

Under Assumption **(L)**, which is stronger than the famous Lipschitz condition, it is well-known that the system (2.2) admits a unique positive solution, for any initial finite condition $\vec{r}(0)$ and all finite t . Then for $f \in \mathcal{C}(\mathbb{L})$ one can define a one-parameter family $\{T(t), t \geq 0\}$, such that

$$T(t)f(\vec{r}(0)) = f(\vec{r}(t)).$$

It is not difficult to see that $\{T(t), t \geq 0\}$ is indeed a strongly continuous contraction semi-group, with generator, say G , which is closed by Theorem A.1. Now the two following properties can be checked at once.

- For all t , $\vec{r}(t) \in \mathcal{H}_2$.
- Taking $f \in \mathcal{C}_2$, then clearly $T(t)f(\vec{r}(0)) \in \mathcal{H}_2$. Since \mathcal{C}_2 is dense in $\mathcal{C}(\mathbb{L})$, it follows by Theorem A.3 that \mathcal{H}_2 is a core for G . Hence $G = \overline{B}$.

The conclusion of the proof is a direct application of Theorem A.4, together with standard results on differential systems, see e.g. [5]. \square

2.3 Some analytic facts

Let, for $|z| \leq 1$,

$$\begin{aligned} r(t, z) &= \sum_{i=1}^{\infty} r_i(t) z^i, & \pi(t, z) &= \sum_{i=1}^{\infty} \pi_i(\vec{r}(t)) z^i, \\ \varphi(z) &= \sum_{i=1}^{\infty} \varphi_i z^i, & \theta(z) &= \sum_{i=0}^{\infty} \theta_i z^i. \end{aligned}$$

In this respect, the model of *Ma Micro Planète* presented in Section 1 corresponds to

$$\theta(z) = pz^{15} + qz^5, \quad \varphi(z) = z^{50}.$$

Taking the summation over i in system (2.2), we get immediately

$$\frac{dr(t, z)}{dt} + \mu \left[1 - \frac{1}{z} \right] r(t, z) = \lambda \varphi(z) - \mu r_1(t) + \alpha [\theta(z) - 1] \pi(t, z). \quad (2.4)$$

The following proposition ensures in particular, for the class of policies considered in Section 2.2, the boundedness of the Cesàro limit of $r(t, 1)$ as $t \rightarrow \infty$, which, from its mathematical definition, takes into account the POIs possibly going to infinity.

Proposition 2.2. *When $\{\theta_k, k \geq 1\}$ is a proper probability distribution, so that $\theta(1) = 1$, we have*

$$\lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t r_1(t) dt = \frac{\lambda}{\mu}, \quad (2.5)$$

Moreover, if $\theta'(1) \cdot \varphi'(1) < \infty$, and for any policy satisfying Assumption **(L)**,

$$\lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t r(t, 1) dt = \frac{\alpha \underline{\pi} \theta'(1) + \lambda \varphi'(1)}{\mu} \leq \frac{\alpha \theta'(1) + \lambda \varphi'(1)}{\mu}, \quad (2.6)$$

where

$$\underline{\pi} = \lim_{s \rightarrow 0} s \pi^* \left(s, \frac{\mu}{\mu + s} \right) \leq 1.$$

Proof. For any arbitrary positive function f , let f^* denote its ordinary Laplace transform

$$f^*(s) \stackrel{\text{def}}{=} \int_0^\infty e^{-st} f(t) dt, \quad \Re(s) \geq 0.$$

Then, by a simple algebra carried out on equation (2.4), we obtain

$$r^*(s, z) = \frac{1}{\mu \left(1 - \frac{1}{z}\right) + s} \left[\frac{\lambda \varphi(z)}{s} + \alpha (\theta(z) - 1) \pi^*(s, z) - \mu r_1^*(s) + r(0, z) \right]. \quad (2.7)$$

Furthermore, applying Cauchy's formula

$$r_1^*(s) = \frac{1}{2i\pi} \int_{\mathcal{C}} \frac{r^*(s, y) dy}{y^2}$$

in equation (2.7), where \mathcal{C} stands for the unit circle centered at the origin, we get

$$\begin{aligned} r_1^*(s) &= \frac{1}{2i\pi} \int_{\mathcal{C}} \frac{\lambda \varphi(y) dy}{s y [(\mu + s)y - \mu]} \\ &\quad + \frac{1}{2i\pi} \int_{\mathcal{C}} \frac{[\alpha (\theta(y) - 1) \pi^*(s, y) - \mu r_1^*(s) + r(0, y)] dy}{y [(\mu + s)y - \mu]}, \end{aligned}$$

which in turn, by a repeated (and elementary) application of Cauchy's residue theorem, allows to extract

$$r_1^*(s) = \frac{\lambda \varphi\left(\frac{\mu}{\mu+s}\right)}{\mu s} + \frac{\alpha}{\mu} \left[\theta\left(\frac{\mu}{\mu+s}\right) - 1 \right] \pi^* \left(s, \frac{\mu}{\mu+s} \right) + \frac{1}{\mu} r\left(0, \frac{\mu}{\mu+s}\right). \quad (2.8)$$

Hence, instantiating the latter expression of $r_1^*(s)$ in (2.7), we obtain finally

$$\begin{aligned} r^*(s, z) &= \frac{1}{\mu \left(1 - \frac{1}{z}\right) + s} \left[\frac{\lambda}{s} \left(\varphi(z) - \varphi\left(\frac{\mu}{\mu+s}\right) \right) \right. \\ &\quad \left. + \alpha (\theta(z) - 1) \pi^*(s, z) + \alpha \left(1 - \theta\left(\frac{\mu}{\mu+s}\right) \right) \pi^* \left(s, \frac{\mu}{\mu+s} \right) \right. \\ &\quad \left. + r(0, z) - r\left(0, \frac{\mu}{\mu+s}\right) \right]. \end{aligned} \quad (2.9)$$

Take now $s > 0$ and let $z \rightarrow 1_-$ by real positive values in (2.9). This yields

$$r^*(s, 1) = \frac{1}{s} \left[\frac{\lambda}{s} \left(1 - \varphi \left(\frac{\mu}{\mu + s} \right) \right) + \alpha \left(1 - \theta \left(\frac{\mu}{\mu + s} \right) \right) \pi^* \left(s, \frac{\mu}{\mu + s} \right) \right], \quad (2.10)$$

where we have used the fact that $\pi^*(s, 1) = 1/s$, since $\pi(t, 1) = 1, \forall t \geq 0$.

Moreover, $\pi^* \left(s, \frac{\mu}{\mu + s} \right) < \pi^*(s, 1) = \frac{1}{s}$, for any $s > 0$, and we are now in a position to apply a Tauberian-type theorem for positive functions, see e.g. [3], just letting $s \rightarrow 0$ by positive values in (2.10), so that

$$\begin{aligned} \lim_{s \rightarrow 0} s r^*(s, 1) &= \lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t r(t, 1) dt \\ &= \lim_{s \rightarrow 0} \left[\frac{\lambda}{s} \left(1 - \varphi \left(\frac{\mu}{\mu + s} \right) \right) + \frac{\alpha}{s} \left(1 - \theta \left(\frac{\mu}{\mu + s} \right) \right) s \pi^*(s, 1) \right] \\ &= \frac{\alpha \pi \theta'(1) + \lambda \varphi'(1)}{\mu}, \end{aligned}$$

which is exactly the asserted equality (2.6). It is worth noting that for the moment there is no need to assume the finiteness of the derivatives $\theta'(1)$ and $\varphi'(1)$, but this will be made more precise in Section 3.

As for (2.5), it is readily obtained from (2.8), using the inequality $s \pi^* \left(s, \frac{\mu}{\mu + s} \right) \leq 1$, and again the above Tauberian theorem. The proof of the proposition is concluded. \square

3 System behavior as $t \rightarrow \infty$

In the sequel, we shall focus our attention solely on the deterministic dynamical system $\vec{r}(t)$ obtained in Theorem 2.1. Of special interest will be the non-degenerate stationary limits

$$\vec{r} = \lim_{t \rightarrow \infty} \vec{r}(t), \quad \text{with } r_i(t) \rightarrow r_i, \forall i \geq 1,$$

and

$$\lim_{z \rightarrow 1_-} \lim_{t \rightarrow \infty} \pi(t, z) = \bar{\pi}. \quad (3.1)$$

Indeed, under Assumption **(L)**, for a given set of parameters, a unique stationary regime of the system exists, but *possibly degenerate* in the sense that $\bar{\pi} < 1$. This follows from the general theory of differential systems (see e.g., [5]). Note that this fact is not easy to prove by a probabilistic argument, since, due to the normalizing condition, system (2.2) cannot be interpreted as forward Kolmogorov's equations for a special birth and death process, where $r(t, z)/r(t, 1)$ would represent the probability distribution function of the score of an arbitrary POI.

Throughout the rest of the paper, we shall speak of *ergodicity*, with a slight abuse of language: this will always refer to a *finite stationary* dynamical system satisfying the conditions

$$r(1) = \sum_{i \geq 1} r_i < \infty, \quad \bar{\pi} = 1. \quad (3.2)$$

3.1 Global conservation equations

Define the generating functions

$$r(z) \stackrel{\text{def}}{=} \sum_{i=1}^{\infty} r_i z^i, \quad \pi(z) \stackrel{\text{def}}{=} \sum_{i=1}^{\infty} \pi_i(\vec{r}) z^i,$$

with the shortened notation $\pi(z) \equiv \pi(z; \vec{r})$, and

$$\frac{1 - \theta(z)}{1 - z} \stackrel{\text{def}}{=} \sum_{n \geq 0} \Theta_n z^n, \quad \frac{1 - \varphi(z)}{1 - z} \stackrel{\text{def}}{=} \sum_{n \geq 0} \Phi_n z^n. \quad (3.3)$$

Two characteristic values are of special interest, namely

- $r(1) = \sum_{i \geq 1} r_i$, the mean number of POIs per player;
- r_1 , since μr_1 is the global death rate of POIs per player.

By (2.4), one sees easily that $r(z)$ satisfies the functional equation

$$\alpha [1 - \theta(z)] \pi(z) + \mu [1 - z^{-1}] r(z) = \lambda \varphi(z) - \mu r_1, \quad (3.4)$$

from which can be deduced the following corollary, which is the direct continuation of Proposition 2.2.

Corollary 3.1.

1. We have always $\lim_{z \rightarrow 1} (1 - z)r(z) = 0$ and $r_1 = \frac{\lambda}{\mu}$.
2. If in addition $\theta'(1) \cdot \varphi'(1) < \infty$, then

$$\frac{r(z)}{z} = \frac{\alpha \Theta(z) \pi(z)}{\mu} + \frac{\lambda \Phi(z)}{\mu}, \quad \forall |z| \leq 1, \quad (3.5)$$

$$r(1) = \frac{\alpha \bar{\pi} \theta'(1) + \lambda \varphi'(1)}{\mu}. \quad (3.6)$$

Proof. Immediate from Proposition 2.2 together with definition (3.3). \square

Assumption (F) *In the rest of the paper, unless otherwise explicitly mentioned, the product $\theta'(1).\varphi'(1) < \infty$ will always be assumed to be finite.*

In the next sections, we examine in detail the effect of specific choices of the policy $\pi(\cdot)$ on the behavior of the system. One of main goals of the study is to find out *the necessary and sufficient conditions* for the system to be *ergodic* in the sense given by the conditions (3.2).

3.2 Selection of POIs according to their scores

In this section, we consider a set of policies where the selection of a POI depends solely on its score. Letting each score $i \geq 1$ be associated with a positive weight a_i , then we recover the policy $\mathbf{P}_{\vec{a}}$ announced in Assumption (L) of Section 2.2, namely

$$\pi_i(\vec{r}) = \frac{a_i r_i}{K}, \quad K \stackrel{\text{def}}{=} \sum_{j=1}^{\infty} a_j r_j. \quad (3.7)$$

In the context of *Ma Micro Planète*, it seems quite realistic to assume that the a_i 's are *increasing with i* , since POIs with high scores should be more attractive. In the sequel, we shall mainly consider this situation of *herding effect*.

From equation (3.5), we get the immediate recursion, valid $\forall n \geq 0$,

$$r_{n+1} = \frac{\alpha}{\mu K} \sum_{i=1}^n a_i r_i \Theta_{n-i} + \frac{\lambda}{\mu} \Phi_n. \quad (3.8)$$

Since the a_i 's are assumed to form an increasing sequence, there exists $\lim_{i \rightarrow \infty} a_i$. When this limit is not finite, the following simple interesting proposition holds.

Proposition 3.2. *If $\lim_{i \rightarrow \infty} a_i = \infty$, then the system is non-ergodic.*

Proof. The relation (3.8) leads to the simple bound

$$r_{n+1} \geq \frac{\alpha}{\mu K} a_n r_n \Theta_0 = \frac{\alpha}{\mu K} a_n r_n,$$

so that no finite non-zero solution to the system can exist when the a_i 's are unbounded. \square

Consequently the only interesting case is when $\{a_i, i \geq 1\}$ is a non-decreasing sequence tending to a *finite limit*, which ad libitum can be taken equal to 1, since the a_i 's are defined up to a constant.

3.3 When scores of selected POIs are exactly increased by one

As it will appear in Sections 3.5 and 5.1, a complete solution of the non-linear equation (3.8) for a general $\theta(z)$ seems to be technically almost untractable. In this section, in order to get ideas about the general behavior of the system, we analyze in detail the simple (but nonetheless not elementary!) case $\theta(z) = z$, for which equation (3.8) becomes the following first order recursive sequence

$$r_{n+1} = \frac{\alpha}{\mu K} a_n r_n + \frac{\lambda}{\mu} \Phi_n. \quad (3.9)$$

A simple algebra carried out on this last equation leads to the formula

$$r_{n+1} = \frac{\lambda}{\mu} A_n \sum_{j=0}^n \frac{\Phi_j}{A_j} \left(\frac{\alpha}{\mu K} \right)^{n-j}, \quad n \geq 0, \quad (3.10)$$

where

$$A_n \stackrel{\text{def}}{=} \prod_{i=1}^n a_i.$$

The constant K can then be recovered as

$$\begin{aligned} K &= \sum_{i \geq 1} a_i r_i = \frac{\lambda}{\mu} \sum_{i=0}^{\infty} a_{i+1} A_i \sum_{j=0}^i \frac{\Phi_j}{A_j} \left(\frac{\alpha}{\mu K} \right)^{i-j} \\ &= \frac{\lambda}{\mu} \sum_{j=0}^{\infty} \frac{\Phi_j}{A_j \left(\frac{\alpha}{\mu K} \right)^j} \sum_{i=j}^{\infty} A_{i+1} \left(\frac{\alpha}{\mu K} \right)^i \\ &= \frac{\lambda K}{\alpha} \sum_{j=0}^{\infty} \frac{\Phi_j}{A_j \left(\frac{\alpha}{\mu K} \right)^j} \sum_{i=j+1}^{\infty} A_i \left(\frac{\alpha}{\mu K} \right)^i, \end{aligned}$$

which leads to the following implicit equation

$$\frac{\alpha}{\lambda} = F\left(\frac{\alpha}{\mu K}\right), \quad (3.11)$$

allowing to determine K , after having set

$$\begin{cases} F(z) \stackrel{\text{def}}{=} \sum_{k=1}^{\infty} u_k z^k, \\ u_k = \sum_{j=0}^{\infty} \frac{A_{j+k} \Phi_j}{A_j}, \quad \forall k \geq 1. \end{cases} \quad (3.12)$$

So, the problem of showing the existence of a *strictly positive and finite constant* K is exactly tantamount to finding a strictly positive real number x , such that

$$\frac{\alpha}{\lambda} = F(x) = \sum_{k=1}^{\infty} u_k x^k.$$

Hence, in order to get effective ergodicity conditions, one has to check the lower and upper bounds of $F(x)$, which is an increasing function of x , for $x \geq 0$. We note first that $F(x)$ is finite for small enough $|x|$ if, and only if,

$$u_1 = \sum_{j=0}^{\infty} a_{j+1} \Phi_j < \infty. \quad (3.13)$$

But, ex hypothesis, $A_{k+1}/A_k = a_k$ increases to 1 from below, so that u_k is a monotone decreasing sequence and inequality (3.13) is plainly equivalent to

$$\varphi'(1) = \sum_{j \geq 0} \Phi_j < \infty. \quad (3.14)$$

As for for the upper bound, we have the immediate inequality

$$u_k \leq \varphi'(1), \quad \forall k \geq 1,$$

and hence the radius of convergence of $F(z)$ is exactly equal to 1. Now, in order to decide whether the consistency relation (3.11) admits a solution, the key point will be to analyze the quantity

$$M = \lim_{x \rightarrow 1_-} F(x).$$

There are two possibilities.

- (i) If M is infinite, then for any value of λ , α , μ , it is possible to find a unique admissible finite K , and we shall say that the system is *ergodic*.
- (ii) Conversely, if $M < \infty$, then it will act as a limiting value for α/λ , and hence a non trivial *ergodicity condition* takes place.

The situation is summarized in the following brief statement.

Theorem 3.3. *Assume that $\lim_{i \rightarrow \infty} a_i \nearrow 1$.*

- (i) *The system is never ergodic when $\varphi'(1) = \infty$.*
- (ii) *When $\varphi'(1) < \infty$, the system is ergodic if, and only if,*

$$\frac{\alpha}{\lambda} \leq \sum_{k=1}^{\infty} \sum_{j=0}^{\infty} \frac{A_{j+k} \Phi_j}{A_j}, \quad (3.15)$$

where the r.h.s. of the inequation above may be infinite.

It is interesting to note a few not so intuitive facts related to (3.15): firstly, this condition does not depend on the value of μ , as long as $\mu > 0$ obviously. Secondly, the ergodicity range increases when the initial score increases in distribution — that is, when the Φ_i 's increase.

To propose a somehow concrete classification, let us concentrate for the remainder of this section on the reasonably general set of sequences satisfying, for some strictly positive constants γ and ν ,

- $a_1 > 0$,
- the a_i 's form an increasing sequence,
- $a_i = 1 - \gamma i^{-\nu} + \mathcal{O}(i^{-\nu-1})$.

The following proposition describes the *phase transition* phenomenon that occurs when the parameters vary.

Proposition 3.4. *Assume the sequence a_i satisfies the above conditions, and that $\theta(z) = z$. Then,*

- if $\nu > 1$, or $\nu = 1$ and $\gamma \leq 1$, the system is always ergodic;
- if $\nu < 1$, or $\nu = 1$ and $\gamma > 1$, there is a finite ergodicity bound M if, and only if, $\sum_{j>0} j^{\nu+1} \varphi_j < \infty$.

Proof. The first step is to obtain a convenient estimate for A_n when n is large. Using the property $0 < a_1 < a_i < 1$, valid for all i , one gets

$$\log A_n = \sum_{i=1}^n \log a_i = -\gamma \sum_{i=1}^n i^{-\nu} + \mathcal{O}(1).$$

The finiteness of M depends strongly on the properties of the series $\sum A_n$, which can be deduced from the following estimates.

- If $\nu > 1$, then the Dirichlet series $\sum_{i \geq 1} i^{-\nu}$ converges to some positive number, so that the series $\sum A_n$ diverges, and then clearly $M = +\infty$.
- If $\nu = 1$, then $A_n \approx C_1 n^{-\gamma}$. When $\gamma \leq 1$, $\sum A_n$ still diverges and $M = +\infty$; conversely, when $\gamma > 1$,

$$\frac{1}{A_j} \sum_{i=j+1}^{\infty} A_i \approx C_2 j,$$

and M is finite whenever the second moment $\sum_{j>0} j^2 \varphi_j$ of the distribution $\{\varphi_k, k \geq 1\}$ exists.

- If $\nu < 1$, then $A_n = \mathcal{O}(e^{-\gamma n^{1-\nu}})$. Using classical techniques (Riemann sums and integration by parts), one gets

$$\sum_{i=j+1}^{\infty} e^{-\gamma i^{1-\nu}} \approx \int_j^{\infty} e^{-\gamma x^{1-\nu}} dx \approx \frac{j^{\nu}}{\gamma(1-\nu)} e^{-\gamma j^{1-\nu}},$$

and hence M is finite whenever $\sum_{j>0} j^{\nu+1} \varphi_j$ is.

This concludes the proof of the proposition, the results of which will be illustrated in Section 4.5. \square

3.4 Picking the largest of c scores

So far, all models which have been considered are based on exogenous parameters a_i , supposed to be known. Actually, it turns out that most of the computations remain valid when these parameters are a *functional of the state* $\vec{r}(t)$. We tackle hereafter such a model, where the ergodicity conditions can be obtained explicitly.

The algorithm is simple: $c > 1$ POIs are selected uniformly, and the one with the largest score will be visited. This model is reminiscent of the queuing model of [9], which proves to be very efficient in thermodynamical limit. The probability of selecting a POI with score less or equal to i is then

$$\left[\frac{1}{r(1)} \sum_{j=1}^i r_j \right]^c,$$

whence we get immediately

$$\pi_i(\vec{r}) = \frac{1}{r(1)^c} \left(\left[\sum_{j=1}^i r_j \right]^c - \left[\sum_{j=1}^{i-1} r_j \right]^c \right). \quad (3.16)$$

From a formal point of view, we are actually in the framework of Section 3.3, as it can be seen by setting

$$a_i = \frac{1}{c r(1)^{c-1}} \frac{1}{r_i} \left(\left[\sum_{j=1}^i r_j \right]^c - \left[\sum_{j=1}^{i-1} r_j \right]^c \right), \quad K = \frac{r(1)}{c}, \quad (3.17)$$

and remarking that the a_i 's are no more exogenous constants as before. To check they form an increasing positive sequence bounded by 1, it suffices to note that x^c is a convex function of x , for any $c \geq 2$, so that

$$0 < \frac{y^c - x^c}{y - x} < cz^{c-1}, \quad \forall x \leq y \leq z,$$

choosing here

$$x = \sum_{j=1}^i r_j, \quad y = \sum_{j=1}^{i-1} r_j, \quad z = r(1).$$

While the equations are similar, the situation is nonetheless very different from Section 3.2. The model is more difficult to solve because π_i 's do not have a simple expression in terms of the score i and a solution like (3.10) is not really usable; however, they can be easily computed by recurrence, since the normalization constant $r(1)$ is given by (3.6). As a consequence, the ergodicity condition is easier to obtain.

Theorem 3.5. *Assume $\theta(z) = z$ and that the POIs are visited according to policy (3.16). Then, according to the definition given in Section 3, the system is ergodic if, and only if,*

$$\frac{\alpha}{\lambda} \leq \frac{\varphi'(1)}{c-1}.$$

Proof. Define

$$R_n \stackrel{\text{def}}{=} \sum_{i=1}^n r_i, \quad B_n \stackrel{\text{def}}{=} \sum_{i=1}^n \Phi_i.$$

Then (3.9) takes the form

$$R_{n+1} - R_n = \frac{\alpha}{\mu} \frac{R_n^c - R_{n-1}^c}{r(1)^c} + \frac{\lambda}{\mu} \Phi_n,$$

which by a direct summation yields

$$R_{n+1} = \frac{\alpha}{\mu} \frac{R_n^c}{r(1)^c} + \frac{\lambda}{\mu} B_n, \quad (3.18)$$

with $R_1 = \lambda/\mu, B_1 = 1$ and $B_n \nearrow B_\infty = \varphi'(1)$. Since we are looking for a sequence R_n satisfying,

$$R_\infty = \lim_{n \rightarrow \infty} R_n = r(1),$$

it will be convenient for the sake of consistency to rewrite (3.18) as

$$R_{n+1} = \left(\frac{\alpha}{\mu K} \right) \frac{R_n^c}{cR_\infty^{c-1}} + \frac{\lambda}{\mu} B_n, \quad (3.19)$$

remembering that $r(1) = cK$ by (3.17). The interest of the form (3.19) is that it allows for a direct graphical analysis of the convergence problem, starting from the following simple facts.

1. The curves $\{y = Ax^c + B_n\}$ and $\{y = Ax^c + B_{n+1}\}$ are convex, parabolic shaped, and obtained from each other by translation of vector $\frac{\lambda}{\mu} \Phi_n$ along the y -axis.
2. The factor cR_∞^{c-1} is the value of the slope at the limit point when it exists.

Hence one sees that (3.19) defines an increasing sequence converging to a unique value R_∞ , if and only if the expansion factor $\alpha/\mu K$ is ≤ 1 . In this case, letting $n \rightarrow \infty$, we get immediately

$$R_\infty = r(1) = \frac{\alpha + \lambda\varphi'(1)}{\mu},$$

which coincides (as expected !) with the result given in (3.6). To conclude the proof of the theorem, it suffices to quote the equivalence

$$\frac{\alpha}{\mu K} \leq 1 \quad \Longleftrightarrow \quad \frac{\alpha}{\lambda} \leq \frac{\varphi'(1)}{c-1}.$$

□

3.5 About the general case: a matrix form solution

The problem becomes more intricate when $\theta(z)$ is an arbitrary probability generating function with $\varphi'(1) < \infty$. Then the following approach, based on the recursion (3.8), involves infinite matrices and seems to be computationally quite acceptable. We sketch hereafter its main lines.

Clearly, equation (3.8) can be rewritten in the vector form

$$\vec{U}_{n+1} = M_n\left(\frac{\alpha}{\mu K}\right)\vec{U}_n + \frac{\lambda}{\mu}\vec{V}_{n+1}, \quad (3.20)$$

where

- \vec{U}_n is the n -column vector $[r_n r_{n-1} \dots r_1]^T$, where T denotes the transpose operation;
- \vec{V}_n is the n -column vector $[\Phi_{n-1}, 0, \dots, 0]^T$;
- $M_n(x)$ is a $(n+1) \times n$ matrix for all $x > 0$. Its first row is the vector $x\vec{B}_n$, where \vec{B}_n denotes the row vector

$$[a_n \Theta_0, a_{n-1} \Theta_1, \dots, a_1 \Theta_{n-1}],$$

and the remaining internal matrix is exactly the $n \times n$ identity matrix denoted by I_n .

By a direct algebra, the solution takes the form

$$\vec{U}_{n+1} = \frac{\lambda}{\mu} \sum_{i=1}^{n+1} Q_{n,i}\left(\frac{\alpha}{\mu K}\right)\vec{V}_i, \quad n \geq 0, \quad (3.21)$$

where

$$\begin{cases} Q_{n,i}(x) = M_n(x)M_{n-1}(x)\dots M_i(x), & i \leq n, \\ Q_{n,n+1}(x) \stackrel{\text{def}}{=} I_{n+1}. \end{cases}$$

The next step is to carry out the scalar product of (3.21) with the row vector

$$\vec{A}_{n+1} \stackrel{\text{def}}{=} [a_{n+1}, \dots, a_1],$$

and then to let $n \rightarrow \infty$. Keeping in mind that $K = \sum_{i \geq 1} a_i r_i$, we get the following final relationship, which is formally similar to the one given by (3.11),

$$\frac{\alpha}{\lambda} = \frac{\alpha}{\mu K} \sum_{n=0}^{\infty} \sum_{i=1}^{n+1} \vec{A}_{n+1} \tilde{Q}_{n,i}\left(\frac{\alpha}{\mu K}\right)\vec{V}_i. \quad (3.22)$$

From the definition of M_n , each term of the double sum in (3.22) is clearly a *decreasing* scalar function of K . Hence, ergodicity conditions could be obtained along the same lines as in Theorem 3.3, by analyzing the series in the right-hand side member of (3.22).

4 Some simple examples and limit cases

For the sake of completeness, we quote hereafter simple peculiar cases, which shed light on some aspects of the behavior of the system.

4.1 No creation of POIs, i.e. $\lambda = 0$

In this case, the system is empty at steady state for any Markovian policy. The argument is elementary, since the vector state $\vec{0}$ is absorbing due to the $M/M/\infty$ character of the score decreasing process.

4.2 The lower bound $\alpha = 0$

Here no POI will see any increase of score upon being visited, and hence one gets a clear lower bound for the model. Solving (3.5) leads to

$$r_i = \frac{\lambda}{\mu} \sum_{k \geq i} \varphi_k, \quad r(1) = \frac{\lambda \varphi'(1)}{\mu}, \quad r'(1) = \frac{\lambda [\varphi''(1) + 2\varphi'(1)]}{2\mu}.$$

Consequently, when $\alpha \neq 0$, we have the following bound, valid for any point selection scheme (i.e. policy) π ,

$$r'(1) \geq \frac{\lambda [\varphi''(1) + 2\varphi'(1)]}{2\mu},$$

and therefore the mean score $r'(1)/r(1)$ of a POI can be finite only when $\varphi''(1)$ is finite. This leads to some unintuitive behavior, since it was shown in Proposition 3.4 that having $\varphi''(1) = \infty$ can at the same time make the system always ergodic.

4.3 Score independent visits

When the POIs are visited uniformly regardless of their score, which amounts to the choice

$$\pi_i(\vec{r}) = \frac{r_i}{\sum_{i \geq 1} r_i},$$

we have $\pi(z) = r(z)/r(1)$. Then equation (3.5) allows to derive immediately the corresponding generating function, which takes the compact explicit form

$$r(z) = \frac{\lambda r(1)(\varphi(z) - 1)}{\alpha(1 - \theta(z)) + \mu r(1)(1 - z^{-1})},$$

where, for $\lambda \neq 0$,

$$r(1) = \frac{\alpha \theta'(1) + \lambda \varphi'(1)}{\mu},$$

and one can check the denominator of $r(z)$ does not vanish for $|z| < 1$. Moreover, $r(1)$ is clearly finite and given by equation (3.6). So, we have

obtained a unique function $r(z)$ analytic in the unit disc and continuous for $|z| = 1$. Consequently the system admits of a unique limit invariant measure $\{r_i, i \geq 1\}$ for any value of the parameters. Intuitively, this is due to the fact that the decrease rate of the scores is somehow proportional to the number of points, like in a $M/M/\infty$ queue. But, when the selection process favors heavily enough the POIs having a big score, proper ergodicity conditions appear, as in the models analyzed in Section 3.2.

4.4 An unstable model with a score dependent policy

A policy that may seem attractive at first sight is to give to each POI a weight proportional to its score

$$\pi_i(\vec{r}) = \frac{ir_i}{\sum_{i \geq 1} ir_i},$$

so that $\pi(z) = zr'(z)/r'(1)$ and (3.5) becomes a differential equation

$$\alpha [1 - \theta(z)] \frac{zr'(z)}{r'(1)} + \mu[1 - z^{-1}]r(z) = \lambda[\varphi(z) - 1].$$

However, the weight $a_i = i$ given to the score i is unbounded, and from Proposition 3.2 we conclude that no choice of parameters can lead to a stable system under this policy. More precisely, *there exists at least one POI having an infinite score.*

4.5 A case study of Proposition 3.4

Assuming the conditions of Proposition 3.4 hold, let us take in addition $\varphi(z) = z$ and, for some $\gamma > 0$,

$$a_i = \left(\frac{i}{i+1}\right)^\gamma.$$

This implies $A_k = 1/(k+1)^\gamma$ and therefore (3.10)–(3.11) become

$$r_{k+1} = \frac{\lambda}{\mu} \frac{1}{(k+1)^\gamma} \left(\frac{\mu K}{\alpha}\right)^k,$$

$$\frac{\alpha}{\lambda} = F(x) = \sum_{k=1}^{\infty} A_k x^k = \Phi(x, \gamma, 1) - 1,$$

where Φ stands for the classical Lerch function [4, formula 9.550]

$$\Phi(x, \gamma, v) \stackrel{\text{def}}{=} \sum_{k=0}^{\infty} \frac{x^k}{(k+v)^\gamma}.$$




Figure 1: Expected score of POIs per player $r'(1)$ as a function of the visit rate α . The parameters $\lambda = \mu = 3$ remain fixed.

When $\varphi(z) = z$, the computation of the expected score of POIs per player $r'(1)$ is easy:

$$r'(1) = \sum_{k=1}^{\infty} k r_k = \sum_{k=1}^{\infty} \frac{\lambda}{\mu} \sum_{k=1}^{\infty} k A_k \left(\frac{\alpha}{\mu K} \right)^k = \frac{\lambda}{\mu} \frac{\alpha}{\mu K} F' \left(\frac{\alpha}{\mu K} \right).$$

The above formulas, allow to represent the expected score per player as a function of α in a parametric way, by letting x vary from 0 to 1 (Figure 1). As expected from the theoretical results obtained above, 3 different situations prevail:

- $\gamma \leq 1$: the system is always ergodic (dashed curves);
- $1 < \gamma \leq 2$: there is an explicit ergodicity condition on α , and $r'(1)$ tends to infinity at the boundary;
- $\gamma > 2$: there is an explicit ergodicity condition on α , but $r'(1)$ remains bounded.

5 Miscellaneous outcomes and open problems

The next three paragraphs present briefly some ideas and mathematical questions concerning the general model.

5.1 About a general solution via an integral equation

Consider the situation of Section 3.2, and let $a(z) \stackrel{\text{def}}{=} \sum_{j \geq 1} a_j z^j$. Then by Hadamard's theorem [7] and definition (3.7), we can write

$$\pi(z) = \frac{1}{2i\pi K} \int_{\mathcal{L}} r(w) a\left(\frac{z}{w}\right) \frac{dw}{w},$$

which yields the non-linear integral equation

$$\frac{\mu r(z)}{z} = \frac{\alpha[1 - \theta(z)]}{2i\pi K(1 - z)} \int_{\mathcal{L}} r(w) a\left(\frac{z}{w}\right) \frac{dw}{w} + \frac{\lambda(1 - \varphi(z))}{1 - z}. \quad (5.1)$$

Methods of solution of (5.1) (e.g. via boundary value problem or Fredholm integral equation) depend heavily on the explicit form of $a(\cdot)$. Alas, tractable explicit expressions can hardly be derived (nor expected!), even when $a(\cdot)$ is a polynomial.

5.2 About transient phenomena

Hereafter, we do not pretend to give exhaustive rigorous proofs of our claims, which can rather be viewed as *very likely true* conjectures, and have been verified by several numerical simulation runs.

Our concern is to extend Theorem 3.3, when there is no solution to (3.15), i.e. when its right-hand side member is finite and α is too large. In fact, it appears that the system has still a stationary regime, but a *finite number of POIs have an infinite score*. This means that, as $t \rightarrow \infty$, the selection policy $\pi(\vec{t})$ becomes *defective*, namely (see equation (3.1)) when

$$\bar{\pi} < 1.$$

This implies exactly

$$\pi_i(\vec{r}) = \frac{a_i r_i}{\sum_{j \geq 1} a_j r_j + \delta},$$

where δ is a positive constant representing the proportion of POIs going to infinity. Obviously, $\delta = 0$ when the system is *normally* ergodic, i.e. no escape of mass to infinity. Then the consistency equation (3.11) must be modified and becomes

$$1 = \frac{\lambda}{\alpha} F\left(\frac{\alpha}{\mu K}\right) + \frac{\delta}{K}, \quad (5.2)$$

which has *always* a solution (K, δ) by monotonicity with respect to K , remembering that $F(1) < \infty$, while $F(1 + \varepsilon) = \infty, \forall \varepsilon > 0$, and $\alpha > \lambda F(1)$. Equation (5.2) is not sufficient to determine the two unknowns K and δ . To get a second equation, we resort to the following heuristic *stochastic least action principle*, which in our opinion should be provable in a very general context.

Definition 5.1. The least action principle. *When a multicomponent irreducible Markovian system ceases to be ergodic, the first component(s) becoming infinite with positive probability) can be identified by continuity with respect to the set of parameters \mathcal{P} defining the system.*

This definition, which might look obscure to the reader, can be rephrased by saying that the phase transition takes place as soon as one touches some regions (hyperplanes or surfaces) in \mathcal{P} . This is for instance the case for the famous Jackson-Kelly stochastic queuing networks...

Applying this principle leads to say that we are looking for the minimal $\delta > 0$ ensuring the system

$$\vec{r} = \lim_{t \rightarrow \infty} \vec{r}(t)$$

is no more ergodic, so that *some POIs go to infinity*. This implies necessarily

$$\frac{\alpha}{\mu K} = 1. \quad (5.3)$$

Then (5.2) and (5.3) yield at once

$$\begin{cases} \delta = \frac{\alpha - \lambda F(1)}{\mu}, \\ \bar{\pi} = \frac{K - \delta}{K} = \frac{\lambda F(1)}{\alpha} < 1, \\ r(1) = \frac{\lambda}{\mu} [F(1) + \varphi'(1)]. \end{cases}$$

A possible way to decide analytically when $\bar{\pi} < 1$ would be to design a converging iterative scheme to solve system (2.2) or (2.4) (see e.g. the polling network analyzed in [1]): this appears to be another interesting, but practically intricate problem.

5.3 The case $\theta'(1) \cdot \varphi'(1) = \infty$

When either $\theta'(1)$ or $\varphi'(1)$ are infinite, we are a priori facing a *transient* phenomenon as $r(1) = \infty$. However, the system can have an interesting behavior provided that $K = \sum_{i \geq 1} a_i r_i < \infty$, which yields necessarily

$$\liminf_{i \rightarrow \infty} a_i = 0,$$

but this would correspond to a *non-herding* system, which is an other story, although most of the mathematical arguments presented in Section 3 could be readily applied.

A Generators, cores and weak convergence

Here we quote the material necessary for the proof of Theorem 2.2. The results are borrowed from [2].

Theorem A.1. *If O is the generator of a strongly continuous semigroup $\{T(t)\}$ on \mathcal{L} , then its domain $\mathcal{D}(O)$ is dense in \mathcal{L} and O is closed.*

Definition A.2. ([2], p. 17) *Let O be a closed linear operator with domain $\mathcal{D}(O)$. A subspace S of $\mathcal{D}(O)$ is said to be a core for O if the closure of the restriction of O to S is equal to O , i.e., if $\overline{O|_S} = O$.*

The next proposition ([2], p. 17), is an important criterion to characterize a core.

Theorem A.3. *Let O be the generator of a strongly continuous contraction semigroup $\{T(t)\}$ on \mathcal{L} . Let \mathcal{D}_0 and \mathcal{D} be dense subspaces of \mathcal{L} with $\mathcal{D}_0 \subset \mathcal{D} \subset \mathcal{D}(O)$. If $T(t) : \mathcal{D}_0 \rightarrow \mathcal{D}$ for all $t \geq 0$, then \mathcal{D} is a core for O .*

The proof of Theorem 2.1 relies heavily on the next general proposition ([2], theorem 6.1, p. 28).

Theorem A.4. *In addition to \mathcal{L} , let $\mathcal{L}_k, k \geq 1$, be a sequence of Banach spaces, $\Pi_k : \mathcal{L} \rightarrow \mathcal{L}_k$ be a bounded linear transformation, subject to the constraint $\sup_k \|\Pi_k\| < \infty$. Let also $\{T_k(t)\}$ and $\{T(t)\}$ be strongly continuous contraction semigroups on \mathcal{L}_k and \mathcal{L} with generators O_k and O . We write $f_k \rightarrow f$ to mean exactly*

$$f \in \mathcal{L}, \quad f_k \in \mathcal{L}_k \quad \text{for } k \geq 1, \quad \text{and} \quad \lim_{k \rightarrow \infty} \|f_k - \Pi_k f\| = 0.$$

Then, if \mathcal{D} is a core for O , the following are equivalent:

- (a) *For each $f \in \mathcal{L}$, $T_k(t)\Pi_k f \rightarrow T(t)f$ for all $t \geq 0$, uniformly on bounded intervals.*
- (b) *For each $f \in \mathcal{L}$, $T_k(t)\Pi_k f \rightarrow T(t)f$ for all $t \geq 0$.*
- (c) *For each $f \in \mathcal{D}$, there exists $f_k \in \mathcal{D}(O_k)$ for each $k \geq 1$, such that $f_k \rightarrow f$ and $O_k f_k \rightarrow O f$.*

References

- [1] F. Delcoigne and G. Fayolle. Thermodynamical limit and propagation of chaos in polling systems. *Markov Processes and Related Fields*, 5(1):89–124, 1999.
- [2] S. N. Ethier and T. G. Kurtz. *Markov processes. Characterization and Convergence*. Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics. John Wiley & Sons Inc., New York, 1986.

- [3] B. A. Fuchs and V. I. Levin. *Functions of a complex variable and some of their applications. Vol. II.* Translated by J. Berry; edited by T. Kövari. Pergamon Press, London, 1961.
- [4] I. S. Gradshteyn and I. M. Ryzhik. *Table of Integrals, Series, and Products.* Academic Press, 7th edition, 2007.
- [5] E. L. Ince. *Ordinary Differential Equations.* Dover Publications, New York, 1956.
- [6] Ma Micro Planète. <http://www.ludigo.net/index.php?rub=3&ssrub=3>, 2011. (in French).
- [7] E. C. Titchmarsh. *The Theory of Functions.* Oxford University Press, 2nd edition, 1939.
- [8] S. Veeraraghavan and L. Debo. Joining longer queues: Information externalities in queue choice. *Manufacturing & Service Operations Management*, 11:543–562, 2009.
- [9] N. D. Vvedenskaya, R. L. Dobrushin, and F. I. Karpelevich. A queueing system with a choice of the shorter of two queues—an asymptotic approach. *Problemy Peredachi Informatsii*, 32(1):20–34, 1996.



**RESEARCH CENTRE
PARIS – ROCQUENCOURT**

Domaine de Voluceau, - Rocquencourt
B.P. 105 - 78153 Le Chesnay Cedex

Publisher
Inria
Domaine de Voluceau - Rocquencourt
BP 105 - 78153 Le Chesnay Cedex
inria.fr

ISSN 0249-6399