



The snapping out Brownian motion

Antoine Lejay

► To cite this version:

| Antoine Lejay. The snapping out Brownian motion. 2012. hal-00781447v2

HAL Id: hal-00781447

<https://inria.hal.science/hal-00781447v2>

Preprint submitted on 6 Jan 2015 (v2), last revised 3 Aug 2015 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The snapping out Brownian motion

Antoine Lejay^{1,2,3,4,5}

January 6, 2015

Abstract

We give a probabilistic representation of a one-dimensional diffusion equation where the solution is discontinuous at 0 with a jump proportional to its flux. This kind of interface condition is usually seen as a semi-permeable barrier. For this, we use a process called here the snapping out Brownian motion, whose properties are studied. As this construction is motivated by applications, for example in brain imaging or in chemistry, a simulation scheme is also provided.

Keywords: Interface condition, elastic Brownian motion, semi-permeable barrier, thin layer, piecing out a Markov process.

1. Introduction

Many diffusion phenomena have to deal with interface conditions. Let D be a diffusivity coefficient which is smooth away from a regular surface S , but presents some discontinuity there. In this case, the solution to the diffusion equation

$$\partial_t u(t, x) = \frac{1}{2} \nabla(D(x) \nabla u(t, x)) = 0 \text{ with } u(0, x) = f(x) \quad (1)$$

¹Université de Lorraine, Institut Élie Cartan de Lorraine, UMR 7502, Vandœuvre-lès-Nancy, F-54500, France

²CNRS, Institut Élie Cartan de Lorraine, UMR 7502, Vandœuvre-lès-Nancy, F-54500, France

³Inria, TOSCA, Villers-lès-Nancy, F-54600, France

⁴Contact: IECL, BP 70238, F-54506 Vandœuvre-lès-Nancy CEDEX, France. Email: Antoine.Lejay@iecn.u-nancy.fr

⁵This work has been supported by the ANR SIMUDMRI (ANR-10-COSI-SIMUDMRI).




Figure 1: The thin layer problem.

has to be understood as a *weak solution*. However, u is smooth away from S and satisfies

$$u(t, x+) = u(t, x-) \text{ and } D(x+)n^+(x) \cdot \nabla u(t, x+) = D(x-)n^-(x) \cdot \nabla u(t, x-), \quad (2)$$

for $x \in S$, when S is assumed to separate locally \mathbb{R}^d into a “+” and a “-” part and where n^\pm is a vector normal to S at x pointing to the “ \pm ” side. The second condition is called the *continuity of the flux*.

Now, let us assume that D takes scalar values, and is constant away from a thin layer of width 2ℓ enclosed between two parallel surfaces S^+ and S^- . When the width ℓ of the layer tends to 0, S^{++} and S^- merge into a single interface located on a surface S .

When the diffusivity D_0 decreases to 0 with ℓ and $D_0/\ell \rightarrow \lambda > 0$, then the solution to (1) converges to a function v satisfying (1) away from S with the interface condition for $x \in S$:

$$\nabla v(t, x+) = \nabla v(t, x-) \text{ and } \frac{\lambda}{2}(v(t, x+) - v(t, x-)) = D(x\pm)\nabla v(t, x\pm). \quad (3)$$

The solution has a continuous flux on S but is discontinuous on S (See for example [32, Chap. 13]). A heuristic explanation is given Figure 1.

If D is smooth on \mathbb{R}^d , it is well known that

$$u(t, x) = \mathbb{E}_x[f(X_t)], \quad (4)$$

where X is the diffusion process generated by $\frac{1}{2}\nabla(D\nabla)$ which is solution under \mathbb{P}_x to the stochastic differential equation (SDE)

$$X_t = x + \int_0^t \sigma(X_s) dB_t + \int_0^t \frac{1}{2} \sum_{i=1}^d \frac{D_{i,\cdot}}{\partial x_i}(X_s) ds \text{ with } \sigma\sigma^T = D \quad (5)$$

for a Brownian motion B .

When D presents some discontinuities, (5) has no longer a meaning. However, a Feller processes $(X, (\mathcal{F}_t)_{t \geq 0}, (\mathbb{P}_x)_{x \in \mathbb{R}})$ is associated to $\frac{1}{2} \nabla(D \nabla \cdot)$ for which (4) holds. In particular, the marginal distributions X_t have a density $p(t, x, \cdot)$ under \mathbb{P}_x , where $p(t, x, y)$ is the fundamental solution to (1) (See [35] for example).

Let us now assume that the dimension of the space is equal to 1 and that D is discontinuous at some separated points $\{x_i\}$ with left and right limit there, and smooth elsewhere. The process X is solution to a SDE with local time. The Itô-Tanaka formula is the key tool to manipulate it, and several simulation algorithms have been proposed (See the references in [25] for example). The process called the *Skew Brownian motion* is the main tool for this construction [23, 24].

Coming back to the thin layer problem, we assume that D is constant and equal to D_1 on $(-\infty, -\ell)$ and (ℓ, ∞) , and to D_0 on $(-\ell, \ell)$. The associated stochastic process is solution to

$$X_t = x + \int_0^t \sqrt{D(X_s)} dB_s + \frac{D_1 - D_0}{D_1 + D_0} L_t^\ell(X) + \frac{D_0 - D_1}{D_1 + D_0} L_t^{-\ell}(X),$$

where $L_t^{\pm\ell}(X)$ is the local time of X at $\pm\ell$ [24].

Letting D_0/ℓ converging to 2κ with $\ell \rightarrow 0$, one may expect that X converges in distribution to a stochastic process Y such that the solution to (1) with the interface condition (3) is given by $v(t, x) = \mathbb{E}_x[f(Y_t)]$.

The article then aims at constructing and giving several properties related to the process Y which we call a *snapping out Brownian motion* (SNOB). This process is Feller on $\mathcal{R} = (-\infty, 0] \cup [0, +\infty)$ but not on \mathbb{R} . The intervals in the definition of \mathcal{R} are disjoint so that 0 corresponds either to $0+$ or $0-$ seen as distinct points.

The behavior of this process is the following: Assume that its starting point is $x \geq 0$. It behaves as a positively reflected Brownian motion until its local time is greater than an independent exponential random variable of parameter 2κ . Then it decides its sign with probability $1/2$ and starts afresh as a new reflected Brownian motion, until its local time is greater than a new exponential random variable, and so on... Using the properties of the exponential random variable, it is equivalent to assert that the particle changes its sign when its local time is greater than an exponential random variable with parameter κ , and behaves like a positively or negatively reflected Brownian motion between these switching times.

Its name is justified by the following fact: As the time at which the particle possibly changes its sign is the same as for the *elastic Brownian motion* [10, 15, 18, 19] (also called the *partially reflected Brownian motion*), it could also be seen as some elastic Brownian motion which is reborn once killed.

The elastic Brownian motion, also called a *partially reflected Brownian motion*, is associated to the Robin boundary condition and has then many applications [8, 15, 34]. This process is the “basic brick” for constructing the SNOB.

The behavior of the SNOB justifies also the old heuristic that the interface condition (3) corresponds to a *semi-permeable barrier*, which arises for example in diffusion Magnetic Resonance Imaging [11] or in chemistry [1, 8]. The interface condition (3) is different from (2), to which is associated a Skew Brownian motion and where the particle crosses the interface when it reaches it, and which corresponds to a *permeable barrier* (See references in [23, 25]).

Here, we work under the condition of a single interface at 0. In short time, it is sufficient to describe the behavior of the process even in a more complex media, since other interface or boundary conditions far enough have “exponentially small” influence on the distribution of the process. This is sufficient for simulation purposes, where particles positions are represented by the stochastic process and move according to its dynamic.

Using similar computations, one may generalize our work to the case where $D(x) = D^+$ if $x \geq 0$, D^- if $x \leq 0$ and an interface condition

$$\nabla u(t, 0+) = \beta \nabla u(t, 0-) \text{ and } \lambda u(t, 0+) - \mu u(t, 0-) = \nabla u(t, 0+)$$

with $\lambda, \mu > 0$. Diffusions on graphs specified by a condition at each vertex could also be considered, which could be of interest in several applications. This process has been described without proof by R. Bobrowski in [6], which have studied its limit behavior when the diffusion coefficients increase.

Although the SNOB may be seen as a diffusion on a graph, it is not a diffusion on a metric graph, where the edges are joint by vertices. Such diffusions have been classified by M. Freidlin and A. Wentzell in [12, 13]. The conditions that are required at the vertices of the graphs are some extension of the possible boundary conditions for a Markov process studied by W. Feller [10]. See also [21] for example for the related problem of pasting diffusions¹.

Our interface condition does not fall in these categories. Our process is best thought as a kind of *random evolution process* which switches back and forth randomly among a collection of processes (See *e.g.* [16, 33]).

Outline. In Sections 2 and 3, we present quickly the main results related to the Elastic Brownian motions and the piecing out procedure. The SNOB is constructed in Section 4 through its resolvent. In Section 5, we show the relationship between the SNOB and the thin layer problem. Finally, in Section 6, we show how to simulate this process.

2. Elastic Brownian motion

Let $(R_t)_{t \geq 0}$ a reflected Brownian motion, and denote by $(L_t)_{t \geq 0}$ its symmetric local time at 0. We add a cemetery point \dagger to \mathbb{R}_+ . For a constant $\kappa > 0$, we consider an

¹The article [31] defines a notion of *semipermeable membrane* which is different from ours, where the solution is continuous with a discontinuous gradient.

exponential random variable ξ independent from B with parameter κ . Set

$$Z_t = \begin{cases} R_t & \text{if } L_t \leq \xi, \\ \dagger & \text{if } L_t > \xi. \end{cases}$$

Thanks to the properties of the local time, this process called the *elastic Brownian motion* (EBM), is still a strong Markov process. It semi-group is

$$P_t^e f(x) = \mathbb{E}_x[\exp(-\kappa L_t) f(X_t)]$$

for f in the set $\mathcal{C}_0(\mathbb{R}_+, \mathbb{R})$ of continuous functions that vanishes at infinity. Closed form expressions of the density transition function are given in [14, 34].

Let \mathfrak{k} be the time at which the EBM is killed, which means $\mathfrak{k} = \inf\{t > 0 \mid L_t \geq \xi\}$. This is a stopping time. Since the local time increases only on the closure of $\mathcal{Z} = \{t > 0 \mid X_t = 0\}$, it holds that $Z_{\mathfrak{k}} = 0$ almost surely. Using standard computations in the inverse of the local time of the Brownian motion,

$$\psi(x, \alpha) = \mathbb{E}_x[\exp(-\alpha \mathfrak{k})] = \frac{\kappa}{\sqrt{2\alpha + \kappa}} \exp(-\sqrt{2\alpha} x). \quad (6)$$

Using the Itô formula, it is easily shown that $u(t, x) = P_t^e f(x)$ is solution to the heat equation with Robin (or third kind) boundary condition [3, 15, 30]

$$\begin{cases} \frac{\partial u(t, x)}{\partial t} = \frac{1}{2} \Delta u(t, x) & \text{on } \mathbb{R}_+^* \times \mathbb{R}_+^* \\ \frac{\partial u(t, 0)}{\partial x} = \kappa u(t, 0). \end{cases}$$

For a Markov process X , let us recall that its resolvent $(G_\alpha)_{\alpha > 0}$ is a family of operators defined by $G_\alpha f(x) = \mathbb{E}_x \left[\int_0^{+\infty} e^{-\alpha s} f(X_s) ds \right]$ for any $f \in \mathcal{C}_0$ and any $\alpha > 0$. It has a density g_α when $G_\alpha f(x) = \int g_\alpha(x, y) f(y) dy$.

Using standard computations on the Green functions, the density $g_\alpha^e(x, y)$ of the resolvent of the EBM is for $x, y \geq 0$,

$$g_\alpha^e(x, y) = \frac{1}{\sqrt{2\alpha}} \begin{cases} \frac{\sqrt{2\alpha - \kappa}}{\sqrt{2\alpha + \kappa}} e^{-\sqrt{2\alpha}(y+x)} + e^{-\sqrt{2\alpha}(x-y)} & \text{for } y \in [0, x], \\ e^{\sqrt{2\alpha}(x-y)} + \frac{\sqrt{2\alpha - \kappa}}{\sqrt{2\alpha + \kappa}} e^{-\sqrt{2\alpha}(x+y)} & \text{for } y \geq x. \end{cases}$$

We extend the EBM to a process on \mathcal{R} by symmetry, so that its resolvent becomes

$$G_\alpha^e f(x) := \mathbb{E}_x \left[\int_0^{\mathfrak{k}} e^{-\alpha s} f(X_s) ds \right] = \int_0^{+\infty} g_\alpha^e(|x|, |y|) f(\text{sgn}(x)y) dy \quad (7)$$

for $x \in \mathcal{R}$. This process evolves either on \mathbb{R}_- or \mathbb{R}_+ but never crosses 0 and could be identified with a process on \mathcal{R} .

3. Piecing out Markov processes

The procedure of piecing out is a way to construct a Markov process from a killed one. We present in this Section a result due to N. Ikeda, M. Nagasawa and S. Watanabe [17] (Similar considerations are given in [28]).

On a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, let $((X_t)_{t \geq 0}, (\mathcal{F}_t)_{t \geq 0}, (\mathbb{P}_x)_{x \in \bar{S}})$ be a right continuous strong Markov process living in state space $\bar{S} = S \cup \{\dagger\}$ with a death point \dagger and a lifetime \mathfrak{k} .

The shift operator associated to X is denoted by $(\theta_t)_{t \geq 0}$.

We also consider a family μ defined on $\Omega \times \bar{S}$ such that $\mu(\omega, \cdot)$ is a probability measure on \bar{S} and for any fixed Borel subset A , $\mu(\cdot, A)$ is $\sigma(X_t, t \geq 0)$ -measurable. We assume additionally that $\mu(\omega, dy) = \delta_{\dagger}(dy)$ when $\mathfrak{k}(\omega) = 0$ and

$$\mathbb{P}_x[\mu(\omega, dy) = \mu(\theta_{\mathfrak{t}(\omega)}\omega, dy), \mathfrak{t}(\omega) < \mathfrak{k}(\omega)] = \mathbb{P}_x[\mathfrak{t} < \mathfrak{k}]$$

for any stopping time \mathfrak{t} . The family μ , called an *instantaneous distribution*, describes the way the process is reborn once killed.

Let $\widehat{\Omega}$ be the product of an infinite, countable, number of copies of $\Omega \times \bar{S}$. We define X on $\widehat{\Omega}$ by

$$X_t(\widehat{\omega}) = \begin{cases} x_t(\omega_1) & \text{if } t \in [0, \mathfrak{k}(\omega_1)), \\ y_1 & \text{if } t = \mathfrak{k}(\omega_1), \\ x_{t-\mathfrak{k}(\omega_1)}(\widetilde{\omega}_2) & \text{if } t \in (\mathfrak{k}(\omega_1), \mathfrak{k}(\omega_1) + \mathfrak{k}(\omega_2)), \\ y_2 & \text{if } t = \mathfrak{k}(\omega_2), \\ \dots & \\ \dagger & \text{if } t \geq \mathfrak{k}(\omega_1) + \dots + \mathfrak{k}_N(\omega_N) \end{cases}$$

with $\widehat{\omega} = (\omega_1, y_1, \omega_2, y_2, \dots) \in \widehat{\Omega}$, and $N = \inf\{k \geq 0 \mid \mathfrak{k}(\omega_k) = 0\}$.

We consider the probability measure

$$\widehat{\mathbb{P}}_x[d\omega_1, dx^1, \dots, d\omega_n, dx^n] = \mathbb{P}_x[d\omega^1]\mu(\omega^1, dx^1)\mathbb{P}_{x^1}[d\omega^2]\mu(\omega^1, dx^2) \cdots \mathbb{P}_{x^n}[d\omega^n]\mu(\omega^n, dx^n).$$

This means that when the path $X(\omega)$ is killed, we let it reborn by placing it at the point x_1 with probability $\mu(\omega, dx_1)$ and then start again...

We left the technical details about the construction of the probability space and the filtration and presents the main result on piecing out Markov process.

Theorem 1 ([17]). *Using the above defined notations, there exists a probability space $(\widehat{\Omega}, \widehat{\mathcal{B}}, \widehat{\mathbb{P}})$ and a filtration $(\widehat{\mathcal{B}}_t)_{t \geq 0}$ on which $(X, (\widehat{\mathcal{B}}_t)_{t \geq 0}, (\widehat{\mathbb{P}}_x)_{x \in S})$ is a strong Markov process on \bar{S} with $\mathbb{P}_{\dagger}[X_t = \dagger, \forall t \geq 0] = 1$.*

4. The snapping out Brownian motion

Definition 1. A *snapping out Brownian motion* (SNOB) X is a strong Markov stochastic process living on \mathcal{R} constructed by making EBM reborn on $0+$ or $0-$ with probability $1/2$ using the piecing-out procedure.

The sign of X changes with probability $1/2$ when its local time L_t at 0 is greater than \mathbf{u}_k with $\mathbf{u}_0 = 0$, $\mathbf{u}_k - \mathbf{u}_{k-1} \sim \exp(\kappa)$ is independent from $(\mathbf{u}_i)_{i \leq k-1}$. From the properties of the exponential and binomial distributions, the sign of X changes when its local time is greater than \mathbf{s}_k with $\mathbf{s}_0 = 0$, $\mathbf{s}_k - \mathbf{s}_{k-1} \sim \exp(\kappa/2)$ is independent from $(\mathbf{s}_i)_{i \leq k-1}$.

It is also immediate that $|X|$ is a Brownian motion, where $|\cdot|$ is the canonical projection of \mathcal{R} onto $[0, +\infty)$.

Proposition 1. *The resolvent family $(G_\alpha)_{\alpha>0}$ of the SNOB is solution to*

$$\left(\alpha - \frac{1}{2}\Delta\right) G_\alpha f(x) = f(x) \text{ for } x \in \mathcal{R}$$

$$\text{with } \nabla G_\alpha f(0+) = \nabla G_\alpha f(0-) \text{ and } \frac{\kappa}{2}(G_\alpha f(0+) - G_\alpha f(0-)) = \nabla G_\alpha f(0),$$

for any bounded, continuous function f on \mathcal{R} that vanishes at infinity.

This proposition identifies the infinitesimal generator of the process X . The points $0+$ and $0-$ could be identified with the sides of a semi-permeable barrier.

Proof. From this very construction and the strong Markov property, for any continuous function f on \mathcal{R} which vanishes at infinity.

$$G_\alpha f(x) = G_\alpha^e f(x) + \frac{\psi(|x|, \alpha)}{2} (G_\alpha f(0+) + G_\alpha f(0-)), \quad (8)$$

where G_α^e is defined by (7).

Using $x = 0+$ and $x = 0-$ in (8) and summing the two equations leads to

$$G_\alpha f(x) = G_\alpha^e f(x) + \frac{\kappa e^{-\sqrt{2\alpha}|x|}}{2\sqrt{2\alpha}} \beta(f) \text{ with } \beta(f) = G_\alpha^e f(0+) + G_\alpha^e f(0-). \quad (9)$$

Then

$$G_\alpha f(x) + G_\alpha f(-x) = G_\alpha^e f(x) + G_\alpha^e f(-x) + \frac{\kappa}{\sqrt{2\alpha}} e^{-\sqrt{2\alpha}|x|} \beta(f), \quad (10)$$

$$G_\alpha f(x) - G_\alpha f(-x) = G_\alpha^e f(x) - G_\alpha^e f(-x). \quad (11)$$

Derivating (10) and setting $x = 0\pm$, since $\nabla G_\alpha^e f(0\pm) = \pm \kappa G_\alpha^e f(0\pm)$,

$$\nabla G_\alpha f(0+) - \nabla G_\alpha f(0-) = 0.$$

Derivating (11),

$$\begin{aligned} 2\nabla G_\alpha f(0\pm) &= \nabla G_\alpha f(0+) + \nabla G_\alpha f(0-) = \nabla G_\alpha^e f(0+) + \nabla G_\alpha^e f(0-) \\ &= \kappa(G_\alpha^e f(0+) - G_\alpha^e f(0-)) = \kappa(G_\alpha f(0+) - G_\alpha f(0-)). \end{aligned}$$

In addition, it is easily seen that $(\alpha - \frac{1}{2}\Delta) G_\alpha f = f$ since $\psi(x, \alpha)$ is solution to $(\alpha - \frac{1}{2}\Delta) \psi(x, \alpha) = 0$. The resolvent is then identified. \square

Proposition 2. *The semi-group $(P_t)_{t \geq 0}$ of the SNOB has the following representation*

$$P_t f(x) = \mathbb{E}_x \left[\left(\frac{1 + e^{-\kappa L_t}}{2} \right) f(\text{sgn}(x)|B_t|) \right] + \mathbb{E}_x \left[\left(\frac{1 - e^{-\kappa L_t}}{2} \right) f(-\text{sgn}(x)|B_t|) \right] \quad (12)$$

for a Brownian motion B .

Proof. Let us decompose a function f as its even and odd parts:

$$\hat{f}(x) = \frac{1}{2}(f(x) + f(-x)) \text{ and } \check{f}(x) = \frac{1}{2}(f(x) - f(-x)).$$

Then $G_\alpha^e \hat{f}(-x) = G_\alpha^e \hat{f}(x)$ and $G_\alpha^e \check{f}(-x) = -G_\alpha^e \check{f}(x)$, so that $\beta(\check{f}) = 0$ for β defined by (9). Thus $G_\alpha \check{f}(x) = G_\alpha^e \check{f}(x)$. In addition, since $\hat{f}(|x|) = \hat{f}(x)$ and the SNOB has the same distribution as the reflected Brownian motion $|B|$,

$$G_\alpha \hat{f}(x) = G_\alpha^r \hat{f}(x) := \mathbb{E}_x \left[\int_0^{+\infty} e^{-\alpha s} \hat{f}(|B_s|) ds \right].$$

This gives an alternative representation for the resolvent of the SNOB: $G_\alpha f(x) = G_\alpha^r \hat{f}(x) + G_\alpha^e \check{f}(x)$. Inverting the resolvent to recover the semi-group $(P_t)_{t \geq 0}$,

$$P_t f(x) = P_t^r \hat{f}(x) + P_t^e \check{f}(x) = \mathbb{E}_x[\hat{f}(|B_t|)] + \mathbb{E}_x[\exp(-\kappa L_t) \check{f}(\text{sgn}(x)|B_t|)].$$

This expression could be arranged as (12). \square

5. The thin layer problem

We now fix $\epsilon > 0$ and we consider the process X^ϵ generated by (See *e.g.* [35] for general considerations on this process)

$$\mathcal{L}^\epsilon = \frac{1}{2} \frac{\partial}{\partial x} \left(a^\epsilon(x) \frac{\partial}{\partial x} \right) \text{ with } a^\epsilon(x) = \begin{cases} 1 & \text{when } x \notin [-\epsilon, \epsilon], \\ \kappa\epsilon & \text{when } x \in [-\epsilon, \epsilon] \end{cases}$$

whose domain is the closure for the operator norm of

$$\left\{ f \in \mathcal{C}(\mathbb{R}) \cap \mathcal{C}^2((-\infty, -\epsilon) \cup (\epsilon, \epsilon) \cup (\epsilon, \infty)) \left| \begin{array}{l} f(\pm\epsilon-) = f(\pm\epsilon+) \\ \nabla f(-\epsilon-) = \kappa\epsilon \nabla f(-\epsilon) \\ \kappa\epsilon \nabla f(\epsilon-) = \nabla f(\epsilon) \end{array} \right. \right\}.$$

This operator is self-adjoint, so that its resolvent density satisfies $g_\alpha^\epsilon(x, y) = g_\alpha^\epsilon(y, x)$. This process is a Feller process, and is a strong solution to the SDE with local time

$$X_t^\epsilon = x + \int_0^t \sqrt{a^\epsilon(X_s^\epsilon)} dB_s + \eta_\epsilon L_t^\epsilon(X^\epsilon) - \eta_\epsilon L_t^{-\epsilon}(X^\epsilon) \text{ with } \eta_\epsilon = \frac{1 - \kappa\epsilon}{1 + \kappa\epsilon},$$

where B is a Brownian motion and $L_t^x(X^\epsilon)$ is the symmetric local time at x of X^ϵ (See *e.g.* [24], and [4, 22] among others for general results on SDEs with local time).

In [10, § 11], the elastic Brownian motion is constructed as the limit of a process which either jumps at ϵ or is killed with probability $\kappa\epsilon$ when it arrives at 0.

Using the piecing out procedure, we construct a strong Markov process Z^ϵ by considering the process X^ϵ which is instantaneously replaced at $-\epsilon$ or ϵ with probability 1/2 when it reaches 0, and then behaving again as X^ϵ until it reaches 0, and so on. This process Z^ϵ could be identified as a process living in \mathcal{R} by defining \mathbb{P}_{0+} as \mathbb{P}_ϵ and \mathbb{P}_{0-} as $\mathbb{P}_{-\epsilon}$, since the process is instantaneously killed when at 0.

Theorem 2. *The process Z^ϵ with $Z_0^\epsilon = x$ converges in distribution to the SNOB starting from x in the Skorohod topology.*

The proof relies on the next two results.

Proposition 3. *Let g_α^ϵ be the resolvent density of X^ϵ . Then $g_\alpha^\epsilon(x, y)$ converges to $g(x, y)$ for any $x, y \neq 0$ and any $\alpha > 0$ as $\epsilon \rightarrow 0$.*

Remark 1. This result follows from classical results in deterministic homogenization theory (See [32] for example) where the convergence holds in Sobolev spaces. Here, we consider a direct computational proof for the convergence of the Green kernel, which we use later.

Proof. We assume that $x > 0$ and we set $\mu := \sqrt{2\alpha}$ for some $\alpha > 0$. The resolvent density g_α^ϵ of X^ϵ has the form, for $x > \epsilon$,

$$g_\alpha^\epsilon(x, y) = \begin{cases} C_\epsilon(x)e^{-\mu y} & \text{for } y > x \\ A_\epsilon(x)e^{-\mu y} + B_\epsilon(x)e^{\mu y} & \text{for } y \in [\epsilon, x], \\ H_\epsilon(x)e^{\mu y/\sqrt{\kappa\epsilon}} + E_\epsilon(x)e^{-\mu y/\sqrt{\kappa\epsilon}} & \text{for } y \in [-\epsilon, \epsilon], \\ F_\epsilon(x)e^{\mu y} & \text{for } y < -\epsilon. \end{cases}$$

By this, we mean that

$$\mathbb{E}_x \left[\int_0^{+\infty} e^{-\alpha t} f(X_s^\epsilon) ds \right] = \int_{\mathbb{R}} g_\alpha^\epsilon(x, y) f(y) dy$$

for any bounded, measurable function f .

With the conditions

$$\begin{aligned} g_\alpha^\epsilon(x, \epsilon+) &= g_\alpha^\epsilon(x, \epsilon-), \quad g_\alpha^\epsilon(x, \epsilon-) = g_\alpha^\epsilon(x, \epsilon+), \\ \nabla_y g_\alpha^\epsilon(x, -\epsilon-) &= \kappa \epsilon \nabla_y g_\alpha^\epsilon(x, -\epsilon+), \quad \kappa \epsilon \nabla_y g_\alpha^\epsilon(x, \epsilon-) = \nabla_y g_\alpha^\epsilon(x, \epsilon+), \\ \nabla_y g_\alpha^\epsilon(x, x+) - \nabla_y g_\alpha^\epsilon(x, x-) &= 2, \end{aligned}$$

With $\mu = \sqrt{2\alpha}$, the coefficients A_ϵ , B_ϵ , C_ϵ , H_ϵ and F_ϵ are expressed as

$$G_\epsilon := \left(2e^{4\frac{\mu\epsilon}{\sqrt{\kappa\epsilon}}} \sqrt{\kappa\epsilon} + e^{4\frac{\mu\epsilon}{\sqrt{\kappa\epsilon}}} \kappa\epsilon + 2\sqrt{\kappa\epsilon} - \kappa\epsilon + e^{4\frac{\mu\epsilon}{\sqrt{\kappa\epsilon}}} - 1 \right) \mu.$$

Since $\epsilon \rightarrow 0$, $G_\epsilon = 4\sqrt{\kappa\epsilon}(1 + \frac{\mu}{\kappa} + O(\kappa\epsilon))\mu$. After tedious computations,

$$\begin{aligned} A_\epsilon(x) &= - \left(e^{4\frac{\mu\epsilon}{\sqrt{\kappa\epsilon}}} \kappa\epsilon - \kappa\epsilon - e^{4\frac{\mu\epsilon}{\sqrt{\kappa\epsilon}}} + 1 \right) e^{\mu(2\epsilon-x)} / G_\epsilon \xrightarrow{\epsilon \rightarrow 0} A_0(x) := -\frac{e^{-\mu x}}{\kappa + \mu}, \\ B_\epsilon(x) &= B_0(x) := -\frac{e^{-\mu x}}{\mu}, \\ C_\epsilon(x) &= -2 \sinh(2\epsilon/\sqrt{\kappa\epsilon}) \left(de^{-\mu x+2\mu\epsilon} - e^{-\mu x+2\mu\epsilon} + de^{\mu x} + e^{-\mu x} \right) e^{\frac{2\mu\epsilon}{\sqrt{\kappa\epsilon}}} / G_\epsilon, \\ &\quad + 4\sqrt{\kappa\epsilon} e^{\mu x} \cosh(2\epsilon/\sqrt{\kappa\epsilon}) e^{\frac{2\mu\epsilon}{\sqrt{\kappa\epsilon}}} / G_\epsilon \xrightarrow{\epsilon \rightarrow 0} C_0(x) := \frac{\kappa e^{\mu x}}{\mu(\kappa + \mu)}, \\ H_\epsilon(x) &= -2e^{\frac{\mu(3\epsilon+\epsilon\sqrt{\kappa\epsilon}-x\sqrt{\kappa\epsilon})}{\sqrt{\kappa\epsilon}}} (1 + \sqrt{\kappa\epsilon}) \sqrt{\kappa\epsilon} / G_\epsilon \xrightarrow{\epsilon \rightarrow 0} H_0(x) := -\frac{\kappa e^{-\mu x}}{2\mu(\kappa + \mu)}, \\ E_\epsilon(x) &= -2e^{\frac{\mu(\epsilon+\epsilon\sqrt{\kappa\epsilon}-x\sqrt{\kappa\epsilon})}{\sqrt{\kappa\epsilon}}} (1 - \sqrt{\kappa\epsilon}) \sqrt{\kappa\epsilon} / G_\epsilon \xrightarrow{\epsilon \rightarrow 0} H_0(x), \\ F_\epsilon(x) &= -4\sqrt{\kappa\epsilon} e^{\frac{\mu(2\epsilon+2\epsilon\sqrt{\kappa\epsilon}-x\sqrt{\kappa\epsilon})}{\sqrt{\kappa\epsilon}}} / G_\epsilon \xrightarrow{\epsilon \rightarrow 0} F_0(x) := \frac{-\kappa e^{-\mu x}}{\mu(\kappa + \mu)} = -C_0(-x). \end{aligned}$$

Let g_α be the function

$$g_\alpha(x, y) := \begin{cases} C_0(x)e^{-\mu y} & \text{if } y > x, \\ A_0(x)e^{-\mu y} + B_0(x)e^{\mu y} & \text{if } y \in [0, x], \\ F_0(x)e^{\mu y} & \text{if } y < 0. \end{cases}$$

A similar work may be performed for $x < 0$. Thus, we easily obtain that $g_\alpha^\epsilon(x, y) \xrightarrow{\epsilon \rightarrow 0} g_\alpha(x, y)$ converges to g_α and that g_α is the density resolvent of the SNOB by checking it satisfies the appropriate conditions at the interface. \square

Proposition 4. Let \mathfrak{h}_0^ϵ be the first hitting time of 0 for X^ϵ .

Under \mathbb{P}_x , \mathfrak{h}_0^ϵ converges in distribution to a random variable \mathfrak{k} distributed as the lifetime of the EBM of parameter κ .

Proof. As in [9, 24], we introduce $\Phi^\epsilon(x)$ as the piecewise linear function defined by

$$\frac{d\Phi^\epsilon}{dx}(x) = \begin{cases} 1/\sqrt{\kappa\epsilon} & \text{if } x \in [-\epsilon, \epsilon], \\ 1 & \text{otherwise.} \end{cases}$$

Set $Y^\epsilon = \Phi^\epsilon(X^\epsilon)$ so that Y^ϵ is solution to the SDE [9, 24]

$$Y_t^\epsilon = \Phi^\epsilon(x) + B_t + \theta^\epsilon L_t^{y^\epsilon}(Y^\epsilon) - \theta^\epsilon L_t^{-y^\epsilon}(Y^\epsilon) \text{ with } \theta^\epsilon = \frac{1 - \sqrt{\kappa\epsilon}}{1 + \sqrt{\kappa\epsilon}} \text{ and } y^\epsilon := \Phi^\epsilon(\epsilon) = \sqrt{\epsilon/\kappa}.$$

The infinitesimal generator of Y^ϵ is $\frac{1}{2}\Delta$ with a domain which is the closure of $f \in \mathcal{C}_b^2(\mathbb{R} \setminus \{-y^\epsilon, y^\epsilon\})$ with $f(\pm y^\epsilon +) = f(\pm y^\epsilon -)$, $(1 - \theta^\epsilon)f'(y^\epsilon -) = (1 + \theta^\epsilon)f'(y^\epsilon +)$ and $(1 + \theta^\epsilon)f'(-y^\epsilon -) = (1 - \theta^\epsilon)f'(-y^\epsilon +)$. From now, we assume that $x > 0$.

The hitting time \mathfrak{h}_0^ϵ is also the first hitting time of zero by Y^ϵ . Since by symmetry $\psi(-x, \alpha) = \psi(x, \alpha)$ for any $x \geq 0$, we consider only that $x \geq 0$.

Since the Feynman-Kac formula is valid for the process Y^ϵ , $\psi^\epsilon(x, \alpha) := \mathbb{E}_x[e^{-\alpha\mathfrak{h}_0^\epsilon}]$ is solution to

$$\begin{cases} \frac{1}{2}\Delta\psi^\epsilon(x, \alpha) = \alpha\psi^\epsilon(x, \alpha) \text{ for } x \neq y^\epsilon, \\ \psi^\epsilon(0, \alpha) = 1, \\ \psi^\epsilon(y^\epsilon -, \alpha) = \psi^\epsilon(y^\epsilon +, \alpha), \\ (1 - \theta^\epsilon)\nabla_x\psi^\epsilon(y^\epsilon -, \alpha) = (1 + \theta^\epsilon)\nabla_x\psi^\epsilon(y^\epsilon +, \alpha), \end{cases}$$

Hence, $\psi^\epsilon(x, \alpha)$ is sought as

$$\psi^\epsilon(x, \alpha) = \begin{cases} \gamma^\epsilon \exp(-\sqrt{2\alpha}x) & \text{if } x > y^\epsilon, \\ \cos(\sqrt{2\alpha}x) + \beta^\epsilon \sin(\sqrt{2\alpha}x) & \text{if } x \in [0, y^\epsilon]. \end{cases}$$

After some computations,

$$\beta^\epsilon = \frac{-\cos(\sqrt{2\alpha}y^\epsilon) + \sqrt{\kappa\epsilon}\sin(\sqrt{2\alpha}y^\epsilon)}{\sin(\sqrt{2\alpha}y^\epsilon) + \sqrt{\kappa\epsilon}\cos(\sqrt{2\alpha}y^\epsilon)} \text{ and } \sqrt{\epsilon}\beta^\epsilon \sim_{\epsilon \rightarrow 0} \frac{-\sqrt{\kappa}}{\kappa + \sqrt{2\alpha}}.$$

Besides,

$$\gamma^\epsilon = e^{\sqrt{2\alpha}y^\epsilon} \sqrt{\kappa\epsilon} (\beta^\epsilon \cos(\sqrt{2\alpha}y^\epsilon) - \beta^\epsilon \sin(\sqrt{2\alpha}y^\epsilon)) \sim_{\epsilon \rightarrow 0} \frac{\kappa}{\kappa + \sqrt{2\alpha}}.$$

Hence, for any $x > 0$,

$$\psi^\epsilon(x, \alpha) \xrightarrow{\epsilon \rightarrow 0} \psi(x, \alpha) := \frac{\kappa}{\kappa + \sqrt{2\alpha}} e^{-\sqrt{2\alpha}x} \quad (13)$$

with ψ defined by (6).

This proves that under \mathbb{P}_x , \mathfrak{h}_0^ϵ converges to a random variable \mathfrak{k} whose Laplace transform is $\psi(x, \alpha)$ under \mathbb{P}_x . This random variable \mathfrak{k} is then the lifetime of an EBM. \square

Proof of Theorem 2. Using the properties of the resolvent, for $\alpha > 0$ and a bounded, measurable function f ,

$$G_\alpha^\epsilon f(x) := \mathbb{E}_x \left[\int_0^{+\infty} e^{-\alpha t} f(X_s^\epsilon) ds \right] = R_\alpha^\epsilon f(x) + \mathbb{E}_x[e^{-\alpha \mathfrak{h}_0^\epsilon}] \frac{1}{2} (G_\alpha^\epsilon f(\epsilon) + G_\alpha^\epsilon f(-\epsilon)).$$

with

$$R_\alpha^\epsilon f(x) := \mathbb{E}_x \left[\int_0^{\mathfrak{h}_0^\epsilon} e^{-\alpha t} f(X_s^\epsilon) ds \right].$$

Since $\psi^\epsilon(x, \alpha) = \psi^\epsilon(-x, \alpha)$,

$$G_\alpha^\epsilon f(x) = R_\epsilon^\alpha f(x) + \frac{\psi^\epsilon(x, \alpha)}{1 - \psi^\epsilon(\epsilon, \alpha)} \frac{R_\epsilon^\alpha f(\epsilon) + R_\epsilon^\alpha f(-\epsilon)}{2}.$$

For the sake of simplicity, we assume that $x > 0$. Using the symmetry properties of \mathcal{L}^ϵ ,

$$R_\alpha^\epsilon f(x) = \int_0^{+\infty} (g_\alpha^\epsilon(x, y) - g_\alpha^\epsilon(x, -y)) f(y) dy.$$

But

$$g_\alpha^\epsilon(x, y) - g_\alpha^\epsilon(x, -y) \xrightarrow{\epsilon \rightarrow 0} g_\alpha(x, y) - g_\alpha(x, -y) = g_\alpha^e(x, y),$$

where $g_\alpha^e(x, y)$ is the resolvent density of the EBM. Thus, $R_\epsilon^\alpha f(x) \xrightarrow{\epsilon \rightarrow 0} G_\alpha^\epsilon f(x)$ for any $x > 0$. It is also easily obtained that

$$R_\epsilon^\alpha f(\epsilon) \xrightarrow{\epsilon \rightarrow 0} G_\alpha^\epsilon f(0+) \text{ and } R_\epsilon^\alpha f(-\epsilon) \xrightarrow{\epsilon \rightarrow 0} G_\alpha^\epsilon f(0-).$$

Using (9) and (13), $G_\alpha^\epsilon f(x) \xrightarrow{\epsilon \rightarrow 0} G_\alpha f(x)$. The Trotter-Kato theorem (See *e.g.* [20, Theorem IX.2.16, p. 504]) and the Markov property imply the convergence in finite-dimensional distributions of Z^ϵ to X under \mathbb{P}_x for $x \geq 0$. By symmetry, this could be extended to $x \leq 0$.

The only remaining point of the tightness. When away from $[-\epsilon, \epsilon]$, X^ϵ behaves like a Brownian motion. Hence, for $0 \leq s \leq t \leq T$, let us set $\mathfrak{f}(s, t) = \inf\{u >$

$s||X_u^\epsilon| = \epsilon\}$ with possibly $\mathfrak{f}(s, t) = +\infty$ and $\mathfrak{l}(s, t) = \sup\{u < t ||X_u^\epsilon| = \epsilon\}$ with possibly $\mathfrak{l}(s, t) = -\infty$.

If $\mathfrak{f}(s, t) \geq t$ and $\mathfrak{l}(s, t) \leq s$, then for $\delta < 1/2$, there exists an integrable random variable $C(\omega)$ such that $|X_t^\epsilon(\omega) - X_s^\epsilon(\omega)| \leq C(\omega)(t - s)^\delta$ for any $0 \leq s \leq t \leq T$.

If $\mathfrak{f}(s, t) \leq t$ and $\mathfrak{l}(s, t) \leq s$, then

$$|X_t^\epsilon - X_s^\epsilon| \leq |X_{\mathfrak{f}(s,t)}^\epsilon - X_s^\epsilon| + |X_t^\epsilon - X_{\mathfrak{f}(s,t)}^\epsilon| \leq C(t - s)^\beta + 2\epsilon$$

since X_t^ϵ belongs to $[-\epsilon, \epsilon]$. A similar analysis could be carried for the other cases, which means that for some integrable random variable C ,

$$\sup_{|t-s|<\delta} |X_t^\epsilon - X_s^\epsilon| \leq C\delta^\beta + 2\epsilon.$$

This proves that $(Z^\epsilon)_{\epsilon>0}$ is tight in the space $\mathcal{D}([0, T]; \mathbb{R})$ of discontinuous functions with the Skorohod topology (See *e.g.* [5]) and then on $\mathcal{D}([0, T]; \mathcal{R})$. Hence, we easily deduce the convergence of Z^ϵ to the SNOB in $\mathcal{D}([0, T]; \mathcal{R})$. \square

6. Simulation of the SNOB

It is easy to simulate a discretized process X in the same way it is easy to simulate the Brownian motion. Following Proposition 2, we draw a random variate with density $p(\Delta t, x, \cdot)$ when x is close enough to 0.

For this, we use a Brownian bridge technique to check if the process reaches $0 \pm$ before Δt (see for example [2] and [25, Sect. B.2] for an example of application and further references) and we then compute the local time using the following representation [26, 27]:

$$(L_t, |B_t|) \stackrel{\text{dist}}{=} (\mathfrak{l}, \mathfrak{l} - H) \text{ where } \mathfrak{l} := \frac{1}{2}(H + \sqrt{V + H^2})$$

with $H \sim \mathcal{N}(0, t)$ and $V \sim \exp(1/2t)$ independent from H .

The density of the inverse Gaussian distribution $\mathcal{IG}(\lambda, \mu)$ is $r_{\mu, \lambda}(x) = \sqrt{\frac{\lambda}{2\pi x^3}} \exp\left(\frac{-\lambda(x-\mu)^2}{2\mu^2 x}\right)$. Random variates with \mathcal{IG} distribution could be simulated by the methods proposed in [7, p. 148] and [29].

The generic algorithm to simulate the process at time Δt when at point x at time 0 is the following:

1. Set $y = x + \sqrt{\Delta t}G$ with G a random variate whose distribution is $\mathcal{N}(0, 1)$.
2. If $|x| \geq 4\sqrt{\Delta t}$, then return y (here, we neglect the exponentially small probability that the process crosses 0 between the times 0 and Δ).
3. If $yx > 0$, then decide with probability $\exp(-2|xy|/\Delta t)$ if the path X has crossed 0.
 - If no crossing occurs, then return y .

- If a crossing occurs, draw $\mathfrak{g} \sim \mathcal{IG}(|x|/|y|, x^2/2\Delta t)$, so that $\mathfrak{z} = \Delta t\mathfrak{g}/(1 + \mathfrak{g})$ is a realization of the first hitting time of 0 for a Brownian bridge with $B_0 = x$ and $B_{\Delta t} = y$. Then go the step 5.
- 4. If $xy < 0$, then draw $\mathfrak{g} \sim \mathcal{IG}(-|x|/|y|, x^2/2\Delta t)$ and set $\mathfrak{z} = \Delta t\mathfrak{g}/(1 + \mathfrak{g})$, the first time the Brownian bridges reaches 0. Go to step 5.
- 5. Set $\mathfrak{r} = \Delta t - \mathfrak{z}$. For three independent random variates $U \sim \mathcal{U}(0, 1)$, $H \sim \mathcal{N}(0, \mathfrak{r})$ and $V \sim \exp(1/2\mathfrak{r})$, set $\mathfrak{l} = (H + \sqrt{V + H^2})/2$.
- 6. Set $\mathfrak{s} := \text{sgn}(x)$ if $\exp(-\kappa\mathfrak{l}) \geq 2U - 1$. Otherwise, set $\mathfrak{s} := -\text{sgn}(x)$.
- 7. Return $\mathfrak{s}(\mathfrak{l} - H)$.

Acknowledgement. The author is indebted to Jing-Rebecca Li and Denis Grebenkov for having proposed this research and interesting discussions about it. The author also wishes to thank warmly Claire Nivlet for having suggested the name of the process.

References

- [1] S. S. Andrews, *Accurate particle-based simulation of adsorption, desorption and partial transmission*, Phys. Biol. **6**, 046015.
- [2] P. Baldi, *Exact asymptotics for the probability of exit from a domain and applications to simulation*, Ann. Probab. **23** (1995), no. 4, 1644–1670.
- [3] Richard F. Bass, Krzysztof Burdzy, and Zhen-Qing Chen, *On the Robin problem in fractal domains*, Proc. Lond. Math. Soc. (3) **96** (2008), no. 2, 273–311, DOI 10.1112/plms/pdm045. MR2396121 (2008m:60158)
- [4] R. F. Bass and Z.-Q. Chen, *Brownian motion with singular drift*, Ann. Probab. **31** (2003), no. 2, 791–817, DOI 10.1214/aop/1048516536. MR1964949 (2004b:60144)
- [5] P. Billingsley, *Convergence of probability measures*, 2nd ed., Wiley Series in Probability and Statistics: Probability and Statistics, John Wiley & Sons Inc., New York, 1999.
- [6] A. Bobrowski, *From Diffusions on Graphs to Markov Chains via Asymptotic State Lumping*, Annales Henri Poincaré, posted on 2012, DOI 10.1007/s00023-012-0158-z.
- [7] L. Devroye, *Non-Uniform Random Variate Generation*, Springer-Verlag, 1986.
- [8] R. Erban and S.J. Chapman, *Reactive boundary conditons for stochastic simulation of reaction-diffusion processes*, Phys. Biol. **4** (2007), 16–28.
- [9] P. Étoré, *On random walk simulation of one-dimensional diffusion processes with discontinuous coefficients*, Electron. J. Probab. **11** (2006), no. 9, 249–275.
- [10] W. Feller, *Diffusion processes in one dimension*, Trans. Amer. Math. Soc. **77** (1954), 1–31.
- [11] E. Fieremans, D. S. Novikov, J. H. Jensen, and J. A. Helpert, *Monte Carlo study of a two-compartment exchange model of diffusion*, NMR in Biomedicine **23** (2010), 711–724, DOI 10.1002/nbm.1577.
- [12] M. I. Freidlin and A. D. Wentzell, *Random perturbations of Hamiltonian systems*, Mem. Amer. Math. Soc. **109** (1994), no. 523.
- [13] ———, *Diffusion processes on graphs and the averaging principle*, Ann. Probab. **21** (1993), no. 4, 2215–2245.

- [14] G. Gallavotti and H. P. McKean, *Boundary conditions for the heat equation in a several-dimensional region*, Nagoya Math. J. **47** (1972), 1–14.
- [15] D. Grebenkov, *Partially Reflected Brownian Motion: A Stochastic Approach to Transport Phenomena*, Focus on Probability Theory (L. R. Velle, ed.), Nova Science Publishers, 2006, pp. 135–169, available at [arXiv:math/0610080v1](https://arxiv.org/abs/math/0610080v1).
- [16] R. J. Griego and A. Moncayo, *Random evolutions and piecing out of Markov processes*, Bol. Soc. Mat. Mexicana (2) **15** (1970), 22–29.
- [17] N. Ikeda, M. Nagasawa, and S. Watanabe, *A construction of Markov processes by piecing out*, Proc. Japan Acad. **42** (1966), 370–375.
- [18] K. Itô and H.P. McKean, *Diffusion processes and their sample paths*, 2nd ed., Springer-Verlag, 1974.
- [19] S. Karlin and H. M. Taylor, *A second course in stochastic processes*, Academic Press Inc. [Harcourt Brace Jovanovich Publishers], New York, 1981.
- [20] T. Kato, *Perturbation theory for linear operators*, 2nd ed., Springer-Verlag, Berlin, 1976.
- [21] B. Kopytko and M. Portento, *The problem of pasting together two diffusion processes and classical potentials*, Theor. Stoch. Proc. **15** (2009), no. 31, 126–139.
- [22] J.-F. Le Gall, *One-dimensional stochastic differential equations involving the local times of the unknown process*, Stochastic analysis and applications (Swansea, 1983), Lecture Notes in Math., vol. 1095, Springer, Berlin, 1984, pp. 51–82, DOI 10.1007/BFb0099122.
- [23] A. Lejay, *On the constructions of the Skew Brownian motion*, Probab. Surv. **3** (2006), 413–466.
- [24] A. Lejay and M. Martinez, *A scheme for simulating one-dimensional diffusion processes with discontinuous coefficients*, Ann. Appl. Probab. **16** (2006), no. 1, 107–139, DOI 10.1214/105051605000000656.
- [25] A. Lejay and G. Pichot, *Simulating diffusion processes in discontinuous media: a numerical scheme with constant time steps*, J. Comput. Phys. **231** (2012), no. 21, 7299–7314, DOI 10.1016/j.jcp.2012.07.011.
- [26] D. Lépine, *Un schéma d'Euler pour équations différentielles stochastiques réfléchies*, C. R. Acad. Sci. Paris Sér. I Math. **316** (1993), no. 6, 601–605.
- [27] ———, *Euler scheme for reflected stochastic differential equations*, Math. Comput. Simulation **38** (1995), no. 1-3, 119–126. Probabilités numériques (Paris, 1992).
- [28] P. A. Meyer, *Renaissance, recollements, mélanges, ralentissement de processus de Markov*, Ann. Inst. Fourier (Grenoble) **25** (1975), no. 3-4.
- [29] J.R. Michael, W.R. Shucany, and R.W. Haas, *Generating random variates using transformations with multiple roots*, American Statistician **30** (1976), no. 2, 88–90.
- [30] V. G. Papanicolaou, *The probabilistic solution of the third boundary value problem for second order elliptic equations*, Probab. Theory Related Fields **87** (1990), no. 1, 27–77, DOI 10.1007/BF01217746.
- [31] N. I. Portenko, *A probabilistic representation for the solution to one problem of mathematical physics*, Ukraïn. Mat. Zh. **52** (2000), no. 9, 1272–1282, DOI 10.1023/A:1010388321016; English transl., Ukrainian Math. J. **52** (2000), no. 9, 1457–1469 (2001).

- [32] E. Sánchez-Palencia, *Non-Homogeneous Media and Vibration Theory*, Lecture Notes in Phys., vol. 127, Springer, Berlin, 1980.
- [33] K. Siegrist, *Random evolution processes with feedback*, Trans. Amer. Math. Soc. **265** (1981), no. 2, 375–392, DOI 10.2307/1999740.
- [34] A. Singer, Z. Schuss, A. Osipov, and D. Holcam, *Partially reflected diffusions*, SIAM J. Appl. Math. **68** (2007/08), no. 3, 844–868, DOI 10.1137/060663258.
- [35] D. W. Stroock, *Diffusion semigroups corresponding to uniformly elliptic divergence form operators*, Séminaire de Probabilités, XXII, Lecture Notes in Math., vol. 1321, Springer, Berlin, 1988, pp. 316–347.