



HAL
open science

External stimulation induces switches between neural oscillations: an illustrative feedback model

Axel Hutt

► **To cite this version:**

Axel Hutt. External stimulation induces switches between neural oscillations: an illustrative feedback model. BMC Neuroscience, 2011, 12 (Suppl 1), pp.P156. 10.1186/1471-2202-12-S1-P156. hal-00784420

HAL Id: hal-00784420

<https://inria.hal.science/hal-00784420>

Submitted on 4 Feb 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

POSTER PRESENTATION

Open Access

External stimulation induces switches between neural oscillations: an illustrative feedback model

Axel Hutt

From Twentieth Annual Computational Neuroscience Meeting: CNS*2011
Stockholm, Sweden. 23-28 July 2011

Oscillatory activity in neural systems plays an important role in information processing. For instance, specific cognitive tasks evoke oscillations in certain frequency bands, which are characteristic for the corresponding experimental paradigm. Examples are the theta-rhythm (4-8Hz) in memory tasks or the gamma-rhythm (30-60Hz) in visual binding tasks. Moreover, changing the external activation of a neural system may lead to the activation of additional oscillations in specific frequency bands or may even yield a switch between specific frequency bands. In the last years, such a switch has been found experimentally in-vivo in some neural systems, such as in the mammalian olfactory bulb [1] and in the primary sensory area of weakly electric fish [2]. Both systems exhibit a feedback topology between two or more interacting populations and an external stimulation that originates from another brain area.

More detailed, the mammalian olfactory bulb is subject to the activation by the receptor cells, which are rhythmically activated by odours due to the breathing cycle. Applying an odour, the Local Field Potentials (LFP) exhibit high gamma-activity in the inhalation phase and switches to high beta-activity in the exhalation phase. In weakly electric fish, the primary sensory area ELL receives input from electro-receptors on the skin of the fish. In the case of spatially uncorrelated noisy stimulation of the skin receptors, single neurons fire mainly in the beta-band, while a strong spatial input correlation switches on an additional spiking rhythm in the gamma-band.

To explain such switches in frequency bands in feedback systems, the current work presents a neural population rate model of a rather general feedback system [3], that allows to derive mathematical conditions for the switch between frequency bands subject to the

spatial correlation of an external input. The proposed model explains the switch between frequency bands as a selective activation of linear spatial oscillation modes. The centre frequency and the width of the bands and the properties of the spatial oscillation modes to be activated result from the intrinsic properties of the feedback system. Such properties are the synaptic time constants and delays between system components. Moreover, the spatial interactions in the feedback system determines the spatial frequency of the oscillation modes. The essential selection of the frequency band happens via the coupling of the spatially extended external input with the spatial modes. For illustration reasons, one may think of a guitar string, whose oscillation frequency depends on the manner how it is stricken.

Published: 18 July 2011

References

1. Buonviso N, Amat C, Litaudon P, Roux S, Royet JP, Farget V, Sicard G: Rhythm sequence through the olfactory bulb layers during the time window of a respiratory cycle. *Europ. J. Neurosci* 2003, **17**:1811-1819.
2. Doiron B, Lindner B, Longtin A, Maler L, Bastian J: Oscillatory activity in electrosensory neurons increases with the spatial correlation of the stochastic input stimulus. *Phys.Rev.Lett* 2004, **93**(4):048101.
3. Hutt A, Sutherland C, Longtin A: Driving neural oscillations with correlated spatial input and topographic feedback. *Phys. Rev. E* 2008, **78**:021911.

doi:10.1186/1471-2202-12-S1-P156

Cite this article as: Hutt: External stimulation induces switches between neural oscillations: an illustrative feedback model. *BMC Neuroscience* 2011 **12**(Suppl 1):P156.

Correspondence: axel.hutt@inria.fr
Equipe Cortex, INRIA Nancy – Grand Est, Villers-les-Nancy, France