



HAL
open science

Irreversible computable functions

Mathieu Hoyrup

► **To cite this version:**

Mathieu Hoyrup. Irreversible computable functions. STACS - 31st Symposium on Theoretical Aspects of Computer Science - 2014, Mar 2014, Lyon, France. hal-00915952v1

HAL Id: hal-00915952

<https://inria.hal.science/hal-00915952v1>

Submitted on 9 Dec 2013 (v1), last revised 14 Jan 2014 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Irreversible computable functions

Mathieu Hoyrup¹

1 Inria, LORIA
615 ru du jardin botanique 54600 Villers-lès-Nancy, France
mathieu.hoyrup@inria.fr

Abstract

The strong relationship between topology and computations has played a central role in the development of several branches of theoretical computer science: foundations of functional programming, computational geometry, computability theory, computable analysis. Often it happens that a given function is not computable simply because it is not continuous. In many cases, the function can moreover be proved to be non-computable in the stronger sense that it does not preserve computability: it maps a computable input to a non-computable output. To date, there is no connection between topology and this kind of non-computability, apart from Pour-El and Richards “First Main Theorem”, applicable to linear operators on Banach spaces only.

In the present paper, we establish such a connection. We identify the discontinuity notion, for the inverse of a computable function, that implies non-preservation of computability. Our result is applicable to a wide range of functions, it unifies many existing *ad hoc* constructions explaining at the same time what makes these constructions possible in particular contexts, sheds light on the relationship between topology and computability and most importantly allows us to solve open problems. In particular it enables us to answer the following open question in the negative: if the sum of two shift-invariant ergodic measures is computable, must these measures be computable as well? We also investigate how generic a point with computable image can be. To this end we introduce a notion of genericity of a point w.r.t. a function, which enables us to unify several finite injury constructions from computability theory.

1998 ACM Subject Classification F.1.1 Models of Computation/Computability theory, F.4.1 Mathematical Logic/Computability theory

Keywords and phrases Computability theory, computable analysis, finite injury, generic set

Digital Object Identifier 10.4230/LIPIcs.xxx.yyy.p

1 Introduction

Many problems in classical computability theory [15] and computable analysis [14, 18] amount to studying the computability of some function f defined on continuous spaces such as the Cantor space or the space of real numbers. One is usually interested in three stronger and stronger notions of computability for f :

- (i) $f(x)$ is computable for every computable x ;
- (ii) $f(x)$ is computable *relative* to x for every x ;
- (iii) $f(x)$ is computable relative to x for every x , *uniformly* in x .

In the first case we say that f is *computably invariant* (terminology introduced in [1]). In the third case we simply say that f is *computable*. It happens that many interesting functions are not computable and even not computably invariant. For instance Braverman and Yampolsky proved in [3] that the function mapping a parameter to the corresponding Julia set does not satisfy (ii); they later strengthened that result in [4] by proving that it does not



© Mathieu Hoyrup;
licensed under Creative Commons License CC-BY

Conference title on which this volume is based on.

Editors: Billy Editor, Bill Editors; pp. 1–17



Leibniz International Proceedings in Informatics

LIPICs Schloss Dagstuhl – Leibniz-Zentrum für Informatik, Dagstuhl Publishing, Germany

satisfy (i) either. By contrast, the function mapping a parameter to the corresponding *filled* Julia set does satisfy condition (ii), while it does not satisfy (iii) because it is discontinuous [5].

While functions that are not computable often fail to be computably invariant, the proof of the former is usually much simpler than the proof of the latter. Indeed, it is often based on the fundamental result that a computable function must be continuous. Hence proving that a function is not computable is often a purely topological argument.

However proving that a function is not computably invariant is usually much more challenging, as a counterexample must be constructed, by encoding the halting set or by using more involved computability-theoretic arguments based on priority methods, e.g. Our point is that topology is still at play in many computability-theoretic constructions¹. Usually the construction of a computable element whose image is not computable implicitly makes use of the discontinuity of the function. Of course mere discontinuity is not sufficient in general to carry out such a construction: there exist discontinuous functions that are computably invariant, such as the floor function or the function that maps a real number to its binary expansion. More is needed and our question is: what discontinuity property is needed to make such a construction possible?

Such discontinuity properties have already been sought by several authors. Pour-El and Richards “First Main Theorem” [14] shows that in the case of linear operators with c.e. closed graph, if the operator is unbounded (i.e., discontinuous) then it is not computably invariant (it is actually an equivalence). Their result subsumes many *ad hoc* constructions, such as Myhill’s differentiable computable function whose derivative is not computable [12]. As part of their open problem no. 7, Pour-El and Richards ask whether their First Main Theorem can be extended to nonlinear operators. A generalization of their theorem to certain algebraic structures was proved by Brattka [1], applicable to operators on the set of compact subsets of \mathbb{R} .

In these results, the underlying algebraic structures enable the authors to provide counterexamples via explicit expressions (such as linear combinations of basic elements with well-chosen weights) by encoding the halting set, which contrasts with many situations in computability theory where explicit constructions are rarely possible and priority methods are often needed to build counterexamples (Friedberg-Muchnik construction of Turing incomparable c.e. sets, e.g.). This observation allows one to hope for stronger results whose proofs involve more complicated, non-explicit constructions.

In this paper we present such a result, applicable to inverses of computable functions. We work on effective topological spaces and effective Polish spaces without additional structure, which makes our result applicable in many situations. We introduce a topological notion, *irreversibility* of a function, whose effective version entails the existence of a non-computable point whose image is computable. We think that this notion is rather simple to verify on particular instances. The proof of the result implicitly uses the priority method with finite injury. We think that our discontinuity notion is rather natural and, in concrete situations, much easier to verify than constructing a computable element whose pre-image is not computable. In other words, our result is not merely an abstract generalization of existing constructions, but a powerful theorem that provides insight into computability theory, as illustrated by the numerous examples we give.

¹ for instance the role of Baire category in computability theory has been revealed by several authors (see [11] e.g.)

This work was originally motivated by the following question, left open in [6]: are there two non-computable shift-invariant ergodic measures whose sum is computable? As an application of our main result, we positively answer this question.

We push our investigation further by studying the following question: how non-computable can a point with a computable image be? We introduce a notion of genericity of a point w.r.t. a function and prove that generic points with computable images exist. The construction unifies several finite injury arguments.

The paper is organized as follows: in Section 2 we introduce basic notions of computable analysis; in Section 3 we introduce a notion of continuous invertibility at a point and prove that for “almost” every point, if a function is computably invertible at that point then it is continuously invertible there (Theorem 7). In Section 4 we introduce the notion of an *irreversible function*, which in substance expresses that a function is topologically hard to inverse. In Section 5 we present our main result: a function that is topologically hard to inverse is computably hard to inverse, in particular it maps a non-computable point to a computable image. In Section 5.1 we present an application of our main result to the non-computability of the ergodic decomposition. In Section 6 we introduce a notion of genericity w.r.t. a function which unifies several finite injury constructions.

A complete version of the paper with all the proofs is available at <http://hal.inria.fr/docs/00/73/57/58/PDF/draft.pdf>. In the present version, we avoid some technical considerations that are necessary for the proofs of the results but not for their understanding. Most of the proofs are given in the appendix.

2 Background and notations

We assume familiarity with basic computability theory on the natural numbers. We implicitly use Weihrauch’s notions of computability on effective topological spaces, based on the standard representation (see [18] for more details), however we do not express them in terms of representations.

2.1 Notations

In a metric space (X, d) , if $x \in X$ and $r \in (0, +\infty)$ then we denote the open ball with center x and radius r by $B(x, r) = \{x' \in X : d(x, x') < r\}$. We denote the corresponding closed ball by $\overline{B}(x, r) = \{x' \in X : d(x, x') \leq r\}$. The Cantor space of infinite binary sequences, or equivalently subsets of \mathbb{N} , is denoted by $2^{\mathbb{N}}$. The halting set, denoted \emptyset' , is the set of numbers of Turing machines that halt. It is a noncomputable set that is computably enumerable (c.e.).

2.2 Effective topology

An *effective topological space* (X, τ, \mathcal{B}) consists of a topological space (X, τ) together with a countable basis $\mathcal{B} = \{B_0, B_1, \dots\}$ numbered in such a way that the finite intersection operator is computable. An open subset $U \subseteq X$ is *effectively open* if $U = \bigcup_{k \in W} B_k$ for some c.e. set $W \subseteq \mathbb{N}$.

To a point $x \in X$ we associate $N(x) = \{n \in \mathbb{N} : x \in B_n\}$. By an *enumeration of $N(x)$* we mean a total function $f : \mathbb{N} \rightarrow \mathbb{N}$ whose range is $N(x)$. A point x is *computable* if $N(x)$ is c.e., i.e. if $N(x)$ has a computable enumeration.

Given points x, y in effective topological spaces X, Y respectively, we say that y is *computable relative to x* if there is an oracle Turing machine M that, given any enumeration

of $N(x)$ as oracle, outputs an enumeration of $N(y)$. We denote it by $M^x = y$. In other words, y is computable relative to x if $N(y)$ is enumeration reducible to $N(x)$. As proved by Selman [16] and pointed out by Miller [10], y is computable relative to x if and only if every enumeration of $N(x)$ computes an enumeration of $N(y)$ (uniformity is not explicitly required, but is a consequence).

A (possibly partial) function $f : X \rightarrow Y$ is **computable** if there is a machine M such that for every $x \in \text{dom}(f)$, $M^x = f(x)$. A computable function is always continuous.

2.3 Effective Polish spaces

An **effective Polish space** is a topological space such that there exists a dense sequence s_0, s_1, \dots of points, called **simple** points and a complete metric d inducing the topology, such that all the real numbers $d(s_i, s_j)$ are computable uniformly in (i, j) . Every effective Polish space can be made an effective topological space, taking as canonical basis the open balls $B(s, r)$ with s simple point and r positive rational together with a standard effective numbering.

In an effective Polish space, a point x is computable if and only if for every $\epsilon > 0$ a simple point s can be computed, uniformly in ϵ , such that $d(s, x) < \epsilon$.

We will be concerned with computability and Baire category, so we will naturally meet the notion of a 1-generic point: a point that does not belong to any “effectively meager set” in the following sense.

► **Definition 1.** $x \in X$ is **1-generic** if x does not belong to the boundary of any effective open set. In other words, for every effective open set U , either $x \in U$ or there exists a neighborhood B of x disjoint from U .

By the Baire category theorem, every Polish space is a Baire space so 1-generic points exist and form a co-meager set.

3 A non-uniform result

Let X be an effective Polish space, Y an effective topological space and $f : X \rightarrow Y$ a computable function.

To introduce informally the results of this section, assume temporarily that f is one-to-one. If f^{-1} is computable, i.e. if every x is computable relative to $f(x)$ *uniformly* in x , then f^{-1} is continuous. As mentioned earlier uniformity is crucial here: that some x is computable relative to $f(x)$ does not imply in general that f^{-1} is continuous at $f(x)$. Theorem 7 below surprisingly shows that a non-uniform version can still be obtained, valid at most points.

Let us now make it precise and formal. We do not assume anymore that f is one-to-one.

When focusing on the problem of inverting a function, one comes naturally to the following basic notions:

- f is *invertible* at x if x is the only pre-image of $f(x)$,
- f is *locally invertible* at x if x is isolated in the pre-image of $f(x)$.

If one has access to x via its image only, then x is determined unambiguously in the first case, with the help of a discrete advice (a basic open set isolating x) in the second case. However, “being uniquely determined” is not sufficient in practice: physically or computationally, one cannot know entirely $f(x)$ in one step, but progressively as a limit of finite approximations. We need to consider stronger, topological versions of the two basic notions of invertibility, expressing that x can be recovered from the knowledge of its image given by finer and finer neighborhoods.

► **Definition 2.** f is *continuously invertible at x* if the pre-images of the neighborhoods of $f(x)$ form a neighborhood basis of x , i.e. for every neighborhood U of x there exists a neighborhood V of $f(x)$ such that $f^{-1}(V) \subseteq U$.

f is *locally continuously invertible at x* if there exists a neighborhood B of x such that the restriction of f to B is continuously invertible at x , i.e. for every neighborhood U of x there exists a neighborhood V of $f(x)$ such that $B \cap f^{-1}(V) \subseteq U$.

Observe that these notions are very natural when investigating the problem of inverting a function: we think that they are not technical *ad hoc* conditions.

The topological space Y is sequential, i.e. continuity notions can be expressed in terms of sequences, which may be more intuitive. We will be particularly interested in the negations of these notions, which we characterize now.

► **Proposition 3.1.** f is not continuously invertible at x if and only if there exist $\delta > 0$ and a sequence x_n such that $d(x, x_n) > \delta$ and $f(x_n)$ converge to $f(x)$.

f is not locally continuously invertible at x if and only if for every $\epsilon > 0$ there exist $\delta > 0$ and a sequence x_n such that $\epsilon > d(x, x_n) > \delta$ and $f(x_n)$ converge to $f(x)$.

Let us illustrate these notions on a few examples.

► **Example 3.** If f is one-to-one then f is continuously invertible at x if and only if f^{-1} is continuous at $f(x)$.

► **Example 4.** The real function $f(x) = x^2$ is continuously invertible exactly at 0, and locally continuously invertible everywhere (for $x \neq 0$ take for B an open interval avoiding 0).

► **Example 5.** The projection $\pi_1 : 2^{\mathbb{N}} \rightarrow 2^{\mathbb{N}}$ which maps $A_1 \oplus A_2 = \{2n : n \in A_1\} \cup \{2n+1 : n \in A_2\}$ to A_1 is not locally continuously invertible anywhere. Indeed, given $A_1, A_2 \in 2^{\mathbb{N}}$, $A_1 \oplus A_2$ is not isolated in the pre-image by π_1 of $A_1 = \pi_1(A_1 \oplus A_2)$.

► **Example 6.** Let X be the Cantor space $2^{\mathbb{N}}$ with the product topology τ generated by the cylinders $[u]$, $u \in 2^*$, Y be the Cantor space with the positive topology τ_{Scott} generated by the sets $\{A \subseteq \mathbb{N} : F \subseteq A\}$ where F varies among the finite subsets of \mathbb{N} . The computable elements of the two effective topological spaces are the computable sets and the c.e. sets respectively. Consider the enumeration operator $\text{Enum} := \text{id} : X \rightarrow Y$. Enum is computable and one-to-one but its inverse is discontinuous. More precisely, (i) it is continuously invertible exactly at \mathbb{N} , (ii) it is locally continuously invertible exactly at the co-finite sets: if A is co-finite then let B be a cylinder specifying all the 0's in A , every cylinder containing A is the intersection of a Scott open set with B .

In general continuous invertibility at a point is strictly stronger than local continuous invertibility. It is not the case for linear operators, where a dichotomy appears. Following Pour-El and Richards [14], by a linear operator $T : X \rightarrow Y$ between Banach spaces we mean a linear function $T : \mathcal{D}(T) \rightarrow Y$ where $\mathcal{D}(T)$ is a subspace of X .

► **Proposition 3.2.** Let X, Y be Banach spaces and $T : X \rightarrow Y$ a one-to-one linear operator.

- If T^{-1} is bounded then T is continuously invertible everywhere.
- If T^{-1} is unbounded then T is nowhere locally continuously invertible.

Proof. The first point simply follows from the fact that T^{-1} is continuous. Assume that T^{-1} is unbounded. There exists a sequence $a_n \in X$ such that $\|a_n\| = 1$ and $\|T(a_n)\| \rightarrow 0$. Let $x \in X$ and $\epsilon > 0$. Take $\delta = \epsilon/3$ and define $x_n = x + 2\delta a_n$: $T(x_n)$ converge to $T(x)$ and $\epsilon > \|x - x_n\| > \delta$ for all n . ◀

Observe that in the case when T is not one-to-one, T is also nowhere locally continuously invertible, with exactly the same proof (one can take $a_n = a$ for some a with $\|a\| = 1$ and $\|T(a)\| = 0$).

We now come to our first result.

► **Theorem 7.** *Let $f : X \rightarrow Y$ be a computable function and $x \in X$ a 1-generic point. If x is computable relative to $f(x)$ then f is locally continuously invertible at x .*

Proof idea. Assume that f is not locally continuously invertible at x and that there is a Turing machine M that computes x on oracle $f(x)$. We show that x belongs to the boundary of an effective open set U , i.e. that x is not 1-generic.

Given a point y , there are two possible ways in which a machine may fail to compute y from $f(y)$: either it diverges, or it outputs something that is incompatible with y . The latter can be recognized in finite time: we then say that $M^{f(y)}$ *positively* fails to compute y . Our effective open set U is the set of points y such that $M^{f(y)}$ positively fails to compute y .

First, if f is not continuously invertible at x , there exists $\delta > 0$ and a sequence x_n such that $d(x_n, x) > \delta$ and $f(x_n)$ converge to $f(x)$. If n is sufficiently large then $f(x_n)$ is arbitrarily close to $f(x)$ so $M^{f(x_n)}$ computes an arbitrarily refined approximation of x . If we take n so large that $M^{f(x_n)}$ computes x at precision $< \delta/2$, then $M^{f(x_n)}$ positively fails to compute x_n so x_n belongs to U .

Now, if f is not *locally continuously invertible at x* then x_n can be taken arbitrarily close to x , so x belongs to the closure of U . ◀

In the sequel we introduce a condition on f which roughly means that f is “almost nowhere” locally continuously invertible and that entails (i) the existence of an x that is not computable relative to $f(x)$ (Theorem 13) and, better, (ii) the existence of a non-computable x such that $f(x)$ is computable (Theorem 20).

4 Reversibility

We define two dual notions for a function: reversible (Section 4.1) and irreversible (Section 4.2). In the sense of Baire category, a reversible function is continuously invertible almost everywhere; an irreversible function is almost nowhere locally continuously invertible.

4.1 Reversible functions

Let X, Y be T_0 topological spaces. For a continuous function $f : X \rightarrow Y$, the following are equivalent:

- f is one-to-one and $f^{-1} : f(X) \rightarrow X$ is continuous,
- the initial topology of f is the topology of X , i.e. for every open set $U \subseteq X$ there exists an open set $V \subseteq Y$ such that $U = f^{-1}(V)$.

A function satisfying these conditions can be *reversed* in the sense that x can be recovered from $f(x)$ for every x : x is not only uniquely determined by $f(x)$, but a neighborhood basis of x can be progressively constructed from a neighborhood basis of $f(x)$.

We first consider a slight weakening of this notion.

► **Definition 8.** We say that f is *reversible* if for every non-empty open set $U \subseteq X$ there is an open set $V \subseteq Y$ such that $\emptyset \neq f^{-1}(V) \subseteq U$.

We say that f is *effectively reversible* if $V = V_U$ can moreover be computed from U (basic open set).

► **Proposition 4.1.** If f is reversible then it is continuously invertible at every point in a dense G_δ -set.

If f is effectively reversible then there is a dense effective G_δ -set D such that $f|_D$ is one-to-one and its inverse is computable on $f(D)$, i.e. x is uniformly computable from $f(x)$ when $x \in D$.

In particular if x is 1-generic then x is computable relative to $f(x)$.

4.2 Irreversible functions

We now consider the dual notion: an *irreversible* function is a function that is not reversible, even locally.

► **Definition 9.** f is *irreversible* if for every open set $B \subseteq X$ the restriction $f|_B : B \rightarrow f(B)$ is not reversible.

Formally, f is irreversible if for every non-empty open set B there exists a non-empty open set $U_B \subseteq B$ such that there is no open set V satisfying $\emptyset \neq f^{-1}(V) \cap B \subseteq U_B$.

In other words, each pre-image of open set that intersects B does so outside U_B . If $x \in U_B$ then we will never know it from $f(x)$, even with the help of the advice $x \in B$.

Observe that one can assume w.l.o.g. that $f^{-1}(V) \cap B \not\subseteq \overline{U_B}$. Indeed, one can replace U_B by some ball $B(s, r)$ such that $\overline{B}(s, r) \subseteq U_B$.

An application of an irreversible function f to x comes with a loss of information about x , that can hardly be recovered. Being irreversible is orthogonal to not being one-to-one: the function $x \mapsto x^2$ is not one-to-one but not irreversible: x can be (continuously or computably) recovered from x^2 ; a one-to-one function can be irreversible if its inverse is dramatically discontinuous (examples of such functions will be encountered in the sequel).

In terms of sequences, f is irreversible if and only if for every B there exists a non-empty open set $U_B \subseteq B$ such that for every $x \in U_B$ there is a sequence $x_n \in B \setminus U_B$ such that $f(x_n)$ converge to $f(x)$.

As announced, the set of points at which an irreversible function is locally continuously invertible is small in the sense of Baire category.

► **Proposition 4.2.** Let f be irreversible. There is a dense G_δ -set D such that f is not locally continuously invertible at any $x \in D$.

In other words, for almost every x the application of f to x comes with a “topological information” loss.

The preceding proposition does not rule out the possibility that the restriction of f to a “large” set be continuously invertible (for instance, the characteristic function of the rational numbers is nowhere continuous, but its restriction to the co-meager set of irrational numbers is continuous). The next assertion shows that it is not possible.

► **Proposition 4.3.** Let f be irreversible and $C \subseteq X$ be such that $f|_C : C \rightarrow f(C)$ is an homeomorphism. C is nowhere dense.

Proof. Assume the closure of C contains a ball B . $U_B \cap C$ is non-empty. Let $x \in U_B \cap C$. There exists a sequence $x_n \in B \setminus \overline{U_B}$ such that $f(x_n)$ converge to $f(x)$. By density of C in B , x_n can be taken in C . As $f|_C$ is an homeomorphism and $f(x_n)$ converge to $f(x)$, x_n should converge to x and eventually enter U_B , which gives a contradiction. ◀

► **Example 10.** Let f be a constant function defined on the Polish space X . f is irreversible if and only if X is perfect, i.e. has no isolated point.

► **Example 11.** The first projection $\pi_1 : 2^{\mathbb{N}} \rightarrow 2^{\mathbb{N}}$ from Example 5 is irreversible. Indeed, to $B = [w]$, associate $U_B = [w00]$. The intersection with $[w]$ of the pre-image of any cylinder cannot be contained in $[w00]$: knowing arbitrarily many bits of $\pi_1(A)$ and the first $|w|$ bits of A does not give any information about the next odd bit of A , so it does not enable one to guess that A belongs to $[w00]$.

In the definition of an irreversible function (Definition 9), B and U_B can be assumed w.l.o.g. to be basic balls.

► **Definition 12.** f is *effectively irreversible* if U_B can be computed from B .

The following result is the effective version of Proposition 4.3.

► **Theorem 13.** *If f is effectively irreversible then for every 1-generic x , x is not computable relative to $f(x)$.*

Proof. The dense G_δ -set provided by Proposition 4.2 is effective when f is effectively irreversible so it contains every 1-generic point. Hence for every 1-generic x , f is not locally continuously invertible at x . We now apply Theorem 7. ◀

In other words, if x is 1-generic then the application of f to x comes with an “algorithmic information” loss. So if f is effectively irreversible then there exists some x that is not computable relative to $f(x)$.

4.3 Examples

Several well-known results in computability theory can be interpreted using Theorem 13 as consequences of the effective irreversibility of some computable function.

► **Example 14.** Consider the enumeration operator of Example 6. Enum is effectively irreversible: to each cylinder $B = [w]$ associate $U_B = [w0]$.

Applying Theorem 13 then gives: if A is 1-generic then A and $\mathbb{N} \setminus A$ have incomparable enumeration degrees. Such an A was first proved to exist by Selman [16].

► **Example 15.** The projection $\pi_1 : 2^{\mathbb{N}} \rightarrow 2^{\mathbb{N}}$ from Examples 5 and 11 is effectively irreversible. Applying Theorem 13 to π_1 and symmetrically to the second projection π_2 gives Jockush and Posner’s result [8] that if $A = A_1 \oplus A_2$ is 1-generic then A_1 and A_2 are Turing incomparable, which implies Kleene-Post theorem, taking a \emptyset' -computable 1-generic set.

► **Example 16.** Jockush [9] proved that every 1-generic $A \in 2^{\mathbb{N}}$ is c.e.a., i.e. A computes some B such that A is c.e. relative to B but not computable relative to B . The proof goes as follows: let $f(A) = \{\langle i, j \rangle : i \in A \wedge \langle i, j \rangle \notin A\}$ (where $\langle \cdot \rangle$ is a computable one-to-one pairing function such that $\langle i, j \rangle > i$). f is computable, if A is 1-generic then A is c.e. in $f(A)$ as $i \in A \iff \exists j, \langle i, j \rangle \in A$. We show that f is effectively irreversible, which by Theorem 13 implies that if A is 1-generic then $A \not\leq_T f(A)$.

First observe that f is not one-to-one: given A and i such that $i \notin A$ and $\langle i, 0 \rangle \notin A$, there exists $\hat{A} \neq A$ such that $f(\hat{A}) = f(A)$. Add $\langle i, 0 \rangle$ to A , and each time some k is added, add all the pairs $\langle k, j \rangle$ that are not already in. One easily checks that $f(\hat{A}) = f(A)$. As a result, given a cylinder $B = [u]$, let $U_B = [u] \cap \{A : i \notin A \text{ and } \langle i, 0 \rangle \notin A\}$. For every $A \in U_B$ there is $\hat{A} \in B \setminus U_B$ such that $f(\hat{A}) = f(A)$, so $f^{-1}f(A)$ intersects $B \setminus U_B$: knowing $f(A)$ and that $A \in B$ does not enable one to know that $A \in U_B$.

Again, linear operators provide a large class of examples. An effective Banach space is a Banach space which is an effective Polish space with the metric induced by the norm, such that 0 is a computable point and the vector space operations are computable functions. Many classical Banach spaces \mathbb{R} , $\mathcal{C}[0, 1]$ (with the uniform norm) or $L^1[0, 1]$ are effective Banach spaces.

► **Proposition 4.4.** Let X, Y be effective Banach spaces and $T : X \rightarrow Y$ a computable linear operator. Assume that either T is not one-to-one or T is one-to-one and T^{-1} is unbounded. T is effectively irreversible.

► **Example 17.** Applying Proposition 4.4 and Theorem 13 to the integration operator that maps $f \in \mathcal{C}[0, 1]$ to $F : x \mapsto \int_0^x f(t) dt$ gives that if $f \in \mathcal{C}[0, 1]$ is 1-generic then f is not computable relative to its primitive F that vanishes at 0.

► **Example 18.** Applying Proposition 4.4 and Theorem 13 to the canonical injection from $\mathcal{C}[0, 1]$ to $L^1[0, 1]$ gives that if $f \in \mathcal{C}[0, 1]$ is 1-generic then it is not computable relative to itself, as an element of $L^1[0, 1]$. In other words, the description of f as an element of $L^1[0, 1]$ contains strictly less algorithmic information than the description of f as an element of $\mathcal{C}[0, 1]$.

► **Example 19.** A function $f : \mathbb{N} \rightarrow \mathbb{N}$ can be described by enumerating its graph or by enumerating the complement of its graph. The former alternative gives in general strictly more information about the function than the latter. Let us make it precise.

Every function F with finite domain induces the cylinder $[F]$ of functions $f : \mathbb{N} \rightarrow \mathbb{N}$ extending F . The product topology on the Baire space \mathbb{B} is generated by the cylinders. The negative topology is generated by the complements of the cylinders, as a subbasis. The identity $\text{id} : (\mathbb{B}, \tau) \rightarrow (\mathbb{B}, \tau_{\text{neg}})$ is computable: from f one can enumerate the cylinders that are incompatible with f , but the converse cannot be done. id is effectively irreversible: to a cylinder $B = [F]$, associate $U_B = [F] \cup \{n \mapsto 0\}$ where n is fresh, i.e. not in the domain of F .

By Theorem 13, if $f : \mathbb{N} \rightarrow \mathbb{N}$ is 1-generic then it is not computable relative to every co-enumeration of its graph.

5 The constructive result

We now present the main result of the paper. It is the constructive version of Theorem 13 as it makes $f(x)$ computable. The construction uses a priority argument with finite injury.

► **Theorem 20.** *If f is effectively irreversible then there exists a non-computable x such that $f(x)$ is computable.*

The proof is given in the appendix. The proof uses the priority method with finite injury, which can be seen as a game between a player, computing $f(x)$, and infinitely many opponents (all the Turing machines) trying to compute x .

5.1 Application to the ergodic decomposition

We now present an application of Theorem 20. Let P be a Borel probability measure P over the Cantor space. P is **computable** if the real numbers $P[w]$ are uniformly computable. P is **shift-invariant** if $P[w] = P[0w] + P[1w]$ for each finite string w . P is **ergodic** if it cannot be written as $P = \frac{1}{2}(P_1 + P_2)$ with $P_1 \neq P_2$ both shift-invariant.

The ergodic decomposition theorem says that every shift-invariant measure can be uniquely decomposed into a convex combination (possibly uncountable) of ergodic measures. Our question is: given a computable shift-invariant measure, can we compute in a sense its ergodic decomposition? This question was implicitly addressed by V'yugin [17] who constructed a counter example: a countably infinite combination of ergodic measures which is computable but not effectively decomposable. In [6] we raised the following question: does the ergodic decomposition become computable when restricting to finite combinations? As an application of Theorem 20, we solve the problem and prove that it is already non-effective in the finite case:

► **Theorem 21.** *There exist two ergodic shift-invariant measures P and Q such that neither P nor Q is computable but $P + Q$ is computable.*

The strategy is as follows: the mapping $(P, Q) \mapsto P + Q$ is computable, two-to-one on the space $\mathcal{E} \times \mathcal{E}$ of pairs of ergodic measures and we prove

► **Theorem 22.** *The function $(P, Q) \mapsto P + Q$ defined on $\mathcal{E} \times \mathcal{E}$ is effectively irreversible.*

which implies the result applying Theorem 20.

6 Genericity

Given an effectively irreversible function f ,

- Theorem 13 tells us that if x is 1-generic then x is not computable relative to $f(x)$,
- Theorem 20 tells us that there exist non-computable x such that $f(x)$ is computable.

The two results are “disjoint” in the sense that a single x cannot at the same time be 1-generic and have a computable image. We raise the following question: is it possible to bring the two results closer together? How far can x be from being computable, given that $f(x)$ is computable? How *generic* can x be?

We now give an answer to that question. We recall that a topological space always comes with an order called the *specialization order*: $x \leq y$ iff every neighborhood of x is also a neighborhood of y . $x \leq y$ means that if one describes x by listing its basic neighborhoods then one can never distinguish x from y . When the space is Hausdorff, the specialization order is trivial. Here \leq denotes the specialization order on the target space Y of f .

► **Definition 23.** x is *f -generic* if x is 1-generic in the subspace $\uparrow_f x := \{x' : f(x) \leq f(x')\}$. In other words, x is f -generic if for every effective open set U , either $x \in U$ or there exists a neighborhood B of x such that $B \cap U \cap \uparrow_f x = \emptyset$.

For instance, taking the first projection $\pi_1 : 2^{\mathbb{N}} \rightarrow 2^{\mathbb{N}}$ of Example 15, $A = A_1 \oplus A_2$ is π_1 -generic iff A_1 is 1-generic relative to A_2 .

Here we focus on a few particular instances of this notion, when f is the identity from a space to itself with two different topologies. We will consider

- i) the enumeration operator $\text{Enum} = \text{id} : (2^{\mathbb{N}}, \tau_{\text{prod}}) \rightarrow (2^{\mathbb{N}}, \tau_{\text{Scott}})$ (Examples 6 and 14),
- ii) $\text{id} : (2^{\mathbb{N}}, \tau_{\text{prod}}) \rightarrow (2^{\mathbb{N}}, \tau_{\text{lex}})$ where τ_{lex} is generated by the sets $\{y \in 2^{\mathbb{N}} : x <_{\text{lex}} y\}$ and
- iii) $\text{id} : (\text{CL}(2^{\mathbb{N}}), \tau_{\text{hit-or-miss}}) \rightarrow (\text{CL}(2^{\mathbb{N}}), \tau_{\text{miss}})$. Here, $\text{CL}(2^{\mathbb{N}})$ is the set of non-empty closed subsets of the Cantor space. τ_{miss} is generated by the sets $\mathcal{U}_u = \{P \in \text{CL}(2^{\mathbb{N}}) : P \cap [u] = \emptyset\}$ where $u \in 2^*$. $\tau_{\text{hit-or-miss}}$ is generated by the sets \mathcal{U}_u together with their complements.

Definition 23 is instantiated as follows:

- **Definition 24.** i) A *generic c.e. set* x is a c.e. set that is 1-generic in the subspace $\{y \in 2^{\mathbb{N}} : x \subseteq y\}$.
 ii) A *generic left-c.e. real* x is a left-c.e. real that is 1-generic in the subspace $\{y \in 2^{\mathbb{N}} : x \leq_{\text{lex}} y\}$.
 iii) A *generic Π_1^0 -class* P is a Π_1^0 -class that is 1-generic in the subspace $\{Q \in \text{CL}(2^{\mathbb{N}}) : Q \subseteq P\}$.

Informally, a generic element belongs to every effective open set that is dense *above* it, for the corresponding specialization order (while a 1-generic elements belongs to every effective open set that is dense *along* it). The next result is the sought combination of Theorems 13 and 20.

- **Theorem 25.** *There exists a co-infinite generic c.e. set, a co-infinite generic left-c.e. real and a generic Π_1^0 -class without isolated point.*

Proof idea (detailed in the appendix). Kurtz built a left-c.e. weakly 1-generic real (see [13] for a proof). The construction actually gives a generic left-c.e. real. The construction of a generic c.e. set and of a generic Π_1^0 -class are exactly the same, replacing the lexicographic order \leq_{lex} by inclusion \subseteq of sets and reverse inclusion of classes respectively, which are the specialization orders of the corresponding topologies. ◀

Theorem 25 is indeed a strengthening of Theorem 20: in Theorem 7, the 1-genericity assumption can actually be weakened to f -genericity (at least in the particular functions under consideration).

- **Proposition 6.1.** In each one of the three cases, if x is generic inside $\uparrow_f x$ and f is not locally continuously invertible at x then x is not computable.

Proof. Using compactness of the space, one can show that f is not locally continuously invertible at x iff x is not isolated in $\uparrow_f x$, i.e. x belongs to the closure of $\uparrow_f x \setminus \{x\}$. If x is computable then the complement of $\{x\}$ is an effective open set, so x cannot be generic inside $\uparrow_f x$. ◀

As the next result shows, Theorem 25 embodies many simple finite injury arguments as Friedberg-Muchnik theorem, e.g.

- **Proposition 6.2.** Let A be a co-infinite generic c.e. set. A is hypersimple, $A = A_1 \oplus A_2$ where A_1 and A_2 are Turing incomparable, A is not autoreducible.

Proof. Same argument as for 1-generic sets, observing that the involved open set is not only dense *along* A , but even *above* A . For instance, to prove that $A_2 \not\leq_T A_1$, given a Turing functional ϕ , let $U = \{A_1 \oplus A_2 : \exists n, \phi^{A_1}(n) = 0 \wedge A_2(n) = 1\}$. If $\phi^{A_1} = A_2$ then replacing a 0 in A_2 by a 1 arbitrarily far gives an element of U arbitrarily close to $A_1 \oplus A_2$ that is *above* (i.e. is a superset of) $A_1 \oplus A_2$. ◀

It happens that the co-infinite generic c.e. sets are exactly the p -generic sets defined by Ingrassia [7].

- **Proposition 6.3.** A generic Π_1^0 -class without isolated point has no computable member.

Proof. Let x be computable. Consider the collection $\mathcal{U} = \{P : x \notin P\}$. \mathcal{U} is an effective open set in the space $(\text{CL}(2^{\mathbb{N}}), \tau_{\text{hit-or-miss}})$ (and even in the topology τ_{miss}). \mathcal{U} is dense and better: for every P without isolated point, there exist $Q \subseteq P$ in \mathcal{U} arbitrarily close to P , so \mathcal{U} is dense below P (here the specialization order is the reverse inclusion). As a result, if P is a generic Π_1^0 -class without isolated point then P belongs to \mathcal{U} , i.e. $x \notin P$. ◀

7 Acknowledgements

The author wishes to thank Peter Gács, Emmanuel Jeandel and Cristóbal Rojas for discussions on the subject and helpful comments on a draft of the paper, and Christopher Porter for suggesting Example 16.

References

- 1 V. Brattka. Computable invariance. *Theor. Comput. Sci.*, 210(1):3–20, 1999.
- 2 V. Brattka. Computable versions of Baire’s category theorem. In J. Sgall, A. Pultr, and P. Kolman, editors, *MFCS*, pages 224–235, London, UK, 2001. Springer-Verlag.
- 3 M. Braverman and M. Yampolsky. Non-computable Julia sets. *Journ. Amer. Math. Soc.*, 19:551–578, 2006.
- 4 M. Braverman and M. Yampolsky. Constructing non-computable Julia sets. In *STOC*, pages 709–716. ACM, 2007.
- 5 M. Braverman and M. Yampolsky. *Computability of Julia Sets*. Springer, 2008.
- 6 M. Hoyrup. Randomness and the ergodic decomposition. In B. Löwe, D. Normann, I. N. Soskov, and A. A. Soskova, editors, *CiE*, volume 6735 of *Lecture Notes in Computer Science*, pages 122–131. Springer, 2011.
- 7 M. Ingrassia. *P-genericity for Recursively Enumerable Sets*. PhD thesis, University of Illinois at Urbana-Champaign, 1981.
- 8 J. Jockusch, Carl G. and D. B. Posner. Double jumps of minimal degrees. *The Journal of Symbolic Logic*, 43(4):pp. 715–724, 1978.
- 9 C. G. Jockusch. Degrees of generic sets. In F. R. Drake and S. S. Wainer, editors, *Recursion Theory: its Generalisations and Applications*, pages 110–139. Cambridge University Press, 1980.
- 10 J. S. Miller. Degrees of unsolvability of continuous functions. *Journal of Symbolic Logic*, 69(2):555–584, 2004.
- 11 J. Myhill. Category methods in recursion theory. *Pac. J. Math.*, 11:1479–1486, 1961.
- 12 J. Myhill. A recursive function, defined on a compact interval and having a continuous derivative that is not recursive. *Michigan Math. J.*, 18(2):97–98, 1971.
- 13 A. Nies. *Computability and randomness*. Oxford logic guides. Oxford University Press, 2009.
- 14 M. B. Pour-El and J. I. Richards. *Computability in Analysis and Physics*. Perspectives in Mathematical Logic. Springer, Berlin, 1989.
- 15 H. J. Rogers. *Theory of Recursive Functions and Effective Computability*. MIT Press, Cambridge, MA, USA, 1987.
- 16 A. L. Selman. Arithmetical reducibilities I. *Mathematical Logic Quarterly*, 17(1):335–350, 1971.
- 17 V. V. V’yugin. Effective convergence in probability and an ergodic theorem for individual random sequences. *SIAM Theory of Probability and Its Applications*, 42(1):39–50, 1997.
- 18 K. Weihrauch. *Computable Analysis*. Springer, Berlin, 2000.
- 19 M. Yasugi, T. Mori, and Y. Tsujii. Effective properties of sets and functions in metric spaces with computability structure. *Theoretical Computer Science*, 219(1-2):467–486, 1999.

A Proofs of simple propositions

A.1 Proposition 4.1

Assume that f is reversible. For each basic ball $U \subseteq X$ there exists $V_U \subseteq Y$ such that $\emptyset \neq f^{-1}(V_U) \subseteq U$. Let W_n be the union of $f^{-1}(V_U)$ over all basic balls U of radius $< 2^{-n}$. W_n is a dense open set. If $x \in W_n$ for all n then f is continuously invertible at x . Indeed, for every n there exists a ball U of radius $< 2^{-n}$ such that $x \in f^{-1}(V_U) \subseteq U$.

If f is effectively irreversible then the sets W_n are uniformly effective open sets and if $x \in \bigcap_n W_n$ then for each n there exists a basic ball U of radius $< 2^{-n}$ such that $f(x) \in V_U$, which can be found from any enumeration of $N(f(x))$. It gives an approximation of x within 2^{-n} .

A.2 Proposition 4.2

Let W_n be the union of U_B for all basic open sets B of radius $< 2^{-n}$. W_n is a dense open set. Let $x \in \bigcap_n W_n$. For each n there is a ball of radius $< 2^{-n}$ such that $x \in U_B$. For every neighborhood V of $f(x)$, $x \in f^{-1}(V) \cap B \neq \emptyset$ so $f^{-1}(V) \cap B \not\subseteq U_B$.

A.3 Proposition 4.4

To a ball $B = B(s, r)$ associate $U_B = B(s, r/2)$. According to the assumption about T , for every ϵ there exists a such that $\|a\| = r/2$ and $\|T(a)\| < \epsilon$. For every $x \in U_B$ and $\epsilon > 0$ there exists $\lambda \in \{-1, 1\}$ such that $r/2 \leq d(x + \lambda a, s) < r$, i.e. $x + \lambda a \in B \setminus U_B$. Indeed, $d(x + a, s) + d(x - a, s) \geq 2\|a\| = r$ and $d(x \pm a, s) < r$. Moreover, $d(T(x \pm \lambda a), T(x)) < \epsilon$.

B Proof of Theorem 20

We fix a one-to-one computable enumeration n_0, n_1, \dots of the halting set \emptyset' . We construct $x \in X$ such that $f(x)$ is computable and \emptyset' is computable relative to x . We construct a shrinking sequence of balls B_n and define x as the unique member of their intersection. Of course, B_n must not be computable otherwise x would be computable. The sequence B_n is constructed in stages: at stage s we define $B_n[s]$ and for each n the sequence $B_n[s]$ is stationary, with limit B_n . For each s , the sequence $B_n[s]$ is shrinking, so the limiting sequence B_n will be shrinking as well. One may imagine, for each s , the sequence $B_n[s]$ as an infinite path in a tree. At stage $s + 1$, n_s is enumerated into A' and the current path branches at depth n_s .

In order to make $f(x)$ computable we enumerate along the construction all its basic neighborhoods into a list L . L is the union of a computable growing sequence of finite lists L_s . At stage s , the current neighborhood of $f(x)$, denoted \mathcal{V}_s is the (finite) intersection of the members of L_s . As $L_s \subseteq L_{s+1}$, $\mathcal{V}_{s+1} \subseteq \mathcal{V}_s$.

In order to construct the list L , we start with a technical point: in the space X , we make an effective change of simple points and basic open sets. We can assume w.l.o.g. that the radius of U_B is at most half the radius of B . Given a basic ball B , consider the computable sequence $U_B^{(n)}$ defined inductively by $U_B^{(0)} = B$ and $U_B^{(n+1)} = U_{U_B^{(n)}}$. $U_B^{(n)}$ is a computable shrinking sequence and the unique member a of $\bigcap_n U_B^{(n)}$ is computable, uniformly in B . The canonical enumeration B_j of basic balls induces a computable dense sequence a_j , which will serve as simple points.

We then change the basic open subsets of X . Let $(V_k)_{k \in \mathbb{N}}$ be the canonical enumeration of the basic open subsets of Y .

► **Lemma 26.** *There is a double-sequence of open sets $O_{k,i} \subseteq X$ such that*

- $O_{k,i} \subseteq O_{k,i+1}$,
- $f^{-1}(V_k) = \bigcup_{i \in \mathbb{N}} O_{k,i}$,
- the predicate $a_j \in O_{k,i}$ is decidable.

Proof. By a standard diagonalization argument (computable Baire theorem on the real numbers), there exists a computable dense sequence of positive real numbers r_n such that $d(s_i, a_j) \neq r_n$ for all i, j, n . The metric balls $B(s_i, r_n)$ form an effective basis and the predicate $a_j \in B(s_i, r_n)$ is decidable. $f^{-1}(V_k)$ can be expressed as an effective union of such balls. Define $O_{k,i}$ as the union of the first i balls enumerated into $f^{-1}(V_k)$. ◀

We now proceed to the construction of the sequence $B_n[s]$ for each stage s . For each s , $B_n[s]$ will be a shrinking sequence, $x[s]$ will be defined as the unique member of their intersection and will be one of the points $\{a_j : j \in \mathbb{N}\}$.

Stage 0. We start with a ball $B_0[0]$ of radius 1, $B_{n+1}[0] = U_{B_n[0]}$ and $\{x[0]\} = \bigcap_n B_n[0]$. Start with $L_0 = \emptyset$ and $\mathcal{V}_0 = Y$. Observe that for each n , $B_n[0] \cap f^{-1}(\mathcal{V}_0)$ is non-empty as it contains $x[0]$.

Stage $s + 1$. First, L_{s+1} is obtained by adding to L_s all the numbers $k \leq s$ such that $x[s] \in O_{k,s}$. Let \mathcal{V}_{s+1} be the intersection of the open sets V_k with $k \in L_{s+1}$.

Let $n = n_s$ be the next element enumerated into the halting set. Let $B_{n+1}[s+1]$ be a ball satisfying $\overline{B_{n+1}[s+1]} \subseteq f^{-1}(\mathcal{V}_{s+1}) \cap B_n[s] \setminus \overline{B_{n+1}[s]}$. Such a ball exists: $f^{-1}(\mathcal{V}_{s+1}) \cap B_n[s]$ is non-empty as it contains $x[s]$, f is irreversible and $B_{n+1}[s] = U_{B_n[s]}$. For $n' \leq n$, let $B_{n'}[s+1] = B_{n'}[s]$. For $n' > n$ define by induction $B_{n'+1}[s+1] = U_{B_{n'}[s+1]}$. Let $\{x[s+1]\} = \bigcap_n B_n[s+1]$.

Verification. By construction one has $\overline{B_{n+1}[s]} \subseteq B_n[s]$ and $B_{n+1}[s] = U_{B_n[s]}$ for sufficiently large n so $B_n[s]$ is a shrinking sequence.

We call the *settling time* of n the minimal number s such that $n_{s'} \geq n$ for all $s' \geq s$.

We say that $n \in \emptyset'$ is a *forward element* if no element $m < n$ is enumerated into \emptyset' after the enumeration stage of n : in other words, the settling time of n coincides with its enumeration stage. As \emptyset' is infinite, it has infinitely many forward elements.

► **Claim.** For each n , $B_n[s]$ is a stationary sequence.

Proof. Let s be the settling time of n : $B_n[s] = B_n[s_0]$ for all $s \geq s_0$. ◀

Let B_n be its limit. B_n is a shrinking sequence as well, let x be the member of its intersection. Observe that the sequence $x[s]$ converges to x : given ϵ , let n be such that B_n has radius $< \epsilon$ and s_0 be the settling time n : for all $s \geq s_0$, $x[s] \in B_n[s] = B_n$ so $d(x[s], x) < \epsilon$.

► **Claim.** $f(x)$ is computable.

Proof. We prove that a basic open set V_k contains $f(x)$ if and only if k is enumerated into the list $L = \bigcup_s L_s$.

If $k \in L_s$ for some s , let n be a forward element which is enumerated at some stage $s' \geq s$. $x \in \overline{B_{n+1}} = \overline{B_{n+1}[s'+1]} \subseteq f^{-1}(\mathcal{V}_{s'+1}) \subseteq f^{-1}(\mathcal{V}_s) \subseteq f^{-1}(V_k)$.

Now let V_k be a basic neighborhood of $f(x)$. Let i_0 be such that $x \in O_{k,i_0}$. As $x[s]$ converge to x there is s such that $x[s] \in O_{k,i_0}$ for all $s' \geq s$. Let $t = \max(s, i_0)$: $x[t] \in O_{k,i_0} \subseteq O_{k,t}$ so k must be added to the list at stage $t + 1$ or earlier. ◀

► **Claim.** \emptyset' is computable relative to x .

Proof. Let p_i be the increasing sequence of forward elements. \emptyset' can be computed from the sequence p_i and the (computable) enumeration of \emptyset' .

From x one can inductively compute the sequence p_i . First, p_0 is the minimal n such that $x \notin B_{n+1}[0]$. Once p_i is known, let s be the stage at which p_i is enumerated into \emptyset' , i.e. $n_s = p_i$. p_{i+1} is the minimal $n > p_i$ such that $x \notin B_{n+1}[s+1]$. ◀

C Proof of Theorem 22

Before proving the theorem, we need some preliminaries so show that $\mathcal{E} \times \mathcal{E}$ is an effective Polish space.

Let X be an effective Polish space. $X' \subseteq X$ is an *effective Polish subspace* if it is an effective Polish space with the induced topology and such that the canonical injection from X' to X to be a computable homeomorphism. Alexandrov theorem gives a way to obtain Polish subspaces of a Polish space, and has an effective version, which we present now.

A set A is an *effective G_δ -set* if there exists a family of uniformly effective open sets U_n such that $A = \bigcap_n U_n$. Such a set is *c.e.* if it contains a dense computable sequence. Examples of such sets are given by the computable Baire theorem [19, 2]: any dense effective G_δ -set is a c.e. effective G_δ -set.

► **Proposition C.1 (Effective Alexandrov Theorem).** Every c.e. effective G_δ -set is an effective Polish subspace of X .

We consider the space $\mathcal{P}(2^\mathbb{N})$ of Borel probability measures over the Cantor space together with the complete metric

$$d(P, Q) = \sum_{w \in \{0,1\}^*} 2^{-|w|} |P[w] - Q[w]|.$$

The finite rational combination of Dirac measures are dense in $\mathcal{P}(2^\mathbb{N})$ and d is computable over them, so $\mathcal{P}(2^\mathbb{N})$ is an effective Polish space. The subset \mathcal{I} of shift-invariant measures is closed so d is complete over \mathcal{I} as well. \mathcal{I} easily contains a dense computable sequence, so \mathcal{I} is an effective Polish subspace of $\mathcal{P}(2^\mathbb{N})$. Let $\mathcal{E} \subseteq \mathcal{I}$ be the set of ergodic shift-invariant measures. d is no more complete over \mathcal{E} , but Proposition C.1 implies that \mathcal{E} is an effective Polish subspace. We work with the basis given by the intersection of the canonical basis of \mathcal{I} with \mathcal{E} .

We now present the proof of Theorem 22. Let $B \subseteq \mathcal{I} \times \mathcal{I}$ be an open set and $(P_0, Q_0) \in B$ with $P_0 \neq Q_0$. Let $\epsilon > 0$ be such that $d(P_0, Q_0) > \epsilon$ and $B(P_0, \epsilon) \times B(Q_0, \epsilon) \subseteq B$. Let $\delta = \epsilon/4$ and $U_B = B(P_0, \delta) \times B(Q_0, \delta) \subseteq B$. Observe that U_B can be effectively obtained from B . We now show how a pair $(P_1, Q_1) \in U_B$ can be moved outside U_B , but still inside B , nearly without changing its sum. By the choice of δ , if $(P_1, Q_1) \in U_B$ then $d(P_1, Q_1) > 2\delta$. For $\lambda \in [0, 1]$, define

$$\begin{aligned} P(\lambda) &= \lambda P_1 + (1 - \lambda) Q_1, \\ Q(\lambda) &= \lambda Q_1 + (1 - \lambda) P_1. \end{aligned}$$

Observe that $P(\lambda) + Q(\lambda) = P_1 + Q_1$ and

$$d(P_1, P(\lambda)) = d(Q_1, Q(\lambda)) = (1 - \lambda)d(P_1, Q_1).$$

As $d(P_1, Q_1) > 2\delta$ there exists $\lambda \in (0, 1)$ such that $(1 - \lambda)d(P_1, Q_1) = 2\delta$. One has

$$d(P_0, P(\lambda)) \leq d(P_0, P_1) + d(P_1, P(\lambda)) < 3\delta < \epsilon$$

and

$$d(P_0, P(\lambda)) \geq d(P_1, P(\lambda)) - d(P_0, P_1) > \delta,$$

and similarly $\delta < d(Q_0, Q(\lambda)) < \epsilon$ so $(P(\lambda), Q(\lambda)) \in B \setminus \overline{U}_B$.

Observe that the shift-invariant measures $P(\lambda)$ and $Q(\lambda)$ are not ergodic. As the ergodic measures are dense in the set of shift-invariant measures, there exist two sequences P_n, Q_n of ergodic measures converging to $P(\lambda)$ and $Q(\lambda)$ respectively. As $(P(\lambda), Q(\lambda))$ belongs to the open set $B \setminus \overline{U}_B$, we can assume w.l.o.g. that $(P_n, Q_n) \in B \setminus \overline{U}_B$ for all n . The mapping $(P, Q) \mapsto P + Q$ is continuous so $P_n + Q_n$ converge to $P(\lambda) + Q(\lambda) = P_1 + Q_1$.

D Proof of Theorem 25

Let us slightly reformulate the three results in a unified way.

Let $X = 2^{\mathbb{N}}$ or the space $\text{CL}(2^{\mathbb{N}})$ of non-empty compact subsets of $2^{\mathbb{N}}$. Both are effective Polish spaces with the usual topologies. Let \leq be an ordering: inclusion or lexicographic ordering on $2^{\mathbb{N}}$, reverse inclusion on $\text{CL}(2^{\mathbb{N}})$. Let X_0 be the sequence of simple points. \leq is computable on X_0 . It is important to observe that given a clopen set $C \subseteq X$ and $s \in X_0$, it is decidable whether C intersects $\uparrow s$.

We say that $x \in X$ is \leq -c.e. if the set $\{s \in X_0 : s < x\}$ is c.e., or equivalently x is the supremum of a computable increasing sequence of simple points. We say that x is \leq -generic if for every effective open set $W \subseteq X$, either $x \in W$ or there exists $\epsilon > 0$ such that $B(x, \epsilon) \cap W \cap \uparrow x = \emptyset$.

► **Theorem 27.** *Let $U_n \subseteq X$ be uniformly effective dense open sets. There exists $x \in \bigcap_n U_n$ which is \leq -c.e. and \leq -generic.*

The notions of generic c.e. set, generic left-c.e. real and generic Π_1^0 -class correspond to the notions of \leq -c.e. and \leq -generic points, when \leq is the corresponding ordering. Being a co-infinite subset of \mathbb{N} or being a closed set without isolated points can both be expressed as an intersection of dense effective open sets in the corresponding space.

Proof. The proof is a finite injury argument. We want to satisfy the requirements

$$R_e : x \in W_e \text{ or } \exists \epsilon, B(x, \epsilon) \cap W_e \cap \uparrow x = \emptyset,$$

where W_e is the effective open set with number e (in the sequel, $W_{e,s}$ will be computable growing clopen sets with union W_e). At stage s , each requirement R_e is assigned a ball $B_e[s]$. They satisfy $\overline{B}_{e+1}[s] \subseteq B_e[s] \cap U_e$. For each e , the sequence $B_e[s]$ is stationary when s grows. At the same time, a computable sequence $x_s \leq x_{s+1}$ is built. The requirement R_e tests whether $B_e[s]$ intersects $W_e \cap \uparrow x_s$.

Stage 0. Start with any ball for $B_0[0]$ and inductively choose $\overline{B}_{e+1}[0] \subseteq B_e[0] \cap U_e$ of radius $< 2^{-e}$. x_0 is the center of $B_0[0]$.

Stage s . Let $e \leq s$ be minimal such that $B_e[s] \cap W_{e,s} \cap \uparrow x_s \neq \emptyset$, if it exists (decidable property). Let $\overline{B}_e[s+1] \subseteq W_{e,s} \cap \uparrow x_s \cap U_{e-1}$ and x_{s+1} be the center of $B_e[s+1]$. Define inductively $\overline{B}_{e'+1}[s+1] \subseteq B_{e'}[s+1] \cap U_{e'}$ of radius $< 2^{-e'}$ for $e' \geq e$.

If e does not exist then let $B_e[s+1] = B_e[s]$ for all e and $x_{s+1} = x_s$.

Verification. By the usual analysis of finite injury arguments, each requirement acts finitely many times, so for each e there is s_0 such that $B_e[s] = B_e[s_0]$ for all $s \geq s_0$. Let $B_e = B_e[s_0]$. One has $\overline{B_{e+1}} \subseteq B_e$. Let x be the unique member of $\bigcap_e B_e$.

► **Claim.** x is c.e. and belongs to $\bigcap_e U_e$.

By construction, $x \in B_{e+1} \subseteq U_e$ so $x \in \bigcap_e U_e$. The sequence x_s converges to x . As $x_s \leq x_{s+1}$, x_s also converges to $\sup_s x_s$, so $x = \sup_s x_s$. As the sequence x_s is computable, x is c.e.

► **Claim.** x is \leq -generic.

Assume $x \notin W_e$. Given e , let s be such that no requirement $e' < e$ acts from stage s on. R_e cannot act at a stage $s' \geq s$, otherwise $x \in B_e = B_e[s' + 1] \subseteq W_e$ which contradicts the assumption about x . As R_e never acts after stage s , it means that $x \in B_e = B_e[s]$ and $B_e[s] \cap W_e \cap \uparrow x = \emptyset$, otherwise there exists $s' \geq s$ such that $B_e[s] \cap W_{e,s'} \cap \uparrow x \neq \emptyset$, hence $B_e[s'] \cap W_{e,s'} \cap \uparrow x_{s'} \neq \emptyset$ as $B_e[s'] = B_e[s]$ and $x_{s'} \leq x$, and then R_e must act at stage s' which is impossible. ◀