



HAL
open science

Observing submesoscale activity in the Bay of Biscay with satellite-derived SST and Chlorophyll concentration

Özge Yelekçi, Guillaume Charria, Daneeja Mawren, Xavier Capet, Gilles Reverdin, Francis Gohin, Joël Sudre, Hussein Yahia

► To cite this version:

Özge Yelekçi, Guillaume Charria, Daneeja Mawren, Xavier Capet, Gilles Reverdin, et al.. Observing submesoscale activity in the Bay of Biscay with satellite-derived SST and Chlorophyll concentration. European Geosciences Union General Assembly 2014, Apr 2014, Vienne, Austria. hal-00942221

HAL Id: hal-00942221

<https://inria.hal.science/hal-00942221v1>

Submitted on 28 Oct 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Observing submesoscale activity in the Bay of Biscay with satellite-derived SST and Chlorophyll concentration

Ozge Yelekci (1), Guillaume Charria (1), Daneeja Mawren (1), Xavier Capet (2), Gilles Reverdin (2), Francis Gohin (3), Joël Sudre (4), and Hussein Yahia (5)

(1) IFREMER, DYNECO/PHYSED, Technôpole de Brest-Iroise, BP 70, F-29280 Plouzané, France (oyelekci@ifremer.fr), (2) LOCEAN/IPSL, CNRS/UPMC/IRD/MNHN, Paris, France, (3) IFREMER, DYNECO/PELAGOS, Technôpole de Brest-Iroise, BP 70, F-29280 Plouzané, France, (4) LEGOS/CNRS, UMR5566, 18 Avenue E. Belin, F-31401 Toulouse Cedex 4, France, (5) GeoStat team, INRIA Bordeaux Sud-Ouest, 351 Cours de La Libération, F-33405 Talence Cedex, France

Submesoscale activity in the upper ocean is linked to vertical velocities that enable transport of nutrients to the euphotic zone and trigger primary production, and yet its contribution to the global production budget is still to be fully investigated. Sampling the submesoscales is a challenge, due to their spatial scales of O(1-10) km and their life spans of a few days. Thus, we utilized high resolution (~ 1 km) satellite images of sea surface temperature (SST) and chlorophyll concentration (Chl) for the purpose of identification and classification of submesoscale activity over the Bay of Biscay continental shelf. For our analysis, we applied an event-based approach. We constructed a large data set of the last ten years' (2003 - 2013) MODIS images of both Aqua and Terra satellites. Due to cloud coverage, number of quality images covering the entire region is limited making the data set discontinuous in time and space. To maximize the portion of usable images covering the temporal and spatial scales we wanted to investigate, we discretized our data set. Given the fact that submesoscale features are highly localized, instead of the entirety of the Bay, we divided the domain into sub-regions that are defined by their hydrological features, such as river plumes. Then, we selected "events" of cloud-free, consecutive daily images of SST and corresponding Chl. For an understanding of the driving mechanisms of submesoscale activity and its outcome as enhanced production, together with satellite images, background environmental conditions such as turbidity, and forcing fields such as wind stress distribution and river runoff before and during these events were analyzed. As a quantitative measure, correlation (or lack thereof) between occurrence and strength of submesoscale features and these factors is presented. As a result of this effort, a classification of the submesoscale features based on their driving mechanisms is proposed.