



HAL
open science

”Trivializing” generalizations of some Izergin-Korepin-type determinants

Tewodros Amdeberhan, Doron Zeilberger

► **To cite this version:**

Tewodros Amdeberhan, Doron Zeilberger. ”Trivializing” generalizations of some Izergin-Korepin-type determinants. *Discrete Mathematics and Theoretical Computer Science*, 2007, Vol. 9 no. 1 (1), pp.203–206. 10.46298/dmtcs.402 . hal-00966504

HAL Id: hal-00966504

<https://inria.hal.science/hal-00966504>

Submitted on 26 Mar 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L’archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d’enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

“Trivializing” Generalizations of some Izergin-Korepin-type Determinants

Tewodros Amdeberhan¹ and Doron Zeilberger²

¹ Tulane Univ., Dept of Math., New Orleans, LA 70118, USA

² Rutgers Univ., Dept of Math., Piscataway, NJ 08854, USA

received May 8, 2007, revised June 13, 2007; July 31, 2007, accepted July 31, 2007.

We generalize (and hence trivialize and routinize) numerous explicit evaluations of determinants and pfaffians due to Kuperberg, as well as a determinant of Tsuchiya. The level of generality of our statements render their proofs easy and routine, by using Dodgson condensation and/or Krattenthaler’s factor exhaustion method.

All our matrices will be assumed to be embedded inside an infinite matrix.

The first theorem adds *parameters* to the determinant formulas found in Kuperberg [Ku, Theorem 15], as well as older determinants, mentioned there, due to Cauchy, Stembridge, Laksov–Lascoux–Thorup, and Tsuchiya [T]. This way, the formulation is suited to the method of [AZ]. Our proofs are much more **succinct** and **automatable**, since their generality enables an easy induction using Dodgson’s rule [D, AZ], or by employing Krattenthaler’s elegant factor exhaustion method [Kr1]. Relevant background for this paper can be found in [Ku], and references thereof.

Theorem 1

$$\det \left(\frac{1}{x_i + y_j + Ax_i y_j} \right)_{i,j}^{1,n} = \frac{\prod_{i < j} (x_j - x_i)(y_j - y_i)}{\prod_{i,j}^n (x_i + y_j + Ax_i y_j)}.$$

$$\det \left(\frac{1}{x_i + y_j} - \frac{1}{1 + x_i y_j} \right)_{i,j}^{1,n} = \frac{\prod_{i < j} (1 - x_i x_j)(1 - y_i y_j)(x_j - x_i)(y_j - y_i)}{\prod_{i,j}^n (x_i + y_j)(1 + x_i y_j)} \cdot \prod_{i=1}^n (1 - x_i)(1 - y_i).$$

$$\det \left(\frac{Ay_j + Bx_i}{y_j + x_i} \right)_{i,j}^{1,n} = (A - B)^{n-1} \frac{\left(A \prod_j^n y_j + (-1)^{n-1} B \prod_i^n x_i \right) \prod_{i < j} (x_i - x_j)(y_i - y_j)}{\prod_{i,j}^n (x_i + y_j)}.$$

$$\det \left(\frac{1 - x_i y_j}{y_j - x_i} \right)_{i,j}^{1,n} = (-1)^{\binom{n}{2}} \frac{\left(\prod_j^n (1 + x_j)(1 - y_j) + \prod_i^n (1 - x_i)(1 + y_i) \right) \prod_{i < j} (x_i - x_j)(y_i - y_j)}{2 \prod_{i,j}^n (y_j - x_i)}.$$

Sketch of Proof: An automatic application of Dodgson Condensation [D, AZ]. \square

Corollary 1 (Cauchy, Stembridge, Laksov–Lascoux–Thorup)

$$\det \left(\frac{1}{x_i + y_j} \right)_{i,j}^{1,n} = \frac{\prod_{i < j} (x_j - x_i)(y_j - y_i)}{\prod_{i,j}^n (x_i + y_j)}.$$

$$\text{Pf}^2 = \det \left(\frac{x_j - x_i}{x_j + x_i} \right)_{i,j}^{1,2n} = \prod_{i < j \leq 2n} \frac{(x_i - x_j)^2}{(x_i + x_j)^2}.$$

$$\text{Pf}^2 = \det \left(\frac{x_j - x_i}{1 - x_i x_j} \right)_{i,j}^{1,2n} = \prod_{i < j \leq 2n} \frac{(x_i - x_j)^2}{(1 - x_i x_j)^2}.$$

Remark 1 Notice that the latter two statements apply only to even-dimensional matrices. An error from [Ku] in the second formula has been corrected here. Pf stands for Pfaffian of a matrix.

The next theorem generalizes, and presents variations of, several of the determinants that appear in Theorems 16 and 17 [Ku] (typos corrected in [Kr2, Theorems 13 and 14]). Below, Z_1, Z_2, Z_3, Z_4, Z_5 are defined by (here $\gamma(a, b) = a - b, \tau(a, b) = a + b$)

$$Z_1(p, q; x, y)_{i,j} = \frac{\gamma(q^{j-i}, x^{j-i})}{\tau(p^{j-i}, y^{j-i})}, \quad Z_2(p, q; x, y)_{i,j} = \frac{\gamma(q^{1+j-i}, x^{1+j-i})}{\tau(p^{1+j-i}, y^{1+j-i})},$$

$$Z_3(p, q; x, y)_{i,j} = \frac{\gamma(q^{-1+j-i}, x^{-1+j-i})}{\tau(p^{-1+j-i}, y^{-1+j-i})}, \quad Z_4(q, q; x, x)_{i,j} = \frac{\gamma(q^{a+j-i}, x^{a+j-i})}{\tau(q^{a+j-i}, x^{a+j-i})},$$

$$Z_5(q, q; x, x)_{i,j} = \frac{\tau(q^{b+j-i}, x^{b+j-i})}{\gamma(q^{b+j-i}, x^{b+j-i})},$$

for $a \in \mathbb{Z}, b = \pm n, \pm(n+1), \dots$. Let $\delta_{e,n} = \frac{1+(-1)^n}{2}$ denote Kronecker's delta function centered at the even integers, and let $\lambda_{i,j} = 1$, if $i \neq j$ and $\lambda_{i,i} = 0$.

Theorem 2 Write $\gamma(a, b) = a - b, \tau(a, b) = a + b$, we have the matrix determinants

$$\det \left(\frac{\gamma(q^{n+j-i}, x^{n+j-i})}{\gamma(p^{n+j-i}, y^{n+j-i})} \right)_{i,j}^{1,n} = (py)^{\binom{n}{3}} \gamma(q, x)^n \frac{\prod_{j>i} \gamma(p^{j-i}, y^{j-i})^2 \gamma(qp^{j-i}, xy^{j-i}) \gamma(xp^{j-i}, qy^{j-i})}{\prod_{i,j} \gamma(p^{n+j-i}, y^{n+j-i})}.$$

$$\det \left(\frac{\tau(q^{j-i}, x^{j-i})}{\tau(p^{j-i}, y^{j-i})} \right)_{i,j}^{1,n} = \frac{\prod_{2|j-i>0} \gamma(p^{j-i}, y^{j-i})^2 \prod_{2|j-i>0} \gamma(qp^{j-i}, xy^{j-i}) \gamma(xp^{j-i}, qy^{j-i})}{(qx)^{\lfloor n^2/4 \rfloor} \prod_{j>i} \tau(p^{j-i}, y^{j-i})^2}.$$

$$\det(Z_1)_{i,j}^{1,n} = \delta_{e,n} \frac{\gamma(q, x)^n (py)^{n/2} \prod_{2|j-i>0} \gamma(p^{j-i}, y^{j-i})^2 \gamma(qp^{j-i}, xy^{j-i}) \gamma(xp^{j-i}, qy^{j-i})}{(qx)^{\lfloor n^2/4 \rfloor} \prod_{j>i} \tau(p^{j-i}, y^{j-i})^2}.$$

$$\det(Z_2)_{i,j}^{1,n} = \frac{\gamma(q, x)^n \prod_{2|j-i>0} \gamma(p^{j-i}, y^{j-i})^2 \gamma(qp^{j-i}, xy^{j-i}) \gamma(xp^{j-i}, qy^{j-i})}{(qx)^{\lfloor (n-1)^2/4 \rfloor} \tau(p^n, y^n)^{1-\delta_{e,n}} \prod_{j>i} \tau(p^{j-i}, y^{j-i})^2}.$$

$$\det(Z_3)_{i,j}^{1,n} = \frac{(-py)^n \gamma(q, x)^n \prod_{2|j-i>0} \gamma(p^{j-i}, y^{j-i})^2 \gamma(qp^{j-i}, xy^{j-i}) \gamma(xp^{j-i}, qy^{j-i})}{(qx)^{\lfloor (n+1)^2/4 \rfloor} \tau(p^n, y^n)^{1-\delta_{e,n}} \prod_{j>i} \tau(p^{j-i}, y^{j-i})^2}.$$

$$\det(Z_4)_{i,j}^{1,n} = 2^{n-1} \frac{q^{na} + (-1)^n x^{na}}{(qx)^{n(n-1)(n+1-3a)/6}} \frac{\prod_{j \neq i} \gamma(q^{|j-i|}, x^{|j-i|})}{\prod_{i,j} \tau(q^{a+j-i}, x^{a+j-i})}.$$

$$\det(Z_5)_{i,j}^{1,n} = (-1)^{\binom{n}{2}} 2^{n-1} \frac{q^{nb} + x^{nb}}{(qx)^{n(n-1)(n+1-3b)/6}} \frac{\prod_{j \neq i} \gamma(q^{|j-i|}, x^{|j-i|})}{\prod_{i,j} \gamma(q^{b+j-i}, x^{b+j-i})}.$$

$$\text{Pf}^2 = \det \left(\lambda_{i,j} \frac{\gamma(q^{j-i}, x^{j-i}) \gamma(r^{j-i}, z^{j-i})}{\gamma(p^{j-i}, y^{j-i})} \right)_{i,j}^{1,2n} = \frac{(yp)^n}{(qxrz)^{n^2}} \frac{\prod_{j \neq i}^{1,n} \gamma(p^{|j-i|}, y^{|j-i|})^2}{\prod_{i,j}^{1,n} \gamma(p^{n+j-i}, y^{n+j-i})^2} \times$$

$$\prod_{i,j}^{1,n} \gamma(qp^{|j-i|}, xy^{|j-i|}) \gamma(xp^{|j-i|}, qy^{|j-i|}) \gamma(rp^{|j-i|}, zy^{|j-i|}) \gamma(zp^{|j-i|}, ry^{|j-i|}).$$

Sketch of Proof: Identities Z_4 and Z_5 are directly amenable to Dodgson’s Condensation technique [AZ]. For the remaining assertions, use the factor exhaustion method [Kr1] (see also [Ku]): the essential idea is to compare zeros and poles on both sides of the equation at hand. We leave the straightforward details to the reader. \square

Acknowledgments

The authors wish to thank Christian Krattenthaler for helpful suggestions.

References

- [AZ] T. Amdeberhan, D. Zeilberger, *Determinants Through the Looking Glass*, Adv. Appl. Math. (Foata special issue) 2/3 **27** (2001), 225–230.
- [D] C.L. Dodgson, *Condensation of Determinants*, Proceedings of the Royal Society of London **15** (1866), 150–155.
- [Kr1] C. Krattenthaler, *Advanced determinant calculus*, Séminaire Lotharingien Combin. **42** (1999), Art. B42q.
- [Kr2] C. Krattenthaler, *Advanced determinant calculus: A Complement*, Linear Alg. Appl. **411** (2005), 68–166.
- [Ku] G. Kuperberg, *Symmetry classes of alternating-sign matrices under one roof*, Ann. of Math. (2) **156** (2002), 835–866.
- [T] O. Tsuchiya, *Determinant formula for the six-vertex model with reflecting end*, J. Math. Phys. **39** (1998), 5946–5951