



HAL
open science

Border Basis relaxation for polynomial optimization

Marta Abril Bucero, Bernard Mourrain

► **To cite this version:**

Marta Abril Bucero, Bernard Mourrain. Border Basis relaxation for polynomial optimization. 2014.
hal-00981546v1

HAL Id: hal-00981546

<https://inria.hal.science/hal-00981546v1>

Preprint submitted on 22 Apr 2014 (v1), last revised 21 Aug 2015 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

BORDER BASIS RELAXATION FOR POLYNOMIAL OPTIMIZATION

MARTA ABRIL BUCERO, BERNARD MOURRAIN

ABSTRACT. A relaxation method based on border basis reduction which improves the efficiency of Lasserre's approach is proposed to compute the optimum of a polynomial function on a basic closed semi algebraic set. A new stopping criterion is given to detect when the relaxation sequence reaches the minimum, using a sparse flat extension criterion. We also provide a new algorithm to reconstruct a finite sum of weighted Dirac measures from a truncated sequence of moments, which can be applied to other sparse reconstruction problems. As an application, we obtain a new algorithm to compute zero-dimensional minimizer ideals and the minimizer points or zero-dimensional G-radical ideals. Experimentations show the impact of this new method on significant benchmarks.

1. INTRODUCTION

Computing the global minimum of a polynomial function f on a semi-algebraic set is a difficult but important problem, with many applications. A relaxation approach was proposed in [11] (see also [27], [31]) which approximates this problem by a sequence of finite dimensional convex optimization problems. These optimization problems can be formulated in terms of linear matrix inequalities on moment matrices associated to the set of monomials of degree $\leq t \in \mathbb{N}$ for increasing values of t . They can be solved by Semi-Definite Programming (SDP) techniques. The sequence of minima converges to the actual minimum f^* of the function under some hypotheses [11]. In some cases, the sequence even reaches the minimum f^* in a finite number of steps [15, 26, 18, 3, 8, 24]. This approach proved to be particularly fruitful in many problems [12]. In contrast with numerical methods such as gradient descent methods, which converge to a local extremum but with no guaranty for the global solution, this relaxation approach can provide certificates for the minimum value f^* in terms of sums of squares representations.

From an algorithmic and computational perspective, some issues need however to be considered.

The size of the SDP problems to be solved is a bottleneck of the method. This size is related to the number of monomials of degree $\leq t$ and is increasing exponentially with the number of variables and the degree t . Many SDP solvers are based on interior point methods which provide an approximation of the optimal moment sequence within a given precision in a polynomial time: namely $\mathcal{O}((ps^{3.5} + cp^2s^{2.5} + cp^3s^{0.5}) \log(\epsilon^{-1}))$ arithmetic operations where $\epsilon > 0$ is the precision of the approximation, s is bounding the size of the moment matrices, p is the number of parameters (usually of the order s^2) and c is the number of constraints [23]. Thus reducing the size s or the number of parameters p can improve significantly the performance of these relaxation methods. Some recent works address this issue, using symmetries (see e.g. [29]) or polynomial reduction (see e.g. [10]). In this paper, we extend this latter approach.

While determining the minimum value of a polynomial function on a semi-algebraic set is important, computing the points where this minimum is reached if they exist, is also critical in many applications. Determining when and how these minimizer points can be computed from the relaxation sequence is a problem that has been adressed for instance

in [9, 25] using full moment matrices. This approach has been used for solving polynomial equations [15, 13, 14, 12].

As an alternative approach, the optimization problem can be reformulated as solving polynomial equations related to the (minimal) critical value of the polynomial f on a semi-algebraic set. Polynomial solvers based for instance on Gröbner basis or border basis computation can then be used to recover the real critical points from the complex solutions of (zero-dimensional) polynomial systems (see e.g. [28, 30, 7]). This type of methods relies entirely on polynomial algebra and univariate root finding. So far, there is no clear comparison of these elimination methods and the relaxation approaches.

Contributions. We propose a new method which combines Lasserre’s SDP relaxation approach with polynomial algebra, in order to increase the efficiency of the optimization algorithm. Border basis computations are considered for their numerical stability [20, 21]. In principle, any graded normal form techniques could be used here.

A new stopping criterion is given to detect when the relaxation sequence reaches the minimum, using a flat extension criterion from [16]. We also provide a new algorithm to reconstruct a finite sum of weighted Dirac measures from a truncated sequence of moments. This reconstruction method can be used in other problems such as tensor decomposition [2] and multivariate sparse interpolation [6].

As shown in [1, 26, 3, 18, 24, 8], an exact SDP relaxation can be constructed for “well-posed” optimization problems. As an application, we obtain a new algorithm to compute zero-dimensional minimizer ideals and the minimizer points, or zero-dimensional G-radical. Experimentations show the impact of this new method compared to the previous relaxation constructions.

Content. The paper is organized as follows. Section 2 describes the minimization problem, the SDP relaxation hierarchies and the concept of optimal linear form. Section 3 describes new algorithms to check exact relaxations and to compute the minimizers for zero-dimensional minimizer ideals. Section 4 analyzes cases for which an exact relaxation can be constructed. Section 5 concludes with the description of the minimization algorithm and experimentation.

Notation. Let $\mathbb{K}[\mathbf{x}]$ be the set of the polynomials in the variables $\mathbf{x} = (x_1, \dots, x_n)$, with coefficients in the field \mathbb{K} . Hereafter, we will choose $\mathbb{K} = \mathbb{R}$ or \mathbb{C} . Let $\overline{\mathbb{K}}$ denotes the algebraic closure of \mathbb{K} . For $\alpha \in \mathbb{N}^n$, $\mathbf{x}^\alpha = x_1^{\alpha_1} \cdots x_n^{\alpha_n}$ is the monomial with exponent α and degree $|\alpha| = \sum_i \alpha_i$. The set of all monomials in \mathbf{x} is denoted $\mathcal{M} = \mathcal{M}(\mathbf{x})$. For a polynomial $f = \sum_\alpha f_\alpha \mathbf{x}^\alpha$, its support is $\text{supp}(f) := \{\mathbf{x}^\alpha \mid f_\alpha \neq 0\}$, the set of monomials occurring with a nonzero coefficient in f .

For $t \in \mathbb{N}$ and $F \subseteq \mathbb{K}[\mathbf{x}]$, we introduce the following sets: F_t is the set of elements of F of degree $\leq t$; $\langle F \rangle = \{ \sum_{f \in F} \lambda_f f \mid f \in F, \lambda_f \in \mathbb{K} \}$ is the linear span of F ; if F is a vector space, F^* is the dual space of linear forms from F to \mathbb{K} ; $(F) = \{ \sum_{f \in F} p_f f \mid p_f \in \mathbb{K}[\mathbf{x}], f \in F \}$ is the ideal in $\mathbb{K}[\mathbf{x}]$ generated by F ; $\langle F \mid t \rangle$ is the vector space spanned by $\{ \mathbf{x}^\alpha f \mid f \in F_t, |\alpha| \leq t - \deg(f) \}$; $F \cdot F := \{ p q \mid p, q \in F \}$; $\Sigma^2(F) = \{ \sum_{i=1}^s f_i^2 \mid f_i \in F \}$ is the set of finite sums of squares of elements of F ; for $F = \{ f_1, \dots, f_m \} \subset \mathbb{K}[\mathbf{x}]$, $\prod(F) = \{ \prod_{i=1}^m f_i^{\epsilon_i} \mid \epsilon_i \in \{0, 1\} \}$.

Hereafter, we will consider $\mathbb{K} = \mathbb{R}$.

2. MINIMIZATION PROBLEM

Let $f, g_1^0, \dots, g_{n_1}^0, g_1^+, \dots, g_{n_2}^+ \in \mathbb{R}[\mathbf{x}]$ be polynomial functions. The minimization problem that we consider all along the paper is the following: compute

$$(1) \quad \begin{aligned} & \inf_{\mathbf{x} \in \mathbb{R}^n} f(\mathbf{x}) \\ \text{s.t. } & g_1^0(\mathbf{x}) = \dots = g_{n_1}^0(\mathbf{x}) = 0 \\ & g_1^+(\mathbf{x}) \geq 0, \dots, g_{n_2}^+(\mathbf{x}) \geq 0 \end{aligned}$$

More precisely, the objectives of the method we are going to describe are to compute the minimum value when f is bounded by below and the points where this minimum value is reached if they exist.

A set of constraints $\mathbf{g} = \{g_1^0, \dots, g_{n_1}^0; g_1^+, \dots, g_{n_2}^+\} \in \mathbb{R}[\mathbf{x}]$ is the union of a finite subset $\mathbf{g}^0 = \{g_1^0, \dots, g_{n_1}^0\}$ of polynomials corresponding to the equality constraints and a finite subset $\mathbf{g}^+ = \{g_1^+, \dots, g_{n_2}^+\}$ corresponding to the non-negativity constraints. We say that $\mathbf{g} \subset \mathbf{g}'$ if $\mathbf{g}^0 \subset \mathbf{g}'^0$ and $\mathbf{g}^+ \subset \mathbf{g}'^+$.

The basic semi-algebraic set defined by the constraints \mathbf{g} will be denoted $\mathcal{S}(\mathbf{g}) = \{\mathbf{x} \in \mathbb{R}^n \mid g_1^0(\mathbf{x}) = \dots = g_{n_1}^0(\mathbf{x}) = 0, g_1^+(\mathbf{x}) \geq 0, \dots, g_{n_2}^+(\mathbf{x}) \geq 0\}$. Hereafter, $S := \mathcal{S}(\mathbf{g})$ will correspond to the basic semi-algebraic set defined by the constraints of our minimisation problem (1). We assume that $S \neq \emptyset$.

When $n_1 = n_2 = 0$, there is no constraint and $S = \mathbb{R}^n$. In this case, we are considering a global minimization problem.

The points $\mathbf{x}^* \in \mathbb{R}^n$ which satisfies $f(\mathbf{x}^*) = \inf_{\mathbf{x} \in S} f(\mathbf{x})$ are called the *minimizers* of f on S . The set of minimizers is denoted $V_{min} = \{\mathbf{x}^* \in S \text{ s.t. } f(\mathbf{x}^*) = f^*\}$. The ideal of $\mathbb{R}[\mathbf{x}]$ defining the set V_{min} is denoted I_{min} .

If the set of minimizers is not empty, we say that the *minimization problem is feasible*. The minimization problem is not feasible means that $V_{min} = \emptyset$ and $I_{min} = \mathbb{R}[\mathbf{x}]$.

2.1. Hierarchies of relaxation problems. In this section, we describe the finite dimensional convex optimization problems that we consider.

Definition 2.1. Given a finite dimensional vector space $E \subset \mathbb{R}[\mathbf{x}]$ and a set of constraints $G \subset \mathbb{R}[\mathbf{x}]$, we define the quadratic module of G on E as

$$\mathcal{Q}_{E,G} = \left\{ \sum_{g \in G^0} g h + \sum_{g' \in G^+} g' h' \mid h \in E, g h \in \langle E \cdot E \rangle, h' \in \Sigma^2(E), g' h' \in \langle E \cdot E \rangle \right\}.$$

If $G^* \subset \mathbb{R}[\mathbf{x}]$ is the set of constraints such that $G^{*0} = G^0$ and $G^{*+} = \prod(G^+)$, the (truncated) quadratic module \mathcal{Q}_{E,G^*} is called the (truncated) preordering of G and denoted $\mathcal{Q}_{E,G}^*$ or $\mathcal{P}_{E,G}$.

By construction, $\mathcal{Q}_{E,G} \subset \langle E \cdot E \rangle$ is a cone of polynomials which are non-negative on the semi-algebraic set S .

We consider now its dual cone.

Definition 2.2. Given a finite dimensional vector space $E \subset \mathbb{R}[\mathbf{x}]$ which contains 1 and a set of constraints $G \subset \mathbb{R}[\mathbf{x}]$, we define

$$\mathcal{L}_{E,G} := \{ \Lambda \in \langle E \cdot E \rangle^* \mid \Lambda(p) \geq 0, \forall p \in \mathcal{Q}_{E,G}, \Lambda(1) = 1 \}.$$

The convex set associated to the preordering $\mathcal{Q}_{E,G}^* = \mathcal{Q}_{E,G^*}$ is denoted $\mathcal{L}_{E,G}^*$.

By this definition, for any element $\Lambda \in \mathcal{L}_{E,G}$ and any $g \in \langle G^0 \rangle \cap E$, we have $\Lambda(g) = 0$.

We introduce now truncated Hankel operators, which will play a central role in the construction of the minimizer ideal of f on S .

Definition 2.3. For a linear form $\Lambda \in \langle E \cdot E \rangle^*$, we define the map $H_\Lambda^E : E \rightarrow E^*$ by $H_\Lambda^E(p)(q) = \Lambda(pq)$ for $p, q \in E$. Thus H_Λ^E is called a truncated Hankel operator, defined on the subspace E .

Its matrix in the monomial and dual bases of E and E^* is usually called the moment matrix of Λ . The kernel of this truncated Hankel operator will be used to compute generators of the minimizer ideal, as we will see.

Definition 2.4. Let $E \subset \mathbb{R}[\mathbf{x}]$ such that $1 \in E$ and a set of constraints $G \subset \mathbb{R}[\mathbf{x}]$. We define the following extrema:

- $f^* = \inf_{\mathbf{x} \in S} f(\mathbf{x})$,
- $f_{E,G}^\mu = \inf \{ \Lambda(f) \text{ s.t. } \Lambda \in \mathcal{L}_{E,G} \}$,
- $f_{E,G}^{sos} = \sup \{ \gamma \in \mathbb{R} \text{ s.t. } f - \gamma \in \mathcal{Q}_{E,G} \}$.

By convention if the sets are empty, $f_{E,G}^{sos} = -\infty$ and $f_{E,G}^\mu = +\infty$.

If $E = \mathbb{R}[\mathbf{x}]_t$ and $G^0 = \langle \mathbf{g}^0 \mid 2t \rangle$, we also denote $f_{E,G}^\mu$ by $f_{t,G}^\mu$ and $f_{E,G}^{sos}$ by $f_{t,G}^{sos}$.

We easily check that $f_{E,G}^{sos} \leq f_{E,G}^\mu$, since if there exists $\gamma \in \mathbb{R}$ such that $f - \gamma = q \in \mathcal{Q}_{E,G}$ then $\forall \Lambda \in \mathcal{L}_{E,G}$, $\Lambda(f - \gamma) = \Lambda(f) - \gamma = \Lambda(q) \geq 0$.

If $S(G) \subset S$, we also have $f_{E,G}^\mu \leq f^*$ since for any $\mathbf{s} \in S$, the evaluation $\mathbf{1}_\mathbf{s} : p \in \mathbb{R}[\mathbf{x}] \mapsto p(\mathbf{s})$ is in $\mathcal{L}_{E,G}$.

Notice that if $E \subset E'$, $G \subset G'$ then $\mathcal{Q}_{E,G} \subset \mathcal{Q}_{E',G'}$, $\mathcal{L}_{E',G'} \subset \mathcal{L}_{E,G}$, $f_{E,G}^\mu \leq f_{E',G'}^\mu$ and $f_{E,G}^{sos} \leq f_{E',G'}^{sos}$.

2.2. Full moment matrix relaxation hierarchy. The relaxation hierarchies introduced in [11] correspond to the case where $E = \mathbb{R}[\mathbf{x}]_t$, $G^0 = \langle \mathbf{g}^0 \mid 2t \rangle$ and $G^+ = \mathbf{g}^+$.

The quadratic module $\mathcal{Q}_{\mathbb{R}[\mathbf{x}]_t, G}$ is denoted $\mathcal{Q}_{t,\mathbf{g}}$ and $\mathcal{L}_{\mathbb{R}[\mathbf{x}]_t, G}$ is denoted $\mathcal{L}_{t,\mathbf{g}}$. Hereafter, we will also call Lasserre hierarchy, the full moment matrix relaxation hierarchy. It corresponds to the sequences

$$\cdots \subset \mathcal{L}_{t+1,\mathbf{g}} \subset \mathcal{L}_{t,\mathbf{g}} \subset \cdots \text{ and } \cdots \subset \mathcal{Q}_{t,\mathbf{g}} \subset \mathcal{Q}_{t+1,\mathbf{g}} \subset \cdots$$

which yield the following increasing sequences for $t \in \mathbb{N}$:

$$\cdots f_{t,\mathbf{g}}^\mu \leq f_{t+1,\mathbf{g}}^\mu \leq \cdots \leq f^* \text{ and } \cdots f_{t,\mathbf{g}}^{sos} \leq f_{t+1,\mathbf{g}}^{sos} \leq \cdots \leq f^*.$$

The foundation of Lasserre's method is to show that these sequences converge to f^* . This is proved under some conditions in [11]. It has also been shown that the limit can even be reached in a finite number of steps in some cases, see e.g. [14, 26, 18, 8, 24, 1]. In this case, the relaxation is said to be *exact*.

2.3. Border basis relaxation hierarchy. In the following we are going to use another type of relaxation ‘‘hierarchies’’, which involves border basis computation. It is aimed at reducing the size of the convex optimization problems solved at each level of the relaxation hierarchy. As we will see in Section 5.2, the impact on the performance of the relaxation approach is significant. We briefly recall the properties of border basis that we need and describe how they are used in the construction of this relaxation hierarchy.

Given a vector space $E \subseteq \mathbb{R}[\mathbf{x}]$, its prolongation $E^+ := E + x_1 E + \dots + x_n E$ is again a vector space.

The vector space E is said to be *connected to 1* if $1 \in E$ and there exists a finite increasing sequence of vector spaces $E_0 \subset E_1 \subset \dots \subset E$ such that $E_0 = \langle 1 \rangle$, $E_{i+1} \subset E_i^+$. For a monomial set $B \subseteq \mathcal{M}$, $B^+ = B \cup x_1 B \cup \dots \cup x_n B$ and $\partial B = B^+ \setminus B$. We easily check that $\langle B \rangle^+ = \langle B^+ \rangle$ and $\langle B \rangle$ is connected to 1 iff $1 \in B$ and for every monomial $m \neq 1$ in B , $m = x_{i_0} m'$ for some $i_0 \in [1, n]$ and some monomial $m' \in B$.

Definition 2.5. Let $B \subset \mathcal{M}$ be connected to 1. A family $F \subset \mathbb{R}[\mathbf{x}] = R$ is a border basis for B in degree $t \in \mathbb{N}$, if $\forall f, f' \in F_t$,

- $\text{supp}(f) \subseteq B^+ \cap R_t$,
- f has exactly **one** monomial in ∂B , denoted $\gamma(f)$ and called the leading monomial of f .
- $\gamma(f) = \gamma(f')$ implies $f = f'$,
- $\forall m \in \partial B \cap R_t, \exists f \in F$ s.t. $\gamma(f) = m$,
- $R_t = \langle B \rangle_t \oplus \langle F|t \rangle$.

A border basis F for B in all degrees t is called a border basis for B . F is graded if moreover $\deg(\gamma(f)) = \deg(f) \forall f \in F$.

There are efficient algorithms to check that a given family F is a border basis for B in degree t and to construct such family from a set of polynomials. We refer to [19, 20, 21, 22] for more details. We will use these tools as “black boxes” in the following.

For a border basis F for B in degree t , we denote by $\pi_{F,B}$ the projection of R_t on $\langle B \rangle_t$ along $\langle F|t \rangle$. We easily check that

- $\forall m \in B_t, \pi_{B,F}(m) = m$,
- $\forall m \in \partial B \cap R_t, \pi_{B,F}(m) = m - f$, where f is the (unique) polynomial in F for which $\gamma(f) = m$, assuming the polynomials $f \in F$ are normalized so that the coefficient of $\gamma(f)$ is 1.

If F is a graded border basis in degree t , one easily verifies that $\deg(\pi_{F,B}(m)) \leq \deg(m)$ for $m \in \mathcal{M}_t$.

Border basis hierarchy. The sequence of relaxation problems that we will use hereafter is defined as follows. For each $t \in \mathbb{N}$, we construct the graded border basis F_{2t} of \mathbf{g}^0 in degree $2t$. Let B be the set of monomials (connected to 1) for which F is a border basis in degree $2t$. We define $E_t := \langle B_t \rangle$, G_t is the set of constraints such that $G_t^0 = \{m - \pi_{B_t, F_{2t}}(m), m \in B_t \cdot B_t\}$ and $G_t^+ = \pi_{B_t, F_{2t}}(\mathbf{g}^+)$, and consider the relaxation sequence

$$(2) \quad \mathcal{Q}_{E_t, G_t} \subset \langle B_t \cdot B_t \rangle \text{ and } \mathcal{L}_{E_t, G_t} \subset \langle B_t \cdot B_t \rangle^*$$

for $t \in \mathbb{N}$. Since the subsets B_t are not necessarily nested, these convex sets are not necessarily included one in the other one. However, by construction of the graded border basis of \mathbf{g} , we have the following inclusions

$$\cdots \subset \langle F_{2t}|2t \rangle \subset \langle F_{2t+2}|2t+2 \rangle \subset \cdots (\mathbf{g}^0),$$

and we can relate the border basis relaxation sequences with the corresponding full moment matrix relaxation hierarchy, using the following proposition:

Proposition 2.6. Let $t \in \mathbb{N}$, $B \subset \mathbb{R}[\mathbf{x}]_{2t}$ be a monomial set connected to 1, $F \subset \mathbb{R}[\mathbf{x}]$ be a border basis for B in degree $2t$, $E := \langle B_t \rangle$, $E' := \mathbb{R}[\mathbf{x}]_t$, G, G' be sets of constraints such that $G^0 = \{m - \pi_{B_t, F}(m), m \in B_t \cdot B_t\}$, $G'^0 = \langle F|2t \rangle$, $G^+ = G'^+$. Then for all $\Lambda \in \mathcal{L}_{E, G}$, there exists a unique $\Lambda' \in \mathcal{L}_{E', G'}$ which extends Λ . Moreover, Λ' satisfies $\text{rank } H_{\Lambda'}^{E'} = \text{rank } H_{\Lambda}^E$ and $\ker H_{\Lambda'}^{E'} = \ker H_{\Lambda}^E + \langle F|t \rangle$.

Proof. As $F \subset \mathbb{R}[\mathbf{x}]$ is a border basis for B in degree $2t$, we have $\mathbb{R}[\mathbf{x}]_{2t} = \langle B \rangle_{2t} \oplus \langle F|2t \rangle$. As $\langle B_t \cdot B_t \rangle \subset \langle B \rangle_{2t} \oplus \langle G^0 \rangle$, $\langle G^0 \rangle \subset \langle G'^0 \rangle = \langle F|2t \rangle$ and $\mathbb{R}[\mathbf{x}]_{2t} = \langle B \rangle_{2t} \oplus \langle F|2t \rangle$, we deduce that for all $\Lambda \in \mathcal{L}_{E, G}$, there exists a unique $\Lambda' \in \mathbb{R}[\mathbf{x}]_{2t}^*$ s.t. $\Lambda'_{|\langle B \rangle_{2t}} = \Lambda$ and $\Lambda'(\langle F|2t \rangle) = 0$.

Let us first prove that $\Lambda' \in \mathcal{L}_{E', G'} = \mathcal{L}_{t, G'}$. As any element q' of $\mathcal{Q}_{E', G'}$ can be decomposed as a sum of an element q of $\mathcal{Q}_{E, G}$ and an element $p \in \langle F|2t \rangle$, we have $\Lambda'(q') = \Lambda'(q) + \Lambda'(p) = \Lambda(q) \geq 0$. This shows that $\Lambda' \in \mathcal{L}_{E', G'}$.

Let us prove now that $\ker H_{\Lambda'}^{E'} = \ker H_{\Lambda}^E + \langle F | t \rangle$ where $E := \langle B_t \rangle$, $E' := \mathbb{R}[\mathbf{x}]_t$. As $E \cdot \langle F | t \rangle \subset \langle F | 2t \rangle = G^{t_0}$, we have $\Lambda'(E \cdot \langle F | t \rangle) = 0$ so that

$$(3) \quad \langle F | t \rangle \subset \ker H_{\Lambda'}^{E'}.$$

For any element $b \in \ker H_{\Lambda}^E$ we have $\forall b' \in E$, $\Lambda(b b') = \Lambda'(b b') = 0$. As $\Lambda'(E \cdot \langle F | t \rangle) = 0$ and $E' = E \oplus \langle F | t \rangle$, for any element $e \in E$, $\Lambda'(b e) = 0$. This proves that

$$(4) \quad \ker H_{\Lambda}^E \subset \ker H_{\Lambda'}^{E'}.$$

Conversely as $E' = E \oplus \langle F | t \rangle$, any element of E' can be reduced modulo $\langle F | t \rangle$ to an element of E , which shows that

$$(5) \quad \ker H_{\Lambda'}^{E'} \subset \ker H_{\Lambda}^E + \langle F | t \rangle.$$

From the inclusions (3), (4) and (5), we deduce that $\ker H_{\Lambda'}^{E'} = \ker H_{\Lambda}^E + \langle F | t \rangle$ and that $\text{rank } H_{\Lambda'}^{E'} = \text{rank } H_{\Lambda}^E$. \square

We deduce from this proposition that $f_{E_t, G_t}^{\mu} = f_{t, \langle F_{2t} | 2t \rangle}^{\mu}$. The sequence of convex sets \mathcal{L}_{E_t, G_t} can be seen as the projections of nested convex sets

$$\cdots \supset \mathcal{L}_{t, \mathbf{g}} \supset \mathcal{L}_{t+1, \mathbf{g}} \supset \cdots$$

so that we have $\cdots \leq f_{E_t, G_t}^{\mu} \leq f_{E_{t+1}, G_{t+1}}^{\mu} \leq \cdots \leq f^*$. We check that similar properties hold for \mathcal{Q}_{E_t, G_t} , $\mathcal{Q}_{t, \mathbf{g}}$ and $f_{E_t, G_t}^{\text{sos}} = f_{t, \mathbf{g}}^{\text{sos}}$, taking the quotient modulo $\langle F_{2t} | 2t \rangle$.

2.4. Optimal linear form. We introduce now the notion of *optimal linear form for f* , involved in the computation of I_{\min} (also called generic linear form when $f = 0$ in [14, 10]):

Definition 2.7. $\Lambda^* \in \mathcal{L}_{E, G}$ is optimal for f if

$$\text{rank } H_{\Lambda^*}^E = \max_{\Lambda \in \mathcal{L}_{E, G}, \Lambda(f) = f_{E, G}^{\mu}} \text{rank } H_{\Lambda}^E.$$

The next result shows that only elements in I_{\min} are involved in the kernel of a truncated Hankel operator associated to an optimal linear form for f .

Theorem 2.8. Let $E \subset \mathbb{R}[\mathbf{x}]$ such that $1 \in E$ and $f \in \langle E \cdot E \rangle$ and let $G \subset \mathbb{R}[\mathbf{x}]$ be a set of constraints with $V_{\min} \subset \mathcal{S}(G)$. If $\Lambda^* \in \mathcal{L}_{E, G}$ is optimal for f and such that $\Lambda^*(f) = f^*$, then $\ker H_{\Lambda^*}^E \subset I_{\min}$.

Proof. The proof is similar e.g. to [10][Theorem 4.9]. \square

Let us describe how optimal linear forms are computed by solving convex optimization problems:

Algorithm 2.1: OPTIMAL LINEAR FORM

Input: $f \in \mathbb{R}[\mathbf{x}]$, $B_t = (\mathbf{x}^{\alpha})_{\alpha \in A}$ a monomial set containing 1 of degree $\leq t$ with $f = \sum_{\alpha \in A+A} f_{\alpha} \mathbf{x}^{\alpha} \in \langle B_t \cdot B_t \rangle$, $G \subset \mathbb{R}[\mathbf{x}]$.

Output: the minimum $f_{t, G}^{\mu}$ of $\sum_{\alpha \in A+A} \lambda_{\alpha} f_{\alpha}$ subject to:

- $H_{\Lambda^*}^{B_t} = (h_{\alpha, \beta})_{\alpha, \beta \in A} \succcurlyeq 0$,
- $H_{\Lambda^*}^{B_t}$ satisfies the Hankel constraints
 $h_{0,0} = 1$, and $h_{\alpha, \beta} = h_{\alpha', \beta'}$ if $\alpha + \beta = \alpha' + \beta'$,
- $\Lambda^*(g^0) = \sum_{\alpha \in A+A} g_{\alpha}^0 \lambda_{\alpha} = 0$ for all $g^0 = \sum_{\alpha \in A+A} g_{\alpha}^0 \mathbf{x}^{\alpha} \in G^0 \cap \langle B_t \cdot B_t \rangle$.
- $H_{g^+, \Lambda^*}^{B_t - w} \succcurlyeq 0$ for all $g^+ \in G^+$ where $w = \lceil \frac{\text{deg}(g^+)}{2} \rceil$.

and $\Lambda^* \in \langle B_t \cdot B_t \rangle^*$ represented by the vector $[\lambda_{\alpha}]_{\alpha \in A+A}$.

This optimization problem is a Semi-Definite Programming problem, corresponding to the optimization of a linear functional on the intersection of a linear subspace with the convex

set of Positive Semi-Definite matrices. It is a convex optimization problem, which can be solved efficiently by SDP solvers. If an Interior Point Method is used, the solution Λ^* is in the interior of a face on which the minimum $\Lambda^*(f)$ is reached so that Λ^* is optimal for f . This is the case for tools such as `csdp`, `sdpa` or `sdpa-gmp`, that we will use in the experimentations.

3. CONVERGENCE CERTIFICATION

To be able to compute the minimizer points from an optimal linear form, we need to detect when the minimum is reached. In this section, we describe a new criterion to check when the kernel of a truncated Hankel operator associated to an optimal linear form for f yields the generators of the minimizer ideal. It involves the flat extension theorem of [16] and applies to polynomial optimization problems where the minimizer ideal I_{min} is zero-dimensional.

3.1. Flat extension criterion.

Definition 3.1. *Given vector subspaces $E_0 \subset E \subset \mathbb{K}[\mathbf{x}]$ and $\Lambda \in \langle E \cdot E \rangle^*$, H_Λ^E is said to be a flat extension of its restriction $H_\Lambda^{E_0}$ if $\text{rank } H_\Lambda^E = \text{rank } H_\Lambda^{E_0}$.*

We recall here a result from [16], which gives a rank condition for the existence of a flat extension of a truncated Hankel operator ¹.

Theorem 3.2. *Let $V \subset E \subset \mathbb{R}[\mathbf{x}]$ be vector spaces connected to 1 with $V^+ \subset E$ and let $\Lambda \in \langle E \cdot E \rangle^*$. Assume that $\text{rank } H_\Lambda^E = \text{rank } H_\Lambda^V = \dim V$. Then there exists a (unique) linear form $\tilde{\Lambda} \in \mathbb{R}[\mathbf{x}]^*$ which extends Λ , i.e., $\tilde{\Lambda}(p) = \Lambda(p)$ for all $p \in \langle E \cdot E \rangle$, satisfying $\text{rank } H_{\tilde{\Lambda}} = \text{rank } H_\Lambda^E$. Moreover, we have $\ker H_{\tilde{\Lambda}} = (\ker H_\Lambda^E)$.*

In other words, the condition $\text{rank } H_\Lambda^E = \text{rank } H_\Lambda^V = \dim V$ implies that the truncated Hankel operator H_Λ^E has a (unique) flat extension to a (full) Hankel operator $H_{\tilde{\Lambda}}$ defined on $\mathbb{R}[\mathbf{x}]$.

Theorem 3.3. *Let $V \subset E \subset \mathbb{R}[\mathbf{x}]$ be finite dimensional vector spaces connected to 1 with $V^+ \subset E$, $G^0 \cdot V \subset \langle E \cdot E \rangle$, $G^+ \cdot V \cdot V \subset \langle E \cdot E \rangle$.*

Let $\Lambda \in \mathcal{L}_{E,G}$ such that $\text{rank } H_\Lambda^E = \text{rank } H_\Lambda^V = \dim V$. Then there exists a linear form $\tilde{\Lambda} \in \mathbb{R}[\mathbf{x}]^$ which is extending Λ and supported on points of $\mathcal{S}(G)$ with positive weights:*

$$\tilde{\Lambda} = \sum_{i=1}^r \omega_i \mathbf{1}_{\xi_i} \text{ with } \omega_i > 0, \xi_i \in \mathcal{S}(G).$$

Moreover, $(\ker H_\Lambda^E) = \mathcal{I}(\xi_1, \dots, \xi_r)$.

Proof. As $\text{rank } H_\Lambda^E = \text{rank } H_\Lambda^V = \dim V$, Theorem 3.2 implies that there exists a (unique) linear function $\tilde{\Lambda} \in \mathbb{R}[\mathbf{x}]^*$ which extends Λ . As $\text{rank } H_{\tilde{\Lambda}} = \text{rank } H_\Lambda^V = |V|$ and $\ker H_{\tilde{\Lambda}} = (\ker H_\Lambda^E)$, any polynomial $p \in \mathbb{R}[\mathbf{x}]$ can be reduced modulo $\ker H_{\tilde{\Lambda}}$ to a polynomial $b \in V$ so that $p - b \in \ker H_{\tilde{\Lambda}}$. Then $\tilde{\Lambda}(p^2) = \tilde{\Lambda}(b^2) = \Lambda(b^2) \geq 0$ since $\Lambda \in \mathcal{L}_{E,G}$. By Theorem 3.14 of [10], $\tilde{\Lambda}$ has a decomposition of the form $\tilde{\Lambda} = \sum_{i=1}^r \omega_i \mathbf{1}_{\xi_i}$ with $\omega_i > 0$ and $\xi_i \in \mathbb{R}^n$.

¹In [16], it is stated with a vector space spanned by a monomial set connected to 1, but its extension to vector spaces connected to 1 is straightforward.

By Lemma 3.5 of [10], V is isomorphic to $\mathbb{R}[\mathbf{x}]/\mathcal{I}(\xi_1, \dots, \xi_r)$ and there exist (interpolation) polynomials $b_1, \dots, b_r \in V$ satisfying $b_i(\xi_j) = 1$ if $i = j$ and $b_i(\xi_j) = 0$ otherwise. We deduce that for $i = 1, \dots, r$ and for all elements $g \in G^0$,

$$\Lambda(b_i g) = 0 = \tilde{\Lambda}(b_i g) = \omega_i g(\xi_i).$$

As $\omega_i > 0$ then $g(\xi_i) = 0$. Similarly, for all $h \in G^+$,

$$\Lambda(b_i^2 h) = \tilde{\Lambda}(b_i^2 h) = \omega_i h(\xi_i) \geq 0$$

and $h(\xi_i) \geq 0$, hence $\xi_i \in \mathcal{S}(G)$.

By Theorem 3.14 of [10] and Theorem 3.2, we have moreover $\ker H_{\tilde{\Lambda}} = \mathcal{I}(\xi_1, \dots, \xi_r) = (\ker H_{\Lambda}^E)$. \square

This theorem applied to an optimal linear form Λ^* for f gives a convergence certificate to check when the minimum f^* is reached and when a generating family of the minimizer ideal is obtained. It generalizes the flat truncation certificate given in [25]. As we will see in the experimentation part, it allows to detect more efficiently when the minimum is reached. Notice that if the test is satisfied, necessarily I_{min} is zero-dimensional.

3.2. Flat extension algorithm. In this section, we describe a new algorithm to check the flat extension property for a linear form for which some moments are known.

Let E be a finite dimensional subspace of $\mathbb{R}[\mathbf{x}]$ connected to 1 and let Λ^* be a linear form defined on $\langle E \cdot E \rangle$ given by its ‘‘moments’’ $\Lambda^*(e_i) := \Lambda_i^*$, where e_1, \dots, e_s is a basis of $\langle E \cdot E \rangle$ (for instance a monomial basis). In the context of global polynomial optimization that we consider here, this linear form is an optimal linear form for f (see Section 2.4) computed by SDP.

We define the linear functional Λ^* from its moments as $\Lambda^* : p = \sum_{i=1}^s p_i e_i \in \langle E \cdot E \rangle \mapsto \sum_{i=1}^s p_i \Lambda_i^*$ and the corresponding inner product:

$$(6) \quad \begin{aligned} E \times E &\rightarrow \mathbb{R} \\ (p, q) &\mapsto \langle p, q \rangle_* := \Lambda^*(pq) \end{aligned}$$

To check the flat extension property, we are going to define inductively vector spaces V_i as follows. Start with $V_0 = \langle 1 \rangle$. Suppose V_i is known and compute a vector space L_i of maximal dimension in V_i^+ such that L_i is orthogonal to V_i : $\langle L_i, V_i \rangle_* = 0$ and $L_i \cap \ker H_{\Lambda^*}^{V_i^+} = \{0\}$. Then we define $V_{i+1} = V_i + L_i$.

Suppose that b_1, \dots, b_{r_i} is an orthogonal basis of V_i : $\langle b_i, b_j \rangle_* = 0$ if $i \neq j$ and $\langle b_i, b_i \rangle_* \neq 0$. Then L_i can be constructed as follows: Compute the vectors

$$b_{i,j} = x_j b_i - \sum_{k=1}^{r_i} \frac{\langle x_j b_i, b_k \rangle_*}{\langle b_k, b_k \rangle_*} b_k,$$

generating V_i^\perp in V_i^+ and extract a maximal orthogonal family $b_{r_i+1}, \dots, b_{r_{i+1}}$ for the inner product $\langle \cdot, \cdot \rangle_*$, that form a basis of L_i . This can be done for instance by computing a QR decomposition of the matrix $[\langle b_{i,j}, b_{i',j'} \rangle_*]_{1 \leq i, i' \leq r_i, 1 \leq j, j' \leq n}$. The process can be repeated until either

- $V_i^+ \not\subset E$ and the algorithm will stop and return **failed**,
- or $L_i = \{0\}$ and $V_i^+ = V_i \oplus \ker H_{\Lambda^*}^{V_i^+}$. In this case, the algorithm stops with **success**.

Here is the complete description of the algorithm:

Algorithm 3.1: DECOMPOSITION

Input: a vector space E connected to 1 and a linear form $\Lambda^* \in \langle E \cdot E \rangle^*$.

- Take $B := \{1\}$; $s := 1$; $r := 1$;
- While $s > 0$ and $B^+ \subset E$ do
 - compute $b_{j,k} := x_k b_j - \sum_{i=1}^r \frac{\langle x_k b_j, b_i \rangle_*}{\langle b_i, b_i \rangle_*} b_i$ for $j = 1, \dots, r$, $k = 1, \dots, n$;
 - compute a maximal subset $B' = \{b'_1, \dots, b'_s\}$ of $\langle b_{j,k} \rangle$ of orthogonal vectors for the inner product $\langle \cdot, \cdot \rangle_*$ and let $B := B \cup B'$, $s = |B'|$ and $r += s$;
- If $B^+ \not\subset E$ then return **failed** else return **success**.

Output: failed or success with

- a basis $B = \{b_1, \dots, b_r\} \subset \mathbb{R}[\mathbf{x}]$;
 - the relations $x_k b_j - \sum_{i=1}^r \frac{\langle x_k b_j, b_i \rangle_*}{\langle b_i, b_i \rangle_*} b_i$, $j = 1 \dots r$ $k = 1 \dots n$.
-

Let us describe the computation performed on the moment matrix, during the main loop of the algorithm. At each step, the moment matrix of Λ^* on V_i^+ is of the form

$$H_{\Lambda^*}^{V_i^+} = \left[\begin{array}{c|c} H_{\Lambda^*}^{B_i, B_i} & H_{\Lambda^*}^{B_i, \partial B_i} \\ \hline H_{\Lambda^*}^{\partial B_i, B_i} & H_{\Lambda^*}^{\partial B_i, \partial B_i} \end{array} \right]$$

where ∂B_i is a subset of $\{b_{i,j}\}$ such that $B_i \cup \partial B_i$ is a basis of $\langle B_i^+ \rangle$. By construction, the matrix $H_{\Lambda^*}^{B_i, B_i}$ is diagonal since B_i is orthogonal for $\langle \cdot, \cdot \rangle_*$. As the polynomials $b_{i,j}$ are orthogonal to B_i , we have $H_{\Lambda^*}^{B_i, \partial B_i} = H_{\Lambda^*}^{\partial B_i, B_i} = 0$. If $H_{\Lambda^*}^{\partial B_i, \partial B_i} = 0$ then the algorithm stops with **success** and all the elements $b_{i,j}$ are in the kernel of $H_{\Lambda^*}^{B_i, B_i}$. Otherwise an orthogonal basis b'_1, \dots, b'_s is extracted. It can then be completed in a basis of $\langle b_{i,j} \rangle$ so that the matrix $H_{\Lambda^*}^{\partial B_i, \partial B_i}$ in this basis is diagonal with zero entries after the $(s+1)^{th}$ index. In the next loop of the algorithm, the basis B_{i+1} contains the maximal orthogonal family b'_1, \dots, b'_s so that the matrix $H_{\Lambda^*}^{B_{i+1}, B_{i+1}}$ remains diagonal and invertible.

Proposition 3.4. *Let $\Lambda^* \in \mathcal{L}_{E,G}$ be optimal for f . If Algorithm 3.1 applied to Λ^* and E stops with **success**, then*

- (1) *there exists a linear form $\tilde{\Lambda} \in \mathbb{R}[\mathbf{x}]^*$ which is extending Λ^* and supported on points in $\mathcal{S}(G)$ with positive weights:*

$$\tilde{\Lambda} = \sum_{i=1}^r \omega_i \mathbf{1}_{\xi_i} \text{ with } \omega_i > 0, \xi_i \in \mathbb{R}^n.$$

- (2) $B = \{b_1, \dots, b_r\}$ is a basis of $\mathcal{A}_{\tilde{\Lambda}} = \mathbb{R}[\mathbf{x}]/I_{\tilde{\Lambda}}$ where $I_{\tilde{\Lambda}} = \ker H_{\tilde{\Lambda}}$,
- (3) $x_k b_j - \sum_{i=1}^r \frac{\langle x_k b_j, b_i \rangle_*}{\langle b_i, b_i \rangle_*} b_i$, $j = 1, \dots, r$, $k = 1, \dots, n$ are generating $I_{\tilde{\Lambda}} = \mathcal{I}(\xi_1, \dots, \xi_r)$,
- (4) $f_{E,G}^\mu = f^*$,
- (5) $V_{min} = \{\xi_1, \dots, \xi_r\}$.

Proof. When the algorithm terminates with **success**, the set B is such that $\text{rank } H_{\Lambda^*}^{B^+} = \text{rank } H_{\Lambda^*}^B = |B|$. By Theorem 3.3, there exists a linear form $\tilde{\Lambda} \in \mathbb{R}[\mathbf{x}]^*$ extending Λ^* and supported on points in $\mathcal{S}(G)$ with positive weights:

$$\tilde{\Lambda} = \sum_{i=1}^r \omega_i \mathbf{1}_{\xi_i} \text{ with } \omega_i > 0, \xi_i \in \mathcal{S}(G).$$

This implies that $\mathcal{A}_{\tilde{\Lambda}}$ is of dimension r and that $I_{\tilde{\Lambda}} = \mathcal{I}(\xi_1, \dots, \xi_r)$. As $H_{\tilde{\Lambda}^*}^B$ is invertible, B is a basis of $\mathcal{A}_{\tilde{\Lambda}}$ which proves the second point.

Let K be the set of polynomials $x_j b_i - \sum_{k=1}^r \frac{\langle x_j b_i, b_k \rangle_{\tilde{\Lambda}^*}}{\langle b_k, b_k \rangle_{\tilde{\Lambda}^*}} b_k$. If the algorithm terminates with **success**, we have $\ker H_{\tilde{\Lambda}^*}^{B^+} = \langle K \rangle$ and by Theorem 3.3, we deduce that $(K) = (\ker H_{\tilde{\Lambda}^*}^{B^+}) = I_{\tilde{\Lambda}}$, which proves the third point.

As $\tilde{\Lambda}(1) = 1$, we have $\sum_{i=1}^r w_i = 1$ and

$$\tilde{\Lambda}(f) = \sum_{i=1}^r \omega_i f(\xi_i) \geq f^*$$

since $\xi_i \in \mathcal{S}(G)$ and $f(\xi_i) \geq f^*$. The relation $f_{E,G}^\mu \leq f^*$ implies that $f(\xi_i) = f^*$ for $i = 1, \dots, r$ and the fourth point is true: $f_{E,G}^\mu = f^*$.

As $f(\xi_i) = f^*$ for $i = 1, \dots, r$, we have $\{\xi_1, \dots, \xi_r\} \subset V_{min}$. By Theorem 2.8, the polynomials of K are in I_{min} so that $V_{min} \subset \mathcal{V}(K) = \{\xi_1, \dots, \xi_r\}$. This shows that $V_{min} = \{\xi_1, \dots, \xi_r\}$ and concludes the proof of this proposition. \square

3.3. Computing the minimizers. The remaining step is the computation of the minimizer points, once Algorithm 3.1 stops with **success** for $\Lambda^* \in \mathcal{L}_{E,G}$ optimal for f . The minimizers can be computed from the eigenvalues of the multiplication operators $M_k : a \in \mathcal{A}_{min} \mapsto x_k a \in \mathcal{A}_{min}$ for $k = 1, \dots, n$ where $\mathcal{A}_{min} = \mathbb{R}[\mathbf{x}]/I_{min}$ and $I_{min} = I_{\tilde{\Lambda}} = \mathcal{I}(\xi_1, \dots, \xi_r)$.

Proposition 3.5. *The matrix of M_k in the basis B of \mathcal{A}_{min} is $[M_k] = (\frac{\Lambda^*(x_k b_i b_j)}{\Lambda^*(b_i b_i)})_{1 \leq i, j \leq r}$. The operators M_k , $k = 1 \dots n$ have r common eigenvectors $\mathbf{u}_1, \dots, \mathbf{u}_r$ which satisfy $M_k \mathbf{u}_i = \xi_{i,k} \mathbf{u}_i$, with $\xi_{i,k}$ the k^{th} coordinate of the minimizer point $\xi_i = (\xi_{i,1}, \dots, \xi_{i,n}) \in S$.*

Proof. By Proposition 3.4 and by definition of the inner-product (6), $B = \{b_1, \dots, b_r\}$ is a basis of $\mathcal{A}_{\tilde{\Lambda}}$ and

$$x_k b_j \equiv \sum_{i=1}^r \frac{\Lambda^*(x_k b_i b_j)}{\Lambda^*(b_i b_i)} b_i \pmod{I_{min}},$$

for $j = 1 \dots r$, $k = 1 \dots n$. This yields the matrix of the operator M_k in the basis B : $[M_k] = (\frac{\Lambda^*(x_k b_i b_j)}{\Lambda^*(b_i b_i)})_{1 \leq i, j \leq r}$.

As the roots of I_{min} are simple, by [4][Theorem 4.23] the eigenvectors of all M_k , $k = 1 \dots n$ are the so-called idempotents $\mathbf{u}_1, \dots, \mathbf{u}_r$ of \mathcal{A}_{min} and the corresponding eigenvalues are $\xi_{1,k}, \dots, \xi_{r,k}$. \square

Algorithm 3.2: MINIMIZER POINTS

Input: B and the relations as output in Algorithm 3.1.

- Compute the matrices $[M_k] = (\frac{\Lambda^*(x_k b_i b_j)}{\Lambda^*(b_i b_i)})_{1 \leq i, j \leq r}$.
- For a generic choice of $l_1, \dots, l_n \in \mathbb{R}$, compute the eigenvectors $\mathbf{u}_1, \dots, \mathbf{u}_r$ of $l_1[M_1] + \dots + l_n[M_n]$.
- Compute $\xi_{i,k} \in \mathbb{R}$ such that $M_k \mathbf{u}_i = \xi_{i,k} \mathbf{u}_i$.

Output: the minimizers $\xi_i = (\xi_{i,1}, \dots, \xi_{i,n})$, $i = 1 \dots r$.

4. FINITE CONVERGENCE

Our approach to compute the minimizers relies on the fact that the border basis relaxation is exact. By Proposition 2.6, the reduced border basis relaxation is exact if and only if the corresponding full moment matrix relaxation is exact.

Though it is not always the case that the full moment matrix relaxation is exact, it is possible to add constraints so that the relaxation becomes exact. In [1], a general strategy to construct exact SDP relaxation hierarchies and to compute the minimizer ideal is described. It applies to the following problems:

Global optimization. Consider the case $n_1 = n_2 = 0$ with $f^* = \min_{\mathbf{x} \in \mathbb{R}^n} f(\mathbf{x})$ reached at a point of \mathbb{R}^n . Taking G such that $G^0 = \{\frac{\partial f}{\partial x_1}, \dots, \frac{\partial f}{\partial x_n}\}$ and $G^+ = \emptyset$, the relaxation associated to the sequence $\mathcal{L}_{t,G}$ is exact and yields I_{min} (see [26, 1]). If I_{min} is finite then the border basis relaxation yields the points and the corresponding border basis.

Regular case. We say that $\mathbf{g} = (g_1^0, \dots, g_{n_1}^0; g_1^+, \dots, g_{n_2}^+)$ is regular if for all points $\mathbf{x} \in \mathcal{S}(\mathbf{g})$ with $\{j_1, \dots, j_k\} = \{j \in [1, n_2] \mid g_j^+(\mathbf{x}) = 0\}$, the vectors $\nabla g_1^0(\mathbf{x}), \dots, \nabla g_{n_1}^0(\mathbf{x}), \nabla g_{j_1}^+(\mathbf{x}), \dots, \nabla g_{j_k}^+(\mathbf{x})$ are linearly independent.

For $\nu = \{j_1, \dots, j_k\} \subset [0, n_2]$ with $|\nu| \leq n - n_1$, let

$$\begin{aligned} A_\nu &= [\nabla f, \nabla g_1^0, \dots, \nabla g_{n_1}^0, \nabla g_{j_1}^+, \dots, \nabla g_{j_k}^+] \\ \Delta_\nu &= \det(A_\nu A_\nu^T) \\ g_\nu &= \Delta_\nu \prod_{j \notin \nu} g_j^+. \end{aligned}$$

Let $G \subset \mathbb{R}[\mathbf{x}]$ be the set of constraints such that $G^0 = \mathbf{g}^0 \cup \{g_\nu \mid \nu \subset [0, n_2], |\nu| \leq n - n_1\}$. Then the relaxation associated to the preordering sequence $\mathcal{L}_{t,G}^*$ is exact and yields I_{min} (see [8, 1] or [24] for \mathbb{C} -regularity and constraints G^0 that involve minors of A_ν).

If I_{min} is non-empty and finite then the border basis relaxation (2) yields the points V_{min} and the border basis of I_{min} .

Boundary Hessian Conditions. If f and \mathbf{g} satisfies the so-called Boundary Hessian Conditions then $f - f^* \in \mathcal{Q}_{t,g}$ and the relaxation associated to $\mathcal{L}_{t,\mathbf{g}}$ is exact and yields I_{min} (see [18]). If moreover I_{min} is finite then the border basis relaxation yields the points V_{min} and the corresponding border basis of I_{min} .

\mathfrak{g}^+ -radical computation. If we optimize $f = 0$ on the set $S = \mathcal{S}(\mathbf{g})$, then all the points of S are minimizers, $V_{min} = S$ and by the Positivstellensatz, I_{min} is equal to

$$\mathfrak{g}^+ \sqrt{\mathbf{g}^0} = \{p \in \mathbb{R}[\mathbf{x}] \mid \exists m \in \mathbb{N} \text{ s.t. } p^{2m} + q = 0, q \in \mathcal{P}_{\mathbb{R}[\mathbf{x}], \mathbf{g}}\}.$$

Here again, the preordering sequence $\mathcal{L}_{t,\mathbf{g}}^*$ is exact. If we assume that $S = \mathcal{S}(\mathbf{g})$ is finite, then the corresponding border basis relaxation yields the points of S and the generators of $\mathfrak{g}^+ \sqrt{\mathbf{g}^0}$. See also [14, 10] for zero dimensional real radical computation and [17].

5. MINIMIZER BORDER BASIS ALGORITHM

In this section we describe the algorithm to compute the minimum of a polynomial on S . It can be seen as a type of border basis algorithm, in which in the main loop we compute the optimal linear form, we check when the minimum is reached and eventually we compute the minimizers points. It is closely connected to the real radical border basis algorithm presented in [10].

5.1. Description.

Algorithm 5.1: MINIMIZATION OF f ON S

Input: A real polynomial function f and a set of constraints $\mathbf{g} \subset \mathbb{R}[\mathbf{x}]$ with V_{min} non-empty finite.

- (1) Take $t = \max(\lceil \frac{\deg(f)}{2} \rceil, d^0, d^+)$ where
 $d^0 = \max_{g^0 \in \mathbf{g}^0}(\lceil \frac{\deg(g^0)}{2} \rceil), d^+ = \max_{g^+ \in \mathbf{g}^+}(\lceil \frac{\deg(g^+)}{2} \rceil)$
- (2) Compute the graded border basis F_{2t} of \mathbf{g}^0 for B in degree $2t$.
- (3) Let B_t be the set of monomials in B of degree $\leq t$.
- (4) Let G_t be the set of constraints such that $G_t^0 = \{m - \pi_{B_t, F_{2t}}(m), m \in B_t \cdot B_t\}$ and $G^+ = \pi_{B_t, F_{2t}}(g^+)$
- (5) $[f_{G_t, B_t}^*, \Lambda^*] := \text{OPTIMAL LINEAR FORM}(f, B_t, G_t)$.
- (6) $[c, B', K] := \text{DECOMPOSITION}(\Lambda^*, B_t)$ where $c = \text{failed}$, $B' = \emptyset, K = \emptyset$ or $c = \text{success}$, B' is the basis and K is the set of the relations.
- (7) if $c = \text{success}$ then $V = \text{MINIMIZER POINTS}(B', K)$
else go to step 2 with $t := t + 1$.

Output: the minimum $f^* = f_{G_t, B_t}^*$, the minimizers $V_{min} = V$, $I_{min} = (K)$ and B' such that K is a border basis for B' .

5.2. Experimentation. In this section, we analyze the practical behavior of the algorithm. In all the examples the minimizer ideal is zero-dimensional hence our algorithm stops in a finite number of steps and yields the minimizer points and generators of the minimizer ideal.

The implementation of the previous algorithm has been performed using the BORDERBASIX package of the MATHEMAGIX² software, which provides a C++ implementation of the border basis algorithm of [22].

For the computation of border basis, we use as a choice function that is tolerant to numerical instability i.e. a choice function that chooses as leading monomial a monomial whose coefficient is maximal among the choosable monomials as described in [21].

The Semi-Definite Programming problems are solved using SDPA³ software. For the link with SDPA we use a file interface since SDPA is not distributed as a library.

Once we have computed the moment matrix, we call the Decomposer Algorithm which is available in the BORDERBASIX package.

The minimizer points are computed from the eigenvalues of the multiplication matrices. This is performed using Lapack routines.

Experiments are made on an Intel Core i5 2.40GHz

In Table 1, we compare this algorithm 5.1 (*bbr*) with the full moment matrix relaxation algorithm (*fmr*) inside the same environment. It reproduces the algorithm described in [12], which is also implemented in the package Gloptipoly of Matlab developed by D. Henrion and J.B. Lasserre. In this table, we record the problem name or the source of the problem, the number of decision variables (v), the number of inequality and equality constraints (c), the maximum degree of the constraints and of the polynomial to minimize (d), the number of minimizer points (sol). For the two algorithms *bbr* and *fmr* we report the CPU time in seconds (t), the order of the relaxation (o), the number of parameters of the SDP problem (p) and the size of the moment matrices (s). The first part of the table contains examples of positive polynomials, which are not some of squares. New equality constraints are added following [1] to compute the minimizer points in the examples marked with \diamond .

²www.mathemagix.org

³http://sdpa.sourceforge.net

problem	v	c	d	sol	t_{bbr}	O_{bbr}	P_{bbr}	S_{bbr}	t_{fmr}	O_{fmr}	P_{fmr}	S_{fmr}
◇ Robinson	2	0	6	8	0.15	4	21	15	*	7	119	36
◇ Motzkin	2	0	6	4	0.17	4	26	15	*	9	189	55
◇ Motzkin perturbed	3	1	6	1	3.78	5	167	56	9.57	5	286	56
◇ [11], Ex. 1	2	0	4	1	0.030	2	8	6	0.050	2	14	6
◇ [11], Ex. 2	2	0	4	1	0.030	2	8	6	0.050	2	14	6
◇ [11], Ex. 3	2	0	6	4	0.432	4	25	15	*	8	152	45
[11], Ex. 5	2	3	2	3	0.045	2	14	6	0.053	2	14	6
[5], Ex. 4.1.4	1	2	4	2	0.024	2	4	3	0.040	2	4	3
[5], Ex. 4.1.6	1	2	6	2	0.027	3	6	4	0.044	3	6	4
[5], Ex. 4.1.7	1	2	4	1	0.023	2	4	3	0.042	2	4	3
[5], Ex. 4.1.8	2	5	4	1	0.060	2	13	6	0.077	2	14	6
[5], Ex. 4.1.9	2	6	4	1	0.20	4	44	15	0.29	4	44	15
[5], Ex. 2.1.1	5	11	2	1	7.60	3	461	56	12.23	3	461	56
[5], Ex. 2.1.2	6	13	2	1	1.00	2	209	26	1.29	2	209	26
[5], Ex. 2.1.3	13	35	2	1	383.97	2	2379	78	417.96	2	2379	78
[5], Ex. 2.1.4	6	15	2	1	1.01	2	209	26	1.48	2	209	26
[5], Ex. 2.1.5	10	31	2	1	29.70	2	1000	66	44.29	2	1000	66
[5], Ex. 2.1.6	10	25	2	1	28.60	2	1000	66	43.68	2	1000	66
** [5], Ex. 2.1.7 (case 1)	20	30	2	1	33219.9	2	10625	231	35310.7	2	10625	231
** [5], Ex. 2.1.7 (case 5)	20	30	2	1	33475.2	2	10625	231	36021.3	2	10625	231
** [5], Ex. 2.1.8	24	58	2	1	3929.23	2	3875	136	>14h	2	20475	325
[5], Ex. 2.1.9	10	11	2	1	12.3	2	714	44	16.76	2	1000	55
[5], Ex. 3.1.3	6	16	2	1	0.96	2	209	26	1.42	2	209	26
[12] cbms1	3	3	3	5	0.16	3	26	17	0.20	3	83	20
[12] rediff3	3	3	2	2	0.07	2	7	7	0.09	2	35	10
[12] quadfor2	4	12	4	2	0.6	3	48	19	0.75	3	210	35
** simplex	15	16	2	1	674.534	2	3059	120	780.371	2	3875	136

TABLE 1. Experimentations

The fourth part of the table contains examples where the real radical $\mathfrak{s}^+\sqrt{\mathbf{g}^0}$ is computed. When there are equality constraints, the border basis computation reduces the size of the moment matrices, as well as the localization matrices associated to the inequalities. This speeds up the SDP computation. In the case where there are only inequalities, the size of the moment matrices is the same but once the optimal linear form is computed using the SDP solver `SDPA`, the `DECOMPOSITION` algorithm which computes the minimizers is more efficient and quicker than the reconstruction algorithm used in the full moment matrix relaxation approach. The performance is not the only issue: numerical problems can also occur due to the bigger size of the moment matrices in the flat extension test and the reconstruction of minimizers. Such examples where the `fmr` algorithm fails because of the numerical rank problems are marked with *. The examples that Gloptipoly cannot treat due to the high number of variables [12] are marked with **.

These experiments show that when the size of the SDP problems becomes significant, most of the time is spent during `sdpa` computation and the border basis time and reconstruction time are negligible. In all the examples, the new border basis relaxation algorithm outperforms the full moment matrix relaxation method.

Acknowledgements. We thank Philippe Trébuchet and Matthieu Dien for their development in the BORDERBASIX package.

REFERENCES

- [1] M. Abril Bucero and B. Mourrain. Exact relaxation for polynomial optimization on semi-algebraic sets. <http://hal.inria.fr/hal-00846977>, 2013.
- [2] J. Brachat, P. Comon, B. Mourrain, and E. Tsigaridas. Symmetric tensor decomposition. *Linear Algebra and Applications*, 433:1851–1872, 2010.
- [3] J. Demmel, J. Nie, and V. Powers. Representations of positive polynomials on noncompact semialgebraic sets via kkt ideals. *Journal of Pure and Applied Algebra*, 209(1):189 – 200, 2007.
- [4] M. Elkadi and B. Mourrain. *Introduction à la résolution des systèmes d'équations algébriques*, volume 59 of *Mathématiques et Applications*. Springer-Verlag, 2007.
- [5] C. A. Floudas, P. M. Pardalos, C. S. Adjiman, W. R. Esposito, Z. H. Gumus, S. T. Harding, J. L. Klepeis, C. A. Meyer, and C. A. Schweiger. *Handbook of Test Problems in Local and Global Optimization*. Kluwer Academic Publishers., 1999. See titan.princeton.edu/Testproblems.
- [6] M. Giesbrecht, G. Labahn, and W.-S. Lee. Symbolic-numeric sparse interpolation of multivariate polynomials. *J. Symb. Comput.*, 44(8):943–959, August 2009.
- [7] A. Greuet and M. Safey El Din. Deciding reachability of the infimum of a multivariate polynomial. In *Proceedings of the 36th international symposium on Symbolic and algebraic computation*, ISSAC '11, pages 131–138, New York, NY, USA, 2011. ACM.
- [8] H. V. Ha and T.S. Pham. Representation of positive polynomials and optimization on noncompact semialgebraic sets. *SIAM Journal on Optimization*, 20(6):3082–3103, 2010.
- [9] D. Henrion and J.B. Lasserre. *Positive Polynomials in Control*, chapter Detecting Global Optimality and Extracting Solutions in GloptiPoly., pages 293–310. Lectures Notes in Control and Information Sciences. Springer, 2005.
- [10] J.-B. Lasserre, M. Laurent, B. Mourrain, P. Rostalski, and P. Trébuchet. Moment matrices, border bases and real radical computation. *Journal of Symbolic Computation*, 2012.
- [11] J.B. Lasserre. Global optimization with polynomials and the problem of moments. *SIAM J. Optim.*, 11:796–817, 2001.
- [12] J.B Lasserre. *Moments, positive polynomials and their applications*. Imperial College Press, 210.
- [13] J.B. Lasserre, M. Laurent, and P. Rostalski. Semidefinite characterization and computation of real radical ideals. *Foundations of Computational Mathematics*, 8(5):607–647, 2008.
- [14] J.B. Lasserre, M. Laurent, and P. Rostalski. A unified approach for real and complex zeros of zero-dimensional ideals. In M. Putinar and S. Sullivant, editors, *Emerging Applications of Algebraic Geometry*, volume 149, pages 125–156. Springer, 2009.
- [15] M. Laurent. Semidefinite representations for finite varieties. *Math. Progr*, 109:1–26, 2007.
- [16] M. Laurent and B. Mourrain. A generalized flat extension theorem for moment matrices. *Arch. Math. (Basel)*, 93(1):87–98, July 2009.
- [17] Y. Ma, Ch. Wang, and Zhi L. A certificate for semidefinite relaxations in computing positive dimensional real varieties. <http://arxiv.org/abs/1212.4924>, 2013.
- [18] M. Marshall. Representations of non-negative polynomials, degree bounds and applications to optimization. *Can. J. Math.*, 61(1):205–221, 2009.
- [19] B. Mourrain. A new criterion for normal form algorithms. In M. Fossorier, H. Imai, Shu Lin, and A. Poli, editors, *Proc. AAECC*, volume 1719 of *LNCS*, pages 430–443. Springer, Berlin, 1999.
- [20] B. Mourrain and P. Trébuchet. Generalized normal forms and polynomials system solving. In M. Kauers, editor, *ISSAC: Proceedings of the ACM SIGSAM International Symposium on Symbolic and Algebraic Computation*, pages 253–260, 2005.
- [21] B. Mourrain and Ph. Trébuchet. Stable normal forms for polynomial system solving. *Theoretical Computer Science*, 409(2):229–240, 2008.
- [22] B. Mourrain and Ph. Trébuchet. Border basis representation of a general quotient algebra. In Joris van der Hoeven, editor, *ISSAC 2012*, pages 265–272, July 2012.
- [23] Y. Nesterov and A. Nemirovski. *Interior-point polynomial algorithms in convex programming*. SIAM, Philadelphia, 1994.
- [24] J. Nie. An exact jacobian SDP relaxation for polynomial optimization. *Mathematical Programming*, pages 1–31, 2011.
- [25] J. Nie. Certifying convergence of Lasserre’s hierarchy via flat truncation. *Mathematical Programming*, pages 1–26, 2012.

- [26] J. Nie, J. Demmel, and B. Sturmfels. Minimizing polynomials via sum of squares over gradient ideal. *Math. Program.*, 106(3):587–606, 2006.
- [27] P.A. Parrilo. Semidefinite programming relaxations for semialgebraic problems. *Mathematical Programming Ser. B*, 96(2):293–320, 2003.
- [28] P.A. Parrilo and B. Sturmfels. Minimizing polynomial functions. In *Proceedings of the DIMACS Workshop on Algorithmic and Quantitative Aspects of Real Algebraic Geometry in Mathematics and Computer Science*, pages 83–100. American Mathematical Society, 2003.
- [29] C. Riener, T. Theobald, L. J. Andr en, and J. B. Lasserre. Exploiting symmetries in SDP-relaxations for polynomial optimization. *Math. Oper. Res.*, 38(1):122–141, February 2013.
- [30] M. Safey El Din. Computing the global optimum of a multivariate polynomial over the reals. In *Proceedings of the twenty-first international symposium on Symbolic and algebraic computation, ISSAC '08*, pages 71–78, New York, NY, USA, 2008. ACM.
- [31] N.Z. Shor. Class of global minimum bounds of polynomial functions. *Cybernetics*, 23:731–734, 1987.