



HAL
open science

Some old and new perspectives for differential games

Lawrence C. Evans

► **To cite this version:**

Lawrence C. Evans. Some old and new perspectives for differential games. NETCO 2014, 2014, Tours, France. hal-01024443

HAL Id: hal-01024443

<https://inria.hal.science/hal-01024443>

Submitted on 16 Jul 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

SOME OLD AND NEW PERSPECTIVES FOR DIFFERENTIAL GAMES

Lawrence C. Evans

Department of Mathematics
University of California, Berkeley

1. ZERO SUM DIFFERENTIAL GAMES

DYNAMICS:

$$\begin{cases} \dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \alpha, \beta) & (0 \leq t \leq \tau) \\ \mathbf{x}(0) = \mathbf{x}, \end{cases}$$

CONTROLS:

$$\begin{cases} \alpha : [0, \tau] \rightarrow A = \text{control for the maximizing player} \\ \beta : [0, \tau] \rightarrow B = \text{control for the minimizing player} \end{cases}$$

PAYOFF:

$$J^{\alpha, \beta}(\mathbf{x}) = \int_0^{\tau} r(\mathbf{x}, \alpha, \beta) dt + g(\mathbf{x}(\tau)).$$

r is the **running payoff** and g is the **terminal payoff**.

MINIMAX CONDITION:

$$\min_{b \in B} \max_{a \in A} \{f(x, a, b) \cdot p + r(x, a, b)\} = \max_{a \in A} \min_{b \in B} \{f(x, a, b) \cdot p + r(x, a, b)\}$$

for all x, p .

VALUE FUNCTION:

$u(x)$ = payoff to max player, if each plays optimally

THEOREM

The value function u , if continuous, solves the **Hamilton–Jacobi–Isaacs (HJI)** equation

$$H(Du, x) = 0$$

in the viscosity sense, for the **game theory Hamiltonian**

$$H(p, x) = - \min_{b \in B} \max_{a \in A} \{f(x, a, b) \cdot p + r(x, a, b)\}.$$

References

- M. Bardi and I. Capuzzo-Dolcetta, *Optimal Control and Viscosity Solutions of Hamilton-Jacobi-Bellman Equations*, Birkhäuser, 1997.
- A. Bressan, *Noncooperative Differential Games. A tutorial*, www.math.psu.edu/bressan/PSPDF/game-lnew.pdf
- P. Cardaliaguet, *Introduction to Differential Games*, www.ceremade.dauphine.fr/~cardalia/CoursJeuxDiff.pdf
- R. Isaacs, *Differential Games. A mathematical theory with applications to warfare and pursuit, control and optimization*, Wiley, 1965.
- J. Lewin, *Differential Games. Theory and methods for solving game problems with singular surfaces*, Springer, 1994.
- A. Melikyan, *Generalized Characteristics of First Order PDEs. Applications in optimal control and differential games*, Birkhäuser, 1998.

Problems for theory applied to “real” examples

- Hamiltonian $H = H(p, x)$ is nonconvex in p .
- Hamiltonian H is not C^1 (and therefore problems with characteristics = optimal trajectories).
- The value function u need not be continuous.
- Boundary conditions are unclear and/or hard to interpret in viscosity sense.

Failure of minimax condition is usually not a problem, at least for examples in Isaacs' and Lewin's books.

AN (UNNAMED) EXAMPLE OF ISAACS

DYNAMICS:

$$\begin{cases} \dot{x} = c(y) + \cos \alpha \\ \dot{y} = 2\beta + \sin \alpha, \end{cases}$$

where $0 \leq \alpha \leq 2\pi$ and $-1 \leq \beta \leq 1$.

$y \mapsto c(y)$ is positive, increasing.

The game is played in the upper half plane $U := \mathbb{R}^2 \cap \{y \geq 0\}$.


PAYOFF:

$J^{\alpha, \beta} = \tau =$ exit time from U along the positive x -axis.

HJI EQUATION: The value function u (= time to exit) solves

$$H(Du, y) = -c(y)u_x - |Du| + 2|u_y| - 1 = 0 \quad \text{in } U.$$

Boundary condition is $u = 0$ on $\{x \geq 0, y = 0\}$.




B = barrier; **ES** = "equivocal surface"

2. NONCONVEX HAMILTON-JACOBI PDE

Study the initial-value problem

$$\begin{cases} u_t + H(Du) = 0 & \text{in } \mathbb{R}^n \times (0, \infty) \\ u = g & \text{on } \mathbb{R}^n \times \{t = 0\}. \end{cases}$$

where H is smooth, but not convex.



The central difficulty for nonconvex Hamilton–Jacobi PDE is that backwards characteristics can hit “shocks”, along which the gradient is discontinuous.

New work of F. Rezakhanlou: Backwards characteristics as martingales.

FORMULAS FOR SOLUTIONS:

- If H is convex,

$$u(x, t) := \inf_{y \in \mathbb{R}^n} \left\{ g(y) + tL\left(\frac{x-y}{t}\right) \right\} \quad (\text{Hopf–Lax})$$

where $L = H^*$

- If g is convex and has linear growth,

$$u(x, t) := \sup_{z \in \mathbb{R}^n} \{ x \cdot z - g^*(z) - tH(z) \} \quad (\text{Hopf})$$

THEOREM (ENVELOPE REPRESENTATION FORMULA)

For \mathcal{L}^{n+1} -a.e. point $(x, t) \in \mathbb{R}^n \times (0, \infty)$ there exists a Radon probability measure $\gamma_{x,t}$ on \mathbb{R}^n such that

$$u(x, t) = \int_{\mathbb{R}^n} g(y) + (x - y) \cdot Dg(y) - tH(Dg(y)) d\gamma_{x,t}.$$



Furthermore,

$$u_t(x, t) = - \int_{\mathbb{R}^n} H(Dg(y)) d\gamma_{x,t}, \quad Du(x, t) = \int_{\mathbb{R}^n} Dg(y) d\gamma_{x,t},$$

and

$$H\left(\int_{\mathbb{R}^n} Dg(y) d\gamma_{x,t}\right) = \int_{\mathbb{R}^n} H(Dg(y)) d\gamma_{x,t}.$$

BARRIERS (= surfaces of discontinuity)



$$H(Du, x) = 0 \quad \text{in } U.$$

APPROXIMATION:

$$-\epsilon \Delta u^\epsilon + H(Du^\epsilon, x) = 0 \quad \text{in } U.$$

Assume:

$$u^\epsilon \leq u^+ + o(1), \quad u^\epsilon \geq u^- - o(1).$$

The **recession function** associated with $H = H(p, x)$ is

$$K(p, x) := \lim_{\lambda \rightarrow \infty} \frac{H(q + \lambda p, x)}{\lambda}.$$

THEOREM

(i)

$$\begin{cases} H(Du^+ + \lambda \nu, x) \leq 0 \\ H(Du^- + \lambda \nu, x) \geq 0 \end{cases} \quad \text{for all } \lambda \geq 0.$$

(ii) *In particular,*

$$K(\nu, x) = 0.$$

This is a PDE for the barrier (Isaacs). We can in principle solve for the location of the discontinuities, before we solve the full HJI equation!

3. NONZERO SUM GAMES: PRINCIPAL-AGENT PROBLEMS

SANNIKOV'S MODEL (simplified)

DYNAMICS:

$$dX = A dt + dB,$$

where B is a one-dimensional Brownian motion.

CONTROLS:

$$\begin{cases} C = \text{principal's control (= payments to agent)} \\ A = \text{agent's control of dynamics.} \end{cases}$$

PAYOFFS:

$$\begin{cases} J^P = \text{principal's payoff} = E \left(\int_0^\infty e^{-t} (A - C) dt \right) \\ J^A = \text{agent's payoff} = E \left(\int_0^\infty e^{-t} (g(C) - h(A)) dt \right) \end{cases}$$

How can we make the agent do what we want?

In particular, how can the principal encourage the agent to perform the action A^* ?

KEY POINT #1: The agent's actions and the Brownian motion are unobservable by the principal; she can only observe X .

DYNAMICS REINTERPRETED:

$$\begin{aligned}dX &= A^* dt + dB \\ &= A dt + dB^A\end{aligned}$$

KEY POINT #2: B is a Brownian motion under the original probability measure P and B^A is a Brownian motion under a new probability measure P^A .

PAYOFFS REINTERPRETED:

$$\begin{cases} J^P = E^A \left(\int_0^\infty e^{-t} (A - C) dt \right) \\ J^A = E^A \left(\int_0^\infty e^{-t} (g(C) - h(A)) dt \right) \end{cases}$$

NEW DYNAMICS: Given $A^* = a(W)$ and $C^* = c(W)$, solve

$$\begin{aligned}dW &= (W - g(C^*) + h(A^*))dt + h'(A^*)dB \\ &= (W - g(C^*) + h(A^*) - A^* h'(A^*))dt + h'(A^*)dX \\ W(0) &= w.\end{aligned}$$

The principal will therefore pay the agent $C^* = c(W)$, depending upon the values of W , which she can compute in terms of X .

THEOREM (Sannikov optimality condition for agent)

Assume

h is convex.

Then it is optimal for the agent to use the control A^ .*

New proof of optimality condition for agent

$$\begin{aligned}d(e^{-t}W) &= e^{-t} [(-g(C^*) + h(A^*))dt + h'(A^*)dB] \\ &= e^{-t} [(-g(C^*) + h(A^*) + h'(A^*)(A - A^*))dt + h'(A^*)dB^A]\end{aligned}$$

Integrate from 0 to ∞ :

$$\begin{aligned}-w &= \int_0^\infty e^{-t}(-g(C^*) + h(A^*) + h'(A^*)(A - A^*)) dt \\ &\quad + \int_0^\infty e^{-t}h'(A^*) dB^A\end{aligned}$$

Take expected value with respect to P^A and rewrite:

$$\begin{aligned}w &= E^A \left(\int_0^\infty e^{-t}(g(C^*) - h(A)) dt \right) \\ &\quad + E^A \left(\int_0^\infty e^{-t}(h(A) - h(A^*) - h'(A^*)(A - A^*)) dt \right)\end{aligned}$$

By convexity, the last term is nonnegative, and is zero for $A = A^*$.



EXERCISE: Use dynamic programming to rederive this optimality condition. What is the agent's value function?

THEOREM (Optimality condition for principal)

The principal's value function $u = u(w)$ solves the HJB equation

$$\max_{a,c} \left\{ \frac{(h'(a))^2}{2} u'' + (w - g(c) + h(a))u' - u + a - c \right\} = 0$$

Select

$$a(w), c(w)$$

giving the max in the HJB equation.

- I.Ekeland, *How to build stable relationships with people who lie and cheat*, www.ceremade.dauphine.fr/~ekeland/Articles/13-03-08-ArticleMilano.pdf

Many fascinating PDE/optimality issues remain open.