



HAL
open science

Bang bang trajectories with a double switching time in the minimum time problem

Laura Poggiolini, Marco Spadini, Gianna Stefani

► **To cite this version:**

Laura Poggiolini, Marco Spadini, Gianna Stefani. Bang bang trajectories with a double switching time in the minimum time problem. NETCO 2014 - New Trends in Optimal Control, Jun 2014, Tours, France. hal-01024733

HAL Id: hal-01024733

<https://inria.hal.science/hal-01024733>

Submitted on 18 Jul 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Bang–bang trajectories with a double switching time in the minimum time problem

L. Poggiolini M. Spadini G. Stefani



NetCo 2014 New trends in optimal control
23-27 June 2014 Tours, France.

The minimum time problem

$$T \rightarrow \min$$

$$\dot{\xi}(t) = f_0(\xi(t)) + \sum_{i=1}^m u_i f_i(\xi(t)),$$

$$\xi(0) \in N_0, \quad \xi(T) \in N_f,$$

$$u = (u_1, \dots, u_m) \in Q := [-1, 1]^m$$

The minimum time problem

$$T \rightarrow \min$$

$$\dot{\xi}(t) = f_0(\xi(t)) + \sum_{i=1}^m u_i f_i(\xi(t)),$$

$$\xi(0) \in N_0, \quad \xi(T) \in N_f,$$

$$u = (u_1, \dots, u_m) \in Q := [-1, 1]^m$$

Assume $(\hat{T}, \hat{\xi}, \hat{u})$ is an admissible triplet satisfying PMP with an adjoint covector

$$\hat{\lambda}: t \in [0, \hat{T}] \mapsto \hat{\lambda}(t) \in T^*M$$

The state space M is a finite dimensional manifold

Problem

Give sufficient second order conditions for $\hat{\xi}$ to be a strong local optimal trajectory

Problem

Give sufficient second order conditions for $\widehat{\xi}$ to be a strong local optimal trajectory

Definition ((time,state)-local optimality)

The trajectory $\widehat{\xi}$ is a (time,state)-local minimiser if there is a neighborhood $\widetilde{\mathcal{U}}$ of its graph such that $\widehat{\xi}$ is a minimiser among the admissible trajectories whose graph is in $\widetilde{\mathcal{U}}$.

Problem

Give sufficient second order conditions for $\widehat{\xi}$ to be a strong local optimal trajectory


Definition ((time,state)-local optimality)

The trajectory $\widehat{\xi}$ is a (time,state)-local minimiser if there is a neighborhood $\widetilde{\mathcal{U}}$ of its graph such that $\widehat{\xi}$ is a minimiser among the admissible trajectories whose graph is in $\widetilde{\mathcal{U}}$.

Definition (state-local optimality)

The trajectory $\widehat{\xi}$ is a state-local minimiser if there are neighborhoods \mathcal{U} of its range $\widehat{\xi}([0, \widehat{T}])$, \mathcal{U}_0 of $\widehat{\xi}(0)$ and \mathcal{U}_f of $\widehat{\xi}(\widehat{T})$ such that $\widehat{\xi}$ is a minimiser among the admissible trajectories whose range is in \mathcal{U} , whose initial point is in $N_0 \cap \mathcal{U}_0$ and whose final point is in $N_f \cap \mathcal{U}_f$.

Example



Hamiltonians associated to the vector fields

$$F_i : \ell = (p, q) \in T^*M \mapsto \langle p, f_i(q) \rangle, \quad i = 0, 1, \dots, m$$

Hamiltonians associated to the vector fields

$$F_i : \ell = (p, q) \in T^*M \mapsto \langle p, f_i(q) \rangle, \quad i = 0, 1, \dots, m$$

Reference Hamiltonian

$$\hat{F}_t = F_0 + \sum_{i=1}^m \hat{u}_i(t) F_i$$

Hamiltonians associated to the vector fields

$$F_i : \ell = (p, q) \in T^*M \mapsto \langle p, f_i(q) \rangle, \quad i = 0, 1, \dots, m$$

Reference Hamiltonian

$$\hat{F}_t = F_0 + \sum_{i=1}^m \hat{u}_i(t) F_i$$

Maximised Hamiltonian

$$F^{\max} : \ell \mapsto \sup_{u \in Q} \left\{ F_0(\ell) + \sum_{i=1}^m u_i F_i(\ell) \right\}$$

Hamiltonian vector fields

$$\vec{F}^{\max}, \vec{F}_t, \vec{F}_i$$

Pontryagin Maximum Principle (PMP)

there exist $\hat{\lambda}: [0, \hat{T}] \rightarrow T^*M$, $\rho_0 \in \{0, 1\}$

such that: $\|\hat{\lambda}(0)\| + \rho_0 \neq 0$

$$\pi \hat{\lambda}(t) = \hat{\xi}(t)$$

$$\frac{d}{dt} \hat{\lambda}(t) = \vec{F}_t(\hat{\lambda}(t))$$

$$\hat{F}_t(\hat{\lambda}(t)) = F^{\max}(\hat{\lambda}(t)) \equiv \rho_0$$

$$\hat{\lambda}(0) \Big|_{T_{\hat{\xi}(0)} N_0} = 0, \quad \hat{\lambda}(\hat{T}) \Big|_{T_{\hat{\xi}(\hat{T})} N_f} = 0$$

Pontryagin Maximum Principle (PMP)

there exist $\hat{\lambda}: [0, \hat{T}] \rightarrow T^*M$, $\rho_0 \in \{0, 1\}$

such that: $\|\hat{\lambda}(0)\| + \rho_0 \neq 0$

$$\pi \hat{\lambda}(t) = \hat{\xi}(t)$$

$$\frac{d}{dt} \hat{\lambda}(t) = \vec{F}_t(\hat{\lambda}(t))$$

$$\hat{F}_t(\hat{\lambda}(t)) = F^{\max}(\hat{\lambda}(t)) \equiv \rho_0$$

$$\hat{\lambda}(0) \Big|_{T_{\hat{\xi}(0)} N_0} = 0, \quad \hat{\lambda}(\hat{T}) \Big|_{T_{\hat{\xi}(\hat{T})} N_f} = 0$$

if $\rho_0 = 1$, $\hat{\lambda}$ is called **normal extremal**

if $\rho_0 = 0$, $\hat{\lambda}$ is called **abnormal extremal**

Hamiltonian approach to (time, state)–local optimality 1/2

- the flow of an over-maximised Hamiltonian

$$H: \ell \in T^*M \mapsto H(\ell) \in \mathbb{R}$$

$$H(\ell) \geq F^{\max}(\ell) \quad H(\widehat{\lambda}(t)) = F^{\max}(\widehat{\lambda}(t))$$

$$\mathcal{H}: (t, \ell) \in [0, \widehat{T}] \times T^*M \mapsto \mathcal{H}_t(\ell) \in T^*M$$

$$\widehat{\lambda}(t) = \mathcal{H}_t(\widehat{\lambda}(0)), \text{ i.e. } \dot{\widehat{\lambda}}(t) = \overrightarrow{H}(\widehat{\lambda}(t))$$

Hamiltonian approach to (time, state)–local optimality 1/2

- the flow of an over-maximised Hamiltonian

$$H: \ell \in T^*M \mapsto H(\ell) \in \mathbb{R}$$

$$H(\ell) \geq F^{\max}(\ell) \quad H(\widehat{\lambda}(t)) = F^{\max}(\widehat{\lambda}(t))$$

$$\mathcal{H}: (t, \ell) \in [0, \widehat{T}] \times T^*M \mapsto \mathcal{H}_t(\ell) \in T^*M$$

$$\widehat{\lambda}(t) = \mathcal{H}_t(\widehat{\lambda}(0)), \text{ i.e. } \dot{\widehat{\lambda}}(t) = \overrightarrow{H}(\widehat{\lambda}(t))$$

- a smooth function $\alpha: M \rightarrow \mathbb{R}$ such that $d\alpha(\widehat{\xi}(0)) = \widehat{\lambda}(0)$

$$\Lambda := \{(d\alpha(x), x)\}$$


Hamiltonian approach to (time, state)–local optimality 2/2

$$\begin{array}{ccc}
 \bullet & [0, \widehat{T}] \times \Lambda & \xrightarrow{\text{id} \times \mathcal{H}} & [0, \widehat{T}] \times T^*M \\
 & & & \downarrow \text{id} \times \pi \\
 & & & [0, \widehat{T}] \times M \\
 & \swarrow (\text{id} \times \pi \mathcal{H})^{-1} & &
 \end{array}$$


Property

$\mathcal{H}^*(p dq - H_t dt)$ is exact on $[0, \widehat{T}] \times \Lambda$


Lifting trajectories



Lifting trajectories



Lifting trajectories



$\mathcal{H}^*(p dq - H_t dt)$ is exact \implies

$$0 = \oint \omega = \int_{(\text{id} \times \pi \mathcal{H})^{-1}(\xi)} \omega + \int_{(\text{id} \times \pi \mathcal{H})^{-1}(\Gamma_f)} \omega - \int_{(\text{id} \times \pi \mathcal{H})^{-1}(\hat{\xi})} \omega + \int_{(\text{id} \times \pi \mathcal{H})^{-1}(\Gamma_0)} \omega$$

$\mathcal{H}^*(p dq - H_t dt)$ is exact \implies

$$0 = \oint \omega = \int_{(\text{id} \times \pi \mathcal{H})^{-1}(\xi)} \omega + \int_{(\text{id} \times \pi \mathcal{H})^{-1}(\Gamma_f)} \omega - \int_{(\text{id} \times \pi \mathcal{H})^{-1}(\hat{\xi})} \omega + \int_{(\text{id} \times \pi \mathcal{H})^{-1}(\Gamma_0)} \omega$$

$$0 \leq p_0(T - \hat{T}) + O((T - \hat{T})^2)$$

If $p_0 = 1 \implies \hat{\xi}$ is a (time, state)-local optimal trajectory

Hamiltonian approach to state–local optimality 1/2

- the flow of an over-maximised Hamiltonian

$$H: \ell \in T^*M \mapsto H(\ell) \in \mathbb{R}$$


$$H(\ell) \geq F^{\max}(\ell) \quad H(\widehat{\lambda}(t)) = F^{\max}(\widehat{\lambda}(t))$$

$$\mathcal{H}: (t, \ell) \in (-\delta, \widehat{T} + \delta) \times T^*M \mapsto \mathcal{H}_t(\ell) \in T^*M$$

$$\widehat{\lambda}(t) = \mathcal{H}_t(\widehat{\lambda}(0))$$

- a smooth function $\alpha: M \rightarrow \mathbb{R}$ such that
 - $d\alpha(\widehat{\xi}(0)) = \widehat{\lambda}(0)$
 - $\Lambda = \{(d\alpha(x), x)\}$ is transverse to $\{H = p_0\}$ in $\widehat{\lambda}(0)$
- $\Lambda_0 := \Lambda \cap \{H = p_0\}$ is a $(n - 1)$ -dim manifold of T^*M


Hamiltonian approach to state–local optimality 2/2




Property

$\mathcal{H}^*(p dq)$ is exact on $(-\delta, \widehat{T} + \delta) \times \Lambda_0$


Lifting trajectories



Lifting trajectories



Lifting trajectories



$\mathcal{H}^*(p dq)$ is exact \implies

$$0 = \oint \omega = \int_{(\pi\mathcal{H})^{-1}(\xi)} \omega + \int_{(\pi\mathcal{H})^{-1}(\gamma_f)} \omega - \int_{(\pi\mathcal{H})^{-1}(\hat{\xi})} \omega + \int_{(\pi\mathcal{H})^{-1}(\gamma_0)} \omega$$

$\mathcal{H}^*(p \, dq)$ is exact \implies

$$0 = \oint \omega = \int_{(\pi\mathcal{H})^{-1}(\xi)} \omega + \int_{(\pi\mathcal{H})^{-1}(\gamma_f)} \omega - \int_{(\pi\mathcal{H})^{-1}(\hat{\xi})} \omega + \int_{(\pi\mathcal{H})^{-1}(\gamma_0)} \omega$$

$$\begin{aligned} p_0(T - \hat{T}) &\geq p_0(t_f - \hat{T}) + \alpha(q_f) - \alpha(\hat{\xi}(0)) + \frac{t_0^2}{2} h_1 \cdot h_1 \cdot \alpha(\bar{q}) \\ &= C(\xi(T)) - C(\hat{\xi}(\hat{T})) + \frac{t_0^2}{2} h_1 \cdot h_1 \cdot \alpha(\bar{q}) \end{aligned}$$

$$C(x) := p_0(\psi^{\mathbb{R}}(x)) + \alpha(\pi\psi^{\wedge_0}(x)) \quad (\psi^{\mathbb{R}}, \psi^{\wedge_0}) := (\pi\mathcal{H})^{-1}$$


$$(t_f, q_f) := (\psi^{\mathbb{R}}, \pi\psi^{\wedge_0})(\xi(T))$$

$$(t_0, q_0) := (\psi^{\mathbb{R}}, \pi\psi^{\wedge_0})(\xi(0))$$

$$\bar{q} = \exp \bar{s} h_1(q_0)$$


Bang-bang controls

$$\hat{u}(t) \in \text{Vertexes}(Q) \quad \forall t \in [0, \hat{T}] \setminus \{\hat{\tau}_j : j = 1, \dots, r\}$$



Simple, double and multiple switches


- **simple switch** $\hat{\tau}$ is a **simple switching time** if only one control component switches at time $\hat{\tau}$
- **double switch** $\hat{\tau}$ is a **double switching time** if two control components switch at time $\hat{\tau}$
- **j -multiplicity switch** $\hat{\tau}$ is a **j multiplicity switching time** if j control components switch at time $\hat{\tau}$



Bang-bang with multiple switches

Simple and double switches

- there exists only double switching time $\hat{\tau}$;
- a certain number of simple switching times $\hat{\theta}_{0j}$ occurs before $\hat{\tau}$ and a certain number of simple switching times $\hat{\theta}_{1j}$ occur after the double switch;



Simplified case

$$T \rightarrow \min,$$

$$\dot{\xi}(t) = f_0(\xi(t)) + u_1(t)f_1(\xi(t)) + u_2(t)f_2(\xi(t)),$$

$$\xi(0) \in N_0, \quad \xi(T) \in N_f,$$

$$|u_i(t)| \leq 1 \quad i = 1, 2, \text{ a.e. } t \in [0, T].$$


Simplified case

$$T \rightarrow \min,$$

$$\dot{\xi}(t) = f_0(\xi(t)) + u_1(t)f_1(\xi(t)) + u_2(t)f_2(\xi(t)),$$

$$\xi(0) \in N_0, \quad \xi(T) \in N_f,$$

$$|u_i(t)| \leq 1 \quad i = 1, 2, \text{ a.e. } t \in [0, T].$$




Simplified case

$$T \rightarrow \min,$$

$$\dot{\xi}(t) = f_0(\xi(t)) + u_1(t)f_1(\xi(t)) + u_2(t)f_2(\xi(t)),$$

$$\xi(0) \in N_0, \quad \xi(T) \in N_f,$$

$$|u_i(t)| \leq 1 \quad i = 1, 2, \text{ a.e. } t \in [0, T].$$



$$H_1(\ell) := \langle \ell, h_1(\pi\ell) \rangle \quad H_2(\ell) := \langle \ell, h_2(\pi\ell) \rangle$$

Assumptions: Regularity along the bang arcs

$$\text{PMP} \implies \begin{aligned} \hat{u}_1(t)F_1(\hat{\lambda}(t)) &= \hat{u}_1(t)\langle \hat{\lambda}(t), f_1(\hat{\xi}(t)) \rangle \geq 0 \\ \hat{u}_2(t)F_2(\hat{\lambda}(t)) &= \hat{u}_2(t)\langle \hat{\lambda}(t), f_2(\hat{\xi}(t)) \rangle \geq 0. \end{aligned}$$

Assumptions: Regularity along the bang arcs

$$\text{PMP} \implies \begin{aligned} \hat{u}_1(t)F_1(\hat{\lambda}(t)) &= \hat{u}_1(t)\langle \hat{\lambda}(t), f_1(\hat{\xi}(t)) \rangle \geq 0 \\ \hat{u}_2(t)F_2(\hat{\lambda}(t)) &= \hat{u}_2(t)\langle \hat{\lambda}(t), f_2(\hat{\xi}(t)) \rangle \geq 0. \end{aligned}$$

Regularity along the bang arcs

If $t \neq \hat{\tau}$, then \hat{u} is the only control that gives the maximised Hamiltonian:

$$\hat{u}_1(t)F_1(\hat{\lambda}(t)) = \hat{u}_1(t)\langle \hat{\lambda}(t), f_1(\hat{\xi}(t)) \rangle > 0$$

$$\hat{u}_2(t)F_2(\hat{\lambda}(t)) = \hat{u}_2(t)\langle \hat{\lambda}(t), f_2(\hat{\xi}(t)) \rangle > 0$$

$$\text{i.e.} \quad \begin{array}{lll} F_1(\hat{\lambda}(t)) < 0 & F_2(\hat{\lambda}(t)) < 0 & \text{if } t \in [0, \hat{\tau}) \\ F_1(\hat{\lambda}(t)) > 0 & F_2(\hat{\lambda}(t)) > 0 & \text{if } t \in (\hat{\tau}, \hat{T}] \end{array}$$

Assumptions: Regularity along the bang arcs

$$\text{PMP} \implies \begin{aligned} \hat{u}_1(t)F_1(\hat{\lambda}(t)) &= \hat{u}_1(t)\langle \hat{\lambda}(t), f_1(\hat{\xi}(t)) \rangle \geq 0 \\ \hat{u}_2(t)F_2(\hat{\lambda}(t)) &= \hat{u}_2(t)\langle \hat{\lambda}(t), f_2(\hat{\xi}(t)) \rangle \geq 0. \end{aligned}$$

Regularity along the bang arcs

If $t \neq \hat{\tau}$, then \hat{u} is the only control that gives the maximised Hamiltonian:

$$\hat{u}_1(t)F_1(\hat{\lambda}(t)) = \hat{u}_1(t)\langle \hat{\lambda}(t), f_1(\hat{\xi}(t)) \rangle > 0$$

$$\hat{u}_2(t)F_2(\hat{\lambda}(t)) = \hat{u}_2(t)\langle \hat{\lambda}(t), f_2(\hat{\xi}(t)) \rangle > 0$$

$$\text{i.e.} \quad \begin{array}{lll} F_1(\hat{\lambda}(t)) < 0 & F_2(\hat{\lambda}(t)) < 0 & \text{if } t \in [0, \hat{\tau}) \\ F_1(\hat{\lambda}(t)) > 0 & F_2(\hat{\lambda}(t)) > 0 & \text{if } t \in (\hat{\tau}, T] \end{array}$$

$$\implies \left. \frac{d}{dt} F_i(\hat{\lambda}(t)) \right|_{t=\hat{\tau}^-} \geq 0 \quad \left. \frac{d}{dt} F_i(\hat{\lambda}(t)) \right|_{t=\hat{\tau}^+} \geq 0$$

Assumptions: Regularity at the switching time 1/2


Regularity at the switching times

$$\left. \frac{d}{dt} F_i(\widehat{\lambda}(t)) \right|_{t=\widehat{\tau}^-} > 0 \quad \left. \frac{d}{dt} F_i(\widehat{\lambda}(t)) \right|_{t=\widehat{\tau}^+} > 0 \quad i = 1, 2.$$

Assumptions: Regularity at the switching time 1/2

Regularity at the switching times

$$\left. \frac{d}{dt} F_i(\hat{\lambda}(t)) \right|_{t=\hat{\tau}^-} > 0 \quad \left. \frac{d}{dt} F_i(\hat{\lambda}(t)) \right|_{t=\hat{\tau}^+} > 0 \quad i = 1, 2.$$



$$h_1 := f_0 - f_1 - f_2$$

$$h_2 := f_0 + f_1 + f_2$$

$$k_1 := f_0 + f_1 - f_2 = h_1 + 2f_1 = h_2 - 2f_2$$

$$k_2 := f_0 - f_1 + f_2 = h_1 + 2f_2 = h_2 - 2f_1$$

$$K_1(\ell) = \langle \ell, k_1(\pi\ell) \rangle$$

$$K_2(\ell) = \langle \ell, k_2(\pi\ell) \rangle$$

Assumptions: Regularity at the switching time 2/2

$$\langle \hat{\lambda}(\hat{\tau}), [h_1, k_1] (\hat{\xi}(\hat{\tau})) \rangle = \sigma \left(\vec{H}_1, \vec{K}_1 \right) (\hat{\lambda}(\hat{\tau})) > 0$$

$$\langle \hat{\lambda}(\hat{\tau}), [h_1, k_2] (\hat{\xi}(\hat{\tau})) \rangle = \sigma \left(\vec{H}_1, \vec{K}_2 \right) (\hat{\lambda}(\hat{\tau})) > 0$$

$$\langle \hat{\lambda}(\hat{\tau}), [k_1, h_2] (\hat{\xi}(\hat{\tau})) \rangle = \sigma \left(\vec{K}_1, \vec{H}_2 \right) (\hat{\lambda}(\hat{\tau})) > 0$$

$$\langle \hat{\lambda}(\hat{\tau}), [k_2, h_2] (\hat{\xi}(\hat{\tau})) \rangle = \sigma \left(\vec{K}_2, \vec{H}_2 \right) (\hat{\lambda}(\hat{\tau})) > 0$$

The maximised flow

- for $\nu = 1, 2$ let $\tau_\nu(\ell)$ be the unique solution to

$$2F_\nu \circ \exp \tau_\nu(\ell) \vec{H}_1(\ell) = (K_\nu - H_1) \circ \exp \tau_\nu(\ell) \vec{H}_1(\ell) = 0$$

in a neighborhood of $(\hat{\tau}, \hat{\lambda}(0))$;

The maximised flow

- for $v = 1, 2$ let $\tau_v(\ell)$ be the unique solution to

$$2F_v \circ \exp \tau_v(\ell) \vec{H}_1(\ell) = (K_v - H_1) \circ \exp \tau_v(\ell) \vec{H}_1(\ell) = 0$$

in a neighborhood of $(\hat{\tau}, \hat{\lambda}(0))$;

- choose

$$\theta_1(\ell) := \min\{\tau_1(\ell), \tau_2(\ell)\};$$

The maximised flow

- for $\nu = 1, 2$ let $\tau_\nu(\ell)$ be the unique solution to

$$2F_\nu \circ \exp \tau_\nu(\ell) \vec{H}_1(\ell) = (K_\nu - H_1) \circ \exp \tau_\nu(\ell) \vec{H}_1(\ell) = 0$$

in a neighborhood of $(\hat{\tau}, \hat{\lambda}(0))$;

- choose

$$\theta_1(\ell) := \min\{\tau_1(\ell), \tau_2(\ell)\};$$

- for $\nu = 1, 2$, let $\tau_\nu^2(\ell)$ be the unique solution to

$$\begin{aligned} 2F_{3-\nu} \circ \exp(\tau_\nu^2(\ell) - \tau_\nu(\ell)) \vec{K}_\nu \circ \exp \tau_\nu(\ell) \vec{H}_1(\ell) \\ = (H_2 - K_\nu) \circ \exp(\tau_\nu^2(\ell) - \tau_\nu(\ell)) \vec{K}_\nu \circ \exp \tau_\nu(\ell) \vec{H}_1(\ell) = 0 \end{aligned}$$

in a neighborhood of $(\hat{\tau}, \hat{\lambda}(0))$;

The maximised flow

- for $\nu = 1, 2$ let $\tau_\nu(\ell)$ be the unique solution to

$$2F_\nu \circ \exp \tau_\nu(\ell) \vec{H}_1(\ell) = (K_\nu - H_1) \circ \exp \tau_\nu(\ell) \vec{H}_1(\ell) = 0$$

in a neighborhood of $(\hat{\tau}, \hat{\lambda}(0))$;

- choose

$$\theta_1(\ell) := \min\{\tau_1(\ell), \tau_2(\ell)\};$$

- for $\nu = 1, 2$, let $\tau_\nu^2(\ell)$ be the unique solution to

$$\begin{aligned} 2F_{3-\nu} \circ \exp(\tau_\nu^2(\ell) - \tau_\nu(\ell)) \vec{K}_\nu \circ \exp \tau_\nu(\ell) \vec{H}_1(\ell) \\ = (H_2 - K_\nu) \circ \exp(\tau_\nu^2(\ell) - \tau_\nu(\ell)) \vec{K}_\nu \circ \exp \tau_\nu(\ell) \vec{H}_1(\ell) = 0 \end{aligned}$$

in a neighborhood of $(\hat{\tau}, \hat{\lambda}(0))$;

- choose

$$\theta_2(\ell) = \begin{cases} \tau_1^2(\ell) & \text{if } \tau_1(\ell) \leq \tau_2(\ell), \\ \tau_2^2(\ell) & \text{if } \tau_2(\ell) < \tau_1(\ell). \end{cases}$$

- $S_0 := \{(t, \ell) : t \in [-\delta, \theta_1(\ell)]\}$

$$\mathcal{F}_t^{\max}(\ell) = \exp t\vec{H}_1(\ell);$$

- $S_1 := \{(t, \ell) : \theta_1(\ell) = \tau_1(\ell), t \in (\theta_1(\ell), \theta_2(\ell))\}$

$$\mathcal{F}_t^{\max}(\ell) = \exp(t - \theta_1(\ell))\vec{K}_1 \circ \exp \theta_1(\ell)\vec{H}_1(\ell);$$

- $S_1^2 := \{(t, \ell) : \theta_1(\ell) = \tau_1(\ell), t \in (\theta_2(\ell), \hat{T} + \delta)\}$

$$\mathcal{F}_t^{\max}(\ell) = \exp(t - \theta_2(\ell))\vec{H}_2(\ell) \circ \exp(\theta_2(\ell) - \theta_1(\ell))\vec{K}_1 \circ \exp \theta_1(\ell)\vec{H}_1(\ell);$$

- $S_2 := \{(t, \ell) : \theta_1(\ell) = \tau_2(\ell), t \in (\theta_1(\ell), \theta_2(\ell))\}$

- $S_2^2 := \{(t, \ell) : \theta_1(\ell) = \tau_2(\ell), t \in (\theta_2(\ell), \hat{T} + \delta)\}$


Finite-dimensional subproblem

Allow for different perturbed switching times of the control components

$$\tau_1 := \hat{\tau} + \varepsilon_1 \quad \tau_2 := \hat{\tau} + \varepsilon_2$$

and for variations of the final time

$$\tau_3 := \hat{T} + \varepsilon_3.$$



if $\varepsilon_1 < \varepsilon_2$

$T \rightarrow \min$

$$\dot{\xi}(t) = \begin{cases} h_1(\xi(t)) & t < \tau_1 \\ k_1(\xi(t)) & \tau_1 < t < \tau_2 \\ h_2(\xi(t)) & \tau_2 < t < \tau_3 \end{cases}$$

$$\xi(0) \in N_0, \quad \xi(T) \in N_f,$$

if $\varepsilon_1 < \varepsilon_2$

$T \rightarrow \min$

$$\dot{\xi}(t) = \begin{cases} h_1(\xi(t)) & t < \tau_1 \\ k_1(\xi(t)) & \tau_1 < t < \tau_2 \\ h_2(\xi(t)) & \tau_2 < t < \tau_3 \end{cases}$$

$$\xi(0) \in N_0, \quad \xi(T) \in N_f,$$

if $\varepsilon_2 < \varepsilon_1$

$T \rightarrow \min$

$$\dot{\xi}(t) = \begin{cases} h_1(\xi(t)) & t < \tau_2 \\ k_2(\xi(t)) & \tau_2 < t < \tau_1 \\ h_2(\xi(t)) & \tau_1 < t < \tau_3 \end{cases}$$

$$\xi(0) \in N_0, \quad \xi(T) \in N_f,$$

Assumption: coercivity of the second order approximations

Coercivity of the second order approximations

Each second order approximation J''_{ν} $\nu = 1, 2$ is coercive on the half-space of the linearised constraints of the finite-dimensional subproblem.

Assumption: coercivity of the second order approximations

Coercivity of the second order approximations

Each second order approximation J''_{ν} $\nu = 1, 2$ is coercive on the half-space of the linearised constraints of the finite-dimensional subproblem.

$$V_{0,\nu}^+ := \left\{ (\delta x, a_1, b, a_2) \in T_{\hat{\xi}(0)} N_0 \times \mathbb{R} \times \mathbb{R}^+ \times \mathbb{R} : \right. \\ \left. \delta x + a_1 g_1(\hat{\xi}(0)) + b j_{\nu}(\hat{\xi}(0)) + a_2 g_2(\hat{\xi}(0)) \in T_{\hat{x}_0} \hat{N}_f \right\}$$

$$J''_{\nu}[\delta x, a_1, b, a_2] = 2 \delta x \cdot (a_1 g_1 + b j_{\nu} + a_2 g_2) \cdot \hat{\beta}(\hat{\xi}(0)) + \\ + (a_1 g_1 + b j_{\nu} + a_2 g_2)^2 \cdot \hat{\beta}(\hat{\xi}(0)) \\ + a_1 b [g_1, j_{\nu}] \cdot \hat{\beta}(\hat{\xi}(0)) + a_1 a_2 [g_1, g_2] \cdot \hat{\beta}(\hat{\xi}(0)) + b a_2 [j_{\nu}, g_2] \cdot \hat{\beta}(\hat{x}_0),$$

$$\hat{N}_f := \hat{S}_{\hat{\tau}}^{-1}(N_f) \quad g_{\nu} := \hat{S}_{\hat{\tau}^*}^{-1} h_{\nu} \circ \hat{S}_{\hat{\tau}} \quad j_{\nu} := \hat{S}_{\hat{\tau}^*}^{-1} k_{\nu} \circ \hat{S}_{\hat{\tau}}$$

$$\hat{\beta}: M \rightarrow \mathbb{R}: d\hat{\beta}(\hat{\xi}(0)) = -\hat{\lambda}(0)$$

The coercivity of J_v'' allows one to prove that there exists a function α such that

- $\Lambda_0 := \{d\alpha(x) : x \in M\} \cap \{H_1 = p_0\}$ and $M_0 := \pi\Lambda_0 = \{x \in M : h_1 \cdot \alpha(x) = p_0\}$ are $(n-1)$ -dimensional manifolds
- For $v = 1, 2$ define


$$M_v := \left\{ \pi \mathcal{F}_{\tau_v(\ell)}^{\max}(\ell) : \ell \in \Lambda_0, \quad \tau_v(\ell) = \theta_1(\ell) \right\},$$

$$M_v^2 := \left\{ \pi \mathcal{F}_{\tau_v^2(\ell)}^{\max}(\ell) : \ell \in \Lambda_0, \quad \tau_v(\ell) = \theta_1(\ell) \right\}.$$


Then these four half-manifolds separate the images of the five smooth pieces of the projected maximised Hamiltonian flow.

Picture for $\nu = 1$

$$M_1 := \left\{ \pi \mathcal{F}_{\tau_1}^{\max}(\ell) : \ell \in \Lambda_0 \right\},$$



$$M_1^2 := \left\{ \pi \mathcal{F}_{\tau_1^2}^{\max}(\ell) : \ell \in \Lambda_0 \right\}.$$



Invertibility of the projected maximised flow

1: The degenerate case $d\tau_1|_{\mathcal{T}_{\hat{\lambda}(0)}\Lambda_0} \equiv d\tau_2|_{\mathcal{T}_{\hat{\lambda}(0)}\Lambda_0}$

- **Clarke's inverse function theorem**

if the convex hull of the Jacobians of each of the five smooth pieces at $(\hat{\tau}, \hat{\lambda}(0))$ is made of invertible matrices only

$\implies \pi_{\mathcal{F}^{\max}}: (-\delta, \hat{\mathcal{T}} + \delta) \times \Lambda_0 \rightarrow M$ is locally one-to-one from a neighborhood of $(\hat{\tau}, \hat{\lambda}(0))$ onto a neighborhood of $\hat{\xi}(\hat{\tau})$

Invertibility of the projected maximised flow

2: The nondegenerate case $d\tau_1|_{T_{\hat{\lambda}(0)}\Lambda_0} \neq d\tau_2|_{T_{\hat{\lambda}(0)}\Lambda_0}$

- Consider the tangent cones to the five sectors where \mathcal{F}^{\max} has a different smooth expression: **the interior of each cone is non-empty**

Invertibility of the projected maximised flow

2: The nondegenerate case $d\tau_1|_{T_{\hat{\lambda}(0)}\Lambda_0} \neq d\tau_2|_{T_{\hat{\lambda}(0)}\Lambda_0}$

- Consider the tangent cones to the five sectors where \mathcal{F}^{\max} has a different smooth expression: **the interior of each cone is non-empty**
- Show that the linearisation of each of the five smooth pieces of $\pi\mathcal{F}^{\max}$ is a proper linear function with positive determinant and degree = 1: for each couple of neighbouring tangent cones take $(\delta t_1, \delta \ell_1)$ and $(\delta t_2, \delta \ell_2)$ in each of the two cones and show that their images under the linearisation of the projected maximised flow do not overlap

Invertibility of the projected maximised flow

2: The nondegenerate case $d\tau_1|_{T_{\hat{\lambda}(0)}\Lambda_0} \neq d\tau_2|_{T_{\hat{\lambda}(0)}\Lambda_0}$

- Consider the tangent cones to the five sectors where \mathcal{F}^{\max} has a different smooth expression: **the interior of each cone is non-empty**
- Show that the linearisation of each of the five smooth pieces of $\pi\mathcal{F}^{\max}$ is a proper linear function with positive determinant and degree = 1: for each couple of neighbouring tangent cones take $(\delta t_1, \delta \ell_1)$ and $(\delta t_2, \delta \ell_2)$ in each of the two cones and show that their images under the linearisation of the projected maximised flow do not overlap
- Show that there exists $\delta y \in T_{\hat{\xi}(\hat{\tau})}M$ such that $(\pi\mathcal{F}^{\max})_*^{-1}(\{\delta y\})$ is a singleton

Invertibility of the projected maximised flow

2: The nondegenerate case $d\tau_1|_{T_{\widehat{\lambda}(0)}}\Lambda_0 \neq d\tau_2|_{T_{\widehat{\lambda}(0)}}\Lambda_0$

- Consider the tangent cones to the five sectors where \mathcal{F}^{\max} has a different smooth expression: **the interior of each cone is non-empty**
- Show that the linearisation of each of the five smooth pieces of $\pi\mathcal{F}^{\max}$ is a proper linear function with positive determinant and degree = 1: for each couple of neighbouring tangent cones take $(\delta t_1, \delta \ell_1)$ and $(\delta t_2, \delta \ell_2)$ in each of the two cones and show that their images under the linearisation of the projected maximised flow do not overlap
- Show that there exists $\delta y \in T_{\widehat{\xi}(\widehat{\tau})}M$ such that $(\pi\mathcal{F}^{\max})_*^{-1}(\{\delta y\})$ is a singleton
- $\implies \deg\left(\pi\mathcal{F}^{\max}, \mathcal{O}(\widehat{\lambda}(0)), \widehat{\xi}(\widehat{\tau})\right) = 1$

Invertibility of the projected maximised flow

2: The nondegenerate case $d\tau_1|_{T_{\hat{\lambda}(0)}\Lambda_0} \neq d\tau_2|_{T_{\hat{\lambda}(0)}\Lambda_0}$

- Consider the tangent cones to the five sectors where \mathcal{F}^{\max} has a different smooth expression: **the interior of each cone is non-empty**
- Show that the linearisation of each of the five smooth pieces of $\pi\mathcal{F}^{\max}$ is a proper linear function with positive determinant and degree = 1: for each couple of neighbouring tangent cones take $(\delta t_1, \delta \ell_1)$ and $(\delta t_2, \delta \ell_2)$ in each of the two cones and show that their images under the linearisation of the projected maximised flow do not overlap
- Show that there exists $\delta y \in T_{\hat{\xi}(\hat{\tau})}M$ such that $(\pi\mathcal{F}^{\max})_*^{-1}(\{\delta y\})$ is a singleton
- $\implies \deg(\pi\mathcal{F}^{\max}, \mathcal{O}(\hat{\lambda}(0)), \hat{\xi}(\hat{\tau})) = 1$
- P-Spadini 2013 (based on Pang-Ralph 1996) $\implies \pi\mathcal{F}^{\max}$ is a Lipschitz homeomorphism from $\mathcal{O}(\hat{\tau}, \hat{\lambda}(0))$ on $\mathcal{O}(\hat{\xi}(\hat{\tau}))$.

The reference trajectory

The reference trajectory

$$\widehat{\xi}: [0, \widehat{T}] \rightarrow M$$

has no self-intersection.

The reference trajectory

The reference trajectory

$$\widehat{\xi}: [0, \widehat{T}] \rightarrow M$$

has no self-intersection.

The Hamiltonian methods can be applied:

- if $p_0 = 1$ then $\widehat{\xi}$ is a state-locally optimal trajectory;
- if $p_0 = 0$ then $\widehat{\xi}$ is isolated among admissible trajectories.

Assumption: coercivity of the second order approximations

Coercivity of the second order approximations

Each second order approximation J''_{ν} $\nu = 1, 2$ is coercive on the half-space of the linearised constraints of the finite-dimensional subproblem **wih zero total variation of the times** .

Assumption: coercivity of the second order approximations

Coercivity of the second order approximations

Each second order approximation J''_{ν} $\nu = 1, 2$ is coercive on the half-space of the linearised constraints of the finite-dimensional subproblem **wih zero total variation of the times** .

$$V_{0,\nu}^+ := \left\{ (\delta x, a_1, b, a_2) \in T_{\hat{\xi}(0)} N_0 \times \mathbb{R} \times \mathbb{R}^+ \times \mathbb{R} : a_1 + b + a_2 = 0 \right. \\ \left. \delta x + a_1 g_1(\hat{\xi}(0)) + b j_{\nu}(\hat{\xi}(0)) + a_2 g_2(\hat{\xi}(0)) \in T_{\hat{x}_0} \hat{N}_f \right\}$$

$$J''_{\nu}[\delta x, a_1, b, a_2] = 2 \delta x \cdot (a_1 g_1 + b j_{\nu} + a_2 g_2) \cdot \hat{\beta}(\hat{\xi}(0)) + \\ + (a_1 g_1 + b j_{\nu} + a_2 g_2)^2 \cdot \hat{\beta}(\hat{\xi}(0)) \\ + a_1 b [g_1, j_{\nu}] \cdot \hat{\beta}(\hat{\xi}(0)) + a_1 a_2 [g_1, g_2] \cdot \hat{\beta}(\hat{\xi}(0)) + b a_2 [j_{\nu}, g_2] \cdot \hat{\beta}(\hat{x}_0),$$

$$\hat{N}_f := \hat{S}_{\hat{T}}^{-1}(N_f) \quad g_{\nu} := \hat{S}_{\hat{\tau}^*}^{-1} h_{\nu} \circ \hat{S}_{\hat{\tau}} \quad j_{\nu} := \hat{S}_{\hat{\tau}^*}^{-1} k_{\nu} \circ \hat{S}_{\hat{\tau}}$$

$$\hat{\beta}: M \rightarrow \mathbb{R} : d\hat{\beta}(\hat{\xi}(0)) = -\hat{\lambda}(0)$$

The coercivity of J'' allows one to prove that there exists a function α such that


$$\text{id} \times \pi \mathcal{F}^{\max}: [0, \widehat{T}] \times \{d\alpha(x) : x \in M\} \rightarrow [0, \widehat{T}] \times M$$

is locally one-to-one at each $(t, \widehat{\lambda}(0))$, $t \in [0, \widehat{T}]$, i.e.

$$\pi \mathcal{F}_t^{\max}: \{d\alpha(x) : x \in M\} \rightarrow M$$

is locally one-to-one at $\widehat{\lambda}(0)$, for each $t \in [0, \widehat{T}]$.

Freeze at $t = \hat{\tau}$



Invertibility of the projected maximised flow

1: The degenerate case $d\tau_1|_{T_{\hat{\lambda}(0)}\Lambda} \equiv d\tau_2|_{T_{\hat{\lambda}(0)}\Lambda}$

Clarke's inverse function theorem

if the convex hull of the Jacobians of each of the five smooth pieces of $\pi\mathcal{F}_{\hat{\tau}}^{\max}$ at $\hat{\lambda}(0)$ is made of invertible matrices only

$\implies \pi\mathcal{F}_{\hat{\tau}}^{\max}: \Lambda \rightarrow M$ is locally one-to-one from a neighborhood of $\hat{\lambda}(0)$ onto a neighborhood of $\hat{\xi}(\hat{\tau})$

Invertibility of the projected maximised flow

2: The nondegenerate case $d\tau_1|_{T_{\hat{\lambda}(0)}\Lambda} \neq d\tau_2|_{T_{\hat{\lambda}(0)}\Lambda}$

- Consider the tangent cones to the five sectors where $\pi\mathcal{F}_{\hat{\tau}}^{\max}$ has a different smooth expression: **the interior of each cone is non-empty**

Invertibility of the projected maximised flow

2: The nondegenerate case $d\tau_1|_{T_{\hat{\lambda}(0)}\Lambda} \neq d\tau_2|_{T_{\hat{\lambda}(0)}\Lambda}$

- Consider the tangent cones to the five sectors where $\pi\mathcal{F}_{\hat{\tau}}^{\max}$ has a different smooth expression: **the interior of each cone is non-empty**
- Show that the linearisation of each of the five smooth pieces of $\pi\mathcal{F}_{\hat{\tau}}^{\max}$ is a proper linear function with positive determinant and degree = 1: for each couple of neighbouring tangent cones take $\delta\ell_1$ and $\delta\ell_2$ in each of the two cones and show that their images under the linearisation of the projected maximised flow do not overlap

Invertibility of the projected maximised flow

2: The nondegenerate case $d\tau_1|_{T_{\hat{\lambda}(0)}\Lambda} \neq d\tau_2|_{T_{\hat{\lambda}(0)}\Lambda}$

- Consider the tangent cones to the five sectors where $\pi\mathcal{F}_{\hat{\tau}}^{\max}$ has a different smooth expression: **the interior of each cone is non-empty**
- Show that the linearisation of each of the five smooth pieces of $\pi\mathcal{F}_{\hat{\tau}}^{\max}$ is a proper linear function with positive determinant and degree = 1: for each couple of neighbouring tangent cones take $\delta\ell_1$ and $\delta\ell_2$ in each of the two cones and show that their images under the linearisation of the projected maximised flow do not overlap
- Show that there exists $\delta y \in T_{\hat{\xi}(\hat{\tau})}M$ such that $(\pi\mathcal{F}_{\hat{\tau}}^{\max})_*^{-1}(\{\delta y\})$ is a singleton

Invertibility of the projected maximised flow

2: The nondegenerate case $d\tau_1|_{T_{\hat{\lambda}(0)}} \wedge \neq d\tau_2|_{T_{\hat{\lambda}(0)}} \wedge$

- Consider the tangent cones to the five sectors where $\pi\mathcal{F}_{\hat{\tau}}^{\max}$ has a different smooth expression: **the interior of each cone is non-empty**
- Show that the linearisation of each of the five smooth pieces of $\pi\mathcal{F}_{\hat{\tau}}^{\max}$ is a proper linear function with positive determinant and degree = 1: for each couple of neighbouring tangent cones take $\delta\ell_1$ and $\delta\ell_2$ in each of the two cones and show that their images under the linearisation of the projected maximised flow do not overlap
- Show that there exists $\delta y \in T_{\hat{\xi}(\hat{\tau})}M$ such that $(\pi\mathcal{F}_{\hat{\tau}}^{\max})_*^{-1}(\{\delta y\})$ is a singleton
- $\implies \deg\left(\pi\mathcal{F}_{\hat{\tau}}^{\max}, \mathcal{O}(\hat{\lambda}(0)), \hat{\xi}(\hat{\tau})\right) = 1$

Invertibility of the projected maximised flow

2: The nondegenerate case $d\tau_1|_{T_{\hat{\lambda}(0)}} \wedge \neq d\tau_2|_{T_{\hat{\lambda}(0)}} \wedge$

- Consider the tangent cones to the five sectors where $\pi\mathcal{F}_{\hat{\tau}}^{\max}$ has a different smooth expression: **the interior of each cone is non-empty**
- Show that the linearisation of each of the five smooth pieces of $\pi\mathcal{F}_{\hat{\tau}}^{\max}$ is a proper linear function with positive determinant and degree = 1: for each couple of neighbouring tangent cones take $\delta\ell_1$ and $\delta\ell_2$ in each of the two cones and show that their images under the linearisation of the projected maximised flow do not overlap
- Show that there exists $\delta y \in T_{\hat{\xi}(\hat{\tau})} M$ such that $(\pi\mathcal{F}_{\hat{\tau}}^{\max})_*^{-1}(\{\delta y\})$ is a singleton
- $\implies \deg(\pi\mathcal{F}_{\hat{\tau}}^{\max}, \mathcal{O}(\hat{\lambda}(0)), \hat{\xi}(\hat{\tau})) = 1$
- P-Spadini 2013 (based on Pang-Ralph 1996) $\implies \pi\mathcal{F}_{\hat{\tau}}^{\max}$ is a Lipschitz homeomorphism from $\mathcal{O}(\hat{\lambda}(0))$ on $\mathcal{O}(\hat{\xi}(\hat{\tau}))$.