



HAL
open science

Astrocytic theory of working memory

Maurizio de Pittà, Eshel Ben-Jacob, Hugues Berry

► **To cite this version:**

Maurizio de Pittà, Eshel Ben-Jacob, Hugues Berry. Astrocytic theory of working memory. BMC Neuroscience, 2014, 15 (Suppl 1), P206 (2 p.). hal-01026524

HAL Id: hal-01026524

<https://inria.hal.science/hal-01026524v1>

Submitted on 21 Jul 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

POSTER PRESENTATION

Open Access


Astrocytic theory of working memory

Maurizio De Pittà^{1,2,3*}, Eshel Ben-Jacob^{4,5}, Hugues Berry^{1,2}

From The Twenty Third Annual Computational Neuroscience Meeting: CNS*2014
Québec City, Canada. 26-31 July 2014

Working memory (WM) is the ability to transiently hold and manipulate goal-related information to forthcoming actions. In primates performing delayed-response tasks, the neural correlate of WM is the emergence of selective persistent activity in the prefrontal cortex (PFC),

during the delay period – i.e., the increase of neuronal firing rates compared to baseline in response of specific stimulatory cues. Neuronal firing during such persistent activity can be highly irregular, with a coefficient of variation (CV) of the interspike interval distribution larger



* Correspondence: maurizio.depitta@gmail.com

¹EPI Beagle, INRIA Rhône-Alpes, Villeurbanne, France

Full list of author information is available at the end of the article

than one. The origin of this increased irregularity is a matter of debate, as it is not known whether it stems from inherent properties of neurons or the plasticity of their synaptic connections or both. Astrocytes, the main type of glial cells in the cortex, have recently emerged as potential active players in synaptic plasticity due to their proposed ability to regulate synaptic neurotransmitter release in response to neuronal activity. Because available WM models do not take into account astrocytes, we investigated the possible role of astrocyte regulation of synaptic plasticity in the emergence of persistent neuronal firing in WM and its statistics. Our modeling study suggests that selective persistent activity could indeed emerge from astrocyte-mediated short-term synaptic facilitation triggered by the brief presentation of a stimulatory cue (Figures 1A,B). Bifurcation analysis revealed that astrocyte-mediated facilitation could outlast the cue and switch to a self-sustained mode where it is maintained by the ongoing synaptic activity, consistent with the existence of bistability of synaptic release (Figures 1C,D). In parallel, mean field analysis showed that such bistable synaptic release resulted in bistability of neural firing rates, with persistent “UP” states characterized by larger irregularity (Figures 1E,F). Altogether, these results suggest a novel astrocyte-based mechanism for WM providing experimentally testable hypotheses for the possible involvement of astrocytes in WM-related cognitive tasks.

Authors' details

¹EPI Beagle, INRIA Rhône-Alpes, Villeurbanne, France. ²LIRIS, Université de Lyon, UMR5205 CNRS-INSA, Villeurbanne, France. ³Department of Statistics, The University of Chicago, 5734 S. University Ave., Chicago, IL, USA. ⁴School of Physics and Astronomy, Tel Aviv University, Ramat Aviv, Israel. ⁵Center for Theoretical Biological Physics, Rice University, Houston, TX, USA.

Published: 21 July 2014

doi:10.1186/1471-2202-15-S1-P206

Cite this article as: De Pittà *et al.*: Astrocytic theory of working memory. *BMC Neuroscience* 2014 15(Suppl 1):P206.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

