



HAL
open science

Normality and Nondegeneracy for Optimal Control Problems with State Constraints

Fernando A.C.C. Fontes, H el ene Frankowska

► **To cite this version:**

Fernando A.C.C. Fontes, H el ene Frankowska. Normality and Nondegeneracy for Optimal Control Problems with State Constraints. 2014. hal-01088903v1

HAL Id: hal-01088903

<https://inria.hal.science/hal-01088903v1>

Preprint submitted on 29 Nov 2014 (v1), last revised 7 Jan 2015 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche franais ou  trangers, des laboratoires publics ou priv es.

Normality and Nondegeneracy for Optimal Control Problems with State Constraints*

Fernando A.C.C. Fontes[†] & H el ene Frankowska[‡]

October 27, 2014

Abstract

In this paper we investigate normal and nondegenerate forms of the maximum principle for optimal control problems with state constraints. We propose new constraint qualifications guaranteeing nondegeneracy and normality that have to be checked on smaller sets of points of an optimal trajectory, than those in known sufficient conditions. In fact, the constraint qualifications proposed impose the existence of an inward pointing velocity just on the instants of time for which the optimal trajectory has an outward pointing velocity.

Keywords. optimal control maximum principle state constraints constraint qualifications normality degeneracy nonsmooth analysis oriented distance.

1 Introduction

We consider optimal control problems with pathwise state constraints. For these problems, we study necessary conditions of optimality in the form of

*The partial supports of projects FP7-ITN-264735-SADCO "Sensitivity Analysis for Deterministic Controller Design" and FCT/FEDER Project PTDC/EEA-CRO/116014/2009 "Optimal Control in Constrained and Hybrid Nonlinear Systems", PTDC/EEI-AUT/1450/2012 "Optimal Control: Health, Energy and Robotics Applications" are gratefully acknowledged.

[†]ISR-Porto, Faculdade de Engenharia, Universidade do Porto, 4200-465 Porto, Portugal

[‡]CNRS, IMJ-PRG, UMR 7586, Sorbonne Universit e, UPMC Univ Paris 06, Univ Paris Diderot, Sorbonne Paris Cit e, Case 247, 4 Place Jussieu, 75252 Paris, France.

a maximum principle that, in some situations, are able to provide more information, by avoiding the abnormality or the degeneracy phenomena.

We say that the maximum principle is *abnormal* when the scalar multiplier associated with the objective function (ahead denoted by λ) is equal to zero. In this case the necessary conditions cannot use the information of the objective function to select minimizers.

The *degeneracy* phenomenon arises when the state constraint is active at the initial time. Then, a possible choice of multipliers are the so-called degenerate multipliers, for which the adjoint multiplier (denoted by q) and therefore the hamiltonian are equal to zero for almost every time. This implies that all conditions in the maximum principle are satisfied for any candidate to solution we might test. Thus, with the degenerate multipliers, the maximum principle is useless to select minimizers.

Nondegenerate and normal forms of the maximum principle are established when the problems satisfy a suitable constraint qualification. In addition to helping, in some situations, to find minimizers or to eliminate some candidates for optimality, the normal maximum principle is also useful to establish regularity properties of the minimizes [6, 12, 17] or to deduce second order optimality conditions [11, 18, 19].

The degeneracy of the maximum principle for state constrained problems has already been well identified and studied in the literature even for non-smooth data, see e.g. [1, 2, 8, 9, 24, 21, 22]. Previous investigations on nondegenerate and normal forms of the maximum principle involved various types of constraint qualifications. The relations between some of them is discussed for instance in [10, 14, 21]. In [22] an integral-type constraint qualification was introduced. There, it was shown that to avoid degeneracy, an inward pointing condition has to be satisfied for some, not all, instants of a neighborhood of the initial time.

In the present paper, we introduce new constraint qualifications that just have to be satisfied for times at which the optimal trajectory has an outward pointing velocity. Furthermore, we remove the convexity assumption on the problem data imposed in [22]. We also provide, in a corollary, additional sufficient conditions to avoid having the adjoint multiplier q equal to zero, or to avoid having q and λ both equal to zero. In this way, we show that a nondegeneracy condition derived previously in a different context in [4] holds true also in our case.

Normality for optimal control problems with state constraints has also been extensively studied, see for instance [10, 13, 16, 20, 24] and references

therein. In the literature it was shown that normality might be guaranteed by assuming constraint qualifications with the inward pointing inequalities imposed on a neighborhood of times for which the optimal trajectory belongs to the boundary of the state constraint. The new constraint qualification **CQn** proposed in Section 3 involves the last instant τ for which the multiplier associated with the state constraint – a measure denoted by μ – is active and, again, the inward pointing inequality has to be satisfied only for times $t < \tau$ from a neighborhood of τ at which the trajectory has an outward pointing velocity.

The constraints qualifications here proposed also help to understand the relations between previous constraints qualifications reported in the literature. Namely, we discuss conditions involving different generalized gradients and, in the smooth case, we compare different types of constraint qualifications.

The outline of the paper is as follows. In Section 2 we recall some notions of nonsmooth analysis, the nonsmooth maximum principle and compare various gradients of distance functions. In Section 3 we state our main results.

2 Preliminaries

2.1 Notations and Definitions

Throughout \mathbb{B} denotes the *closed unit ball* in \mathbb{R}^n , S^{n-1} the *unit sphere* in \mathbb{R}^n , $B(x, r)$ the *closed ball* in \mathbb{R}^n of center x and radius r , $p \cdot v$ the usual *scalar product* of $p, v \in \mathbb{R}^n$ and $|\cdot|$ the *Euclidean norm*. The sets $\text{co } K$, $\overline{\text{co}} K$, $\text{bdy } K$, K^c and $\text{int } K$ stand for the *convex hull*, *closed convex hull*, *boundary*, *complement* and *interior* of a set $K \subset \mathbb{R}^n$, respectively.

For a Borel measure μ on $[0, 1]$ we denote by $\text{supp}\{\mu\}$ its support and we use $\ell(I)$ for the Lebesgue measure of a Lebesgue measurable set $I \subset \mathbb{R}$. The product σ -algebra generated by the Lebesgue subsets \mathcal{L} of $[0, 1]$ and the Borel subsets of \mathbb{R}^m is denoted by $\mathcal{L} \times \mathcal{B}^m$ and the norm in the space of essentially bounded functions from $[0, 1]$ into \mathbb{R}^n by $\|\cdot\|_{L^\infty}$. Finally, $W^{1,1}([0, 1]; \mathbb{R}^n)$ denotes the space of absolutely continuous functions from $[0, 1]$ into \mathbb{R}^n .

Let Θ be a metric space. For a family of subsets $A_\tau \subset \mathbb{R}^n$ with $\tau \in \Theta$ and any $\bar{\tau} \in \Theta$, the upper set limit of A_τ at $\bar{\tau}$ is defined by

$$\text{Limsup}_{\tau \rightarrow \bar{\tau}} A_\tau := \{v \in \mathbb{R}^n : \exists \tau_i \rightarrow \bar{\tau}, v_i \in A_{\tau_i} \text{ such that } \lim_{i \rightarrow \infty} v_i = v\}.$$

We recall next the notion of *limiting normal cone* to a closed set $K \subset \mathbb{R}^n$ at $\bar{x} \in K$. Define first the contingent cone to K at x by

$$T_K(x) = \text{Limsup}_{r \rightarrow 0^+} \frac{1}{r}(K - x)$$

and consider its negative polar

$$\widehat{N}_K(x) = \{p \mid p \cdot v \leq 0, \forall v \in T_K(x)\}.$$

The limiting normal cone to K at x is defined by

$$N_K(\bar{x}) := \text{Limsup}_{x \rightarrow \bar{x}} \widehat{N}_K(x).$$

The negative polar of $N_K(\bar{x})$ is the Clarke tangent cone to K at \bar{x} :

$$\overline{T}_K(\bar{x}) := \{y \in \mathbb{R}^n : y \cdot p \leq 0 \text{ for all } p \in N_K(\bar{x})\}.$$

Given a lower semicontinuous function $f : \mathbb{R}^n \rightarrow \mathbb{R} \cup \{\infty\}$, the *limiting subdifferential* of f at a point $\bar{x} \in \mathbb{R}^n$ with $f(\bar{x}) < +\infty$ is the set

$$\partial f(\bar{x}) := \{\eta \in \mathbb{R}^n : (\eta, -1) \in N_{\text{epi } f}(\bar{x}, f(\bar{x}))\},$$

where $\text{epi } f := \{(x, \alpha) : \alpha \geq f(x)\}$. If f is differentiable at x , then we denote by $\nabla f(x)$ its gradient.

For a locally Lipschitz $h : \mathbb{R}^n \rightarrow \mathbb{R}$ the reachable gradient of h at x is defined by

$$\partial^* h(x) := \text{Limsup}_{y \rightarrow x} \{\nabla h(y)\}.$$

Recall that $\text{co } \partial^* h(x)$ is equal to the generalized gradient $\partial^C h(x)$ of h at x and that $\text{co } \partial h(x) = \partial^C h(x)$, see for instance [26, Proposition 4.7.6].

The *reachable hybrid subdifferential* of h at x is defined as

$$\partial^{*>} h(x) := \{\zeta \mid \exists x_i \rightarrow x \text{ such that } h(x_i) > 0 \forall i \text{ and } \lim_{i \rightarrow \infty} \nabla h(x_i) = \zeta\} \subset \partial h(x)$$

and the *hybrid subdifferential* $\partial^> h(x)$ is defined as

$$\partial^> h(x) := \text{co } \partial^{*>} h(x).$$

Observe that $\partial^{*>} h(x) \subset \partial^* h(x)$.

Similarly, for a locally Lipschitz $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ and $x \in \mathbb{R}^n$ the reachable Jacobian is defined by

$$\partial^* f(x) = \text{Limsup}_{y \rightarrow x} \{f'(y)\},$$

where $f'(y)$ denotes the Jacobian of f at y (which exists a.e. in \mathbb{R}^n).

We refer to [3, 26] for further concepts of nonsmooth and set-valued analysis, the last one using notations similar to those of the present paper.

2.2 Generalized Gradients of Distance Functions

For a closed nonempty set $K \subset \mathbb{R}^n$ and $x \in \mathbb{R}^n$, the distance $d_K(x)$ from x to K is given by $d_K(x) := \min_{y \in K} |x - y|$.

For K different from \mathbb{R}^n , the oriented distance function is defined by

$$d(x) := d_K(x) - d_{K^c}(x).$$

When $K = \mathbb{R}^n$ we set $d(x) = 0$ for every $x \in \mathbb{R}^n$.

These two functions are Lipschitz continuous with Lipschitz constant equal to one.

Therefore, we get $\partial d(x) \subset \mathbb{B}$. Also, $\partial^{*\rangle} d_K(\cdot) = \partial^{*\rangle} d(\cdot)$. By [25, Example 8.53] we know that

$$\partial d_K(x) = N_K(x) \cap \mathbb{B}, \quad \forall x \in K. \quad (1)$$

Proposition 2.1 *Let $x \in \text{bdy } K$. If $0 \notin \partial^{\rangle} d_K(x)$, then $\overline{T}_K(x)$ has a nonempty interior and is equal to the negative polar cone of $\partial^{*\rangle} d_K(x)$.*

Proof. If $0 \notin \partial^{\rangle} d_K(x)$, then the set $\mathbb{R}_+(\partial^{\rangle} d_K(x))$ is closed, see for instance [25, 3.48 (a)]. Consequently $\mathbb{R}_+(\partial^{\rangle} d_K(x))$ is a closed convex cone spanned by a convex compact set not containing zero. For this reason its negative polar cone has a nonempty interior.

By [5, Proposition 2.4 and Corollary 2.5], $\partial^C d_K(x) = \text{co}\{0, \partial^{*\rangle} d_K(x)\}$. Thus from (1) we deduce that

$$\partial^{\rangle} d_K(x) \subset \text{co } \partial d_K(x) = \text{co}(N_K(x) \cap \mathbb{B}) = \partial^C d_K(x) = \{\lambda \partial^{\rangle} d_K(x) \mid \lambda \in [0, 1]\}.$$

This implies that

$$\mathbb{R}_+(\partial^{\rangle} d_K(x)) = \text{co } N_K(x). \quad (2)$$

Since $\overline{T}_K(x)$ is equal to the negative polar of $\text{co } N_K(x)$ the proof follows.

Lemma 2.2 *Let $x \in \text{bdy } K$ be such that $\text{int } \overline{T}_K(x) \neq \emptyset$. Then*

$$\partial^* d_K(x) = \partial^{*\rangle} d_K(x) \cup \{0\}$$

and

$$\partial^{\rangle} d_K(x) \subset \partial^C d_K(x) = \{\lambda \partial^{\rangle} d_K(x) \mid \lambda \in [0, 1]\} \subset \mathbb{B}.$$

Furthermore,

$$N_K(x) \cap S^{n-1} = \partial^{*>} d_K(x) = \partial d_K(x) \cap S^{n-1}$$

and

$$\partial^* d(x) = \partial^{*>} d_K(x) = \partial^{*>} d(x).$$

In particular, $\partial^C d(x) = \partial^{>} d_K(x)$.

Proof. Fix any $v \in \text{int } \overline{T}_K(x)$. By [3], there exists $\varepsilon > 0$ such that

$$y + [0, \varepsilon](v + \varepsilon\mathbb{B}) \subset K, \quad \forall y \in K \cap B(x, \varepsilon).$$

Thus for all small $s > 0$ and any $b \in \mathbb{B}$, $d_K(y + s(v + \varepsilon b)) = 0$. This implies that $\langle \nabla d_K(y), v + \varepsilon b \rangle = 0$ for any $b \in \mathbb{B}$ whenever d_K is differentiable at $y \in K \cap B(x, \varepsilon)$. Hence for any such y we have $\nabla d_K(y) = 0$. Consequently, if $x_i \rightarrow x$ are such that $d_K(x_i) = 0$ and $\nabla d_K(x_i)$ exist and converge to some $\zeta \in \mathbb{R}^n$, then $\zeta = 0$.

Clearly, $\partial^{*>} d_K(x) \subset \partial^* d_K(x)$. Let $x_i \in \text{int } K$ converge to x . Then $\nabla d_K(x_i) = 0$ and therefore $0 \in \partial^* d_K(x)$.

Pick any $\zeta \in \partial^* d_K(x)$ and let $x_i \rightarrow x$ be such that $\nabla d_K(x_i)$ exist and converge to ζ when $i \rightarrow \infty$. If $d_K(x_{i_k}) = 0$ for a subsequence $\{x_{i_k}\}$, then $\zeta = 0$ by the first part of the proof. Otherwise $\zeta \in \partial^{*>} d_K(x)$. This implies the first equality of our Lemma.

The second relation can be easily deduced from the first one.

Let $\zeta \in \partial^{*>} d_K(x)$ and $x_i \rightarrow x$ be such that $x_i \notin K$ and $\nabla d_K(x_i) \rightarrow \zeta$. Then for some $y_i \in \text{bdy } K$, $\nabla d_K(x_i) \in \widehat{N}_K(y_i) \cap S^{n-1}$ and we deduce that $\zeta \in N_K(x) \cap S^{n-1}$. On the other hand, for any $\zeta \in \partial^{>} d_K(x) \setminus \partial^{*>} d_K(x)$ we have $|\zeta| < 1$, because $\partial^{*>} d_K(x) \subset S^{n-1}$ and \mathbb{B} is a strictly convex set. Hence $\partial^C d_K(x) \cap S^{n-1} = \partial^{*>} d_K(x)$. Consequently, by (1),

$$\partial^{*>} d_K(x) \subset N_K(x) \cap S^{n-1} \subset \partial d_K(x) \cap S^{n-1} \subset \partial^C d(x) \cap S^{n-1} = \partial^{*>} d_K(x).$$

To prove the last statement, observe that $\partial^{*>} d_K(x) = \partial^{*>} d(x) \subset \partial^* d(x)$. Let $x_i \in K$ be such that $d(\cdot)$ is differentiable at x_i and x_i converge to x while $\nabla d(x_i)$ converge to some ζ . Then there exists a unique projection y_i of x_i on $\text{bdy } K$. Using that for all large i , $\text{int } \overline{T}_K(y_i) \neq \emptyset$, applying the same proof as the one of Proposition 5 in [15] on a neighborhood of x , we deduce that $\nabla d(y_i) \in \widehat{N}_K(y_i) \cap S^{n-1}$ for all large i . Hence $\zeta \in N_K(x) \cap S^{n-1} = \partial^{*>} d_K(x)$.

2.3 Maximum Principle

Consider the following optimal control problem with state constraints:

$$(P) \quad \begin{cases} \text{Minimize} & g(x(1)) \\ \text{subject to} & \\ & \dot{x}(t) = f(t, x(t), u(t)), \quad u(t) \in U(t) \quad t \in [0, 1] \\ & x(0) = x_0, \quad x(1) \in K_1 \\ & x(t) \in K \quad \text{for all } t \in [0, 1]. \end{cases}$$

The data for this problem comprise functions $g : \mathbb{R}^n \rightarrow \mathbb{R}$, $f : [0, 1] \times \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^n$, an initial state $x_0 \in \mathbb{R}^n$, sets $K, K_1 \subset \mathbb{R}^n$ of the state and the end-point constraints, and a set-valued map $U : [0, 1] \rightrightarrows \mathbb{R}^m$. The set of *control functions* for (P) is

$$\mathcal{U} := \{u : [0, 1] \rightarrow \mathbb{R}^m : u \text{ is Lebesgue measurable, } u(t) \in U(t) \text{ a.e. } t \in [0, 1]\}.$$

We say that a trajectory-control pair (\bar{x}, \bar{u}) of the above system is a local minimizer of (P) if for some $\delta > 0$ and every trajectory-control pair (x, u) satisfying $\|x - \bar{x}\|_{L^\infty} < \delta$ we have $g(\bar{x}(1)) \leq g(x(1))$.

Using the distance or the oriented distance functions, the state constraint $x(t) \in K$ can be written equivalently as the inequality constraint

$$d(x(t)) \leq 0 \quad \text{for all } t \in [0, 1]$$

or as

$$d_K(x(t)) \leq 0 \quad \text{for all } t \in [0, 1].$$

We impose the following hypotheses which refer to a fixed local minimizer (\bar{x}, \bar{u}) and a δ' neighborhood of \bar{x} .

H1 The function $(t, u) \rightarrow f(t, x, u)$ is $\mathcal{L} \times \mathcal{B}^m$ measurable for each x and $U(\cdot)$ is measurable with closed nonempty images.

H2 There exists $C_f \geq 0$ such that

$$|f(t, x, u) - f(t, x', u)| \leq C_f |x - x'|$$

for $x, x' \in \bar{x}(t) + \delta'\mathbb{B}$, $u \in U(t)$ and a.e. $t \in [0, 1]$.

H3 There exists $C_u \geq 0$ such that $|f(t, x, u)| \leq C_u$ for $x \in \bar{x}(t) + \delta'\mathbb{B}$, $u \in U(t)$, and $t \in [0, 1]$.

H4 The function g is Lipschitz on $\bar{x}(1) + \delta'\mathbb{B}$.

H5 The sets K and K_1 are closed.

The maximum principle for problems with state constraints has been reported in different versions with different degrees of generality (see e.g. [23, 7, 27]). The version in the next theorem can be deduced from [26, pp. 329 and 204] applied to the state constraints described by an inequality involving the oriented distance function $d(\cdot)$.

Theorem 2.3 *Let (\bar{x}, \bar{u}) be a local minimizer and assume **H1-H5**. Then, there exist $p \in W^{1,1}([0, 1]; \mathbb{R}^n)$, a non-negative (finite) Borel measure μ on $[0, 1]$ and a scalar $\lambda \in \{0, 1\}$ such that*

$$\mu([0, 1]) + \|p\|_{L^\infty} + \lambda \neq 0, \quad (3)$$

$$-\dot{p}(t) \in q(t) \operatorname{co} \partial_x^* f(t, \bar{x}(t), \bar{u}(t)) \quad a.e. \ t \in [0, 1], \quad (4)$$

$$-q(1) \in \lambda \partial g(\bar{x}(1)) + N_{K_1}(\bar{x}(1)),$$

$$\gamma(t) \in \partial^> d(\bar{x}(t)) \quad \mu - a.e.,$$

$$\operatorname{supp}\{\mu\} \subset \{t \in [0, 1] \mid \bar{x}(t) \in \operatorname{bdy} K\},$$

and, for almost every $t \in [0, 1]$, and all $u \in U(t)$

$$q(t) \cdot (f(t, \bar{x}(t), \bar{u}(t)) - f(t, \bar{x}(t), u)) \geq 0, \quad (5)$$

where

$$q(t) := \begin{cases} p(t) + \int_{[0,t]} \gamma(s) \mu(ds), & t \in [0, 1) \\ p(1) + \int_{[0,1]} \gamma(s) \mu(ds), & t = 1. \end{cases}$$

The relation $\gamma(t) \in \partial^> d(\bar{x}(t))$ is equivalent to $\gamma(t) \in \partial^> d_K(\bar{x}(t))$. It follows from Lemma 2.2 that at every $t \in [0, 1]$ such that $\operatorname{int} \bar{T}_K(\bar{x}(t))$ is nonempty, it can be equivalently written as $\gamma(t) \in \partial^C d(\bar{x}(t))$.

Since μ is a finite Borel measure on $[0, 1]$, it is regular and therefore $q(\cdot)$ is left continuous on $(0, 1)$.

2.4 Abnormality and Degeneracy Phenomena

The above maximum principle is called normal if $\lambda = 1$.

In this paper we say that a maximum principle is nondegenerate if

$$\mu((0, 1]) + \|q\|_{L^\infty} + \lambda \neq 0.$$

The degeneracy phenomenon arises when the state constraint is active at the initial time, i.e. when $x_0 \in \text{bdy } K$. Then a possible choice of multipliers, here called degenerate multipliers, is

$$\lambda = 0, \quad \mu = \delta_{\{0\}}, \quad p \equiv -\xi, \quad \text{with } \xi \in \partial^> d(\bar{x}(0)),$$

where $\delta_{\{0\}}$ is the Dirac unit measure concentrated at 0. We note that with such multipliers, the expression $p(t) + \int_{[0,t)} \xi \mu(ds)$, which features in the conditions of the maximum principle, vanishes for almost all times. So, the maximum principle holds true for any pair (\bar{x}, \bar{u}) we might test. Such multipliers are useless to identify minimizers.

Previous works on the nondegeneracy of the maximum principle mainly differ by the constraint qualification used, in addition to assumptions on the data of the problem. We can identify, in the literature, four types of constraint qualifications (CQ), which we adapt here to the context of state constraints $x(t) \in K$ for all $t \in [0, 1]$:

1. Inward pointing velocity CQ.

If $x_0 \in \text{bdy } K$, then there exist $\delta > 0, \epsilon > 0$ and a control function $\hat{u} \in \mathcal{U}$ such that

$$\max_{\gamma \in \partial^> d_K(x_0)} \gamma \cdot f(t, x_0, \hat{u}(t)) < -\delta \quad \text{a.e. } t \in [0, \epsilon].$$

2. CQ involving the optimal control.

If $x_0 \in \text{bdy } K$, then there exist $\delta > 0, \epsilon > 0$ and a control function $\hat{u} \in \mathcal{U}$ such that

$$\max_{\gamma \in \partial^> d_K(x_0)} \gamma \cdot (f(t, x_0, \hat{u}(t)) - f(t, x_0, \bar{u}(t))) < -\delta \quad \text{a.e. } t \in [0, \epsilon].$$

3. Integral-type CQ.

If $x_0 \in \text{bdy } K$, then there exist $\delta > 0, \epsilon > 0$ and a control function $\hat{u} \in \mathcal{U}$ such that

$$\int_0^t \max_{\gamma \in \partial^> d_K(x_0)} \gamma \cdot (f(s, x_0, \hat{u}(s)) - f(s, x_0, \bar{u}(s))) ds < -\delta t \quad \forall t \in [0, \epsilon].$$

4. CQ involving a strictly feasible initial trajectory.

If $x_0 \in \text{bdy}K$, then there exists $\epsilon > 0$ such that

$$\bar{x}(t) \in \text{int} K \quad \forall t \in (0, \epsilon).$$

The relation between the first two types of constraint qualification is discussed for instance in [10, 13, 21], where some bibliographical references for each type of constraint qualifications are provided. The third type is described in [22]. The fourth type appears, for instance, in [8].

Let us underline that, by Proposition 2.1, the constraint qualifications 1, 2, 3 imply that $\text{int} \bar{T}_K(x_0)$ is nonempty. Thus, by Lemma 2.2, the first three constraints qualifications can be written in an equivalent form by replacing $\partial^>d_K(x_0)$ by $\partial^*d(x_0)$ or by $\partial^C d(x_0)$.

3 Nondegeneracy and Normality

At the end of this section, we provide sufficient conditions for normality of the maximum principle. But first we show how the previous maximum principle can be strengthened to avoid the degeneracy by introducing a new type of constraint qualification.

CQd (*Constraint Qualification to avoid degeneracy*) If $x_0 \in \text{bdy}K$, then there exist $\epsilon > 0$, $\delta > 0$ satisfying

$$\inf_{u \in U(t)} \max_{\gamma \in \partial^*d(x_0)} \gamma \cdot (f(t, x_0, u) - f(t, x_0, \bar{u}(t))) < -\delta, \quad (6)$$

for a.e. $t \in \{r \in [0, \epsilon] : \max_{\xi \in \partial^*d(\bar{x}(r))} \xi \cdot f(r, \bar{x}(r), \bar{u}(r)) \geq 0\}$.

Remark 3.1 *If the set $\{r \in [0, \epsilon] : \max_{\xi \in \partial^*d(\bar{x}(r))} \xi \cdot f(r, \bar{x}(r), \bar{u}(r)) \geq 0\}$ has a positive Lebesgue measure, then **CQd** implies that the interior of $\bar{T}_K(x_0)$ is nonempty. Indeed, by (6), there exists $t \in [0, 1]$ such that*

$$\inf_{u \in U(t)} \max_{\gamma \in \partial^*d(x_0)} \gamma \cdot (f(t, x_0, u) - f(t, x_0, \bar{u}(t))) < -\delta.$$

Hence, by Proposition 2.1, $\text{int} \bar{T}_K(x_0) \neq \emptyset$.

Consequently, by Lemma 2.2, we can replace $\partial^*d(x_0)$ in (6) by $\partial^{*>}d_K(x_0)$. Since the scalar product is bilinear, **CQd** takes then the form of a more classical constraint qualification :

CQd' Suppose that the set $\{r \in [0, \alpha] : \max_{\xi \in \partial^* d(\bar{x}(r))} \xi \cdot f(r, \bar{x}(r), \bar{u}(r)) \geq 0\}$ has a positive Lebesgue measure for any $\alpha > 0$.

If $x_0 \in \text{bdy}K$, then there exist $\epsilon > 0$, $\delta > 0$ such that

$$\inf_{u \in U(t)} \max_{\gamma \in \partial^* d_K(x_0)} \gamma \cdot (f(t, x_0, u) - f(t, x_0, \bar{u}(t))) < -\delta$$

for a.e. $t \in \{r \in [0, \epsilon] : \max_{\xi \in \partial^* d_K(\bar{x}(r))} \xi \cdot f(r, \bar{x}(r), \bar{u}(r)) \geq 0\}$.

We would like to underline that if for some $\alpha > 0$ the Lebesgue measure of the set $\{r \in [0, \alpha] : \max_{\xi \in \partial^* d(\bar{x}(r))} \xi \cdot f(r, \bar{x}(r), \bar{u}(r)) \geq 0\}$ is equal to zero, then the proof of Theorem 3.2 below concerning the nondegenerate maximum principle does not use any constraint qualification at all, beside the assumption $\text{int } \bar{T}_K(x_0) \neq \emptyset$. For this reason the additional requirement imposed in **CQd'** is not restrictive to derive a nondegenerate maximum principle.

We note that **CQd'** requires less than the constraint qualifications of type **2** from Section 2.3 because the relevant inequality does not have to be satisfied *a.e.* in the whole time interval $[0, \epsilon]$, but just for *a.e.* instant for which the optimal trajectory has an outward pointing velocity, i.e. when $\max_{\xi \in \partial^* d_K(\bar{x}(r))} \xi \cdot f(r, \bar{x}(r), \bar{u}(r)) \geq 0$.

We state our first main result.

Theorem 3.2 Let (\bar{x}, \bar{u}) be a local minimizer, $\text{int } \bar{T}_K(x_0) \neq \emptyset$ and assume that **H1-H5**, **CQd** are satisfied. Then, the maximum principle of Theorem 2.3 holds true with the nontriviality condition (3) strengthened to

$$\mu((0, 1]) + \|q\|_{L^\infty} + \lambda > 0. \quad (7)$$

Remark 3.3 Since $q(\cdot)$ is left continuous on $(0, 1)$, if $q(\cdot) = 0$ a.e., then $q(t) = 0$ for all $t \in (0, 1)$. In this case, the maximum principle (5) holds true for all $t \in (0, 1)$ and $u \in U(t)$. Furthermore, from the adjoint equation it follows that $p(\cdot) \equiv p(0)$. Since $p(0) + \int_{[0, t]} \gamma(s) \mu(ds) = 0$ for all $t \in (0, 1)$, we deduce that for all $0 < t_1 < t_2 < 1$, $\int_{[t_1, t_2]} \gamma(s) \mu(ds) = 0$ and, because μ is regular, that for all $0 < t_1 < t_2 < 1$

$$\int_{[t_1, t_2]} \gamma(s) \mu(ds) = 0. \quad (8)$$

Assume that $\text{int } \bar{T}_K(\bar{x}(t)) \neq \emptyset$ for every $t \in [0, 1]$ satisfying $\bar{x}(t) \in \text{bdy } K$. If $\mu((0, 1)) > 0$, then the above implies that $\|q\|_{L^\infty} \neq 0$. Indeed, if $\|q\|_{L^\infty} = 0$, then, using Lemma 2.2, that μ is nonnegative and that $\partial^> d(\bar{x}(\cdot))$ is upper semicontinuous on the compact interval $[0, 1]$, we derive a contradiction with (8) for a choice of $0 < t_1 < t_2 < 1$.

We next observe that if moreover $\mu((0, 1]) > 0$ and

$$\text{co}N_K(\bar{x}(1)) \cap (-N_{K_1}(\bar{x}(1))) = \{0\}, \quad (9)$$

then $\|q\|_{L^\infty} + \lambda \neq 0$. Indeed, assume for a moment that $\|q\|_{L^\infty} + \lambda = 0$. Then $\mu((0, 1)) = 0$ and, therefore, $\mu(\{1\}) > 0$. We deduce that $p(\cdot) \equiv -\gamma(0)\mu(\{0\})$ and therefore

$$q(1) = p(1) + \gamma(0)\mu(\{0\}) + \gamma(1)\mu(\{1\}) = \gamma(1)\mu(\{1\}).$$

Since $\gamma(1) \in \partial^> d(\bar{x}(1))$, Lemma 2.2 and (2) yield $q(1) \in \text{co}N_K(\bar{x}(1))$. On the other hand, $-q(1) \in N_{K_1}(\bar{x}(1))$ in contradiction with (9). This proves our claim.

Furthermore, $\|q\|_{L^\infty} \neq 0$ if in addition

$$\text{co}N_K(\bar{x}(1)) \cap (-\partial g(\bar{x}(1)) - N_{K_1}(\bar{x}(1))) = \emptyset. \quad (10)$$

Indeed, otherwise $\lambda = 1$, $\mu((0, 1)) = 0$, $-q(1) \in \partial g(\bar{x}(1)) + N_{K_1}(\bar{x}(1))$. In the same way as before we show that $q(1) = \gamma(1)\mu(\{1\}) \in \text{co}N_K(\bar{x}(1))$, which contradicts (10).

Hence we have proved the following corollary.

Corollary 3.4 *Let (\bar{x}, \bar{u}) be a local minimizer. Assume **H1-H5**, **CQd**, (9) and that $\text{int } \bar{T}_K(\bar{x}(t)) \neq \emptyset$ for every $t \in [0, 1]$ satisfying $\bar{x}(t) \in \text{bdy } K$. Then, the maximum principle of Theorem 2.3 holds true with the nontriviality condition (3) strengthened to*

$$\|q\|_{L^\infty} + \lambda \neq 0. \quad (11)$$

Moreover, if also (10) is satisfied, then $\|q\|_{L^\infty} \neq 0$.

Another main result in this section provides sufficient conditions for the maximum principle to be normal. Namely, under the constraint qualification **CQn** proposed below, the nondegenerate maximum principle is in fact normal whenever $\bar{x}(1) \in \text{int } K_1$.

CQn (*Constraint Qualification to guarantee normality*) Consider μ as in the maximum principle and define

$$\tau := \inf \{t \in [0, 1] : \mu\{[t, 1]\} = 0\}.$$

Assume that there exist $\epsilon > 0$, $\delta > 0$ such that

$$\inf_{u \in U(t)} \max_{\gamma \in \partial^* d(\bar{x}(\tau))} \gamma \cdot (f(t, \bar{x}(\tau), u) - f(t, \bar{x}(\tau), \bar{u}(t))) < -\delta, \quad (12)$$

for a.e. $t \in \{r \in (\tau - \epsilon, \tau] \cap [0, 1] : \max_{\xi \in \partial^* d(\bar{x}(r))} \xi \cdot f(r, \bar{x}(r), \bar{u}(r)) \geq 0\}$.

Remark 3.5 *In the proof of Theorem 3.6 below we show that for a nondegenerate multiplier we always have $\tau > 0$ and that the set $\{r \in [\tau - \epsilon, \tau] : \max_{\xi \in \partial^* d(\bar{x}(r))} \xi \cdot f(r, \bar{x}(r), \bar{u}(r)) \geq 0\}$ has a positive Lebesgue measure. Thus, as in Remark 3.1, **CQn** implies that $\text{int } \bar{T}_K(\bar{x}(\tau))$ is nonempty. Then, by Lemma 2.2, we can replace $\partial^* d(\bar{x}(\tau))$ in (12) by $\partial^> d_K(\bar{x}(\tau))$ and to obtain a more classical constraint qualification :*

CQn' Consider μ as in the nondegenerate maximum principle and define

$$\tau := \inf \{t \in [0, 1] : \mu\{[t, 1]\} = 0\}.$$

Assume that there exist $\epsilon > 0$, $\delta > 0$ such that

$$\inf_{u \in U(t)} \max_{\gamma \in \partial^> d_K(\bar{x}(\tau))} \gamma \cdot (f(t, \bar{x}(\tau), u) - f(t, \bar{x}(\tau), \bar{u}(t))) < -\delta,$$

for a.e. $t \in \{r \in [\tau - \epsilon, \tau] \cap [0, 1] : \max_{\xi \in \partial^> d_K(\bar{x}(r))} \xi \cdot f(r, \bar{x}(r), \bar{u}(r)) \geq 0\}$.

We note that **CQn'** is less restrictive than the constraint qualification proposed in [20, (8)] because the relevant inequality does not have to be satisfied *a.e.* in the whole time interval $[\tau - \epsilon, \tau]$, but just for *a.e.* instant for which the optimal trajectory has an outward pointing velocity, i.e. when $\max_{\xi \in \partial^> d(\bar{x}(r))} \xi \cdot f(r, \bar{x}(r), \bar{u}(r)) \geq 0$.

Theorem 3.6 *Let (\bar{x}, \bar{u}) be a local minimizer. Assume **H1** - **H5** and that the maximum principle of Theorem 2.3 holds true with the strengthened non-triviality condition (7). If **CQn** is satisfied, then $\lambda + |q(1)| \neq 0$. In particular, if $\bar{x}(1) \in \text{int } K_1$, then $\lambda = 1$.*

Remark 3.7 When the oriented distance $d(\cdot)$ is continuously differentiable on a neighborhood of x_0 , then $\partial^>d(x_0) = \{\nabla d(x_0)\}$ is the outer unit normal to K at x_0 and we can establish interesting connections between the various types of constraint qualifications discussed previously.

Namely, consider the following constraint qualification:

CQd1 If $x_0 \in \text{bdy}K$, then there exist $\epsilon > 0$, $\delta > 0$ and a control $\hat{u} \in \mathcal{U}$ satisfying

$$\nabla d(x_0) \cdot f(t, x_0, \hat{u}(t)) < -\delta, \quad (13)$$

for a.e. $t \in \{r \in [0, \epsilon] : \nabla d(\bar{x}(r)) \cdot f(r, \bar{x}(r), \bar{u}(r)) \geq 0\}$.

This constraint qualification involving an inward pointing inequality (13) and not depending on the optimal control is akin to the constraint qualifications of type 1 mentioned above.

Using Lipschitz continuity of $d(\cdot)$ and **H2**, **H3**, we deduce that for a sufficiently small $\epsilon > 0$ and for a.e. $t \in \{r \in [0, \epsilon] : \nabla d(\bar{x}(r)) \cdot f(r, \bar{x}(r), \bar{u}(r)) \geq 0\}$ we have

$$\nabla d(\bar{x}(t)) \cdot f(t, \bar{x}(t), \hat{u}(t)) < -\delta/2.$$

Thus for a.e. $t \in \{r \in [0, \epsilon] : \nabla d(\bar{x}(r)) \cdot f(r, \bar{x}(r), \bar{u}(r)) \geq 0\}$,

$$\nabla d(\bar{x}(t)) \cdot (f(t, \bar{x}(t), \hat{u}(t)) - f(t, \bar{x}(t), \bar{u}(t))) < -\delta/2.$$

By the measurable selection theorem, **CQd** is also equivalent to the following constraint qualification

CQd2 If $x_0 \in \text{bdy}K$, then there exist $\epsilon > 0$, $\delta > 0$, and a control $\hat{u} \in \mathcal{U}$ satisfying

$$\nabla d(x_0) \cdot (f(t, x_0, \hat{u}(t)) - f(t, x_0, \bar{u}(t))) < -\delta, \quad (14)$$

for a.e. $t \in \{r \in [0, \epsilon] : \nabla d(\bar{x}(r)) \cdot f(r, \bar{x}(r), \bar{u}(r)) \geq 0\}$.

We deduce that in the case when the oriented distance function $d(\cdot)$ is continuously differentiable on a neighborhood of x_0 and either **CQd1** or **CQd2** holds true, then we can write the maximum principle with the stronger non-triviality condition (7).

Recalling that the support of μ is contained in the set of times for which the optimal trajectory is on the boundary of K , for the normality results, we may consider the following two constraint qualifications

CQn1 Let μ be as in the nondegenerate maximum principle, $\mu((0, 1]) > 0$ and define

$$\tau := \inf \{t \in [0, 1] : \mu\{[t, 1]\} = 0\} > 0.$$

Assume that $d(\cdot)$ is continuously differentiable on a neighborhood of $\bar{x}(\tau)$ and that there exist $\epsilon > 0$, $\delta > 0$ such that

$$\inf_{u \in U(t)} \nabla d(\bar{x}(\tau)) \cdot f(t, \bar{x}(\tau), u) < -\delta,$$

for a.e. $t \in \{r \in [\tau - \epsilon, \tau] \cap [0, 1] : \nabla d(\bar{x}(r)) \cdot f(r, \bar{x}(r), \bar{u}(r)) \geq 0\}$

and

CQn2 Let μ , τ be as in **CQn1**. Assume that $d(\cdot)$ is continuously differentiable on a neighborhood of $\bar{x}(\tau)$ and that there exist $\epsilon > 0$, $\delta > 0$ such that

$$\inf_{u \in U(t)} \nabla d(\bar{x}(\tau)) \cdot (f(t, \bar{x}(\tau), u) - f(t, \bar{x}(\tau), \bar{u}(t))) < -\delta,$$

for a.e. $t \in \{r \in [\tau - \epsilon, \tau] \cap [0, 1] : \nabla d(\bar{x}(r)) \cdot f(r, \bar{x}(r), \bar{u}(r)) \geq 0\}$.

Using Lipschitz continuity of the functions involved with arguments as above, we can show that the constraint qualification **CQn1** implies **CQn2**.

Also, since the support of μ is contained in the set of times for which the optimal trajectory is on the boundary of K , we have $\bar{x}(\tau) \in \partial K$.

We can, therefore, write a constraint qualification that is less explicitly dependent on the optimal control and the corresponding multipliers (and might be simpler to verify whether it is satisfied).

CQn1 For every $\tau \in (0, 1]$ such that $\bar{x}(\tau) \in \partial K$ there exist $\epsilon > 0$, $\delta > 0$, and a control $\hat{u} \in \mathcal{U}$ such that

$$\nabla d(\bar{x}(\tau)) \cdot f(t, \bar{x}(\tau), \hat{u}(t)) < -\delta,$$

for a.e. $t \in \{r \in [\tau - \epsilon, \tau] \cap [0, 1] : \nabla d(\bar{x}(r)) \cdot f(r, \bar{x}(r), \bar{u}(r)) \geq 0\}$.

We have established as a corollary of Thm. 3.6 that, under the hypotheses of Thm. 3.2 and $\bar{x}(1) \in \text{int } K_1$, if we assume that either **CQn1**, **CQn2**, or **CQn1** are satisfied, then the maximum principle is normal.

References

- [1] A. V. Arutyunov, *Optimality Conditions: Abnormal and Degenerate Problems*, Kluwer, Boston, 2000.
- [2] A. V. Arutyunov, S. M. Aseev, Investigation of the degeneracy phenomenon of the maximum principle for optimal control problems with state constraints, *SIAM Journal on Control and Optimization* 35 (1997) 930–952.
- [3] J. P. Aubin, H. Frankowska, *Set-Valued Analysis*, Birkhäuser, Boston, 1990.
- [4] A. Cernea, H. Frankowska (2006) *A connection between the maximum principle and dynamic programming for constrained control problems*, *SIAM J. on Control*, 44, 673-703
- [5] F. Clarke, Generalized gradients and applications, *Trans. Amer. Math. Soc.* 205 (1975) 247-262.
- [6] M. M. A. Ferreira, On the regularity of optimal controls for a class of problems with state constraints, *International Journal of Systems Science* 37 (8) (2006) 495–502.
- [7] A. Ya. Dubovitskii and A.A. Milyutin, Extremum problems under constraints, *Dokl. Akad. Nauk SSSR*, 149 (1963), 759762.
- [8] M. M. A. Ferreira, R. B. Vinter, When is the maximum principle for state constrained problems nondegenerate?, *Journal of Mathematical Analysis and Applications* 187 (2) (1994) 438–467.
- [9] M. M. A. Ferreira, F. A. C. C. Fontes, R. B. Vinter, Nondegenerate necessary conditions for nonconvex optimal control problems with state constraints, *Journal of Mathematical Analysis and Applications* 233 (1999) 116–129.
- [10] H. Frankowska, Normality of the maximum principle for absolutely continuous solutions to Bolza problems under state constraints, *Control and Cybernetics* 38 (4B) (2009) 1327–1340.

- [11] H. Frankowska, D. Hoehener, D. Tonon, A second-order maximum principle in optimal control under state constraints, *Serdica Mathematical Journal*, 39 (3-4) (2013) 233-270.
- [12] H. Frankowska, E. M. Marchini, Lipschitzianity of optimal trajectories for the Bolza optimal control problem, *Calculus of Variations and Partial Differential Equations* 27 (4) (2006) 467–492.
- [13] H. Frankowska, M. Mazzola, Optimal synthesis and normality of the maximum principle for optimal control problems with pure state constraints, in: *Proceedings of 9th IEEE International Conference on Control and Automation (ICCA)*, Santiago, Chile, (2011) 945–950.
- [14] H. Frankowska, M. Mazzola, On relations of the adjoint state to the value function for optimal control problems with state constraints, *Nonlinear Differential Equations and Applications* 20 (2013) 361–383.
- [15] H. Frankowska, M. Mazzola Discontinuous solutions of Hamilton-Jacobi-Bellman equation under state constraints, *Calculus of Variations and Partial Differential Equations* 46 (2013) 725-747.
- [16] H. Frankowska, D. Tonon, Inward pointing trajectories, normality of the maximum principle and the non occurrence of the Lavrentieff phenomenon in optimal control under state constraints, *Journal of Convex Analysis* 20 (4) (2013) 1147–1180.
- [17] G. N. Galbraith, R. B. Vinter, Lipschitz continuity of optimal controls for state constrained problems, *SIAM J. Control Optim.* 42 (5) (2003) 1727–1744.
- [18] D. Hoehener, Variational approach to second-order optimality conditions for control problems with pure state constraints. *SIAM J. Control Optim.* 50 (2012), 1139–1173.
- [19] D. Hoehener, Feasible perturbations of control systems with pure state constraints and applications to second-order optimality conditions. *Appl. Math. Optim.* 68 (2013), 219–253.
- [20] S. O. Lopes, F. A. C. C. Fontes, Normal forms of necessary conditions for dynamic optimization problems with pathwise inequality constraints, *Journal of Mathematical Analysis and Applications* 399 (2013) 27–37.

- [21] S. O. Lopes, F. A. C. C. Fontes, M. R. de Pinho, On constraint qualifications for nondegenerate necessary conditions of optimality applied to optimal control problems, *Discrete and Continuous Dynamical System-A* 29 (2) (2011) 559–576.
- [22] S. O. Lopes, F. A. C. C. Fontes, M. R. de Pinho, An integral-type constraint qualification to guarantee nondegeneracy of the maximum principle for optimal control problems with state constraints, *Systems and Control Letters* 62 (2013) 686–692.
- [23] Pontryagin, L. S., V. G. Boltyanskii, and R. V. Gamkrelidze, *The mathematical theory of optimal processes*, Wiley, New York, 1962.
- [24] F. Rampazzo, R. B. Vinter, Degenerate optimal control problems with state constraints, *SIAM Journal of Control and Optimization* 39 (2000) 989–1007.
- [25] R. T. Rockafellar, R. B. Wets, *Variational Analysis*, Berlin: Grundlehren Math. Wiss. 317, Springer-Verlag, 1998.
- [26] R. B. Vinter, *Optimal control*, Birkhäuser, Boston, 2000.
- [27] R. B. Vinter, G. Pappas, A maximum principle for nonsmooth optimal-control problems with state constraints, *Journal of Mathematical Analysis and Applications* 89.1 (1982): 212-232.