



HAL
open science

Genericity of weakly computable objects

Mathieu Hoyrup

► **To cite this version:**

| Mathieu Hoyrup. Genericity of weakly computable objects. 2014. hal-01095864v1

HAL Id: hal-01095864

<https://inria.hal.science/hal-01095864v1>

Preprint submitted on 16 Dec 2014 (v1), last revised 9 Dec 2016 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Genericity of weakly computable objects

Mathieu Hoyrup

December 16, 2014

Abstract

In computability theory many results state the existence of objects that in many respects lack algorithmic structure but at the same time are effective in some sense. Friedberg and Muchnik’s answer to Post problem is one of the most celebrated results in this form. The main goal of the paper is to develop a general result that embodies a large number of these particular constructions, capturing the essential idea that is common to all of them, and expressing it in topological terms.

To do so, we introduce the effective topological notions of *irreversible function* and *directional genericity* and provide two main results that identify situations when such constructions are possible, clarifying the role of topology in many arguments from computability theory. We apply these abstract results to particular situations, illustrating their strength and deriving new results.

This paper is an extended version of the conference paper [Hoy14] with detailed proofs and new results.

Contents

1	Introduction	2
2	Background and notations	3
2.1	Notations	4
2.2	Effective topology	4
2.3	Effective Polish spaces	4
3	Irreversible functions	6
3.1	A non-uniform result	6
3.2	Irreversible functions	8
3.3	The constructive result	10
3.4	Application to the ergodic decomposition	13

4 Directional genericity	14
4.1 Being generic from above	15
4.2 Genericity for c.e. sets	19
4.3 Genericity for left-c.e. reals	20
4.4 Genericity for Π_1^0 -classes	23
4.5 Genericity for regular Π_1^0 -classes	24

1 Introduction

One of the main goals of computability theory is to understand and classify the algorithmic content of infinite objects, which can be expressed as the difficulty of computing them or as their ability to help solving problems. In establishing this classification one is often led to separate classes of algorithmic complexity and the construction of counter-examples is usually a hard task that requires the use of advanced techniques, one of the simplest being the so-called priority method with finite injury. The difficulty in carrying out such constructions is that the built object should meet two types of requirements going in opposite directions: (i) it should lack algorithmic content but at the same time (ii) it should be constructible in some way. In other words, these objects live somewhere between *generic* objects (objects with no structure) and *computable* objects (the most constructible objects). While computability theory provides formal notions of genericity, these ones are always incompatible with computability, so the two prescriptions are conflicting.

In this paper we propose an abstract view on this topic by providing general results that, when applied properly implement the sought constructions, and provide a shortcut for producing objects with prescribed properties. We expect that checking the conditions of these theorems is much easier than writing down the complete construction which is often merely an adaptation of known techniques to new situations.

The first main result enables one to construct objects that (i) are not computable, but (ii) have a computable image under some given function. As an illustration, we prove a negative result about the computability of the ergodic decomposition. This part is developed in Section 3 and is the version with detailed proofs of the results already presented in [Hoy14].

The second main result enables one to construct objects that (i) are generic in some sense, but (ii) are computable in a weak sense. It is done by introducing a new notion of genericity, called *directional genericity*, that has two advantages: it is formally close to plain genericity, being defined

as a slight variation of 1-genericity, and we prove that it is compatible with various weak forms of computability. More precisely each weak notion of computability, such as being a c.e. set, a left-c.e. real, or a Π_1^0 -class, implicitly comes with an ordering which we exploit to define a notion of directional genericity. In addition to introducing these new notions, our main contribution is the result that they are compatible with weak computability. This result has important consequences: many ad hoc existing constructions are subsumed and unified by this result, new results can be obtained whenever the new notion of genericity captures the sought properties, and the result clarifies the role of topology in computability theory. We then illustrate these notions of genericity in several contexts, showing evidence that they clarify existing results, and we apply our main theorem to derive a separation result. This part is developed in Section 4 and is a significant extension of what appears in [Hoy14].

The proofs of our main results use the priority method with finite injury. We observe that while this technique has been massively used in many situations, no result has convincingly clarified its scope and the contexts in which it can be applied, and each particular construction is yet another adaptation of the same argument, based on the same idea. We hope that the results presented here shed light on this subject.

The paper is organized as follows. In Section 2 we present the needed background on computable analysis, providing along the way a few results of independent interest on effective Polish spaces. In Section 3 we introduce the notion of irreversible function and prove the first main result: if a computable function is effectively irreversible then it maps a non-computable point to a computable image. The results of this section come from [Hoy14], we present here their detailed proofs. In Section 4 we introduce a notion of genericity that is compatible with weak computability. We prove the second main result of the paper: under suitable assumptions, a point exists that is at the same time “generic” and “weakly computable”. We then apply this result to four classes of objects: c.e. sets, left-c.e. reals, Π_1^0 -classes and regular Π_1^0 -classes.

2 Background and notations

We assume familiarity with basic computability theory on the natural numbers. We implicitly use Weihrauch’s notions of computability on effective topological spaces, based on the standard representation (see [Wei00] for more details), however we do not express them in terms of representations.

2.1 Notations

In a metric space (X, d) , if $x \in X$ and $r \in (0, +\infty)$ then we denote the open ball with center x and radius r by $B(x, r) = \{x' \in X : d(x, x') < r\}$. We denote the corresponding closed ball by $\overline{B}(x, r) = \{x' \in X : d(x, x') \leq r\}$. The Cantor space of infinite binary sequences, or equivalently subsets of \mathbb{N} , is denoted by $2^{\mathbb{N}}$. The halting set, denoted \emptyset' , is the set of numbers of Turing machines that halt. It is a noncomputable set that is computably enumerable (c.e.).

2.2 Effective topology

An *effective topological space* (X, τ, \mathcal{B}) consists of a topological space¹ (X, τ) together with a countable basis $\mathcal{B} = \{B_0, B_1, \dots\}$ numbered in such a way that the finite intersection operator is computable. An open subset $U \subseteq X$ is *effectively open* if $U = \bigcup_{k \in W} B_k$ for some c.e. set $W \subseteq \mathbb{N}$.

To a point $x \in X$ we associate $N(x) = \{n \in \mathbb{N} : x \in B_n\}$. By an *enumeration of $N(x)$* we mean a total function $f : \mathbb{N} \rightarrow \mathbb{N}$ whose range is $N(x)$. A point x is *computable* if $N(x)$ is c.e., i.e. if $N(x)$ has a computable enumeration.

Given points x, y in effective topological spaces X, Y respectively, we say that y is *computable relative to x* if there is an oracle Turing machine M that, given any enumeration of $N(x)$ as oracle, outputs an enumeration of $N(y)$. We denote it by $M^x = y$. In other words, y is computable relative to x if $N(y)$ is enumeration reducible to $N(x)$. As proved by Selman [Sel71] and pointed out by Miller [Mil04], y is computable relative to x if and only if every enumeration of $N(x)$ computes an enumeration of $N(y)$ (uniformity is not explicitly required, but is a consequence).

A (possibly partial) function $f : X \rightarrow Y$ is *computable* if there is a machine M such that for every $x \in \text{dom}(f)$, $M^x = f(x)$. A computable function is always continuous.

2.3 Effective Polish spaces

An *effective Polish space* is a topological space such that there exists a dense sequence s_0, s_1, \dots of points, called *simple* points and a complete metric d inducing the topology, such that all the reals numbers $d(s_i, s_j)$ are computable uniformly in (i, j) . Every effective Polish space can be made an effective topological space, taking as canonical basis the open balls $B(s, r)$

¹Such spaces are usually assumed to be T_0 , however this assumption is not necessary

with s simple point and r positive rational together with a standard effective numbering. It is sometimes useful to use another basis: we say that a family $\mathcal{B} = \{B_i\}_{i \in \mathbb{N}}$ of open sets is an *effective basis* if the identity between the effective topological spaces induced by \mathcal{B} and by the canonical basis is computable in the two directions. There always exists an effective basis \mathcal{B} that makes the predicate $s_i \in B_j$ computable in i, j .

In an effective Polish space, a point x is computable if and only if for every $\epsilon > 0$ a simple point s can be computed, uniformly in ϵ , such that $d(s, x) < \epsilon$.

We present a few simple observations on effective Polish spaces which seem to be new and are of independent interest. Let X be an effective Polish space. $X' \subseteq X$ is an *effective Polish subspace* if it is an effective Polish space with the induced topology and such that the canonical injection from X' to X to be a computable homeomorphism. Alexandrov theorem gives a way to obtain Polish subspaces of a Polish space, and has an effective version, which we present now.

A set A is an *effective G_δ -set* if there exists a family of uniformly effective open sets U_n such that $A = \bigcap_n U_n$. The following result unifies two known results: every closed set whose hit set (see statement below) is c.e. contains a dense computable sequence; every dense effective G_δ -set contains a dense computable sequence (computable Baire theorem [YMT99, Bra01]).

Proposition 2.1. *Let A be an effective G_δ -subset of an effective Polish space. The hit set $\{i \in \mathbb{N} : B_i \cap A \neq \emptyset\}$ is c.e. if and only if A contains a computable sequence dense in A . We then say that A is **c.e.***

Proof. We prove the non-trivial direction. Given a ball B_0 intersecting A , one can compute a sequence of balls B_n of radius $< 2^{-n}$ such that $\overline{B_{n+1}} \subseteq B_n \subseteq U_n$. The intersection $\bigcap_n B_n$ contains one point that is computable and belongs to A . The induction hypothesis on B_n is that it intersects A . Once B_n has been built, there must be some ball B of radius $< 2^{-n-1}$ such that $\overline{B} \subseteq B_n \cap U_n$ and B intersects A , and such a B can be effectively computed. Let $B_{n+1} = B$. To obtain a dense computable sequence, start from any ball B_0 intersecting A . \square

Proposition 2.2 (Effective Alexandrov Theorem). *Let X be an effective Polish space. Every c.e. effective G_δ -set is an effective Polish subspace of X .*

Proof. Let $A = \bigcap_n U_n$ be a c.e. effective G_δ -set. Let d be a complete computable metric on X . Let $d_n : X \rightarrow [0, +\infty)$ be uniformly computable

functions such that $d_n(x) > 0 \iff x \in U_n$. d_n is a computable version of the distance to the complement of U_n . For $x \in U_n$, let $f_n(x) = \frac{1}{d_n(x)}$. For $x, y \in A$, let

$$d'(x, y) = d(x, y) + \sum_n 2^{-n} \frac{|f_n(x) - f_n(y)|}{1 + |f_n(x) - f_n(y)|}.$$

The function $d' : A \times A \rightarrow \mathbb{R}$ is computable. On A it is a complete metric that induces the same topology as d . The computable sequence which is dense in A can serve as special points in A . The canonical injection is 1-Lipschitz, hence has a computable modulus of uniform continuity and is computable on the special points of A , so it is computable. \square

The theorem is actually an equivalence.

We will be concerned with computability and Baire category, so we will naturally meet the notion of a 1-generic point: a point that does not belong to any “effectively meager set” in the following sense.

Definition 2.1. $x \in X$ is **1-generic** if x does not belong to the boundary of any effective open set. In other words, for every effective open set U , either $x \in U$ or there exists a neighborhood B of x disjoint from U .

By the Baire category theorem, every Polish space is a Baire space so 1-generic points exist and form a co-meager set.

3 Irreversible functions

3.1 A non-uniform result

Let X be an effective Polish space, Y an effective topological space and $f : X \rightarrow Y$ a (total) computable function.

To introduce informally the results of this section, assume temporarily that f is one-to-one. If f^{-1} is computable, i.e. if every x is computable relative to $f(x)$ *uniformly* in x , then f^{-1} is continuous. As mentioned earlier uniformity is crucial here: that some x is computable relative to $f(x)$ does not imply in general that f^{-1} is continuous at $f(x)$. Theorem 3.1 below surprisingly shows that a non-uniform version can still be obtained, valid at most points.

Let us now make it precise and formal. We do not assume anymore that f is one-to-one.

When focusing on the problem of inverting a function, one comes naturally to the following basic notions:

- f is *invertible* at x if x is the only pre-image of $f(x)$,
- f is *locally invertible* at x if x is isolated in the pre-image of $f(x)$.

If one has access to x via its image only, then x is determined unambiguously in the first case, with the help of a discrete advice (a basic open set isolating x) in the second case. However, “being uniquely determined” is not sufficient in practice: physically or computationally, one cannot know entirely $f(x)$ in one step, but progressively as a limit of finite approximations. We need to consider stronger, topological versions of the two basic notions of invertibility, expressing that x can be recovered from the knowledge of its image given by finer and finer neighborhoods.

Definition 3.1. Let $f : X \rightarrow Y$ be a function.

We say that f is ***continuously invertible at x*** if the pre-images of the neighborhoods of $f(x)$ form a neighborhood basis of x , i.e. for every neighborhood U of x there exists a neighborhood V of $f(x)$ such that $f^{-1}(V) \subseteq U$.

We say that f is ***locally continuously invertible at x*** if there exists a neighborhood B of x such that the restriction of f to B is continuously invertible at x , i.e. for every neighborhood U of x there exists a neighborhood V of $f(x)$ such that $B \cap f^{-1}(V) \subseteq U$.

Observe that these notions are very natural when investigating the problem of inverting a function: we think that they are not technical *ad hoc* conditions.

Every effective topological space Y has a countable basis hence is sequential, i.e. continuity notions can be expressed in terms of sequences, which may be more intuitive. We will be particularly interested in the negations of these notions, which we characterize now.

Proposition 3.1. *f is not continuously invertible at x if and only if there exist $\delta > 0$ and a sequence x_n such that $d(x, x_n) > \delta$ and $f(x_n)$ converges to $f(x)$.*

f is not locally continuously invertible at x if and only if for every $\epsilon > 0$ there exist $\delta > 0$ and a sequence x_n such that $\epsilon > d(x, x_n) > \delta$ and $f(x_n)$ converges to $f(x)$.

We now come to our first result.

Theorem 3.1 (Computability implies continuity, pointwise). *Let $f : X \rightarrow Y$ be a computable function and $x \in X$ a 1-generic point.*

If x is computable relative to $f(x)$ then f is locally continuously invertible at x .

Proof. Assume that x is computable relative to $f(x)$ and f is not locally continuously invertible at x . We show that x belongs to the boundary of an effective open set U , i.e. that x is not 1-generic.

Intuitively, for a point y , there are two possible ways in which a Turing machine may fail to compute y from $f(y)$: either it diverges, or it outputs something that is incompatible with y . The latter can be recognized in finite time: we then say that $M^{f(y)}$ *positively* fails to compute y . Our effective open set U will be the set of points y such that $M^{f(y)}$ positively fails to compute y .

Let us make it more precise. As x is computable relative to $f(x)$, there exist uniformly effective open sets $V_n \subseteq Y$ such that for all i , $x \in B_n \iff f(x) \in V_n$, where B_n is the canonical basis of the topology on X . We then define

$$U = \bigcup_n f^{-1}(V_n) \setminus \overline{B_n}$$

which is an effective open set. By definition of V_n , x does not belong to U .

Let us show that x belongs to the closure of U . Let B be a neighborhood of x and U_B another neighborhood coming from the fact that f is not locally continuously invertible at x . Let B_n be a neighborhood of x such that $\overline{B_n} \subseteq U_B$. The set V_n is a neighborhood of $f(x)$, so $f^{-1}(V_n)$ intersects $B \setminus U_B \subseteq B \setminus \overline{B_n}$. As a result, B intersects U . This is true of every neighborhood B of x , so x belongs to the closure of U . \square

In the sequel we introduce a condition on f which roughly means that f is “almost nowhere” locally continuously invertible and that entails (i) the existence of an x that is not computable relative to $f(x)$ (Theorem 3.2) and, better, (ii) the existence of a non-computable x such that $f(x)$ is computable (Theorem 3.3).

3.2 Irreversible functions

We now consider the following notion: an *irreversible* function is locally continuously invertible at almost no point, in the sense of Baire category.

Definition 3.2. f is *irreversible* if for every non-empty open set B there exists a non-empty open set $U_B \subseteq B$ such that there is no open set V satisfying $\emptyset \neq f^{-1}(V) \cap B \subseteq U_B$.

In other words, if the pre-image of an open set intersects B then it intersects $B \setminus U_B$.

Intuitively, in a game between a player progressively describing $f(x)$ for some $x \in U_B$ and an opponent trying to progressively guess x , the opponent can never guess that $x \in U_B$ even knowing that $x \in B$.

Observe that one can assume w.l.o.g. that U_B is a basic ball and $f^{-1}(V) \cap B \not\subseteq \overline{U}_B$. Indeed, one can replace U_B by some ball $B(s, r)$ such that $\overline{B}(s, r) \subseteq U_B$.

An application of an irreversible function f to x comes with a loss of information about x , that can hardly be recovered. Being irreversible is orthogonal to not being one-to-one: the function $x \mapsto x^2$ is not one-to-one but not irreversible: x can be (continuously or computably) recovered from x^2 ; a one-to-one function can be irreversible if its inverse is dramatically discontinuous (examples of such functions will be encountered in the sequel).

In terms of sequences, f is irreversible if and only if for every B there exists a non-empty open set $U_B \subseteq B$ such that for every $x \in U_B$ there is a sequence $x_n \in B \setminus U_B$ such that $f(x_n)$ converges to $f(x)$.

As announced, the set of points at which an irreversible function is locally continuously invertible is small in the sense of Baire category.

Proposition 3.2. *Let f be irreversible. There is a dense G_δ -set D such that f is not locally continuously invertible at any $x \in D$.*

Proof. Let W_n be the union of U_B for all basic open sets B of radius $< 2^{-n}$. W_n is a dense open set. Let $x \in \bigcap_n W_n$. For each n there is a ball of radius $< 2^{-n}$ such that $x \in U_B$. For every neighborhood V of $f(x)$, $x \in f^{-1}(V) \cap B \neq \emptyset$ so $f^{-1}(V) \cap B \not\subseteq U_B$. \square

In other words, for almost every x the application of f to x comes with a “topological information” loss.

The preceding proposition does not rule out the possibility that the restriction of f to a “large” set be continuously invertible (for instance, the characteristic function of the rational numbers is nowhere continuous, but its restriction to the co-meager set of irrational numbers is continuous). The next assertion shows that this is not possible.

Proposition 3.3. *Let f be irreversible and $C \subseteq X$ be such that $f|_C : C \rightarrow f(C)$ is an homeomorphism. Then C is nowhere dense.*

Proof. Assume that the closure of C contains a ball B . Let $x \in U_B \cap C$. There exists a sequence $x_n \in B \setminus \overline{U}_B$ such that $f(x_n)$ converges to $f(x)$. By density of C in B , x_n can be taken in C . As $f|_C$ is an homeomorphism and $f(x_n)$ converges to $f(x)$, x_n should converges to x and eventually enter U_B , which gives a contradiction. \square

In the definition of an irreversible function (Definition 3.2), B and U_B can be assumed w.l.o.g. to be basic balls.

Definition 3.3. f is *effectively irreversible* if U_B can be computed from B in Definition 3.2.

The following result is the effective version of Proposition 3.3.

Theorem 3.2. *If f is computable and effectively irreversible then for every 1-generic x , x is not computable relative to $f(x)$.*

Proof. The dense G_δ -set provided by Proposition 3.2 is effective when f is effectively irreversible so it contains every 1-generic point. Hence for every 1-generic x , f is not locally continuously invertible at x . We now apply Theorem 3.1. \square

In other words, if x is 1-generic then the application of f to x comes with an “algorithmic information” loss. So if f is effectively irreversible then there exists some x that is not computable relative to $f(x)$.

3.3 The constructive result

We now present the main result of the paper. It is the constructive version of Theorem 3.2 as it makes $f(x)$ computable. The construction uses a priority argument with finite injury.

Theorem 3.3. *If f is computable and effectively irreversible then there exists a non-computable x such that $f(x)$ is computable.*

The proof uses the priority method with finite injury, which can be seen as a game between a player, computing $f(x)$, and infinitely many opponents (all the Turing machines) trying to compute x . The remainder of this section is devoted to the detailed proof of Theorem 3.3.

We fix a one-to-one computable enumeration n_0, n_1, \dots of the halting set \emptyset' . We construct $x \in X$ such that $f(x)$ is computable and \emptyset' is computable relative to x . We construct a shrinking sequence of balls B_n and define x as the unique member of their intersection. Of course, B_n must not be computable otherwise x would be computable. The sequence B_n is constructed in stages: at stage s we define $B_n[s]$ and for each n the sequence $B_n[s]$ is stationary, with limit B_n . For each s , the sequence $B_n[s]$ is shrinking, so the limiting sequence B_n will be shrinking as well. One may imagine, for each s , the sequence $B_n[s]$ as an infinite path in a tree. At stage $s + 1$, n_s is enumerated into A' and the current path branches at depth n_s .

In order to make $f(x)$ computable we enumerate along the construction the indices of all its basic neighborhoods into a list $L \subseteq \mathbb{N}$. L is the union of a computable growing sequence of finite lists L_s . At stage s , the current neighborhood of $f(x)$, denoted \mathcal{V}_s is the (finite) intersection of the basic open sets indexed by L_s . As $L_s \subseteq L_{s+1}$, $\mathcal{V}_{s+1} \subseteq \mathcal{V}_s$.

We need to consider two technical points. First we use a particular set of special points, induced by the effective irreversibility of f , obtained as follows. We can assume w.l.o.g. that the radius of U_B is at most half the radius of B . Given a basic ball B , consider the computable sequence $U_B^{(n)}$ defined inductively by $U_B^{(0)} = B$ and $U_B^{(n+1)} = U_{U_B^{(n)}}$. $U_B^{(n)}$ is a computable shrinking sequence and the unique member a of $\bigcap_n U_B^{(n)}$ is computable, uniformly in B . The canonical enumeration B_j of basic balls induces a computable dense sequence a_j , which will serve as simple points.

We then come to the second technical point. Let $(V_k)_{k \in \mathbb{N}}$ be the canonical enumeration of the basic open subsets of Y . We assume that the effective open sets $f^{-1}(V_k)$ come with growing enumerations $f^{-1}(V_k)[s]$ such that the predicate $a_i \in f^{-1}(V_k)[s]$ is decidable in i, k, s . Such enumerations exist as there exists an effective basis \mathcal{B}' of X that makes the predicate $a_i \in B'_j$ decidable, and the effective open sets $f^{-1}(V_k)$ can be expressed as effective unions of elements of \mathcal{B}' .

We now proceed to the construction of the sequence $B_n[s]$ for each stage s . For each s , $B_n[s]$ will be a shrinking sequence, $x[s]$ will be defined as the unique member of their intersection and will be one of the points $\{a_j : j \in \mathbb{N}\}$.

Stage 0. We start with a ball $B_0[0]$ of radius 1, $B_{n+1}[0] = U_{B_n[0]}$ and $\{x[0]\} = \bigcap_n B_n[0]$. Start with $L_0 = \emptyset$ and $\mathcal{V}_0 = Y$. Observe that for each n , $B_n[0] \cap f^{-1}(\mathcal{V}_0)$ is non-empty as it contains $x[0]$.

Stage $s + 1$. First, L_{s+1} is obtained by adding to L_s all the numbers $k \leq s$ such that $x[s] \in O_{k,s}$. Let \mathcal{V}_{s+1} be the intersection of the open sets V_k with $k \in L_{s+1}$.

Let $n = n_s$ be the next element enumerated into the halting set. Let $B_{n+1}[s+1]$ be a ball satisfying $\overline{B_{n+1}[s+1]} \subseteq f^{-1}(\mathcal{V}_{s+1}) \cap B_n[s] \setminus \overline{B_{n+1}[s]}$. Such a ball exists: $f^{-1}(\mathcal{V}_{s+1}) \cap B_{n+1}[s]$ is non-empty as it contains $x[s]$, f is irreversible and $B_{n+1}[s] = U_{B_n[s]}$. For $n' \leq n$, let $B_{n'}[s+1] = B_{n'}[s]$. For $n' > n$ define by induction $B_{n'+1}[s+1] = U_{B_{n'}[s+1]}$. Let $\{x[s+1]\} = \bigcap_n B_n[s+1]$.

Verification. By construction one has $\overline{B_{n+1}[s]} \subseteq B_n[s]$ and $B_{n+1}[s] = U_{B_n[s]}$ for sufficiently large n so $B_n[s]$ is a shrinking sequence.

We call the *settling time* of n the minimal number s such that $n_{s'} \geq n$ for all $s' \geq s$.

We say that $n \in \emptyset'$ is a *forward element* if no element $m < n$ is enumerated into \emptyset' after the enumeration stage of n : in other words, the settling time of n coincides with its enumeration stage. As \emptyset' is infinite, it has infinitely many forward elements.

Claim 1. For each n , $B_n[s]$ is a stationary sequence.

Proof. Let s be the settling time of n : $B_n[s] = B_n[s_0]$ for all $s \geq s_0$. \square

Let B_n be its limit. B_n is a shrinking sequence as well, let x be the member of its intersection. Observe that the sequence $x[s]$ converges to x : given ϵ , let n be such that B_n has radius $< \epsilon$ and s_0 be the settling time n : for all $s \geq s_0$, $x[s] \in B_n[s] = B_n$ so $d(x[s], x) < \epsilon$.

Claim 2. $f(x)$ is computable.

Proof. We prove that a basic open set V_k contains $f(x)$ if and only if k is enumerated into the list $L = \bigcup_s L_s$.

If $k \in L_s$ for some s , let n be a forward element which is enumerated at some stage $s' \geq s$. $x \in \overline{B}_{n+1} = \overline{B}_{n+1}[s' + 1] \subseteq f^{-1}(\mathcal{V}_{s'+1}) \subseteq f^{-1}(\mathcal{V}_s) \subseteq f^{-1}(V_k)$.

Now let V_k be a basic neighborhood of $f(x)$. Let i_0 be such that $x \in O_{k,i_0}$. As $x[s]$ converges to x there is s such that $x[s] \in O_{k,i_0}$ for all $s' \geq s$. Let $t = \max(s, i_0)$: $x[t] \in O_{k,i_0} \subseteq O_{k,t}$ so k must be added to the list at stage $t + 1$ or earlier. \square

Claim 3. \emptyset' is computable relative to x .

Proof. Let p_i be the increasing sequence of forward elements. \emptyset' can be computed from the sequence p_i and the (computable) enumeration of \emptyset' .

From x one can inductively compute the sequence p_i . First, p_0 is the minimal n such that $x \notin B_{n+1}[0]$. Once p_i is known, let s be the stage at which p_i is enumerated into \emptyset' , i.e. $n_s = p_i$. p_{i+1} is the minimal $n > p_i$ such that $x \notin B_{n+1}[s + 1]$. \square

In the proof \emptyset' is encoded in x . The argument relativizes: given a set $A \subseteq \mathbb{N}$, there exists x_A such that A computes $f(x_A)$ and the pair (x_A, A) computes A' . All the reductions are uniform, so computing the jump operator can be reduced to computing the inverse of f (when f is one-to-one). The notion capturing this idea is Weihrauch reducibility [Wei92, BG11].

Corollary 3.1. *If f is one-to-one, computable and effectively irreversible then the jump operator is Weihrauch reducible to f^{-1} .*

3.4 Application to the ergodic decomposition

We now present an application of Theorem 3.3. Let P be a Borel probability measure P over the Cantor space. P is **computable** if the real numbers $P[w]$ are uniformly computable. P is **shift-invariant** if $P[w] = P[0w] + P[1w]$ for each finite string w . P is **ergodic** if it cannot be written as $P = \frac{1}{2}(P_1 + P_2)$ with $P_1 \neq P_2$ both shift-invariant.

The ergodic decomposition theorem says that every shift-invariant measure can be uniquely decomposed into a convex combination (possibly uncountable) of ergodic measures. Our question is: given a computable shift-invariant measure, can we compute in a sense its ergodic decomposition? This question was implicitly addressed by V'yugin [V'y97] who constructed a counter example: a countably infinite combination of ergodic measures which is computable but not effectively decomposable. In [Hoy11] we raised the following question: does the ergodic decomposition become computable when restricting to finite combinations? As an application of Theorem 3.3, we solve the problem and prove that it is already non-effective in the finite case:

Theorem 3.4. *There exist two ergodic shift-invariant measures P and Q such that neither P nor Q is computable but $P + Q$ is computable.*

The strategy is as follows: the mapping $(P, Q) \mapsto P + Q$ is computable, two-to-one on the space $\mathcal{E} \times \mathcal{E}$ of pairs of ergodic measures and we prove

Theorem 3.5. *The function $(P, Q) \mapsto P + Q$ defined on $\mathcal{E} \times \mathcal{E}$ is effectively irreversible.*

which implies Theorem 3.4 applying Theorem 3.3.

Before proving the theorem, we need some preliminaries so show that $\mathcal{E} \times \mathcal{E}$ is an effective Polish space.

We consider the space $\mathcal{P}(2^{\mathbb{N}})$ of Borel probability measures over the Cantor space together with the complete metric

$$d(P, Q) = \sum_{w \in \{0,1\}^*} 2^{-|w|} |P[w] - Q[w]|.$$

The finite rational combination of Dirac measures are dense in $\mathcal{P}(2^{\mathbb{N}})$ and d is computable over them, so $\mathcal{P}(2^{\mathbb{N}})$ is an effective Polish space. The subset

\mathcal{I} of shift-invariant measures is closed so d is complete over \mathcal{I} as well. \mathcal{I} easily contains a dense computable sequence (take the Markovian measures with rational coefficients), so \mathcal{I} is an effective Polish subspace of $\mathcal{P}(2^{\mathbb{N}})$. Let $\mathcal{E} \subseteq \mathcal{I}$ be the set of ergodic shift-invariant measures. The metric d is no more complete over \mathcal{E} , but \mathcal{E} is an effective G_δ -set that is c.e. so Proposition 2.2 implies that \mathcal{E} is an effective Polish subspace (see [Par61] for results on the Baire category of the set of ergodic measures). We work with the basis given by the intersection of the canonical basis of \mathcal{I} with \mathcal{E} .

We now present the proof of Theorem 3.5.

Proof of Theorem 3.5. Let $B \subseteq \mathcal{I} \times \mathcal{I}$ be an open set and $(P_0, Q_0) \in B$ with $P_0 \neq Q_0$. Let $\epsilon > 0$ be such that $d(P_0, Q_0) > \epsilon$ and $B(P_0, \epsilon) \times B(Q_0, \epsilon) \subseteq B$. Let $\delta = \epsilon/4$ and $U_B = B(P_0, \delta) \times B(Q_0, \delta) \subseteq B$. Observe that U_B can be effectively obtained from B . We now show how a pair $(P_1, Q_1) \in U_B$ can be moved outside U_B , but still inside B , nearly without changing its sum. By the choice of δ , if $(P_1, Q_1) \in U_B$ then $d(P_1, Q_1) > 2\delta$. For $\lambda \in [0, 1]$, define

$$\begin{aligned} P(\lambda) &= \lambda P_1 + (1 - \lambda)Q_1, \\ Q(\lambda) &= \lambda Q_1 + (1 - \lambda)P_1. \end{aligned}$$

Observe that $P(\lambda) + Q(\lambda) = P_1 + Q_1$ and

$$d(P_1, P(\lambda)) = d(Q_1, Q(\lambda)) = (1 - \lambda)d(P_1, Q_1).$$

As $d(P_1, Q_1) > 2\delta$ there exists $\lambda \in (0, 1)$ such that $(1 - \lambda)d(P_1, Q_1) = 2\delta$. One has

$$d(P_0, P(\lambda)) \leq d(P_0, P_1) + d(P_1, P(\lambda)) < 3\delta < \epsilon$$

and

$$d(P_0, P(\lambda)) \geq d(P_1, P(\lambda)) - d(P_0, P_1) > \delta,$$

and similarly $\delta < d(Q_0, Q(\lambda)) < \epsilon$ so $(P(\lambda), Q(\lambda)) \in B \setminus \overline{U}_B$.

Observe that the shift-invariant measures $P(\lambda)$ and $Q(\lambda)$ are not ergodic. As the ergodic measures are dense in the set of shift-invariant measures, there exist two sequences P_n, Q_n of ergodic measures converging to $P(\lambda)$ and $Q(\lambda)$ respectively. As $(P(\lambda), Q(\lambda))$ belongs to the open set $B \setminus \overline{U}_B$, we can assume w.l.o.g. that $(P_n, Q_n) \in B \setminus \overline{U}_B$ for all n . The mapping $(P, Q) \mapsto P + Q$ is continuous so $P_n + Q_n$ converges to $P(\lambda) + Q(\lambda) = P_1 + Q_1$. \square

4 Directional genericity

Given an effectively irreversible function f ,

- Theorem 3.2 tells us that if x is 1-generic then x is not computable relative to $f(x)$,
- Theorem 3.3 tells us that there exist non-computable x such that $f(x)$ is computable.

The two results are “disjoint” in the sense that in general a single x cannot at the same time be 1-generic and have a computable image, except for some particular functions like constant functions. We raise the following question: is it possible to bring the two results closer together? How far can x be from being computable, given that $f(x)$ is computable? How *generic* can x be? In this section we give an answer to these questions, introducing a notion of genericity that is compatible with a weak form of computability.

For the sake of simplicity, we will assume that f is the identity. We fix a set X , endowed with an effective Polish topology τ and a weaker effective topology τ' . In doing so we lose no generality, as a function $f : (X, \tau_X) \rightarrow (Y, \tau_Y)$ can always be thought as the identity from (X, τ_X) to (X, τ') where τ' is the initial topology of f whose open sets are the preimages of τ_Y -open sets.

4.1 Being generic from above

Let (X, τ) be an effective Polish space and τ' be an effective topology on X that is effectively weaker than τ : the basic τ' -open sets are effective τ -open sets, uniformly. In other words, we require the identity function from (X, τ) to (X, τ') to be computable.

Our general goal is to build elements of X that are to some extent generic in the topology τ but still computable in the topology τ' . The latter condition is usually weaker than being computable in the topology τ , and will be our weak notion of computability. We now have to define a suitable notion of genericity.

The topology τ' induces a pre-order on X , called the *specialization* pre-order \leq :

$$x \leq y \iff \text{every } \tau'\text{-neighborhood of } x \text{ is a } \tau'\text{-neighborhood of } y.$$

$x \leq y$ means that if one describes x by listing its basic neighborhoods then one can never distinguish x from y . Observe that when τ' is T_0 , \leq is an order, and when τ' is T_1 , \leq is the trivial ordering (equality).

Definition 4.1. To $x \in X$ we associate

$$S_x = \{y \in X : x \leq y\}$$

which is the intersection of all the τ' -neighborhoods of x .

S_x is the set of elements that cannot be distinguished from x when describing x in the topology τ' . If τ' is T_1 then $S_x = \{x\}$ for all x .

In a game where the player describes an element x in the τ' -topology, the player enumerates the basic τ' -neighborhoods of x . Each enumerated basic open set is a commitment: if V is enumerated then x must belong to V . Each commitment reduces the degrees of freedom of the player to fool the opponent. However some free space is always left, and this space is precisely S_x : at any moment the player is allowed to move into S_x (and then change x at the same time). As a consequence, during the computation of x in the topology τ' , the player is able to make x as generic as possible, *inside the subset S_x* . This motivates the following definition.

Definition 4.2. Let (X, τ) be an effective Polish space, $A \subseteq X$ and $x \in A$. We say that x is **generic inside** A if for every effective open set $U \subseteq X$,

- either $x \in U$,
- or there exists a neighborhood B of x such that $B \cap U \cap A = \emptyset$.

If τ' is a weaker topology on X then we say that x is **generic from above** if x is generic inside $S_x = \{y \in X : x \leq y\}$, where \leq is the specialization pre-order induced by τ' .

Every 1-generic element is generic inside any set containing it. Let us give a few examples illustrating these notions. For $A = X$, being generic inside A is the same as being 1-generic. Every element x is vacuously generic inside $\{x\}$. In the product space $X \times X$, (x, y) is generic in $\{x\} \times X$ if and only if y is 1-generic relative to x (effective open subsets of X relative to x are the same as the sections at x of effective open subsets of $X \times X$).

Informally, x is generic from above means that x belongs to every effective open set that is dense *above* it, for the specialization order induced by τ' (while a 1-generic elements belongs to every effective open set that is dense *along* it).

With this notion in hand we obtain the sought combination of Theorems 3.2 and 3.3. For this we need a reasonable technical assumption on the bases \mathcal{B} and \mathcal{B}' of the topologies τ and τ' respectively.

Assumption 1 The non-emptiness of finite conjunctions and disjunctions of basic open sets from both \mathcal{B} and \mathcal{B}' is computable.

Theorem 4.1 (Generic and computable). *Let U_n be uniformly effective dense τ -open sets. Under assumption 1, there exists $x \in \bigcap_n U_n$ such that*

- x is generic inside S_x ,
- x is τ' -computable.

Observe that the theorem is only interesting when τ' is not T_1 , otherwise $S_x = \{x\}$ and the first condition is vacuously satisfied for every x .

Proof. Let a_i be a dense computable sequence of simple points in (X, τ) . Again we can assume w.l.o.g. that the basic open sets $V_k \in \mathcal{B}'$ come with a computable enumeration $V_k[s]$ ($V_k[s]$ is a finite union of elements of \mathcal{B}) such that the predicate $a_i \in V_k[s]$ is decidable in i, k, s .

The proof is a finite injury argument. We want to satisfy the requirements

$$R_e : x \in W_e \text{ or } \exists \epsilon, B(x, \epsilon) \cap W_e \cap S_x = \emptyset,$$

where W_e is the effective open set with number e (in the sequel, $W_{e,s}$ will be computable growing finite unions of elements of \mathcal{B} with union W_e). At stage s , each requirement R_e is assigned a ball $B_e[s]$. They satisfy $\overline{B_{e+1}[s]} \subseteq B_e[s] \cap U_e$. For each e , the sequence $B_e[s]$ is stationary when s grows. At the same time, a list $L \subseteq \mathbb{N}$ is enumerated containing exactly the indices of the basic \mathcal{B}' -neighborhoods of x . L is obtained as the union of a growing computable sequence of finite sets L_s . We denote by \mathcal{V}_s the finite intersection of elements of \mathcal{B}' whose indices are given by L_s . \mathcal{V}_s will be a neighborhood of x in the topology τ' . The requirement R_e tests whether $B_e[s]$ intersects $W_{e,s} \cap \mathcal{V}_s$.

Stage 0. Let $B_0[0]$ be any ball of radius $< 2^{-0}$ and inductively choose $\overline{B_{e+1}[0]} \subseteq B_e[0] \cap U_e$ of radius $< 2^{-e-1}$. Let x_0 be the center of $B_0[0]$ and $L_0 = \emptyset$.

Stage $s+1$. Let $e \leq s$ be minimal such that $B_e[s] \cap W_{e,s} \cap \mathcal{V}_s \neq \emptyset$ and R_e is not already declared satisfied, if it exists (decidable property). Let $B_{e'}[s+1] = B_{e'}[s]$ for $e' \leq e$, let $\overline{B_{e+1}[s+1]} \subseteq B_e[s] \cap W_{e,s} \cap \mathcal{V}_s \cap U_e$ have radius $< 2^{-e-1}$ and x_{s+1} be the center of $B_{e+1}[s+1]$. Define inductively $\overline{B_{e'+1}[s+1]} \subseteq B_{e'}[s+1] \cap U_{e'}$ of radius $< 2^{-e'-1}$ for $e' > e$. We say that R_e acts and R_e is declared satisfied. All $R_{e'}$ with $e' > e$ are initialized, which means that all of them are declared unsatisfied.

If e does not exist then let $B_e[s+1] = B_e[s]$ for all e and $x_{s+1} = x_s$.

Put $k \leq s$ in L_{s+1} iff $x_{s+1} \in V_k[s]$.

Verification. By the usual analysis of finite injury arguments, each requirement acts finitely many times, so for each e there is s_0 such that $B_e[s] = B_e[s_0]$ for all $s \geq s_0$. Let $B_e = B_e[s_0]$. One has $\overline{B_{e+1}} \subseteq B_e$ and B_e has radius $< 2^{-e}$. Let x be the unique member of $\bigcap_e B_e$.

Claim 4. The sequence x_s converges to x .

For each k , and all sufficiently large s , only requirements R_e with $e > k$ act, so $x_{s+1} \in B_{e+1}[s+1] \subseteq B_k[s+1] = B_k$. As $x \in B_k$ and B_k has radius $< 2^{-k}$, $d(x_{s+1}, x) < 2^{-k+1}$ for all sufficiently large s .

Claim 5. $x \in \bigcap_e U_e$.

By construction, $B_{e+1} \subseteq U_e$ for all e , hence $x \in \bigcap_e B_{e+1} \subseteq \bigcap_e U_e$.

Claim 6. L lists exactly the elements of \mathcal{B}' containing x , hence x is τ' -computable.

First, x belongs to each \mathcal{V}_s so L lists only (indices of) neighborhoods of x . Assume that V_k is a neighborhood of x . It implies that some $V_k[s_0]$ is a neighborhood of x . As x_s converges to x , x_s belongs to $V_k[s_0]$ for all s larger than some s_1 . As a result, for $s > \max(s_0, s_1)$, k is listed in L_s .

Claim 7. x is generic inside S_x .

Let $e \in \mathbb{N}$ be such that $x \notin W_e$. Let s be such that no requirement $e' < e$ acts from stage s on. R_e cannot act at a stage $s' \geq s$, otherwise $x \in B_{e+1} = B_{e+1}[s'+1] \subseteq W_e$ which contradicts the assumption $x \notin W_e$. In the same way, R_e cannot be declared satisfied at stage s , otherwise $x \in B_{e+1} = B_{e+1}[s'+1] \subseteq W_e$. As R_e never acts after stage s , it means that $x \in B_e = B_e[s]$ and $B_e[s] \cap W_e \cap S_x = \emptyset$, otherwise there exists $s' \geq s$ such that $B_e[s] \cap W_{e,s'} \cap S_x \neq \emptyset$, hence $B_e[s'] \cap W_{e,s'} \cap \mathcal{V}_{s'} \neq \emptyset$ as $B_e[s'] = B_e[s]$ and $S_x \subseteq \mathcal{V}_{s'}$, and then R_e must act at stage s' as it is not declared satisfied. This is a contradiction. \square

The point x provided by Theorem 4.1 actually satisfies a stronger notion of genericity.

Lemma 4.1. *For each effective open set $W_e = \bigcup_s W_e[s]$,*

- *either $x \in W_e$,*
- *or there exists a neighborhood B of x such that $B \cap (\bigcup_s W_e[s] \cap \mathcal{V}_s) = \emptyset$.*

Proof. Indeed, if $x \notin W_e$ then for when all $R_{e'}$ with $e' \leq e$ have settled, $B_e[s] = B_e$ and $B_e[s] \cap W_e[s] \cap \mathcal{V}_s = \emptyset$ otherwise R_e will act and force x to fall into W_e , so one can take $B = B_e$. \square

In other words, x satisfies the condition of 1-genericity not for every effective open set W_e , but for every the effective open set $\bigcup_s W_e[s] \cap \mathcal{V}_s$. This genericity condition has the consequence that x behaves in some respects as a 1-generic point, as illustrated by the following two results.

Corollary 4.1. *The point x provided by Theorem 4.1 is low, i.e. the set $\{e \in \mathbb{N} : x \in W_e\}$ is Δ_2^0 (or \emptyset' -computable, or limit of a computable sequence).*

Proof. For each e , the computable predicate $B_e[s] \cap W_e[s] \cap \mathcal{V}_s \neq \emptyset$ converges to the predicate $x \in W_e$. Indeed, if $B_e[s] \cap W_e[s] \cap \mathcal{V}_s \neq \emptyset$ for infinitely many s then R_e will eventually act and never be injured later, so x will be forced to fall into W_e . If $x \in W_e$ then for sufficiently large s , $B_e[s] = B_e$ and $x \in W_e[s]$ so $B_e[s] \cap W_e[s] \cap \mathcal{V}_s \neq \emptyset$. \square

On compact spaces, Theorem 4.1 is indeed a strengthening of Theorem 3.3.

Corollary 4.2. *Assume that (X, τ) is compact. If $\text{id} : (X, \tau) \rightarrow (X, \tau')$ is effectively irreversible then the point x provided by Theorem 4.1 can be taken non-computable.*

Proof. One can take x in the dense effective G_δ given by Proposition 3.2, so that id is not locally continuously invertible at x .

If x is computable then there exists e such that $W_e = X \setminus \{x\}$. As $x \notin W_e$ there exists B such that $B \cap \bigcup_s W_e[s] \cap \mathcal{V}_s = \emptyset$. Let $\bar{U}_B \subseteq B$ come from the local continuous non-invertibility of id at x . As W_e covers the compact set $X \setminus U_B$, there exists s such that $W_e[s]$ already covers that set. As \mathcal{V}_s is a τ' -neighborhood of x , $B \cap \mathcal{V}_s \setminus U_B \neq \emptyset$ so $B \cap \mathcal{V}_s \cap W_e[s] \neq \emptyset$, which contradicts the choice of B . \square

We now illustrate directional genericity in several situations and show how Theorem 4.1 embodies many constructions encountered in computability theory. It means that in many situations, in order to construct an object satisfying a given set of requirements, one only has to find the suitable topologies τ and τ' that make directionally generic objects have the sought properties. Theorem 4.1 can then be directly applied, instead of explicitly constructing the object by means of a finite injury argument.

4.2 Genericity for c.e. sets

We consider the Cantor space X of subsets of \mathbb{N} . Here τ is the Cantor topology and τ' is the Scott topology. For a set $A \subseteq \mathbb{N}$, $S_A = \{B \subseteq \mathbb{N} : A \subseteq B\}$ is the class of supersets of A .

Definition 4.3. A set $A \subseteq \mathbb{N}$ is *generic from above* if it is 1-generic inside S_A , which means that for every effective open class \mathcal{U} , either $A \in \mathcal{U}$ or there exists n such that $[A \upharpoonright_n] \cap \mathcal{U} \cap S_A = \emptyset$.

In other words, A is generic from above if it belongs to every effective open class that is dense *above* A in the subset ordering. This should be compared to the notion of 1-generic set, which belongs to every effective open class that is dense *along* it.

As a direct application of Theorem 4.1 we obtain:

Corollary 4.3. *There exists a co-infinite c.e. set $A \subseteq \mathbb{N}$ that is generic from above.*

Proof. The class of co-infinite sets is a dense effective G_δ -set. □

As the next result shows, Theorem 4.1 embodies simple finite injury arguments as Friedberg-Muchnik theorem, e.g. Interestingly co-infinite sets that are generic from above inherit many properties of 1-generic sets, with the same arguments (observing that the effective open classes appearing in those arguments are not only dense *along* the set, but also *above* it).

Proposition 4.1. *Let A be co-infinite and generic from above. $\mathbb{N} \setminus A$ is hyperimmune, $A = A_1 \oplus A_2$ where A_1 and A_2 are Turing incomparable, A is not autoreducible.*

Proof. Same argument as for 1-generic sets, observing that the involved open set is not only dense *along* A , but even *above* A . For instance, to prove that $A_2 \not\leq_T A_1$, given a Turing functional ϕ , let $U = \{A_1 \oplus A_2 : \exists n, \phi^{A_1}(n) = 0 \wedge A_2(n) = 1\}$. If $\phi^{A_1} = A_2$ then replacing a 0 in A_2 by a 1 arbitrarily far gives an element of U arbitrarily close to $A_1 \oplus A_2$ that is *above* (i.e. is a superset of) $A_1 \oplus A_2$. □

It happens that the co-infinite sets that are generic from above are exactly the p -generic sets defined by Ingrassia [Ing81].

4.3 Genericity for left-c.e. reals

We consider the unit real interval $[0, 1]$. τ is the Euclidean topology, τ' is the topology induced by the semi-lines $(x, 1]$. The specialization order is the natural ordering on real numbers. For a real $x \in [0, 1]$, $S_x = [x, 1]$.

Definition 4.4. A real $x \in [0, 1]$ is *generic from the right* if it is 1-generic inside $[x, 1]$, which means that for every effective open set $\mathcal{U} \subseteq [0, 1]$, either $x \in \mathcal{U}$ or there exists $\epsilon > 0$ such that $[x, x + \epsilon) \cap \mathcal{U} = \emptyset$.

Again x is generic from the right if it belongs to every effective open set that is dense *above* x in the real ordering.

Kurtz built a left-c.e. weakly 1-generic real (see [Nie09] for a proof). The construction even gives a left-c.e. real that is generic from the right. One can think of the proof of Theorem 4.1 as a kind of generalization of this argument (replacing the lexicographic order used in the proof appearing in [Nie09] by the specialization pre-order induced by τ').

Genericity from the right easily lies between two classical notions of genericity.

Proposition 4.2. *Every 1-generic is generic from the right. Every generic from the right is weakly-1-generic. The implications are strict.*

Proof. An open set that is dense is dense on the right of x . An open set that is dense on the right of x is dense along x . Right-c.e. reals cannot be generic on the right, but there exists a right-c.e. weakly-1-generic real. Left-c.e. reals cannot be 1-generic, but there exists a left-c.e. real that is generic on the right. \square

In Section 4.5 we will separate genericity from the right from weakly-1-genericity among the left-c.e. reals.

Solovay reducibility vs. cl-reducibility. If A is a subset of \mathbb{N} then we denote by x_A the real number whose binary expansion is A . In [DHL04] it is proved that there exist two sets A, B such that A is a c.e. set, x_B is a left-c.e. real and $A \leq_{\text{cl}} B$ but $x_A \not\leq_S x_B$. Here \leq_{cl} stands for computably Lipschitz reducibility and \leq_S stand for Solovay reducibility. The construction is a finite injury argument, which again is captured by Theorem 4.1.

Let us recall that $A \leq_{\text{cl}} B$ means that there is a Turing functional computing A with oracle B , reading the first $n + c$ bits of B to compute the n first bits of A , for some constant c and all n . $x_A \leq_S x_B$ means that there exists a constant $c \in \mathbb{N}$ and computable sequences $a_i \nearrow x_A$ and $b_i \nearrow x_B$ such that $x_A - a_i \leq c(x_B - b_i)$ for all i , or equivalently that $cx_B - x_A$ is left-c.e.

Theorem 4.2. *Let x_B be left-c.e. and generic on the right and $A = \{w \in 2^{<\mathbb{N}} : w <_{\text{lex}} B\}$. One has $A \leq_{\text{cl}} B$ but $x_A \not\leq_S x_B$.*

Note that we identify A with a subset of \mathbb{N} by using a computable bijection between $2^{<\mathbb{N}}$ and \mathbb{N} (we will assume that the string represented by a number n has length at most n).

Proof. The reduction $A \leq_{\text{cl}} B$ is obvious: to know whether $w <_{\text{lex}} B$, one only needs to know the $|w|$ first bits of B .

Assume that $x_A \leq_S x_B$. It implies the existence of a one-to-one computable enumeration w_i of A , a computable sequence $b_i \nearrow x_B$ and a computable order $h : \mathbb{N} \rightarrow \mathbb{N}$ such that for all $i \in \mathbb{N}$, if $x_B - b_i < 2^{-h(n)}$ then $|w_i| > n$. Let B_i be the maximal element of $\{w_0, \dots, w_i\}$ in the lexicographic ordering. If $x_B - b_i < 2^{-h(n)}$ then $B_i \upharpoonright_n = B \upharpoonright_n$: indeed, the string $B \upharpoonright_n$ belongs to A so it must be w_j for some j , which must be less than i .

For $w \in 2^{<\mathbb{N}}$, let $[w]$ be the interval containing all real numbers having a binary expansion starting with w , namely $[w] = [0.w, 0.w + 2^{-|w|}]$.

Let $U = \bigcup_{n,i} (b_i, b_i + 2^{-h(n)}) \setminus [B_i \upharpoonright_n]$. U is an effective open set. U does not contain x_B . Indeed, if $x_B - b_i < 2^{-h(n)}$ then $B_i \upharpoonright_n = B \upharpoonright_n$ so $x_B \in [B_i \upharpoonright_n]$. We now prove that U is dense on the right of x_B , which contradicts the assumption that x_B is generic on the right. As x_B is generic on the right it is weakly 1-generic, so there exist infinitely many n such that B contains all the natural numbers from n to $h(n)$. In other words for infinitely many n , x_B is very close to the right endpoint of $[B \upharpoonright_n]$, namely at distance $< 2^{-h(n)}$. For such n and sufficiently large i , $(b_i, b_i + 2^{-h(n)}) \setminus [B_i \upharpoonright_n]$ is a non-empty subset of the interval $(x_B, x_B + 2^{-h(n)})$. \square

We actually prove more: there is no computable order h and no computable sequences $a_i \nearrow x_A$, $b_i \nearrow x_B$ such that $x_B - b_i \leq 2^{-h(n)}$ implies $x_A - a_i \leq 2^{-n}$, which would be a generalization of Solovay reducibility.

Left-c.e. reals with only computable presentations. Downey and LaForte [DL02] proved the existence of non-computable left-c.e. reals x all of whose presentations are computable: each prefix-free c.e. set A of finite binary strings satisfying $\sum_{w \in A} 2^{-|w|} = x$ is actually a computable set. A corollary of a result of Stephan and Wu [SW05] is that any such real is weakly 1-random, i.e. it must belong to every effective open set of measure one. Actually it must be weakly-1-generic and even generic from the right.

Proposition 4.3. *If x is a non-computable left-c.e. real all of whose presentations are computable then x is generic from the right.*

Proof. Let U be an effective open set that does not contain x : we must find $y > x$ such that $[x, y)$ is disjoint from U . First replace U by $V = U \cup [0, x)$. Let A be a prefix-free c.e. set such that $V = \bigcup_{w \in A} [w]$. The set $A_{<x} = \{w \in A : w <_{\text{lex}} x\}$ is a presentation of x hence it is computable, so $A_{>x} = \{w \in A : w >_{\text{lex}} x\} = A \setminus A_{<x}$ is c.e. As a result, $y := \inf \bigcup_{w \in A_{>x}} [w]$

is right-c.e. As x is not computable and $x \leq y$, one has $x < y$ and we get the result as $[x, y)$ is disjoint from U . \square

4.4 Genericity for Π_1^0 -classes

We work on the set $\text{CL}(2^{\mathbb{N}})$ of non-empty closed subsets of the Cantor space, endowed with the so-called *hit-or-miss* topology τ_{hm} . τ_{hm} is generated by the *miss* sets $\mathcal{U}_u = \{P \in \text{CL}(2^{\mathbb{N}}) : P \cap [u] = \emptyset\}$ where $u \in 2^{<\mathbb{N}}$, together with their complements (the *hit* sets). We obtain an effective Polish space $(\text{CL}(2^{\mathbb{N}}), \tau_{hm})$. A computable element of this space is usually called a computable closed set, and is the set of infinite branches of a computable tree without dead-ends.

Proposition 4.4. *In the space $(\text{CL}(2^{\mathbb{N}}), \tau_{hm})$, every weakly-1-generic element contains only weakly-1-generic sequences.*

Proof. Let $U \subseteq 2^{\mathbb{N}}$ be a dense effective open set. Let $\mathcal{U} = \{P \in \text{CL}(2^{\mathbb{N}}) : P \subseteq U\}$. \mathcal{U} is a dense effective open set in the space $\text{CL}(2^{\mathbb{N}})$. Hence every weakly-1-generic closed set P belongs to \mathcal{U} , i.e. is contained in U . \square

We consider a weaker topology τ_m called the *miss* topology, generated by the miss sets \mathcal{U}_u with $u \in 2^{<\mathbb{N}}$. A Π_1^0 -class is a computable member of $(\text{CL}(2^{\mathbb{N}}), \tau_m)$. The specialization pre-order induced by τ_m is the reverse inclusion, so that for each non-empty closed set P one has $\mathcal{S}_P = \{Q \in \text{CL}(2^{\mathbb{N}}) : Q \subseteq P\}$ (being “above” P in this pre-order means being *inside* P). Definition 4.2 is instantiated as follows.

Definition 4.5. A non-empty closed set $P \subseteq 2^{\mathbb{N}}$ is **generic from inside** if P is 1-generic inside \mathcal{S}_P , which means that for every effective open set $\mathcal{U} \subseteq \text{CL}(2^{\mathbb{N}})$, either $P \in \mathcal{U}$ or there exists a τ_{hm} -neighborhood \mathcal{N} of P such that $\mathcal{N} \cap \mathcal{U} \cap \mathcal{S}_P = \emptyset$.

One can easily see that a closed set that is generic from inside has empty interior, i.e. has a dense complement. As a result, no member of a Π_1^0 -class P that is generic from inside can be weakly-1-generic. However all the elements of P are weakly-1-generic *inside* P : if U is an effective open subset of the Cantor space such that $P \cap U$ is dense in P then U contains every member of P , i.e. U contains P .

Proposition 4.5. *If P is generic from inside then every member of P is weakly-1-generic inside P .*

Proof. Let $U \subseteq 2^{\mathbb{N}}$ be an effective open set. Consider the set $\mathcal{U} = \{P : P \subseteq U\}$. \mathcal{U} is an effective open set in the space $(\text{CL}(2^{\mathbb{N}}), \tau_{hm})$ (and even in the topology τ_m). If $P \cap U$ is dense in P then there exists $Q \subseteq P \cap U$ arbitrarily τ_{hm} -close to P (let Q be a finite set of points from $P \cap U$ whose ϵ -neighborhood covers P , for arbitrarily small ϵ), so P belongs to the closure of $\mathcal{U} \cap \mathcal{S}_P$. If P is generic from inside then P must belong to \mathcal{U} , so $P \subseteq U$. \square

In particular,

Corollary 4.4. *A perfect closed set that is generic from inside has no computable member.*

Proof. If x is computable then $U = 2^{\mathbb{N}} \setminus \{x\}$ is an effective open set. If P has no isolated point then $P \cap U$ is dense in P . By the previous result, P is then contained in U , i.e. P does not contain x . \square

Now, Theorem 4.1 can be instantiated as follows.

Corollary 4.5. *There exists a perfect Π_1^0 -class that is generic from inside.*

Proof. Being perfect, or having no isolated point is a dense effective G_δ -property in the space $(\text{CL}(2^{\mathbb{N}}), \tau_{hm})$. \square

4.5 Genericity for regular Π_1^0 -classes

We know from Proposition 4.2 that every real $x \in [0, 1]$ that is generic on the right is weakly-1-generic, but not the converse. Here we prove the existence of *left-c.e.* reals that are weakly-1-generic but not generic from the right.

To this end we need to construct a Π_1^0 -set P such that (i) its leftmost element x is weakly-1-generic, and (ii) the complement of P is dense on the right of x . The first condition requires the class to have non-empty interior, and even that the interior of P be dense along x . Together with the second condition, it implies that x should not be isolated in the boundary of P .

The class P that we build will actually satisfy these conditions *at every point of its boundary*: P is regular (it coincides with the closure of its interior), its boundary is perfect (has no isolated point) and contains only weakly-1-generic points.

A suitable way of describing a regular closed set C is by giving approximations of C in the hit-or-miss topology and at the same time enumerating its interior. This can be formalized by introducing a new topology τ on $\text{CL}([0, 1])$ that is stronger than the hit-or-miss topology τ_{hm} . It is generated by the hit-or-miss open sets together with the sets

$$\{C \in \text{CL}([0, 1]) : [a, b] \subseteq \text{int}(C)\},$$

where $a < b$ are rational numbers and $\text{int}(C)$ is the interior of C . A canonical enumeration of the rational numbers gives a numbered basis for the topology τ , which makes $(\text{CL}([0, 1]), \tau)$ an effective topological space.

Intuitively, describing a closed set C in the topology τ consists in giving approximations of C in the hit-or-miss topology and at the same time enumerating the interior of C , which is equivalent to giving approximations of both C and $(\text{int}(C))^c$ in the hit-or-miss topology.

Proposition 4.6. *The space $(\text{CL}([0, 1]), \tau)$ is an effective Polish space.*

Proof. The space can be embedded into $\text{CL}([0, 1]) \times \text{CL}([0, 1])$ endowed with the product of the hit-or-miss topology, which is an effective Polish space. Indeed, the space is computably homeomorphic to the subset $\{(C, (\text{int}(C))^c) : C \in \text{CL}([0, 1])\}$ of $\text{CL}([0, 1]) \times \text{CL}([0, 1])$. We show that this subset is a c.e. effective G_δ -set, which implies that it is an effective Polish space by Proposition 2.2.

Claim 8. The set $\{(A, B) : (\text{int}(A))^c \subseteq B\}$ is Π_1^0 .

Indeed, $(\text{int}(A))^c \subseteq B$ is equivalent to $A^c \subseteq B$, which holds iff every rational interval $[a, b]$ that intersects A^c intersects B .

Claim 9. The set $\{(A, B) : B \subseteq (\text{int}(A))^c\}$ is Π_2^0 .

Indeed, $B \subseteq (\text{int}(A))^c$ iff every rational interval (a, b) that intersects B also intersects A^c .

Now it is c.e. The collection of pairs $(C, (\text{int}(C))^c)$ where C ranges over the finite unions of closed rational intervals is dense in it. \square

The Polish topology τ induces a notion of 1-genericity that fits with our objectives.

Proposition 4.7. *In the space $(\text{CL}([0, 1]), \tau)$, every 1-generic element is regular and its boundary contains only weakly 1-generic reals.*

Proof. First, the set of regular closed sets is a dense effective G_δ -set. Indeed, C is regular iff every rational interval (a, b) is disjoint from C or intersects the interior of C . The collection of finite unions of closed rational intervals (regular sets) is dense.

Let $U \subseteq [0, 1]$ be a dense effective open set. Let $\mathcal{U} \subseteq \text{CL}([0, 1])$ be the collection of closed sets whose boundary is contained in U . It is an effective open set in the topology τ : the boundary of C is contained in U iff $\text{int}(C) \cup C^c \cup U$ covers $[0, 1]$, which is semi-decidable from a description of C . \mathcal{U} is moreover dense. \square

Theorem 4.3. *Let P be a non-empty regular closed set that is τ_{reg} -generic from inside. The boundary of P contains only weakly-1-generic points.*

Proof. Let $U \subseteq [0, 1]$ be a dense effective open set. The class \mathcal{U} of regular closed sets whose boundary is contained in U is an effective open class in the topology τ_{reg} . Now let P be a regular closed set. There exists a sequence P_n of finite union of closed intervals contained in $\text{int}(P)$ and converging to P in the topology τ_{reg} . As U is dense, the endpoints of the intervals constituting P_n can be taken in U . Each P_n is contained in P (i.e. P_n belongs to S_P), and belongs to \mathcal{U} , so \mathcal{U} is dense below P . If P is τ_{reg} -generic from inside then P must belong to \mathcal{U} , i.e. its boundary must be contained in U . \square

Theorem 4.1 directly gives the following result.

Corollary 4.6. *In the space $(CL([0, 1]), \tau)$ there exists a regular Π_1^0 -class that is generic from inside and whose boundary is perfect.*

Proof. Having a perfect boundary is again a dense effective G_δ -property in the topology τ . \square

As a result, the leftmost element of this set is a left-c.e. real that is weakly-1-generic but not generic from the right.

Acknowledgements

The author wishes to thank Peter Gács, Emmanuel Jeandel and Cristóbal Rojas for discussions on the subject.

References

- [BG11] Vasco Brattka and Guido Gherardi. Weihrauch degrees, omniscience principles and weak computability. *J. Symb. Log.*, 76(1):143–176, 2011.
- [Bra99] Vasco Brattka. Computable invariance. *Theor. Comput. Sci.*, 210(1):3–20, 1999.
- [Bra01] Vasco Brattka. Computable versions of Baire’s category theorem. In Jiri Sgall, Ales Pultr, and Petr Kolman, editors, *MFCS*, pages 224–235, London, UK, 2001. Springer-Verlag.

- [BY06] M. Braverman and M. Yampolsky. Non-computable Julia sets. *Journ. Amer. Math. Soc.*, 19:551–578, 2006.
- [BY07] Mark Braverman and Michael Yampolsky. Constructing non-computable Julia sets. In *STOC*, pages 709–716. ACM, 2007.
- [BY08] Mark Braverman and Michael Yampolsky. *Computability of Julia Sets*. Springer, 2008.
- [DHL04] Rodney G. Downey, Denis R. Hirschfeldt, and Geoffrey LaForte. Randomness and reducibility. *J. Comput. Syst. Sci.*, 68(1):96–114, 2004.
- [DL02] Rodney G. Downey and Geoffrey LaForte. Presentations of computably enumerable reals. *Theor. Comput. Sci.*, 284(2):539–555, 2002.
- [Hoy11] Mathieu Hoyrup. Randomness and the ergodic decomposition. In Benedikt Löwe, Dag Normann, Ivan N. Soskov, and Alexandra A. Soskova, editors, *CiE*, volume 6735 of *Lecture Notes in Computer Science*, pages 122–131. Springer, 2011.
- [Hoy14] Mathieu Hoyrup. Irreversible computable functions. In Ernst W. Mayr and Natacha Portier, editors, *31st International Symposium on Theoretical Aspects of Computer Science (STACS 2014), STACS 2014, March 5-8, 2014, Lyon, France*, volume 25 of *LIPICs*, pages 362–373. Schloss Dagstuhl - Leibniz-Zentrum fuer Informatik, 2014.
- [Ing81] M.A. Ingrassia. *P-genericity for Recursively Enumerable Sets*. PhD thesis, University of Illinois at Urbana-Champaign, 1981.
- [Mil04] Joseph S. Miller. Degrees of unsolvability of continuous functions. *Journal of Symbolic Logic*, 69(2):555–584, 2004.
- [Myh61] J. Myhill. Category methods in recursion theory. *Pac. J. Math.*, 11:1479–1486, 1961.
- [Myh71] J. Myhill. A recursive function, defined on a compact interval and having a continuous derivative that is not recursive. *Michigan Math. J.*, 18(2):97–98, 1971.
- [Nie09] André Nies. *Computability and randomness*. Oxford logic guides. Oxford University Press, 2009.

- [Par61] K.R. Parthasarathy. On the category of ergodic measures. *Ill. J. Math.*, 5:648–656, 1961.
- [PER89] Marian B. Pour-El and J. Ian Richards. *Computability in Analysis and Physics*. Perspectives in Mathematical Logic. Springer, Berlin, 1989.
- [Rog87] Hartley Jr. Rogers. *Theory of Recursive Functions and Effective Computability*. MIT Press, Cambridge, MA, USA, 1987.
- [Sel71] Alan L. Selman. Arithmetical reducibilities I. *Mathematical Logic Quarterly*, 17(1):335–350, 1971.
- [SW05] Frank Stephan and Guohua Wu. Presentations of K-trivial reals and Ksolmogorov complexity. In *CiE*, pages 461–469, Berlin, Heidelberg, 2005. Springer-Verlag.
- [V’y97] Vladimir V. V’yugin. Effective convergence in probability and an ergodic theorem for individual random sequences. *SIAM Theory of Probability and Its Applications*, 42(1):39–50, 1997.
- [Wei92] Klaus Weihrauch. The TTE-interpretation of three hierarchies of omniscience principles. Informatik Berichte 130, FernUniversität Hagen, 1992.
- [Wei00] Klaus Weihrauch. *Computable Analysis*. Springer, Berlin, 2000.
- [YMT99] Mariko Yasugi, Takakazu Mori, and Yoshiki Tsujii. Effective properties of sets and functions in metric spaces with computability structure. *Theoretical Computer Science*, 219(1-2):467–486, 1999.