



HAL
open science

A methodological framework for drug development in rare diseases

Patrice Nony, Polina Kurbatova, Agathe Bajard, Salma Malik, Charlotte Castellan, Sylvie Chabaud, Vitaly Volpert, Nathalie Eymard, B. Kassai, Catherine Cornu

► To cite this version:

Patrice Nony, Polina Kurbatova, Agathe Bajard, Salma Malik, Charlotte Castellan, et al.. A methodological framework for drug development in rare diseases. *Orphanet Journal of Rare Diseases*, 2014, 9 (164), pp.1-10. 10.1186/s13023-014-0164-y . hal-01097224

HAL Id: hal-01097224

<https://inria.hal.science/hal-01097224>

Submitted on 20 May 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

THERAPIES special issue number 2 march april 2020

HEADING: DRUG AND ORPHAN DISEASES

A methodological framework for drug development in rare diseases.

The CRESim program: epilogue and perspectives

The CRESim program: methodological framework for drug development in rare diseases

Patrice Nony^{a,b,*}, Behrouz Kassai^{a,b,c}, Catherine Cornu^{b,c}, for the CRESim and Epi-CRESim research groups*

^a *Service hospitalo-universitaire de pharmacotoxicologie (SHUPT), hospices civils de Lyon, 69424 Lyon, France*

^b *Laboratoire de biométrie et biologie évolutive (LBBE), UMR 5558 CNRS, University Lyon 1, Lyon, 69376, France*

^c *EPICIME-CIC 1407 de Lyon, hospices civils de Lyon, Inserm, 69677 Bron, France*

Received September 16, 2019; accepted November 15, 2019

***Corresponding author.** Service hospital-universitaire de pharmacotoxicologie, hospices civils de Lyon, 69424 Lyon, France.

E-mail address: patrice.nony@univ-lyon1.fr

*This article has been written as part of the Child-Rare-Euro-Simulation (CRESim) and Epilepsy-CRESim (Epi-CRESim) projects.

Members of the CRESim and Epi-CRESim project groups: Leon Aarons, Corinne Alberti, Agathe Bajard, Pascal Benquet, Yves Bertrand, Frank Bretz, Daan Caudri, Charlotte Castellan, Sylvie Chabaud, Catherine Chiron, Catherine Cornu, Frank Dufour, Nathalie Eymard, Roland Fisch, Renzo Guerrini, Vincent Jullien, Behrouz Kassai, Polina Kurbatova, Salma Malik, Rima Nabbout,

Patrice Nony, Kayode Ogungbenro, David Pérol, Gérard Pons, Anna Rosati, Harm Tiddens, Fabrice Wendling.

Summary

Based on the ‘European Child-Rare-Euro-Simulation’ (CRESim) project, this article proposes a generalizable strategy utilizing datasets analysis in combination with modeling and simulation, in order to optimize the clinical drug development applied in the field of rare diseases. The global process includes the simulation of a realistic virtual population of patients (modeled from a real dataset of patients), the modeling of disease pathophysiological components and of pharmacokinetic-pharmacodynamic relations of the drug(s) of interest, the modeling of several randomized controlled clinical trials (RCTs) designs and the analysis of the results (multi-dimensional approach for RCTs durations and precision of the estimation of the treatment effect). However, whereas modeling and numerical simulation may provide supplementary tools for drug development, they cannot be considered as a substitute for RCTs performed in ‘real’ patients.

KEY WORDS

Integrative modelling; Numerical simulation; Randomized controlled clinical trials; Drug development; Rare diseases

Abbreviations

CRESim: Child-Rare-Euro-Simulation

FDA: Food and Drug Administration

IO: input-output (model)

M&S: modeling and simulation

ODE: ordinary differential equations

PDE: partial differential equations

PK/PD: pharmacokinetic/pharmacodynamic

PrioMed Child: priority medicines for children

RCT: randomized controlled trials

Introduction

Drug development is a long – often more than 10 years – and costly – more than 1 billion Euros each – machinery [1-3], requiring experiments in animals and humans. More recent alternative approaches, such as computer modeling and simulation (M&S) provide supplementary tools for developing drugs [4-8]. These M&S approaches, successfully applied in a variety of industries (aircraft and car manufacturing, nuclear power plants, etc.) will represent a major paradigm shift and factor of progress in a close future for drug development. EU Avicenna roadmap [9], and FDA recent position [10] are important signs of these coming changes.

Such an approach may be of particular interest in the field of rare diseases where conducting clinical trials may be very challenging and the level of available evidence from clinical studies remains insufficient. Recently, the CRESim consortium [11] tested whether *in silico* clinical trial simulation could help to choose the most appropriate design for drugs in three rare disease situations. The results of this M&S program have however to be further discussed and some additional perspectives can be proposed for future research in this field.

Specificities of orphan drug development

There is a need to facilitate the development and availability of high-quality, ethically sound, and appropriately authorized medicines in rare diseases, without subjecting patients (especially children) to unnecessary trials [12, 13]. Conducting adequately powered randomized controlled trials (RCTs) may not be possible because of the small number of potential participants. Small clinical trials have a low statistical power and also lack of generalizability. New approaches are needed for designing valid trials with small sample sizes that can assess the potential therapeutic effect of drugs, biological products, medical devices, and other medical interventions. Therefore, the aim of the CRESim program was to propose a generalizable strategy utilizing datasets analysis in combination with modeling and simulation, in order to optimize the clinical drug development in the field of rare diseases.

The CRESim program

The “European Child-Rare-Euro-Simulation” project (CRESim project no. 40-41800-98-008) was funded in 2011 by the European Union’s ERA-Net PrioMedChild (priority medicines for children). Briefly, the main objective was to investigate and compare RCT designs for drug evaluation in children with rare diseases using *in silico* experiments (i.e. RCTs modeling and simulation). For demonstration purposes, three diseases were studied: Dravet syndrome (myoclonic epilepsy in infancy), cystic fibrosis, and T-cell lymphoblastic lymphoma. Several RCTs designs have been modeled, i.e. the parallel-group, cross-over, randomized withdrawal, early-escape and N-of-1 trial designs. Various performance criteria were simulated for comparison purpose in terms of: 1) trial durations, this criterion represents the ‘burden’ of the trial for the patient, the investigator and the sponsor, and 2) precision of the estimation of the treatment effect, a criterion evaluating the “performance” of the trial. Eleven scientific articles referenced in PubMed [14-24] showed the feasibility and the interest of such an approach. The successive dimensions that were considered are described in Fig. 1.

Basic principles

For each disease of interest, up-to-date biological and biomedical knowledge must be synthesized through mathematical models to simulate 1) the function of one or several systems in human (and possibly in some animal species), their dysfunctions leading to one or several components of the disease, and 2) the impact of current (in the same indication or in repositioning, i.e. in a new indication -rare disease especially-) and new treatments on these dysfunctions [19, 22, 25, 26]. Such a process results from a close collaboration between relevant disciplines: applied mathematics, computational modeling, systems biology and pharmacology, pharmacokinetic and pharmacodynamic modeling, clinical pharmacology and medical specialities. Drug safety has also to be taken into account as far as possible, in order to assess a net clinical benefit [27].

Existing and new mathematical models are developed to capture variability of *in vivo*

parameters and system(s) functioning, in individuals, female and male of different ages (children, teenagers, young and mature adults, elderly). This development is possible through retrospective databases analyses and/or specifically devoted prospective cross-sectional studies of healthy volunteers and patients.

Using these mathematical models realistic virtual individuals and populations will be generated and will be randomised in virtual clinical trial simulations [28]. These populations are called “realistic” because of the similarities between these virtual individuals and real individuals, their ability to simulate and predict biomarkers evolution and/or the occurrence of some clinical endpoints in patients [29,30] and to reproduce the therapeutic impact of drugs.

Validation process of mathematical models remains a key issue and must involve several aspects: 1) Usual internal validation process for model building: the model is built with a set of available data, and validated using a separate set of data from the same database; 2) Confrontation of predicted and observed results of specific biomarkers in human cohorts of diseased patients; 3) Ability of the model to predict results from already performed clinical trials (if any) in terms of clinical events, in both treatment groups (placebo and tested drug); 4) possibly, ability of the model to explain discrepancies between animal and human experiments.

Proposal for specific points to consider

Databases

Depending on the existence of clinical databases, the corresponding strategy could be considered as follows:

- Using existing clinical databases (epidemiological and RCTs databases if any), their use is necessary for: 1) Identifying prognostic and predictive biomarkers: prognostic biomarkers are biomarkers which are linked with a specific favourable or unfavourable evolution of the disease, and predictive biomarkers are associated with the response of patients to specific drugs; the validation of a predictive biomarker requires specific statistical approaches [31, 32]; 2) Constructing realistic virtual population(s) of patients: sets of virtual patients are matrixes of

variables and covariates representing individuals; they are realistic when the interrelations between variables are controlled to realistically mimic patients (using parametric methods and/or resampling techniques and/or Bayesian networks); 3) Model validation: the validation process includes an internal validation (with the use of a training set), temporal validation [evaluation of the performance of a model on subsequent patients from the same centre(s)/dataset], an external validation [i.e. generalizability of the model by testing the performance on other centre(s)/dataset] and sensitivity analysis (the study of how variability in the output of a model can be attributed to different sources of variability in the model input [33]).

- If appropriate, generating new prospective cross-sectional studies involving healthy subjects and / or patients for: 1) Identification of numerical values of model parameters; 2) Assessment of intra and inter-individual parameter variability; 3) External model validation (replication of biomarkers results using specific individual patient characteristics).

Progressive and integrative model building

This step is a way to incorporate each new piece of information in a relevant edifice. Mathematical language enables prediction and establishing dynamic links between: 1) Inputs, such as drug dosage, route of administration, posology, treatment duration and patient compliance to the drug regimen [34–37]; 2) Individual characteristics that modulate the relationships as covariates; 3) Relationships between entities such as dose and concentration, concentration and effect on biomarkers/surrogate outcomes, impact of biomarkers on final clinical outcomes. These mathematical models are initially simple and become more and more sophisticated and accurate when incorporating more information. They become sometimes more complex, to take into account: 1) Specificities in drug distribution with several virtual compartments, or more numerous compartments mimicking the reality of different organs in which drug diffuses (physiology-based pharmacokinetic models) or 2) Several biomarkers/surrogates resulting in indirect and/or delayed effects, or a chain of causal mechanisms, or feedback loops mimicking counter-regulation mechanisms. They become also more and more fit to individual profiles, in order to capture the different characteristics that explain the between individual variability.

Modeling steps

The modeling process consists in:

- Construction of specific (sub)models for: 1) Disease(s) characteristics/spontaneous evolution using statistical and/or mechanistic models [most often ordinary differential equations (ODE) and/or partial differential equations (PDE) systems]; 2) Drug(s) effect(s) [38] using pharmacokinetic-pharmacodynamic (PK-PD) models [39, 40, 41]; 3) RCT characteristics (experimental designs [42, 43, 44]) and their corresponding statistical methods of analysis (e.g. frequentist versus Bayesian approaches [45, 46], parametric versus non parametric approaches, meta-analysis).
- Prospective validation of the disease and drug-effect models (model verification and validation, comparison between predictions with observations, sensitivity analyses).
- Connection of submodels together into a meta-model for simulation purposes.

RCTs modeling

This dimension involves:

- RCT simulation using: 1) An input-output (IO) submodel; 2) A covariate distribution submodel; 3) An execution submodel; 4) A specific statistical analysis according to each considered design; 5) An additional exploratory search for predictive biomarkers.
- When required, (e.g. in ultra-rare diseases), simulations of cohorts of virtual patients can be performed for: 1) Formal external indirect comparisons approaches; 2) Investigation of new prognostic biomarkers of potential interest; 3) Construction of prognostic scores for model calibration purposes.

Simulation strategies

This results from exposing realistic virtual populations to a variety of simulated clinical trials with a considered drug. The drug impact is directly observed from the comparison of the disease evaluation in the same population simulated after drug or placebo exposure. A large number of scenarios regarding RCT designs, drug dosages, drug associations, selection of patients, length of exposure, etc. can be explored through this approach. The main steps are as follows:

- Samples of patients randomly drawn from the ‘validated’ virtual population for inclusion in *in silico* RCTs.
- Applying a pre-specified simulation strategy to optimize and limit the number of simulations.
- Use of the same specific statistical analysis for each simulated RCT results.

Simulation results analysis

A multidimensional analysis must be finally performed to put into perspective the simulation results. At the end of the simulation process, an example of the expression of results is given below:

- the median of the estimations of the treatment effect and the medians of the lower/upper limits of their 95% CI;
- an evaluation of the precision of the estimation of the treatment effect, e.g. the coefficient of variation (corresponding to the ratio between the standard deviation of treatment effect and the corresponding median).

The corresponding graphical representation may use for example the radar plot approach. Each axis of radar plot represents an evaluation of one component of interest. The furthest value from the center corresponds to the ‘ideal’ value for clinical trial, e.g. a high power (%), a low variability (%), a short patient duration (days), short investigator and sponsor durations (years), and a high number of patients under active treatment.

Other possible applications of simulation

Their main interest is to compare results from real RCTs with the results from simulated similar RCTs that would have been designed if the mathematical models were available and used. In addition, case studies can specifically assess the potential of the innovative approach to achieve better level of evidence at lesser costs, to reduce the number of human (and animal) test-subjects, e.g. through optimising experimental design, or explaining and preventing failures in drug development.

Comprehensive overview of the overall process

The global process may be summarized in Fig. 2 that includes the simulation of a virtual population of patients (modeled from a real dataset of patients), the modeling of disease components and PK-PD of the drug of interest, the modeling of several RCTs designs and the analysis of the results (multi-dimensional approach for RCTs durations and precision of the estimation of the treatment effect).

Limitations

Some possible barriers have to be mentioned and further taken into account:

- conservatism of pharmaceutical industry, regulatory bodies, pharmacologists, methodologists and physicians;
- need to involve different disciplines that do not work usually together;

- time consuming nature of the process (results are not ‘handy’ and their integration into the development plan is still not well defined);
- the cost-effectiveness of the modeling approach should be demonstrated;
- the regulatory framework has to be better defined.

Perspectives

Recent advances in *data sharing* and in clinical trial methodology offer new perspective to clinical trial simulation. The development of databases which are either open access or on demand access facilitates formal indirect external comparisons for prediction issued by modeling, and building datasets of virtual patients who become more and more realistic.

Similarly, *biomarkers* previously identified using bioinformatic approaches could be first validated in silico.

The same approach could allow the in silico evaluation of the interest of coupling therapeutic and diagnostic (*theranostic*) strategies.

Extensive use of databases could allow the application of specific *enrichment strategies*, i.e. the selection of patients whose characteristics are likely to optimize the chances of success of a trial. This strategy could also be applied to the field of rare diseases whenever possible [47, 48].

The application of modeling and simulation in the field of *4P medicine* (predictive, personalized, preventive and participatory) offers the perspective for each new patient to have the results of a given trial reanalyzed according to his/her own characteristics / preferences (e.g. interest of use of generalized multiple pairwise comparisons for trial re-analysis [49, 50]).

Academic and industrial partners should have free access to dedicated *RCT modeling and simulation platforms*.

Conclusion

M&S will speed the availability of well documented drugs on the market, positioning adequately new drugs. It has the potential to reduce costs of drug development, including the number of needed individuals to be recruited in clinical trials, by avoiding useless trials and optimizing precision through trial designs [51–54].

All stakeholders (regulators, academics, firms, patients associations) should be involved as soon as possible in each new research program to allow their expectations to be taken into account, and in return to be informed on the program advances and for regulators to prepare guidelines.

Lastly, whereas M&S results obtained from simulated RCTs performed in ‘virtual’ patients can improve the methodological approach of drug evaluation in the field of rare diseases (particularly in children), they cannot be considered as substitute for RCTs performed in ‘real’ patients.

Disclosure of interest

Authors have no competing interest to declare.

References

- [1] Morgan S, Grootendorst P, Lexchin J, Cunningham C, Greyson D. The cost of drug development: A systematic review. *Health Policy (New York)* 2011;100(1):4-17.
- [2] DiMasi JA, Grabowski HG, Hansen RW. Innovation in the pharmaceutical industry: New estimates of R&D costs. *J Health Econ* 2016;47:20-33.
- [3] Assistant Secretary of Planning and Evaluation (ASPE). Examination of clinical trial costs and barriers for drug development | ASPE. July 2014. <https://aspe.hhs.gov/report/examination-clinical-trial-costs-and-barriers-drug-development>. [Accessed December 20, 2019].
- [4] Boissel JP, Ribba B, Grenier E, Chapuisat G, Dronne MA. Modelling methodology in physiopathology. *Prog Biophys Mol Biol* 2008;97(1):28-39.
- [5] Holford N, Ma SC, Ploeger BA. Clinical trial simulation: a review. *Clin Pharmacol Ther* 2010;88(2):166-82.
- [6] Medicines Agency E. Human medicines development and evaluation EFPIA-EMA modelling and simulation workshop report EMA-EFPIA modelling and simulation workshop. 2019. <http://www.ema.europa.eu>. [Accessed December 20, 2019].
- [7] Food and Drug Administration. FDA guidances. 2019. <http://www.fda.gov/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/default.htm>. [Accessed December 20, 2019].
- [8] Dorsey ER, Venuto C, Venkataraman V, Harris DA, Kieburz K. Novel Methods and technologies for 21st-Century clinical trials. *JAMA Neurol* 2015;72(5):582.
- [9] Avicenna Roadmap | Avicenna. <https://avicenna-alliance.com/about-us/avicenna-roadmap/>. 2019. [Accessed December 20, 2019].
- [10] Food and Drug Administration. Reporting of computational modeling studies in medical device submissions. September 2016. <https://www.fda.gov/regulatory-information/search-fda-guidance-documents/reporting-computational-modeling-studies-medical-device-submissions>. [Accessed December 20, 2019].

- [11] UMR CNRS 5558 Laboratoire de biométrie et biologie évolutive - CRESIM. 2019. <https://lbbe.univ-lyon1.fr/-CRESIM-.html>. [Accessed December 20, 2019].
- [12] Ahmed MA, Okour M, Brundage R, Kartha R V. Orphan drug development: the increasing role of clinical pharmacology. *J Pharmacokinet Pharmacodyn* July 2019. 2019 Oct;46(5):395-409.
- [13] van Hasselt JGC, van Eijkelenburg NKA, Beijnen JH, Schellens JHM, Huitema ADR. Optimizing drug development of anti-cancer drugs in children using modelling and simulation. *Br J Clin Pharmacol* 2013;76(1):30-47.
- [14] Eymard N, Volpert V, Kurbatova P, Volpert V, Bessonov N, Ogungbenro K, et al. Mathematical model of T-cell lymphoblastic lymphoma: disease, treatment, cure or relapse of a virtual cohort of patients. *Math Med Biol* 2018;35(1):25-47.
- [15] Cornu C, Kassai B, Fisch R, Chiron C, Alberti C, Guerrini R, et al. Experimental designs for small randomised clinical trials: an algorithm for choice. *Orphanet J Rare Dis* 2013;8(1):48.
- [16] Nony P, Kurbatova P, Bajard A, Malik S, Castellan C, Chabaud S, et al. A methodological framework for drug development in rare diseases. *Orphanet J Rare Dis* 2014;9(1):164.
- [17] Ogungbenro K, Aarons L, CRESim & Epi-CRESim Project Groups. Physiologically based pharmacokinetic modelling of methotrexate and 6-mercaptopurine in adults and children. Part 2: 6-mercaptopurine and its interaction with methotrexate. *J Pharmacokinet Pharmacodyn* 2014;41(2):173-85.
- [18] Ogungbenro K, Aarons L. Physiologically based pharmacokinetic modelling of methotrexate and 6-mercaptopurine in adults and children. Part 1: methotrexate. *J Pharmacokinet Pharmacodyn* 2014;41(2):159-71.
- [19] Kurbatova P, Bessonov N, Volpert V, Tiddens HA, Cornu C, Nony P, et al. Model of mucociliary clearance in cystic fibrosis lungs. *J Theor Biol* 2015;372:81-8.
- [20] Bajard A, Chabaud S, Cornu C, Castellan C, Malik S, Kurbatova P, et al. An in silico approach helped to identify the best experimental design, population, and outcome for future randomized clinical trials. *J Clin Epidemiol* 2016;69:125-36.
- [21] Ogungbenro K, Aarons L, CRESim & Epi-CRESim Project Groups. Physiologically based pharmacokinetic model for 6-mercaptopurine: exploring the role of genetic polymorphism in TPMT enzyme activity. *Br J Clin Pharmacol* 2015;80(1):86-100.


- [22] Kurbatova P, Wendling F, Kaminska A, Rosati A, Nabbout R, Guerrini R, et al. Dynamic changes of depolarizing GABA in a computational model of epileptogenic brain: Insight for Dravet syndrome. *Exp Neurol* 2016;283(Pt A):57-72.
- [23] Ogungbenro K, Aarons L, CRESim & Epi-CRESim Project Groups. A physiologically based pharmacokinetic model for valproic acid in adults and children. *Eur J Pharm Sci* 2014;63:45-52.
- [24] Ogungbenro K, Aarons L, CRESim & Epi-CRESim Project Groups. A physiologically based pharmacokinetic model for clobazam and stiripentol in adults and children. *Pharm Res* 2015;32(1):144-57.
- [25] Eymard N, Volpert V, Quere I, Lajoinie A, Nony P, Cornu C. A 2D computational model of lymphedema and of its management with compression device. Bocharov G, Simakov S, Vassilevski Y, Volpert V, eds. *Math Model Nat Phenom* 2017;12(5):180-95.
- [26] Tallon J, Couenne F, Bordes C, Tayakout-Fayolle M. Immune system modelling in case of a septic shock. *Comput Aided Chem Eng* 2019;46:1093-8.
- [27] Gal J, Milano G, Ferrero JM, Saâda-Bouزيد E, Viotti J, Chabaud S, et al. Optimizing drug development in oncology by clinical trial simulation: Why and how? *Brief Bioinform* 2018;19(6):1203-17.
- [28] Teutonico D, Musuamba F, Maas HJ, Acius A, Yang S, Danhof M, et al. Generating virtual patients by multivariate and discrete re-sampling techniques. *Pharm Res* 2015;32(10):3228-37.
- [29] Marchant I, Boissel JP, Nony P, Gueyffier F. High risk versus proportional benefit: modelling equitable strategies in cardiovascular prevention. *PLoS One*. 2015;10(11):e0140793.
- [30] Marchant I, Nony P, Cucherat M, Boissel JP, Thomas SR, Bejan-angoulvant T, et al. The global risk approach should be better applied in French hypertensive patients: a comparison between simulation and observation studies. *PLoS One* 2011;6(3):e17508.
- [31] Gerds TA, Cai T, Schumacher M. The performance of risk prediction models. *Biom J* 2008;50(4):457-479.
- [32] Gail M, Simon R. Testing for qualitative interactions between treatment effects and patient subsets. *Biometrics* 1985;41(2):361-72.

- [33] Altman DG, Vergouwe Y, Royston P, Moons KGM. Prognosis and prognostic research: validating a prognostic model. *BMJ* 2009;338(may28 1):b605-b605.
- [34] Nony P, Boissel JP. Use of sensitivity functions to characterise and compare the forgiveness of drugs. *Clin Pharmacokinet* 2002;41(5):371-80.
- [35] Nony P, Cucherat M, Boissel JP. Revisiting the effect compartment through timing errors in drug administration. *Trends Pharmacol Sci* 1998;19(2):49-54.
- [36] Boissel JP, Nony P. Using pharmacokinetic- pharmacodynamic relationships to predict the effect of poor compliance. *Clin Pharmacokinet* 2002;41(1):1-6.
- [37] Blesius A, Chabaud S, Cucherat M, Mismetti P, Boissel JP, Nony P. Compliance-guided therapy. *Clin Pharmacokinet* 2006;45(1):95-104.
- [38] Wang H, Boissel JP, Nony P. Revisiting the relationship between baseline risk and risk under treatment. *Emerg Themes Epidemiol* 2009;6(1):1.
- [39] Duffull SB, Chabaud S, Nony P, Laveille C, Girard P, Aarons L. A pharmacokinetic simulation model for ivabradine in healthy volunteers. *Eur J Pharm Sci* 2000;10(4):285-94.
- [40] Chabaud S, Girard P, Nony P, Boissel JP, HERapeutic MOdeling and Simulation Group. Clinical trial simulation using therapeutic effect modeling: application to ivabradine efficacy in patients with angina pectoris. *J Pharmacokinet Pharmacodyn* 2002;29(4):339-63.
- [41] Colquitt RB, Colquhoun DA, Thiele RH. In silico modelling of physiologic systems. *Best Pract Res Clin Anaesthesiol* 2011;25(4):499-510.
- [42] Bajard A, Chabaud S, Pérol D, Boissel JP, Nony P. Revisiting the level of evidence in randomized controlled clinical trials: A simulation approach. *Contemp Clin Trials* 2009;30(5):400-10.
- [43] Abbas I, Rovira J, Cobo E, Casanovas J. Simulation models for optimizing the design of clinical trials. *Qual Reliab Eng Int* 2006;22(6):683-91.
- [44] Abbas I, Rovira J, Casanovas J. Clinical trial optimization: Monte Carlo simulation Markov model for planning clinical trials recruitment. *Contemp Clin Trials* 2007;28(3):220-31.
- [45] CDRH. Guidance for industry and FDA staff guidance for the use of Bayesian statistics in medical device clinical trials preface public comment. 2010. <http://www.regulations.gov>. [Accessed December 20, 2019].

- [46] Kadane JB, ed. Bayesian methods and ethics in a clinical trial design. Hoboken, NJ, USA: John Wiley & Sons, Inc.; 1996. ISBN 0-471-84680-5.
- [47] FDA, CDER, CBER. Enrichment strategies for clinical trials to support determination of effectiveness of human drugs and biological products guidance for industry. 2019. <https://www.fda.gov/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/default.htm> and/or <https://www.fda.gov/BiologicsBloodVaccines/GuidanceComplianceRegulatoryInformation/Guidances/default.htm>. [Accessed December 20, 2019 (45 pp.)].
- [48] Janes H, Brown MD, Huang Y, Pepe MS. An approach to evaluating and comparing biomarkers for patient treatment selection. *Int J Biostat* 2014;10(1):99-121.
- [49] Vickers AJ, Kattan MW, Sargent DJ. Method for evaluating prediction models that apply the results of randomized trials to individual patients. *Trials* 2007;8(1):14.
- [50] Buyse M. Generalized pairwise comparisons of prioritized outcomes in the two-sample problem. *Stat Med* 2010;29(30):3245-57.
- [51] Claxton K, Posnett J. An economic approach to clinical trial design and research priority-setting. *Health Econ* 1996;5(6):513-24.
- [52] Chilcott J, Brennan A, Booth A, Karnon J, Tappenden P. The role of modelling in prioritising and planning clinical trials. *Health Technol Assess* 2003;7(23):iii, 1-125.
- [53] Eckermann S, Willan AR. Optimal global value of information trials: better aligning manufacturer and decision maker interests and enabling feasible risk sharing. *Pharmacoeconomics* 2013;31(5):393-401.
- [54] DiMasi JA. The value of improving the productivity of the drug development process: faster times and better decisions. *Pharmacoeconomics* 2002;20 Suppl 3(Supplement 3):1-10.

Figure 1. Modeling steps for RCT simulation.

Construction of a virtual population of patients from epidemiologic databases, modeling disease prognosis and therapeutic effect from pathophysiology and PK-PD relations, RCTs simulations using different experimental designs, comparison of design performances




PK/PD: pharmacokinetic/pharmacodynamic; RCT: randomized controlled trials

Adapted from [16] Nony P, Kurbatova P, Bajard A, Malik S, Castellan C, Chabaud S, et al. A methodological framework for drug development in rare diseases. *Orphanet J Rare Dis* 2014;9(1):164. With courtesy Springer Nature (M Mohammad Suleik).

Figure 2. Flow diagram of the overall process.

Using data of a real population of patients (1), a virtual patient dataset is built (2) and connected with a disease pathophysiological model (3a) and PK-PD models of drug(s) of interest (3b). Samples of virtual patients are randomized in virtual RCTs with different experimental designs (4). Simulations results are compared using a multi-dimensional approach (5).



PK/PD: pharmacokinetic/pharmacodynamic; RCT: randomized controlled trials