



HAL
open science

Detectability and state estimation for linear age-structured population diffusion models

Karim Ramdani, Marius Tucsnak, Julie Valein

► **To cite this version:**

Karim Ramdani, Marius Tucsnak, Julie Valein. Detectability and state estimation for linear age-structured population diffusion models. 2015. hal-01140166v1

HAL Id: hal-01140166

<https://inria.hal.science/hal-01140166v1>

Preprint submitted on 9 Jun 2015 (v1), last revised 21 Oct 2016 (v4)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Detectability and state estimation for linear age-structured population diffusion models

Karim Ramdani*

Marius Tucsnak^{†‡}

Julie Valein^{†‡}

Abstract

We investigate a state estimation problem for an infinite dimensional system appearing in population dynamics. More precisely, given a linear model for age-structured populations with spatial diffusion, we assume the initial distribution to be unknown and that we have at our disposal an observation locally distributed in both age and space. Using Luenberger observers, we solve the inverse problem of recovering asymptotically in time the distribution of population. The observer is designed using a finite dimensional stabilizing output injection operator, yielding an effective reconstruction method. Numerical experiments are provided showing the feasibility of the proposed reconstruction method.

Keywords Inverse problems; Observers; Stabilization; Population Dynamics; Spatial Diffusion

1 Introduction and problem setting

We consider a linear age-structured population model with spatial diffusion described by the following system:

$$\left\{ \begin{array}{ll} \partial_t p(a, x, t) + \partial_a p(a, x, t), & a \in (0, a^*), x \in \Omega, t > 0, \\ = -\mu(a)p(a, x, t) + k\Delta p(a, x, t), & \\ p(a, x, t) = 0, & a \in (0, a^*), x \in \partial\Omega, t > 0, \\ p(a, x, 0) = p_0(a, x), & a \in (0, a^*), x \in \Omega, \\ p(0, x, t) = \int_0^{a^*} \beta(a)p(a, x, t) da, & x \in \Omega, t > 0. \end{array} \right. \quad (1.1)$$

In the above equations:

- $\Omega \subset \mathbb{R}^n$, $n \geq 1$, denotes a smooth bounded domain, k is a positive constant diffusion coefficient and Δ the laplacian with respect to the space variable x .

*Inria, Villers-lès-Nancy, F-54600, France.

[†]Université de Lorraine, Institut Élie Cartan de Lorraine, UMR 7502, Vandœuvre-lès-Nancy, F-54506, France.

[‡]CNRS, Institut Élie Cartan de Lorraine, UMR 7502, Vandœuvre-lès-Nancy, F-54506, France.

- $p(a, x, t)$ denotes the distribution density of the population of age a at spatial position $x \in \Omega$ at time t ;
- p_0 denotes the initial distribution;
- $a^* \in (0, +\infty)$ is the maximal life expectancy;
- $\beta(a)$ and $\mu(a)$ are positive functions denoting respectively the birth and death rates, which are supposed to be independent of x (see Figure 5 for typical graphs of these functions).

The last equation in (1.1) describing the birth process is the so-called *renewal equation*. We assume here homogeneous Dirichlet boundary conditions (in space) which model a hostile habitat at the boundary $\partial\Omega$, but other type of boundary conditions (in space) can also be considered (typically Neumann homogeneous boundary conditions which correspond to isolated population, i.e. vanishing incoming and outgoing flux).




Figure 1: How to estimate $p(T)$ as $T \rightarrow +\infty$ from the knowledge of $y(t)$ for $t \in (0, T)$?

In this paper, we investigate the following inverse problem (see Figure 1):

Assuming the initial age distribution p_0 to be unknown, but knowing the age distribution

$$y(a, x, t) := p(a, x, t), \quad t \in (0, T), \quad a \in (a_1, a_2), \quad x \in \mathcal{O},$$

where \mathcal{O} is some given subset of Ω and $0 \leq a_1 < a_2 \leq a^$, is it possible, and if so how, to estimate when $T \rightarrow +\infty$ the age distribution $p(a, x, T)$, for $x \in \Omega$ and $a \in (0, a^*)$?*

We provide a positive answer to this question and we propose an effective reconstruction algorithm. Based on a semigroup formulation, our method crucially uses two main ingredients: the Luenberger observers [44] and the fact that the problem has finitely many unstable eigenvalues (i.e. eigenvalues with positive real part). Let us make this statement more precise by outlining in a formal way the main ideas of our reconstruction method.

Using a semigroup formulation, we first rewrite problem (1.1) in the abstract form (throughout the paper, the dot denotes the derivative with respect to time)

$$\begin{cases} \dot{p}(t) = Ap(t), & t \in (0, T), \\ p(0) = p_0, \end{cases}$$

where $A : \mathcal{D}(A) \rightarrow X$ is the generator of a C_0 -semigroup on a Hilbert space X . Similarly, the available observation can also be reformulated using a bounded observation operator $C \in \mathcal{L}(X, Y)$, where Y is another Hilbert space:

$$y(t) = Cp(t), \quad t \in (0, T).$$

We define the Luenberger type observer as the solution \hat{p} of the following system

$$\begin{cases} \dot{\hat{p}}(t) = A\hat{p}(t) + L(C\hat{p}(t) - y(t)), & t \in (0, T), \\ \hat{p}(0) = 0, \end{cases} \quad (1.2)$$

where $L \in \mathcal{L}(Y, X)$ is a linear operator to be chosen in such way that $A + LC$ generates an exponentially stable semigroup on X . Indeed, since the error $e := \hat{p} - p$ satisfies by construction

$$e(t) = (A + LC)e(t), \quad t \in (0, T),$$

this condition ensures that $e(T)$ converges exponentially to 0, i.e. that ($\|\cdot\|$ denotes the norm on X)

$$\|\hat{p}(T) - p(T)\| \leq Me^{-\omega T} \|p_0\|,$$

showing that $\hat{p}(T)$ constitutes an approximation of $p(T)$ as $T \rightarrow +\infty$. Let us emphasize that $\hat{p}(T)$ can be computed (by solving (1.2)) exclusively from the knowledge of the observation $y(t)$, provided we have determined a stabilizing output injection operator L .

To compute this operator, we use the second main ingredient of our method: the fact that A has finitely many unstable eigenvalues. Following Triggiani [59, 60], we will show that the operator L can be easily deduced from the finite dimensional operator L^+ stabilizing the finite dimensional system (A^+, C^+) , where A^+ is the restriction of A to the subspace X^+ of X spanned by the unstable (generalized) eigenfunctions of A and C^+ is the corresponding observation operator.

For an overview on population dynamics models, we refer the reader to the monograph of Webb [63] and to the introductory paper [64]. Gurtin [27] introduced the first model with spatial diffusion (see also Gurtin and MacCamy [28]). For linear models, existence of a semigroup in L^2 and large time asymptotics have been developed by Chan and Guo in the case of homogeneous Dirichlet boundary (in [13] for age dependent birth and death rates and in [26] for age and space dependent birth and death rates) and by Huyer [33, 34] in the case of homogeneous Neumann boundary conditions using perturbation techniques. The L^1 setting has been investigated by Rhandi [53] for a linear age-dependent model with spatial diffusion involving age and space dependent birth and death rates. More recently, Walker [62] proposed an abstract functional analytic framework to study the large time behavior for an age-structured linear diffusive model. Existence and large time behavior for non linear models have been studied by Langlais [41, 42]. Exact and approximate controllability issues for age-structured dynamic systems have been studied in Ainseba and Langlais [3], Ainseba and Iannelli [1, 2], Traore [57], Traore and Kavian [37].

The literature on inverse problems for population dynamics models is less rich, especially when spatial diffusion is assumed. Without spatial diffusion, Pilant and Rundel [50], Rundell [54], Engl, Rundell and Scherzer [23] studied the determination of the death rate, Gyllenberg, Osipov and Päivärinta [29] investigated uniqueness for the reconstruction of birth and death rates. State estimation problems, like the one considered here, have been considered only recently. Di Blasio and Lorenzi [21, 22] studied the well-posedness of

the inverse problem of reconstructing the initial state for a linear age-dependent model. Traore [58] considered a data assimilation problem similar to the one studied in this work, in which the state has to be recovered from locally distributed observation in space and full observation in age. However, he used a completely different approach, based on a null controllability result.

Unless otherwise stated, we will make the following classical assumptions on β and μ :

- **(H $_{\beta}$)** $\beta \in L^{\infty}(0, a^*)$, $\beta \geq 0$ a.e. in $(0, a^*)$;
- **(H $_{\mu}$)** $\mu \in L^1_{\text{loc}}(0, a^*)$, $\mu \geq 0$ a.e. in $(0, a^*)$ and

$$\lim_{a \rightarrow a^*} \int_0^{a^*} \mu(s) ds = +\infty. \quad (1.3)$$

We also introduce the function

$$\Pi(a) := \exp\left(-\int_0^{a^*} \mu(s) ds\right)$$

which represents the probability to survive at age a (also known as the life table function). Hence, it is a decreasing function satisfying (thanks to condition (1.3))

$$\lim_{a \rightarrow a^*} \Pi(a) = 0. \quad (1.4)$$

Section 2 gathers some well-known facts about the population semigroup associated to (1.1) and the spectral properties of its generator A . In particular, we will see that A has finitely many unstable eigenvalues. In Section 3, we provide some abstract results about the detectability (see Definition 3.1) of infinite dimensional linear systems (A, C) where $A : \mathcal{D}(A) \rightarrow X$ is the generator of a C_0 -semigroup on a Hilbert space X having finitely many unstable eigenvalues, and $C \in \mathcal{L}(X, Y)$ is a bounded observation operator. In particular, we show the existence of a finite dimensional stabilizing output injection operator for such systems, provided (A, C) satisfy a Hautus test (this follows from combining Theorem 3.4 and Proposition 3.5). In Section 4, we show that problem (1.1) fits into the general framework described in Section 3. This allows us to design a Luenberger type observer for (1.1) providing a solution to our estimation inverse problem. The last two sections of the paper are devoted to the implementation of our reconstruction method. In Section 5 we present finite difference full discretizations of the open loop system (used for synthetic data generation) and the observer. In Section 6, we provide numerical experiments illustrating the theoretical results and showing the efficiency of the reconstruction method.

2 Some background on the population semigroup

We collect below some of the existing results on the population semigroup for the linear age-structured model without (§. 2.1) and with (§. 2.2) spatial diffusion. In particular, we will recall the spectrum's structure of the semigroup generators in these two cases.

2.1 The diffusion free population dynamics

The diffusion free case is described by the so-called McKendrick–Von Foerster model (see [56] for the proofs of the results given below):

$$\begin{cases} \partial_t p(a, t) + \partial_a p(a, t) = -\mu(a)p(a, t), & a \in (0, a^*), t > 0, \\ p(a, 0) = p_0(a), & a \in (0, a^*), \\ p(0, t) = \int_0^{a^*} \beta(a)p(a, t) da, & t > 0. \end{cases}$$

The population operator A_0 corresponding to the above system is defined as follows

$$\begin{aligned} \mathcal{D}(A_0) &= \left\{ \varphi \in H^1(0, a^*) \mid \varphi(0) = \int_0^{a^*} \beta(a)\varphi(a) da; -\frac{d\varphi}{da} - \mu\varphi \in L^2(0, a^*) \right\} \\ A_0\varphi &= -\frac{d\varphi}{da} - \mu\varphi, \quad \forall \varphi \in \mathcal{D}(A_0). \end{aligned} \quad (2.1)$$

Theorem 2.1. *The operator A_0 defined by (2.1) has compact resolvent and its spectrum is constituted of a countable (infinite) set of isolated eigenvalues with finite algebraic multiplicity. The eigenvalues $(\lambda_n^0)_{n \geq 1}$ of A_0 (counted without multiplicity) are the solutions of the characteristic equation*

$$F(\lambda) := \int_0^{a^*} \beta(a)e^{-\lambda a} \Pi(a) da = 1. \quad (2.2)$$

The eigenvalues $(\lambda_n^0)_{n \geq 1}$ are of geometric multiplicity one, the eigenspace associated to λ_n^0 being the one-dimensional subspace of $L^2(0, a^)$ generated by the function*

$$\varphi_n^0(a) = e^{-\lambda_n^0 a} \Pi(a) = e^{-\lambda_n^0 a - \int_0^a \mu(s) ds}.$$

Finally, every vertical strip of the complex plane $\alpha_1 \leq \operatorname{Re}(z) \leq \alpha_2$, $\alpha_1, \alpha_2 \in \mathbb{R}$, contains a finite number of eigenvalues of A_0 .

Theorem 2.2. *The operator A_0 defined by (2.1) has a unique real eigenvalue λ_1^0 . Moreover, we have the following properties*

1. λ_1^0 is of algebraic multiplicity one;
2. $\lambda_1^0 > 0$ (reps. $\lambda_1^0 < 0$) if and only if $F(0) > 1$ (resp. $F(0) < 1$);
3. λ_1^0 is a real dominant eigenvalue:

$$\lambda_1^0 > \operatorname{Re}(\lambda_n^0), \quad \forall n \geq 2.$$

2.2 The population dynamics with diffusion

Chan and Guo [13] showed the existence of a semigroup on $L^2((0, a^*) \times \Omega)$ for linear age-structured population model with (constant) diffusion coefficient and age dependent birth and death rates in [13]. They extended this result to the of age and space dependent birth

and death rates. For reader's convenience, we sketch their proof here in this last case (which is more general than the one under consideration in this paper).

Let $X := L^2((0, a^*) \times \Omega)$ and define the following unbounded operator A on X :

$$\mathcal{D}(A) = \left\{ \varphi \in X \mid \begin{aligned} &\varphi \in C([0, a^*]; L^2(\Omega)) \cap L^2([0, a^*]; H_0^1(\Omega)); \\ &\beta\varphi \in L^1([0, a^*]; L^2(\Omega)); \quad -\partial_a\varphi - \mu\varphi + k\Delta\varphi \in X; \\ &\varphi(0, x) = \int_0^{a^*} \beta(a, x)\varphi(a, x) da \text{ for almost all } x \in \Omega \end{aligned} \right\} \quad (2.3)$$

$$A\varphi = -\partial_a\varphi - \mu\varphi + k\Delta\varphi, \quad \forall \varphi \in \mathcal{D}(A). \quad (2.4)$$

Assume that β and μ satisfy the following assumptions

- $\beta \in L^\infty([0, a^*] \times \Omega)$, $\beta \geq 0$ a.e. in $(0, a^*) \times \Omega$ and

$$\text{meas} \{a \in [0, a^*] \mid \inf_{x \in \overline{\Omega}} \beta(a, x) > 0\} > 0;$$

- μ is measurable and non negative on $[0, a] \times \overline{\Omega}$ for every $a \in (0, a^*)$;
- the function $a_0 \mapsto \sup_{a \in [0, a_0]} \int_{\Omega} \mu^2(a, x) dx$ is continuous with respect to $a \in [0, a^*)$;
- the functions $\underline{\mu}(s) = \inf_{x \in \overline{\Omega}} \mu(s, x)$ and $\overline{\mu}(s) = \sup_{x \in \overline{\Omega}} \mu(s, x)$ satisfy

$$\lim_{a \rightarrow a^*} \int_0^a \underline{\mu}(s) ds = +\infty, \quad \overline{\mu} \in L^1[0, a], \quad a \in [0, a^*].$$

Proposition 2.3. *A generates a C^0 semigroup on X .*

Proof. We just check that $\lambda I - A$ is onto for λ large enough. First, let us recall that from standard regularity for parabolic equations, given $s \geq 0$ and $\psi \in L^2(\Omega)$, the initial and boundary value problem of unknown $v(a, x)$:

$$\begin{cases} \partial_a v(a, x) = k\Delta v(a, x) - \mu(a, x)v(a, x), & s \leq a < a^*, \quad x \in \Omega, \\ v(a, x) = 0, & s \leq a < a^*, \quad x \in \partial\Omega, \\ v(s, x) = \psi(x), & x \in \Omega, \end{cases}$$

admits a unique solution

$$v \in C([s, a^*]; L^2(\Omega)) \cap L^2([s, a^*]; H_0^1(\Omega)).$$

In the sequel, we denote the solution operator by $\mathcal{F}(a, s)$, in the sense that

$$v(a, x) = \mathcal{F}(a, s)\psi(x), \quad 0 \leq s \leq a \leq a^*, \quad x \in \Omega.$$

Now, given $\lambda \in \mathbb{R}$, $f \in X$ and $\psi \in L^2(\Omega)$, the initial and boundary value problem

$$\begin{cases} \partial_a v(a, x) = k\Delta v(a, x) - (\mu(a, x) + \lambda)v(a, x) + f(a, x), & 0 \leq a < a^*, \quad x \in \Omega, \\ v(a, x) = 0, & 0 \leq a < a^*, \quad x \in \partial\Omega, \\ v(0, x) = \psi(x), & x \in \Omega, \end{cases}$$

admits a unique solution

$$v \in C\left([0, a^*]; L^2(\Omega)\right) \cap L^2\left([0, a^*]; H_0^1(\Omega)\right),$$

given by

$$v(a, \cdot) = e^{-\lambda a} \mathcal{F}(a, 0)\psi + \int_0^a e^{-\lambda(a-s)} \mathcal{F}(a, s) f(s, \cdot) ds \quad (0 \leq a < a^*). \quad (2.5)$$

It follows that

$$\begin{aligned} v(0, x) - \int_0^{a^*} \beta(a, x) v(a, x) da &= \psi(x) - \int_0^{a^*} e^{-\lambda a} \beta(a, x) \mathcal{F}(a, 0)\psi(x) da \\ &\quad - \int_0^{a^*} \beta(a, x) \int_0^a e^{-\lambda(a-s)} \mathcal{F}(a, s) f(s, x) ds da. \end{aligned} \quad (2.6)$$

On the other hand, consider the operator \mathcal{B}_λ defined by

$$(\mathcal{B}_\lambda \psi)(x) = \int_0^{a^*} e^{-\lambda a} \beta(a, x) \mathcal{F}(a, 0)\psi(x) da \quad (\psi \in L^2(\Omega)).$$

We clearly have that $\mathcal{B}_\lambda \in \mathcal{L}(L^2(\Omega))$ for every $\lambda \in \mathbb{R}$ and that $\lim_{\lambda \rightarrow \infty} \|\mathcal{B}_\lambda\|_{\mathcal{L}(L^2(\Omega))} = 0$. This implies that $I - \mathcal{B}_\lambda$ is invertible for λ large enough. Taking

$$\psi = (I - \mathcal{B}_\lambda)^{-1} \int_0^{a^*} \beta(a, \cdot) \int_0^a e^{-\lambda(a-s)} \mathcal{F}(a, s) f(s, \cdot) ds da,$$

and using (2.6), it follows that v defined by (2.5) lies in $\mathcal{D}(A)$ and $\lambda v - Av = f$. We have thus shown that A is a semigroup generator. \square

Under assumptions $(\mathbf{H}_\beta) - (\mathbf{H}_\mu)$, problem (1.1) obviously fits into the above framework. Hence, defining A by (2.3)-(2.4), we can write problem (1.1) in the abstract form:

$$\begin{cases} \dot{p}(t) := Ap(t), & t > 0 \\ p(0) = p_0. \end{cases} \quad (2.7)$$




Figure 2: The spectra of the diffusion free operator A_0 (green crosses) and of $-k\Delta$ (red circles).

The generator A of the population semigroup is the sum of a population operator without diffusion $-d/da - \mu I$ and a spatial diffusion term $k\Delta$. It turns out that spectral properties of A can be easily obtained from those of these two operators, as it can be seen from the following result, proved by Chan and Guo [13].

Theorem 2.4. *Let $0 < \lambda_1^D < \lambda_2^D \leq \lambda_3^D \leq \dots$ be the increasing sequence of eigenvalues of $-k\Delta$ with Dirichlet boundary conditions and let $(\varphi_n^D)_{n \geq 1}$ be a corresponding orthonormal basis of $L^2(\Omega)$. Let $(\lambda_n^0)_{n \geq 1}$ and $(\varphi_n^0)_{n \geq 1}$ be respectively the sequence of eigenvalues and eigenfunctions of the free diffusion operator A_0 defined by (2.1) (see Theorem 2.1). Then the following assertions hold:*

1. *The operator A has compact resolvents and its eigenvalues are given by*

$$\sigma(A) = \left\{ \lambda_i^0 - \lambda_j^D \mid i, j \in \mathbb{N}^* \right\}.$$

2. *A has a real dominant eigenvalue:*

$$\lambda_1 = \lambda_1^0 - \lambda_1^D > \operatorname{Re}(\lambda), \quad \forall \lambda \in \sigma(A), \lambda \neq \lambda_1.$$

Moreover, λ_1 is a simple eigenvalue, the corresponding eigenspace being generated by

$$\varphi_1(a, x) := \varphi_1^0(a) \varphi_1^D(x) = e^{-\lambda_1^0 a} \Pi(a) \varphi_1^D(x).$$

3. *The eigenspace associated to an eigenvalue λ of A is given by*

$$\operatorname{Span} \left\{ \varphi_i^0(a) \varphi_j^D(x) = e^{-\lambda_i^0 a} \Pi(a) \varphi_j^D(x) \mid \lambda_i^0 - \lambda_j^D = \lambda \right\}.$$

We also have the following useful result (see [13, Theorem 2]):

Proposition 2.5. *The semigroup e^{tA} generated on X by A is compact for $t \geq a^*$.*

According to Zabczyk [65, Section 2], this implies in particular that

$$\omega_a(A) = \omega_0(A)$$

where $\omega_a(A) := \lim_{t \rightarrow +\infty} t^{-1} \ln \|e^{tA}\|$ denotes the growth bound of the semigroup e^{tA} and $\omega_0(A) := \sup \{\operatorname{Re} \lambda \mid \lambda \in \sigma(A)\}$ the spectral bound of A . It is worth noticing that the above condition ensures that the exponential stability of e^{tA} is equivalent to the condition $\omega_0(A) < 0$.

3 Detectability of infinite-dimensional systems with finitely many unstable eigenvalues

3.1 General case

Let $A : \mathcal{D}(A) \rightarrow X$ be a linear operator with compact resolvents on a Hilbert space X generating a C_0 -semigroup in X , and let $C \in \mathcal{L}(X, Y)$, where Y is another Hilbert space. We assume that A satisfies the following assumptions

- (A1.)** *A admits M eigenvalues (counted without multiplicities) with real part greater than 0. More precisely we can reorder the eigenvalues $(\lambda_n)_{n \in \mathbb{N}^*}$ of A so that the sequence $(\operatorname{Re} \lambda_n)_{n \in \mathbb{N}^*}$ is nonincreasing, and we suppose that $M \in \mathbb{N}^*$ is such that*

$$\dots \leq \operatorname{Re} \lambda_{M+1} < 0 \leq \operatorname{Re} \lambda_M \leq \dots \leq \operatorname{Re} \lambda_2 \leq \operatorname{Re} \lambda_1.$$

(A2.) We have the equality

$$\omega_a(A) = \omega_0(A)$$

where $\omega_a(A) := \lim_{t \rightarrow +\infty} t^{-1} \ln \|e^{tA}\|$ denotes the growth bound of e^{tA} and $\omega_0(A) := \sup \{\operatorname{Re} \lambda \mid \lambda \in \sigma(A)\}$ the spectral bound of A .

Note that assumption (A2.) is in particular satisfied if e^{tA} is an analytic semigroup or if e^{tA} is compact for some t (see [65, p. 252]).

Definition 3.1. *The pair (A, C) is detectable if there exists $L \in \mathcal{L}(Y, X)$ such that $(A + LC)$ generates an exponentially stable semigroup. Such an operator L is called a stabilizing output injection operator for (A, C) .*

In this section, our goal is to show that the detectability of the infinite dimensional system (A, C) can be deduced from the detectability of the finite dimensional system corresponding to the unstable part of A (namely the projection on the finite dimensional space associated with the unstable eigenvalues $\lambda_1, \dots, \lambda_M$) provided a Hautus type assumption is satisfied. The results of this section can be seen as dual to those obtained in [59] for the stabilization of infinite-dimensional systems having finitely many unstable eigenvalues. Later on, this approach has been successfully used for the control and stabilization of parabolic systems [61, 17], especially in fluid dynamics [12, 52, 9, 10].

We set $\Sigma_+ := \{\lambda_1, \dots, \lambda_M\}$ and let Γ_+ be a positively oriented curve enclosing Σ_+ but no other point of the spectrum $\sigma(A)$ of A . Let $P_+ : X \rightarrow X$ be the projection operator defined by

$$P_+ := -\frac{1}{2\pi i} \int_{\Gamma_+} (\xi - A)^{-1} d\xi. \quad (3.1)$$

We set $X_+ := P_+X$ and $X_- := (I - P_+)X$, and then P_+ provides the following decomposition of X

$$X = X_+ \oplus X_-.$$

Moreover X_+ and X_- are invariant subspaces under A (since A and P_+ commute) and the spectra of the restricted operators $A|_{X_+}$ and $A|_{X_-}$ are respectively Σ_+ and $\sigma(A) \setminus \Sigma_+$ (see [36]). We also set

$$A_+ := A|_{\mathcal{D}(A) \cap X_+} : \mathcal{D}(A) \cap X_+ \rightarrow X_+ \quad \text{and} \quad A_- := A|_{\mathcal{D}(A) \cap X_-} : \mathcal{D}(A) \cap X_- \rightarrow X_-.$$

Remark 3.2. *The space X_+ is the finite dimensional space spanned by the generalized eigenfunctions of A associated to the unstable eigenvalues (i.e. the eigenvalues contained in Σ_+):*

$$X_+ = \bigoplus_{k=1}^M \operatorname{Ker} (A - \lambda_k)^{m_k^P}$$

where m_k^P is the multiplicity of the pole λ_k in the resolvent. If $m_k^A := \dim \operatorname{Ker} (A - \lambda_k)^{m_k^P}$ denotes the algebraic multiplicity of λ_k , $k = 1, \dots, M$, then the dimension of X_+ is the sum of the algebraic multiplicities:

$$\dim X_+ = \sum_{k=1}^M m_k^A. \quad (3.2)$$

Remark 3.3. The operator P_+ is not necessarily an orthogonal projection. More precisely, if we define the projection operator corresponding to A^* :

$$P_+^* := -\frac{1}{2\pi i} \int_{\Gamma_+} (\xi - A^*)^{-1} d\xi, \quad (3.3)$$

and if we set $X_+^* := P_+^* X$ and $X_-^* := (I - P_+^*) X$, then we have

$$(X_+)^{\perp} = X_-^* \quad \text{and} \quad (X_+^*)^{\perp} = X_-.$$

We state now the main result of this section.

Theorem 3.4. Let $Q_+ : Y \rightarrow Y_+ := CX_+$ be the orthogonal projection operator from Y to Y_+ , $i_{X_+} : X_+ \rightarrow X$ be the injection operator from X_+ into X and let

$$C_+ = Q_+ C i_{X_+} \in \mathcal{L}(X_+, Y_+). \quad (3.4)$$

Assume that the finite dimensional projected system (A_+, C_+) is detectable through a stabilizing output injection operator $L_+ \in \mathcal{L}(Y_+, X_+)$. Then, the infinite dimensional system (A, C) is detectable through the stabilizing output injection operator

$$L = i_{X_+} L_+ Q_+ \in \mathcal{L}(Y, X). \quad (3.5)$$

Proof. Throughout the proof, we denote by K a positive constant that may change from line to line. For $L \in \mathcal{L}(Y, X)$, consider the system

$$\dot{z}(t) = Az(t) + LCz(t). \quad (3.6)$$

If we write $z = z_+ + z_-$ where $z_+ := P_+ z$ and $z_- := (I - P_+) z$, by applying P_+ and $(I - P_+)$ to (3.6), there is a corresponding splitting of (3.6) into two equations satisfied by z_+ and z_- respectively

$$\dot{z}_+(t) = A_+ z_+(t) + P_+ LCz(t) \quad \text{and} \quad \dot{z}_-(t) = A_- z_-(t) + (I - P_+) LCz(t).$$

Using (3.5) and the facts that $P_+ i_{X_+} = \text{Id}_{X_+}$ and $(I - P_+) i_{X_+} = 0$, we obtain

$$\dot{z}_+(t) = A_+ z_+(t) + L_+ Q_+ C z(t) \quad \text{and} \quad \dot{z}_-(t) = A_- z_-(t).$$

It follows from assumption **(A2.)** that z_- is exponentially stable:

$$\|z_-(t)\| \leq K e^{-\omega_- t} \|z(0)\| \quad (3.7)$$

where $\omega_- = -\text{Re } \lambda_{M+1} > 0$. On the other hand, by using (3.4) and since $i_{X_+} z_+ = z_+$, we have

$$\begin{aligned} \dot{z}_+(t) &= A_+ z_+(t) + L_+ Q_+ C(z_+(t) + z_-(t)) \\ &= A_+ z_+(t) + L_+ Q_+ C i_{X_+} z_+(t) + L_+ Q_+ C z_-(t) \\ &= (A_+ + L_+ C_+) z_+(t) + L_+ Q_+ C z_-(t). \end{aligned}$$

Using Duhamel's formula, we get

$$z_+(t) = \mathbb{T}_t^+ z_+(0) + \int_0^t \mathbb{T}_{t-s}^+ L_+ Q_+ C z_-(s) ds,$$

where \mathbb{T}_t^+ is the semigroup generated by $(A_+ + L_+C_+)$, which is exponentially stable by the detectability assumption, i.e. there exists $\omega_+ > 0$ such that

$$\left\| \mathbb{T}_t^+ x \right\| \leq K e^{-\omega_+ t} \|x\| \quad \forall x \in X_+, \forall t > 0.$$

Combined with (3.7), this yields

$$\|z_+(t)\| \leq K \left\{ e^{-\omega_+ t} \|z_+(0)\| + \|L_+\| \|C\| \int_0^t e^{-\omega_+(t-s)} e^{-\omega_- s} \|z_-(0)\| ds \right\},$$

and consequently

$$\|z_+(t)\| \leq K \left(e^{-\omega_+ t} + \|L_+\| \|C\| \frac{e^{-\omega_+ t} - e^{-\omega_- t}}{\omega_- - \omega_+} \right) \|z_0\|.$$

It is then sufficient to choose ω_+ small enough such that $0 < \omega_+ < \omega_-$ to have the exponential decay of $t \mapsto z_+(t)$:

$$\|z_+(t)\| \leq K e^{-\omega_+ t} \|z_0\|, \quad t > 0. \quad (3.8)$$

Relations (3.7) and (3.8) yield immediately the exponential decay of $z = z_+ + z_-$. \square

The following result provides a Hautus type sufficient condition for the detectability of the finite dimensional projected system (A_+, C_+) .

Proposition 3.5. *If the Hautus test*

$$(\varphi \in \mathcal{D}(A) \mid A\varphi = \lambda\varphi \text{ for } \lambda \in \Sigma_+ \text{ and } C\varphi = 0) \implies \varphi = 0 \quad (3.9)$$

is satisfied, then (A_+, C_+) is detectable.

Proof. Since $C_+z_+ = Cz_+$ for any $z_+ \in X_+$ by definition of Y_+ and Q_+ , if the Hautus criterion (3.9) is satisfied, then it is clear that the following Hautus test is also satisfied:

$$(\varphi \in \mathcal{D}(A) \cap X_+ \mid A_+\varphi = \lambda\varphi \text{ and } C_+\varphi = 0) \implies \varphi = 0.$$

As the above system is finite dimensional, (A_+, C_+) is detectable. \square

Combining Theorem 3.4 and Proposition 3.5 shows that (A, C) is detectable via the stabilizing output injection operator L defined by (3.5) provided that the Hautus test (3.9) is satisfied.

Remark 3.6. *Classically, the stabilizing output injection operator L^+ of the finite dimensional projected system (A^+, C^+) can be determined by solving a (finite dimensional) Riccati equation. It is worth noticing that the matrices A^+ and C^+ are in practice of small size (their dimensions are respectively $\dim X_+ \times \dim X_+$ and $\dim Y_+ \times \dim X_+$, with $\dim Y_+ \leq \dim X_+$), making the computation of the Riccati matrix affordable.*

Remark 3.7. *A natural question is to determine the minimal number of effective outputs needed to stabilize an infinite dimensional system A having a finite number of unstable eigenvalues. This issue has been investigated for the dual problem of stabilization of parabolic infinite dimensional systems by finite dimensional controls. For the boundary control of a linear parabolic equation, Triggiani has shown in [61] that this number is $\max_{k=1,\dots,M} m_k^G$ provided that the family $(C\varphi_{k,i})_{1 \leq k \leq M, 1 \leq i \leq m_k^G}$ is linearly independent (here $(\varphi_{k,i})_{i \in \{1,\dots,m_k^G\}}$ denotes a basis of $\text{Ker}(A - \lambda_k)$ and m_k^G is the geometric multiplicity of λ_k). More recently, Badra and Takahashi [9, 10] have proved for nonlinear parabolic infinite dimensional systems, that this number is indeed $\max_{k=1,\dots,M} m_k^G$ (even when the linear independence assumption fails). In particular, for an infinite dimensional system having simple eigenvalues, a one dimensional stabilizing feedback can be constructed.*

3.2 The diagonalizable case

The rest of this section is devoted to the particular but important case where the unstable system is diagonalizable. Some of the results given here can be found in Barbu and Triggiani [12]. We assume that $A_+ := A|_{\mathcal{D}(A) \cap X_+}$ is diagonalizable, i.e. $m_k^G = m_k^A$ for all positive eigenvalues λ_k of A . This implies in particular that the unstable space is $X_+ = \bigoplus_{k=1}^M \text{Ker}(A - \lambda_k)$, since $m_k^P = 1$, for all $k = 1, \dots, M$.

We denote by N the number of unstable eigenvalues of A counted **with** multiplicities. For the sake of simplicity, these unstable eigenvalues are still denoted λ_k , $k = 1, \dots, N$. We denote then by $(\varphi_k)_{1 \leq k \leq N}$ a basis of X_+ . The eigenvalues of A^* are given by the complex conjugates $\overline{\lambda_k}$ of the eigenvalues λ_k of A and have the same algebraic and geometric multiplicity. Denote by ψ_k an eigenfunction of A^* corresponding to the unstable eigenvalue $\overline{\lambda_k}$ ($1 \leq k \leq N$). It can be shown (see [12, p. 1453]) that the family $(\psi_k)_{1 \leq k \leq N}$ can be chosen such that $(\varphi_k)_{1 \leq k \leq N}$ and $(\psi_k)_{1 \leq k \leq N}$ form bi-orthogonal sequences, in the sense that $(\varphi_k, \psi_m)_X = \delta_{km}$. It follows then that the projection operator $P_+ \in \mathcal{L}(X, X_+)$ defined in (3.1) can be expressed as

$$P_+z = \sum_{k=1}^N (z, \psi_k)_X \varphi_k \quad (z \in X).$$

Since

$$X_+ = P_+X = \text{Span} \{\varphi_k, 1 \leq k \leq N\},$$

it follows that

$$Y_+ = CX_+ = \text{Span} \{C\varphi_k, 1 \leq k \leq N\}.$$

Assume now that the family

$$(C\varphi_k)_{1 \leq k \leq N} \text{ is linearly independent in } X. \quad (3.10)$$

According to Claim 3.3. of [12, p. 1458], this assumption holds true for an infinite dimensional system described by an evolution PDE with internal observation (this assertion will be proved in Lemma 4.3 in the setting of our population problem). Also note that it implies that the Hautus test (3.9) is necessarily satisfied and that:

$$\dim Y_+ = \dim X_+ = N.$$

We denote by \mathcal{C} the Hermitian matrix of size $N \times N$ defined by

$$\mathcal{C} = ((C\varphi_i, C\varphi_j)_Y)_{1 \leq i \leq N, 1 \leq j \leq N}. \quad (3.11)$$

It is not difficult to prove that (3.10) is equivalent to the fact that \mathcal{C} is invertible. We can then express the orthogonal projection operator $Q_+ : Y \rightarrow Y_+ := CX_+$ as follows (note that $(C\varphi_k)_{1 \leq k \leq N}$ is not an orthogonal family).

Lemma 3.8. *Let \mathcal{C} be the matrix defined by (3.11) and assume that property (3.10) holds true. Then, for any $y \in Y$, Q_+y is defined by*

$$Q_+y = \sum_{i=1}^N (y, \eta_i)_Y C\varphi_i, \quad (3.12)$$

where

$$\eta_i = \sum_{j=1}^N \alpha_{ij} C\varphi_j \quad (3.13)$$

and

$$(\alpha_{ij})_{1 \leq i \leq N, 1 \leq j \leq N} = \mathcal{C}^{-1}. \quad (3.14)$$

Proof. Recall that $Q_+y \in Y^+$ is characterized by

$$\forall y \in Y, \forall n \in \{1, \dots, N\}, \quad (Q_+y - y, C\varphi_n)_Y = 0. \quad (3.15)$$

Noting that $(\overline{\alpha_{ij}})_{1 \leq i \leq N, 1 \leq j \leq N}$ is hermitian (because so is \mathcal{C}), we obtain

$$\begin{aligned} \left(\sum_{i=1}^N (y, \eta_i)_Y C\varphi_i, C\varphi_n \right) &= \sum_{j=1}^N \left(\sum_{i=1}^N \overline{\alpha_{ij}} \mathcal{C}_{in} \right) (y, C\varphi_j) \\ &= \sum_{j=1}^N \left(\sum_{i=1}^N \alpha_{ji} \mathcal{C}_{in} \right) (y, C\varphi_j) \\ &= \sum_{j=1}^N \delta_{jn} (y, C\varphi_j) \\ &= (y, C\varphi_n) \end{aligned}$$

and (3.12) is thus proved. \square

The (finite dimensional) operator $C_+ \in \mathcal{L}(X_+, Y_+)$ defined by (3.4) satisfies $C_+\varphi_k = C\varphi_k$ for any $k \in \{1, \dots, N\}$. Note that C_+ is nothing but the identity matrix when we choose as basis for X_+ and Y_+ respectively $(\varphi_k)_{1 \leq k \leq N}$ and $(C\varphi_k)_{1 \leq k \leq N}$. Therefore, using these bases, $A_+ + L_+C_+$ is a Hurwitz matrix provided $\text{diag}(\lambda_1, \dots, \lambda_N) + L_+$ is Hurwitz. It is thus sufficient to take $L_+ = -\sigma I$ with

$$\sigma > \text{Re } \lambda_1$$

to ensure the stability of $A_+ + L_+C_+$.

The corresponding operator $L \in \mathcal{L}(Y, X)$ defined by (3.5) satisfies, for every $y \in Y$

$$Ly = L_+Q_+y = L_+ \left(\sum_{i=1}^N (y, \eta_i)_Y C\varphi_i \right) = -\sigma \sum_{i=1}^N (y, \eta_i)_Y \varphi_i,$$

and Theorem 3.4 implies that $A + LC$ generates an exponentially stable semigroup. We summarize this useful result in the following Corollary.

Corollary 3.9. *Assume that $A_+ := A|_{\mathcal{D}(A) \cap X_+}$ is diagonalizable and let N be the number of unstable eigenvalues of A counted with multiplicities. We denote by $(\varphi_i)_{1 \leq i \leq N}$ a basis of X_+ and we assume that the family $\{C\varphi_i, 1 \leq i \leq N\}$ is linearly independent in X . Given $\sigma > \operatorname{Re} \lambda_1$, let $L \in \mathcal{L}(Y, X)$ be defined by*

$$Ly = -\sigma \sum_{i=1}^N (y, \eta_i)_Y \varphi_i, \quad (3.16)$$

where $\eta_i, i = 1, \dots, N$, is defined in Lemma 3.8. Then, $A + LC$ generates an exponentially stable semigroup on X .

4 Application : detectability and observer design for the population dynamics system

Going back to the population system (1.1), our aim is to recover the distribution density of the population $p(a, x, t)$ for arbitrary a and x and for t large enough, from the knowledge of $p(a, x, t)$ for arbitrary t but only for an age interval (a_1, a_2) (where $0 \leq a_1 < a_2 \leq a^*$) and a subdomain $\mathcal{O} \subset \Omega$. In other words, the available output is $y(t) = p|_{(a_1, a_2) \times \mathcal{O}}$ where $t \in (0, T)$ (T has to be chosen large enough).

To achieve this, we will construct an observer for system (1.1). Observers have been introduced by Luenberger to estimate the state of a finite dimensional dynamical system from the knowledge of an output (of course assuming that the initial state is unknown). Roughly speaking, an observer is an auxiliary dynamical system that uses as inputs the available measurements (that is the output of the original system) and converges asymptotically (in time) towards the state of the original system. The generalization of observers (also called estimators or filters in the stochastic framework) to infinite dimensional systems goes to the eighties (see for instance Baras *et al.* [11, 35], Chapman and Pritchard [15], Curtain and Zwart [19] and Hidayat *et al.* [32] for a brief overview of observers in the context of distributed systems). Observers have been successfully used to solve data assimilation problems in geophysics (see Auroux and Blum [5], Auroux and Nodet [6], Auroux, Bansart and Blum [4]), initial data reconstruction (see Fridman [24], Haine [30], Haine and Ramdani [31], Ramdani, Tucsnak and Weiss [51]) and joint state and/or parameter identification by Curtain, Demetriou and Ito [18], Moireau *et al.* [14, 46, 47], Krstic *et al.* [40, 55].

Using the notation of Section 2, let A be defined by (2.3)-(2.4). We consider the abstract system (2.7), i.e.

$$\dot{p}(t) = Ap(t), \quad t \in (0, T),$$

with the output

$$y(t) = Cp(t), \quad t \in (0, T),$$

where the observation operator $C \in \mathcal{L}(X, Y)$, with $Y := L^2((a_1, a_2) \times \mathcal{O})$ is defined by

$$C\varphi = \varphi|_{(a_1, a_2) \times \mathcal{O}} \quad (\varphi \in X). \quad (4.1)$$

We introduce the observer

$$\begin{cases} \dot{\hat{p}}(t) = A\hat{p}(t) + LC\hat{p}(t) - Ly(t), & t \in (0, T), \\ \hat{p}(0) = 0, \end{cases} \quad (4.2)$$

where $L \in \mathcal{L}(Y, X)$ is a linear operator to be defined. As explained in Section 1, the error $e := \hat{p} - p$ satisfies

$$\begin{cases} \dot{e}(t) = (A + LC)e(t), & t \in (0, T), \\ e(0) = -p(0). \end{cases} \quad (4.3)$$

According to Theorem 2.1, Theorem 2.4 and Proposition 2.5, assumptions **(A1.)** and **(A2.)** (see the beginning of Section 3) are satisfied for the above system. Therefore, the problem of determining the stabilizing output injection operator L for (A, C) fits into the framework described in Section 3 and we can apply the approach developed therein. Denoting by Σ_+ the set of the eigenvalues of A with positive real part, it only remains to verify the assumption of Proposition 3.5, and in particular the following lemma shows that the Hautus test of Proposition 3.5 is satisfied for our system (A, C) .

Lemma 4.1. *If $\varphi \in \mathcal{D}(A)$ satisfies $A\varphi = \lambda\varphi$ for $\lambda \in \Sigma_+$ and $C\varphi = 0$, then φ vanishes identically.*

Proof. Let λ be an unstable eigenvalue of A (i.e. $\operatorname{Re} \lambda \geq 0$) and let $\varphi \in \mathcal{D}(A)$ satisfying $A\varphi = \lambda\varphi$. According to assertion 3 in Theorem 2.4, we have

$$\varphi(a, x) = \sum_{i,j|\lambda=\lambda_i^0-\lambda_j^D} \alpha_j e^{-\lambda_i^0 a} \Pi(a) \varphi_j^D(x).$$

The fact that $C\varphi = 0$, with φ an eigenvector of A corresponding to the eigenvalue λ , writes

$$\sum_{i,j|\lambda=\lambda_i^0-\lambda_j^D} \alpha_j e^{-\lambda_i^0 a} \Pi(a) \varphi_j^D|_{\mathcal{O}} = 0, \quad a \in (a_1, a_2).$$

As the eigenvalues $(\lambda_i^0)_{i \in \mathbb{N}}$ are distinct, this equation reduces to

$$\sum_{j|\lambda+\lambda_j^D \in \sigma(A_0)} \alpha_j e^{-\lambda_{i(j)}^0 a} \varphi_j^D|_{\mathcal{O}}(x) = 0, \quad a \in (a_1, a_2),$$

where $i(j) \in \mathbb{N}^*$ is the unique index such that $\lambda_i^0 = \lambda + \lambda_j^D$. Since the eigenfunctions of $-k\Delta$ with Dirichlet boundary conditions are analytic, we immediately obtain that $\varphi = 0$. \square

Applying Theorem 3.4 and Proposition 3.5 to (4.3) immediately leads to the following result.

Theorem 4.2. *Let $p_0 \in X$ and p the solution of (1.1). Assume that $y(t) = p|_{(a_1, a_2) \times \mathcal{O}}$ ($t > 0$) is known. Let \hat{p} the observer defined by (4.2), where $L \in \mathcal{L}(Y, X)$ is the (finite dimensional) stabilizing output injection operator defined by (3.5). Then, there exists $C > 0$ and $\omega > 0$ such that*

$$\|\hat{p}(t) - p(t)\| \leq C e^{-\omega t} \|p_0\|, \quad t > 0.$$

To conclude this Section, we go back to the particular but important case where the unstable system is diagonalizable (see §.3.2), i.e. we assume that $A_+ := A|_{\mathcal{D}(A) \cap X_+}$ is diagonalizable. For the sake of simplicity, we denote here by M the number of unstable eigenvalues of A counted **without** multiplicities (still denoted λ_k , $k = 1, \dots, M$) and we

denote by $(\varphi_{k,i})_{i \in \{1, \dots, m_k^G\}}$ a basis of $\text{Ker}(A - \lambda_k)$ where m_k^G is the geometric multiplicity of λ_k ($k = 1, \dots, M$). In order to show that the stabilizing output injection operator L can be rewritten as (3.16), it remains to verify that assumption (3.10) holds. This is done in the following lemma, which is exactly Part (A) of Claim 3.3. in [12, p. 1458]. The proof is recalled here for reader's convenience.

Lemma 4.3. *The family $(C\varphi_{k,i})_{1 \leq k \leq M, 1 \leq i \leq m_k^G}$ is linearly independent in X .*

Proof. Step 1. It is easy to prove that the Hautus test (3.9), which is verified in Lemma 4.1 for A and C respectively defined by (2.3)-(2.4) and (4.1), is equivalent to the fact that

$$\forall k \in \{1, \dots, M\}, \text{ the family } (C\varphi_{k,i})_{1 \leq i \leq m_k^G} \text{ is linearly independent in } X.$$

Step 2. Consider now the case of two distinct eigenvalues, say λ_1 and λ_2 . Assume by contradiction that the corresponding system of eigenfunctions

$$\left\{ C\varphi_{1,i}, C\varphi_{2,j} \mid 1 \leq i \leq m_1^G, 1 \leq j \leq m_2^G \right\}$$

is linearly dependent and assume, for instance, that

$$C\varphi_{2,m_2^G} = \sum_{i=1}^{m_1^G} \alpha_i C\varphi_{1,i} + \sum_{j=1}^{m_2^G-1} \beta_j C\varphi_{2,j},$$

i.e. that

$$\varphi_{2,m_2^G} = \sum_{i=1}^{m_1^G} \alpha_i \varphi_{1,i} + \sum_{j=1}^{m_2^G-1} \beta_j \varphi_{2,j} \quad \text{on } (a_1, a_2) \times \mathcal{O}. \quad (4.4)$$

Applying the differential operator $A = -\partial_a - \mu \cdot + k\Delta$ defined by (2.4), we obtain that on $(a_1, a_2) \times \mathcal{O}$:

$$0 = A \left(\varphi_{2,m_2^G} - \sum_{i=1}^{m_1^G} \alpha_i \varphi_{1,i} - \sum_{j=1}^{m_2^G-1} \beta_j \varphi_{2,j} \right) = \lambda_2 \varphi_{2,m_2^G} - \lambda_1 \sum_{i=1}^{m_1^G} \alpha_i \varphi_{1,i} - \lambda_2 \sum_{j=1}^{m_2^G-1} \beta_j \varphi_{2,j},$$

which implies, using again (4.4) that

$$(\lambda_2 - \lambda_1) \sum_{i=1}^{m_1^G} \alpha_i \varphi_{1,i} = 0 \quad \text{on } (a_1, a_2) \times \mathcal{O}.$$

Using the fact that $\lambda_1 \neq \lambda_2$ and Step 1, we find that $\alpha_i = 0$ for all $i \in \{1, \dots, m_1^G\}$.

Therefore $\varphi_{2,m_2^G} = \sum_{j=1}^{m_2^G-1} \beta_j \varphi_{2,j}$ on $(a_1, a_2) \times \mathcal{O}$, which contradicts Step 1.

Step 3. Finally we prove the statement of Lemma 4.3 by induction with respect to the number $q \leq M$ of distinct eigenvalues λ_k . Assume that the statement holds for $q-1$ distinct eigenvalues $\lambda_1, \dots, \lambda_{q-1}$ and consider the system $(C\varphi_{k,i})_{1 \leq k \leq q, 1 \leq i \leq m_k^G}$. By contradiction, assume that this system is linearly dependent. As in Step 2, we may assume, for instance, that

$$\varphi_{q,m_q^G} = \sum_{k=1}^{q-1} \sum_{i=1}^{m_k^G} \alpha_{ki} \varphi_{k,i} + \sum_{j=1}^{m_q^G-1} \beta_j \varphi_{q,j} \quad \text{on } (a_1, a_2) \times \mathcal{O}.$$

Applying the differential operator A and proceeding as in Step 2, we obtain that

$$\sum_{k=1}^{q-1} (\lambda_q - \lambda_k) \sum_{i=1}^{m_k^G} \alpha_{ki} \varphi_{k,i} = 0 \quad \text{on } (a_1, a_2) \times \mathcal{O}.$$

As the eigenvalues are distinct, by the inductive hypothesis, this implies that $\alpha_{ki} = 0$ for any $k \in \{1, \dots, q-1\}$ and $i \in \{1, \dots, m_k^G\}$. Therefore

$$\varphi_{q, m_q^G} = \sum_{j=1}^{m_q^G - 1} \beta_j \varphi_{q,j} \quad \text{on } (a_1, a_2) \times \mathcal{O},$$

which contradicts Step 1 and ends the proof. \square

The results in this section can be extended to other observation operators, as shown in the two following remarks.

Remark 4.4. *Given a_1, a_2 with $0 \leq a_1 < a_2 \leq a^*$, we can take $Y = L^2(\mathcal{O})$ and $C \in \mathcal{L}(X, Y)$ defined by*

$$(C\varphi)(x) = \int_{a_1}^{a_2} \varphi(a, x) da, \quad \varphi \in X, x \in \mathcal{O}. \quad (4.5)$$

Indeed, using the third assertion in Theorem 2.4, it follows that if φ satisfies $A\varphi = \lambda\varphi$ for some $\lambda \in \mathbb{C}$ and $C\varphi = 0$ then there exist complex numbers α_j such that

$$\sum_{j|\lambda + \lambda_j^D \in \sigma(A_0)} \alpha_j \varphi_j^D(x) \int_{a_1}^{a_2} e^{-\lambda_{i(j)}^0 a} \Pi(a) da = 0, \quad x \in \mathcal{O},$$

where $i(j) \in \mathbb{N}^$ is the unique index such that $\lambda_i^0 = \lambda + \lambda_j^D$. Using again the analyticity of the functions φ_j^D , combined with the fact that $e^{-\lambda_{i(j)}^0 a} \Pi(a) > 0$ for $a \in [a_1, a_2]$, we conclude that $\varphi = 0$. The observation operator defined in (4.5) is clearly of biological interest since measuring $y = Cp$ signifies counting the population having an age in the interval (a_1, a_2) and localized in the spatial domain \mathcal{O} .*

Remark 4.5. *Another possible choice of the observation operator, which can be seen as a limit case of the operator defined in (4.1) is*

$$(C\varphi)(x) = \varphi(a_0, x), \quad \varphi \in X, x \in \mathcal{O}, \quad (4.6)$$

where $a_0 \in (0, a^)$ is a fixed age. In this case, the conditions $A\varphi = \lambda\varphi$ for some $\lambda \in \mathbb{C}$ and $C\varphi = 0$ write*

$$-\partial_a \varphi - \mu\varphi + k\Delta\varphi = \lambda\varphi, \quad \varphi \in \mathcal{D}(A), \quad (4.7)$$

$$\varphi(a_0, x) = 0, \quad x \in \mathcal{O}. \quad (4.8)$$

*Looking to the first equation in (4.7) as a parabolic equation with a in the role of the time variable and applying a well-known unique continuation result for parabolic equations (see, for instance, Phung and Wang [49, Proposition 2.2]) we obtain that $\varphi = 0$. Note that the argument used to show the observability of eigenfunctions in this case (and, consequently, for the observation operator (4.1)) can be used, without any modification, in the case $\mu = \mu(a, x)$ and $\beta = \beta(a, x)$, as assumptions **(A1.)** and **(A2.)** are satisfied in this case (see for instance [26]).*

5 Discretization

From now on, we assume for the sake of simplicity that Ω is the interval $(0, \ell)$, that $A_+ := A|_{\mathcal{D}(A) \cap X_+}$ is diagonalizable and that A has N unstable eigenvalues $\lambda_1, \dots, \lambda_N$ counted with multiplicities.

Our goal in the last two sections is to show how to implement the proposed method to estimate the final state from partial observations. More precisely, assuming that p_0 is an *unknown* initial data, we want to estimate p at time $t = T$ where p solves the following open loop system:

$$\begin{cases} \partial_t p(a, x, t) + \partial_a p(a, x, t) + \mu(a)p(a, x, t) - k\partial_{xx}p(a, x, t) = 0, & a \in (0, a^*), x \in (0, \ell), t > 0, \\ p(a, 0, t) = p(a, \ell, t) = 0, & a \in (0, a^*), t > 0, \\ p(a, x, 0) = p_0(a, x), & a \in (0, a^*), x \in (0, \ell), \\ p(0, x, t) = \int_0^{a^*} \beta(a)p(a, x, t) da, & x \in (0, \ell), t > 0, \end{cases} \quad (5.1)$$

provided we know the observation

$$y(t) = p(t)|_{(a_1, a_2) \times (\ell_1, \ell_2)}, \quad t \in (0, T),$$

where $0 \leq a_1 < a_2 \leq a^*$ and $0 \leq \ell_1 < \ell_2 \leq \ell$.

This leads to define the observation operator $C \in \mathcal{L}(X, Y)$

$$C\varphi = \varphi|_{(a_1, a_2) \times (\ell_1, \ell_2)}, \quad \forall \varphi \in X$$

where $X = L^2((0, a^*) \times (0, \ell))$ and $Y = L^2((a_1, a_2) \times (\ell_1, \ell_2))$.

In order to estimate $p(T)$, we use the observer designed in Section 4. As the unstable space is the N -dimensional space

$$X_+ = \bigoplus_{k=1}^N \text{Ker}(A - \lambda_k) = \text{Span} \{ \varphi_k, 1 \leq k \leq N \} = \text{Span} \left\{ \varphi_i^0(a) \varphi_j^D(x), \lambda_i^0 - \lambda_j^D = \lambda_k, 1 \leq k \leq N \right\}$$

(see Section 2), the observer defined by (4.2) involves the stabilizing output injection operator L defined by (3.16) (see §. 3.2)

$$Ly = -\sigma \sum_{i=1}^N (y, \eta_i)_Y \varphi_i \quad (y \in Y),$$

where $\sigma > \lambda_1$ (gain coefficient) and

$$\eta_i = \sum_{j=1}^N \alpha_{ij} C \varphi_j \quad \text{and} \quad (\alpha_{ij})_{1 \leq i \leq N, 1 \leq j \leq N} = \mathcal{C}^{-1}.$$

(see (3.13)-(3.14)).

The observer solves then the following system

$$\left\{ \begin{array}{l} \partial_t \widehat{p}(a, x, t) + \partial_a \widehat{p}(a, x, t) + \mu(a) \widehat{p}(a, x, t) - k \partial_{xx} \widehat{p}(a, x, t) \\ \quad + \sigma \sum_{i=1}^N (C \widehat{p}, \eta_i)_Y \varphi_i(a, x) = \sigma \sum_{i=1}^N (y, \eta_i)_Y \varphi_i(a, x), \quad a \in (0, a^*), x \in (0, \ell), t > 0, \\ \widehat{p}(a, 0, t) = \widehat{p}(a, \ell, t) = 0, \quad a \in (0, a^*), t > 0, \\ \widehat{p}(a, x, 0) = 0, \quad a \in (0, a^*), x \in (0, \ell), \\ \widehat{p}(0, x, t) = \int_0^{a^*} \beta(a) \widehat{p}(a, x, t) da, \quad x \in (0, \ell), t > 0. \end{array} \right. \quad (5.2)$$

Numerical approximation of population models with spatial diffusion (typically of the form (5.1)) has been considered in several papers. Lopez and Trigiante proposed in [43] an adaptative finite difference scheme to handle the singularity of the mortality function μ as $a \rightarrow a^*$. Milner [45] and Kim [38] proposed schemes using finite difference along the characteristic age-time direction and finite elements in space or mixed finite element [39]. Huyer [33] developed a semigroup approach to discretize the problem using a Galerkin approximation involving Laguerre polynomials in age and spectral decomposition in space. Ayati [7] proposed a method allowing variable time steps and independent age and time discretizations. Ayati and Dupont [8] proposed Galerkin approximations in both age and space for a population model with nonlinear diffusion. Let us also mention the contributions of Gerardo-Giorda *et al.* who proposed approximations using finite difference in age and time and finite element in space for an age structured model with space and age dependent diffusion in [20] and density dependent diffusion in [25]. We refer the interested reader to the PhD of Pelovska [48] for more references.

Below, we successively present finite difference discretizations of systems (5.1) and (5.2). For both problems, one of the main difficulties comes from the discretization of the renewal equation $p(0, x, t) = \int_0^{a^*} \beta(a) p(a, x, t) da$ appearing in the domain of the operator. Indeed, this non local equation couples the unknowns corresponding to different ages for a given time and at a given point of space. In [33], Huyer overcame this problem by proposing a reformulation of the system in which this condition is taken into account by adding a Dirac term in the evolution PDE defining the dynamics. Here, we use the same approach as in [20], in which the renewal equation is simplified via the time discretization: the density of population at age 0 for a given time t^n is evaluated using the values of the density for all ages at the previous time t^{n-1} (see Figure 3).

5.1 Discretization of the open loop problem : data generation

5.1.1 Rescaling the problem

In order to overcome the difficulties due to singular behavior of the coefficient μ (see equation (1.3)), we follow [20] and introduce the auxiliary variable

$$u(a, x, t) = \frac{p(a, x, t)}{\Pi(a)} = \exp\left(\int_0^{a^*} \mu(s) ds\right) p(a, x, t).$$

One can easily check that u satisfies

$$\begin{cases} \partial_t u(a, x, t) + \partial_a u(a, x, t) - k \partial_{xx} u(a, x, t) = 0, & a \in (0, a^*), x \in (0, \ell), t > 0, \\ u(a, 0, t) = u(a, \ell, t) = 0, & a \in (0, a^*), t > 0, \\ u(a, x, 0) = u_0(a, x), & a \in (0, a^*), x \in (0, \ell), \\ u(0, x, t) = \int_0^{a^*} m(a) u(a, x, t) da, & x \in (0, \ell), t > 0, \end{cases}$$

where we have set $u_0(a, x) = p_0(a, x)/\Pi(a)$ and where $m(a) = \beta(a)\Pi(a)$ stands for the maternity function.

5.1.2 Finite difference discretization in time

Let $u^n(a, x)$ be an approximation of $u(a, x, t^n)$, where $t^n = n\Delta t$, $0 \leq n \leq N_t$, $\Delta t = T/N_t$ is a discretization of $(0, T)$. Starting from $u^0(a, x) = u_0(a, x)$, we construct u^n for $n \geq 1$ using an Euler's backwards scheme

$$\begin{cases} \frac{1}{\Delta t} (u^n(a, x) - u^{n-1}(a, x)) + \partial_a u^n(a, x) - k \partial_{xx} u^n(a, x) = 0, & a \in (0, a^*), x \in (0, \ell), \\ u^n(a, 0) = u^n(a, \ell) = 0, & a \in (0, a^*), \\ u^0(a, x) = u_0(a, x), & a \in (0, a^*), x \in (0, \ell), \\ u^n(0, x) = \int_0^{a^*} m(a) u^{n-1}(a, x) da, & x \in (0, \ell). \end{cases}$$

5.1.3 Finite difference discretization in space

Denoting by $u_i^n(a)$ an approximation of $u^n(x_i, a)$ (where $x_i = ih = i\ell/(N_x + 1)$, with $0 \leq i \leq N_x + 1$) and using a classical centered approximation for the second order derivative in space, the above system yields

$$\begin{cases} \frac{d\mathbf{U}^n}{da}(a) + \frac{1}{h^2} \mathbb{K} \mathbf{U}^n(a) + \frac{1}{\Delta t} \mathbf{U}^n(a) = \frac{1}{\Delta t} \mathbf{U}^{n-1}(a), \\ \mathbf{U}^n(0) = \int_0^{a^*} m(a) \mathbf{U}^{n-1}(a) da, \\ \mathbf{U}^0(a) = \mathbf{U}_0(a), \end{cases} \quad (5.3)$$

where

$$\mathbf{U}^n(a) := \begin{pmatrix} u_1^n(a) \\ \vdots \\ \vdots \\ u_{N_x}^n(a) \end{pmatrix}, \quad \mathbf{U}_0(a) = \begin{pmatrix} u_0(a, x_1) \\ \vdots \\ \vdots \\ u_0(a, x_{N_x}) \end{pmatrix}, \quad \mathbb{K} = k \begin{pmatrix} 2 & -1 & & & \\ -1 & 2 & -1 & & 0 \\ & -1 & 2 & -1 & \\ & & \ddots & \ddots & \ddots \\ 0 & & & -1 & 2 & -1 \\ & & & & -1 & 2 \end{pmatrix}.$$

5.1.4 Finite difference discretization in age

We use a Crank-Nicholson scheme to approximate the solution of problem (5.3). Denoting by $u_i^{n,k}$ an approximation of $u_i^n(a^k)$, where $a^k = k\Delta a$, $0 \leq k \leq N_a$, $\Delta a = a^*/N_a$, and by

$$\mathbf{U}^{n,k} := \begin{pmatrix} u_1^{n,k} \\ \vdots \\ \vdots \\ u_{N_x}^{n,k} \end{pmatrix}$$

an approximation of $\mathbf{U}^n(a^k)$, we move from age a^{k-1} to age a^k following the scheme

$$\begin{aligned} \frac{1}{\Delta a} (\mathbf{U}^{n,k} - \mathbf{U}^{n,k-1}) + \frac{1}{h^2} \mathbb{K} \left(\frac{\mathbf{U}^{n,k} + \mathbf{U}^{n,k-1}}{2} \right) + \frac{1}{\Delta t} \left(\frac{\mathbf{U}^{n,k} + \mathbf{U}^{n,k-1}}{2} \right) \\ = \frac{1}{\Delta t} \left(\frac{\mathbf{U}^{n-1,k} + \mathbf{U}^{n-1,k-1}}{2} \right), \end{aligned}$$

with the initial conditions

$$\begin{cases} \mathbf{U}^{0,k} = \mathbf{U}_0(a^k), & \forall k = 0, \dots, N_a, \\ \mathbf{U}^{n,0} = \sum_{k=0}^{N_a} \omega_k m(a^k) \mathbf{U}^{n-1,k} \simeq \int_0^{a^*} m(a) \mathbf{U}^{n-1}(a) da, \end{cases}$$

where ω_k are the weights of the quadrature formula used.

5.1.5 Algorithm for the open loop system




Figure 3: Step n of the algorithm. The unknown $\mathbf{U}^{n,k}$ at each node (k, n) is the vector containing the solution at the different points of the space discretization.

1. For $n = 0$: Initialization of $\mathbf{U}^{0,k}$ via the formula

$$\mathbf{U}^{0,k} = \begin{bmatrix} u_0(k\Delta a, x_1) \\ \vdots \\ u_0(k\Delta a, x_{N_h}) \end{bmatrix}.$$

2. For $n = 1, \dots, N_t$:

- $k = 0$: Initialization of $\mathbf{U}^{n,0}$ using the values of $(\mathbf{U}^{n-1,j})_{j=0}^{N_a}$:

$$\mathbf{U}^{n,0} = \sum_{k=0}^{N_a} \omega_k m(a^k) \mathbf{U}^{n-1,k}$$

- For $k = 1, \dots, N_a$, $\mathbf{U}^{n,k} = \begin{pmatrix} u_1^{n,k} \\ \vdots \\ u_{N_x}^{n,k} \end{pmatrix}$ solves the linear system

$$\mathbb{A} \mathbf{U}^{n,k} = \mathbf{b}^{n,k}$$

where

$$\mathbb{A} = \left(\Delta t + \frac{1}{2} \Delta a \right) \mathbb{I} + \frac{1}{2} \frac{\Delta t \Delta a}{h^2} \mathbb{K}$$

$$\mathbf{b}^{n,k} = \frac{1}{2} \Delta a \left(\mathbf{U}^{n-1,k} + \mathbf{U}^{n-1,k-1} \right) + \left[\left(\Delta t - \frac{1}{2} \Delta a \right) \mathbb{I} - \frac{1}{2} \frac{\Delta t \Delta a}{h^2} \mathbb{K} \right] \mathbf{U}^{n,k-1}$$

- $\mathbf{P}^{n,k} = \begin{pmatrix} p_1^{n,k} \\ \vdots \\ p_{N_x}^{n,k} \end{pmatrix}$ where $p_i^{n,k} = \Pi(a^k) u_i^{n,k}$.

- $\mathbf{Y}^{n,k} = \begin{pmatrix} y_1^{n,k} \\ \vdots \\ y_{N_x}^{n,k} \end{pmatrix}$ where $y_i^{n,k} = p_i^{n,k}$ if $\ell_1 \leq ih \leq \ell_2$ and $y_i^{n,k} = 0$ otherwise.

5.2 Discretization of the closed loop system : observer design

In order to discretize (5.2), we follow the same procedure as in the previous subsection for the open loop system. The main differences in the dynamics are the presence of a source term coming from the observation y and the extra term in the evolution equation

$$\sum_{i=1}^N (C \hat{p}_i, \eta_i)_Y \varphi_i.$$

5.2.1 Rescaling the problem

First of all, in order to overcome the difficulties due to singular behavior of the coefficient μ , we introduce as we did for the open loop system, the auxiliary variable

$$\hat{u}(a, x, t) = \frac{\hat{p}(a, x, t)}{\Pi(a)} = \exp\left(\int_0^{a^*} \mu(s) ds\right) \hat{p}(a, x, t),$$

where \hat{p} solves (5.2).

One can easily check that \hat{u} satisfies

$$\left\{ \begin{array}{l} \partial_t \hat{u}(a, x, t) + \partial_a \hat{u}(a, x, t) - k \partial_{xx} \hat{u}(a, x, t) \\ \quad + \sigma \sum_{i=1}^N (C\Pi \hat{u}, \eta_i)_Y v_i(a, x) = \sigma \sum_{i=1}^N (y(t), \eta_i)_Y v_i(a, x), \quad a \in (0, a^*), \quad x \in (0, \ell), \quad t > 0, \\ \hat{u}(a, 0, t) = \hat{u}(a, \ell, t) = 0, \quad a \in (0, a^*), \quad t > 0, \\ \hat{u}(a, x, 0) = 0, \quad a \in (0, a^*), \quad x \in (0, \ell), \\ \hat{u}(0, x, t) = \int_0^{a^*} m(a) \hat{u}(a, x, t) da, \quad x \in (0, \ell), \quad t > 0, \end{array} \right. \quad (5.4)$$

where we have set $v_i(a, x) = \varphi_i(a, x)/\Pi(a)$ ($1 \leq i \leq N$).

In order to discretize the term $(C\Pi \hat{u}, \eta_i)_Y$ in (5.4), let us introduce the new (scalar) unknowns

$$\alpha_i(t) = (C\Pi \hat{u}, \eta_i)_Y \quad (1 \leq i \leq N).$$

Using the first equation in (5.4) and the fact that, for any $i, j \in \{1, \dots, N\}$,

$$(C\Pi v_j, \eta_i)_Y = (C\varphi_j, \eta_i)_Y = \sum_{k=1}^N \overline{\alpha_{ik}} (C\varphi_j, C\varphi_k)_Y = \sum_{k=1}^N \overline{\alpha_{ik} (C\varphi_k, C\varphi_j)_Y} = \delta_{ij},$$

(by (3.13)-(3.14)), we remark that α_i satisfies

$$\dot{\alpha}_i(t) = - (C\Pi \partial_a \hat{u}, \eta_i)_Y + k (C\Pi \partial_{xx} \hat{u}, \eta_i)_Y - \sigma \alpha_i(t) + \sigma (y, \eta_i)_Y.$$

Consequently, problem (5.4) can be reformulated as follows

$$\left\{ \begin{array}{l} \dot{\alpha}_i(t) = - (C\Pi \partial_a \hat{u}(t), \eta_i)_Y + k (C\Pi \partial_{xx} \hat{u}(t), \eta_i)_Y \\ \quad - \sigma \alpha_i(t) + \sigma (y(t), \eta_i)_Y, \quad t > 0, \quad 1 \leq i \leq N, \\ \partial_t \hat{u}(a, x, t) + \partial_a \hat{u}(a, x, t) - k \partial_{xx} \hat{u}(a, x, t) + \sigma \sum_{i=1}^N \alpha_i(t) v_i(a, x) \\ \quad = \sigma \sum_{i=1}^N (y(t), \eta_i)_Y v_i(a, x), \quad a \in (0, a^*), \quad x \in (0, \ell), \quad t > 0, \\ \alpha_i(0) = 0, \quad 1 \leq i \leq N, \\ \hat{u}(a, 0, t) = \hat{u}(a, \ell, t) = 0, \quad a \in (0, a^*), \quad t > 0, \\ \hat{u}(a, x, 0) = 0, \quad a \in (0, a^*), \quad x \in (0, \ell), \\ \hat{u}(0, x, t) = \int_0^{a^*} m(a) \hat{u}(a, x, t) da, \quad x \in (0, \ell), \quad t > 0. \end{array} \right.$$

5.2.2 Finite difference discretization in time

Let $\hat{u}^n(a, x)$ (resp. $\alpha_i^n, y^n(a, x)$) be an approximation of $\hat{u}(a, x, t^n)$ (resp. $\alpha_i(t^n), y(a, x, t^n)$), where $t^n = n\Delta t$, $0 \leq n \leq N_t$, $\Delta t = T/N_t$ is a discretization of $(0, T)$. Starting from $\alpha_i^0 = 0$ and $\hat{u}^0(a, x) = 0$, we construct α_i^n and \hat{u}^n for $n \geq 1$ using an Euler's backwards scheme

$$\left\{ \begin{array}{l} \frac{1}{\Delta t} (\alpha_i^n - \alpha_i^{n-1}) = - (C\Pi\partial_a\hat{u}^{n-1}, \eta_i)_Y + k (C\Pi\partial_{xx}\hat{u}^{n-1}, \eta_i)_Y \\ \quad - \sigma\alpha_i^n + \sigma (y^n, \eta_i)_Y, \quad 1 \leq i \leq N, \\ \frac{1}{\Delta t} (\hat{u}^n(a, x) - \hat{u}^{n-1}(a, x)) + \partial_a\hat{u}^n(a, x) - k\partial_{xx}\hat{u}^n(a, x) \\ \quad + \sigma \sum_{i=1}^N \alpha_i^n v_i(a, x) = \sigma \sum_{i=1}^N (y^n, \eta_i)_Y v_i(a, x), \quad a \in (0, a^*), x \in (0, \ell), \\ \alpha_i^0 = 0, \quad 1 \leq i \leq N, \\ \hat{u}^n(a, 0) = \hat{u}^n(a, \ell) = 0, \quad a \in (0, a^*), \\ \hat{u}^0(a, x) = 0, \quad a \in (0, a^*), x \in (0, \ell), \\ \hat{u}^n(0, x) = \int_0^{a^*} m(a)\hat{u}^{n-1}(a, x) da, \quad x \in (0, \ell). \end{array} \right.$$

5.2.3 Finite difference discretization in space

Denoting by $\hat{u}_i^n(a)$ an approximation of $\hat{u}^n(x_i, a)$ (where $x_i = ih = i\ell/(N_x + 1)$, with $0 \leq i \leq N_x + 1$) and using a classical centered approximation for the second order derivative in space, the above system yields

$$\left\{ \begin{array}{l} \frac{1}{\Delta t} \alpha_i^n = \frac{1}{\Delta t} \alpha_i^{n-1} - \sigma\alpha_i^n - h \left(\int_{a_1}^{a_2} \Pi(a)\boldsymbol{\eta}_i(a)^T \partial_a \hat{\mathbf{U}}^{n-1}(a) da \right) \\ \quad - \frac{1}{h} \left(\int_{a_1}^{a_2} \Pi(a)\boldsymbol{\eta}_i(a)^T \mathbb{K} \hat{\mathbf{U}}^{n-1}(a) da \right) + \sigma h \left(\int_{a_1}^{a_2} \boldsymbol{\eta}_i(a)^T \mathbf{Y}^n(a) da \right), \\ \frac{d\hat{\mathbf{U}}^n}{da}(a) + \frac{1}{h^2} \mathbb{K} \hat{\mathbf{U}}^n(a) + \frac{1}{\Delta t} \hat{\mathbf{U}}^n(a) + \sigma \sum_{i=1}^N \alpha_i^n \mathbf{V}_i(a) = \frac{1}{\Delta t} \hat{\mathbf{U}}^{n-1}(a) \\ \quad + \sigma h \sum_{i=1}^N \left(\int_{a_1}^{a_2} \boldsymbol{\eta}_i(a)^T \mathbf{Y}^n(a) da \right) \mathbf{V}_i(a), \quad (5.5) \\ \alpha_i^0 = 0, \\ \hat{\mathbf{U}}^n(0) = \int_0^{a^*} m(a)\hat{\mathbf{U}}^{n-1}(a) da, \\ \hat{\mathbf{U}}^0(a) = 0, \end{array} \right.$$

where

$$\hat{\mathbf{U}}^n(a) := \begin{pmatrix} \hat{u}_1^n(a) \\ \vdots \\ \hat{u}_{N_x}^n(a) \end{pmatrix}, \quad \mathbf{V}_i(a) := \begin{pmatrix} v_i(a, x_1) \\ \vdots \\ v_i(a, x_{N_x}) \end{pmatrix}, \quad \boldsymbol{\eta}_i(a) := \begin{pmatrix} \eta_i(a, x_1) \\ \vdots \\ \eta_i(a, x_{N_x}) \end{pmatrix}, \quad \mathbf{Y}^n(a) := \begin{pmatrix} y^n(a, x_1) \\ \vdots \\ y^n(a, x_{N_x}) \end{pmatrix}.$$

We take here $y^n(a, x_i) = 0$ for all $i \in \{1, \dots, N_x\}$ such that $x_i \notin (\ell_1, \ell_2)$, for all $a \in (0, a^*)$ and all $n \geq 1$.

5.2.4 Finite difference discretization in age

We use a Crank-Nicholson scheme to approximate the solution of problem (5.5). Denoting by $\widehat{u}_i^{n,k}$ an approximation of $\widehat{u}_i^n(a^k)$, where $a^k = k\Delta a$, $0 \leq k \leq N_a$, $\Delta a = a^*/N_a$, $N_{a_1} = \lfloor a_1/\Delta a \rfloor$, $N_{a_2} = \lceil a_2/\Delta a \rceil$ and by

$$\widehat{\mathbf{U}}^{n,k} := \begin{pmatrix} \widehat{u}_1^{n,k} \\ \vdots \\ \widehat{u}_{N_x}^{n,k} \end{pmatrix}$$

an approximation of $\widehat{\mathbf{U}}^n(a^k)$, we move from age a^{k-1} to age a^k following the scheme

$$\begin{aligned} \frac{1}{\Delta t} \alpha_i^n &= \frac{1}{\Delta t} \alpha_i^{n-1} - \sigma \alpha_i^n - h \Delta a \left(\sum_{k=N_{a_1}}^{N_{a_2}} \Pi(k\Delta a) (\boldsymbol{\eta}_i^k)^\mathbf{T} \left(\frac{\widehat{\mathbf{U}}^{n-1,k} - \widehat{\mathbf{U}}^{n-1,k-1}}{\Delta a} \right) \right) \\ &\quad - \frac{\Delta a}{h} \left(\sum_{k=N_{a_1}}^{N_{a_2}} \Pi(k\Delta a) (\boldsymbol{\eta}_i^k)^\mathbf{T} \mathbb{K} \widehat{\mathbf{U}}^{n-1,k} \right) + \sigma h \Delta a \left(\sum_{k=N_{a_1}}^{N_{a_2}} (\boldsymbol{\eta}_i^k)^\mathbf{T} \mathbf{Y}^{n,k} \right), \end{aligned}$$

and

$$\begin{aligned} \frac{1}{\Delta a} \left(\widehat{\mathbf{U}}^{n,k} - \widehat{\mathbf{U}}^{n,k-1} \right) &+ \frac{1}{h^2} \mathbb{K} \left(\frac{\widehat{\mathbf{U}}^{n,k} + \widehat{\mathbf{U}}^{n,k-1}}{2} \right) + \frac{1}{\Delta t} \left(\frac{\widehat{\mathbf{U}}^{n,k} + \widehat{\mathbf{U}}^{n,k-1}}{2} \right) + \sigma \sum_{i=1}^N \alpha_i^n \mathbf{V}_i^k \\ &= \frac{1}{\Delta t} \left(\frac{\widehat{\mathbf{U}}^{n-1,k} + \widehat{\mathbf{U}}^{n-1,k-1}}{2} \right) + \sigma h \Delta a \sum_{i=1}^N \left(\sum_{j=N_{a_1}}^{N_{a_2}} (\boldsymbol{\eta}_i^j)^\mathbf{T} \mathbf{Y}^{n,j} \right) \mathbf{V}_i^k, \end{aligned}$$

with the initial conditions

$$\begin{cases} \alpha_i^0 = 0, & \forall i = 1, \dots, N, \\ \widehat{\mathbf{U}}^{0,k} = 0, & \forall k = 0, \dots, N_a, \\ \widehat{\mathbf{U}}^{n,0} = \sum_{k=0}^{N_a} \omega_k m(a^k) \widehat{\mathbf{U}}^{n-1,k}. \end{cases}$$

Here

$$\mathbf{V}_i^k := \begin{pmatrix} v_i(k\Delta a, x_1) \\ \vdots \\ v_i(k\Delta a, x_{N_x}) \end{pmatrix}, \quad \boldsymbol{\eta}_i^k := \begin{pmatrix} \eta_i(k\Delta a, x_1) \\ \vdots \\ \eta_i(k\Delta a, x_{N_x}) \end{pmatrix}, \quad \mathbf{Y}^{n,k} := \begin{pmatrix} y^n(k\Delta a, x_1) \\ \vdots \\ y^n(k\Delta a, x_{N_x}) \end{pmatrix}.$$

5.2.5 Algorithm for the observer

1. For $n = 0$: Initialization of α_i^0 ($1 \leq i \leq N$) and $\widehat{\mathbf{U}}^{0,k}$ ($k = 0, \dots, N_a$) at 0.
2. For $n = 1, \dots, N_t$:




Figure 4: Step n of the algorithm. First, α_i^n is computed using α_i^{n-1} and $\widehat{\mathbf{U}}^{n-1,k}$, for $k = N_{a_1} - 1, \dots, N_{a_2}$ (pink arrows). Next, $\widehat{\mathbf{U}}^{n,0}$ is deduced from $\widehat{\mathbf{U}}^{n-1,k}$, for $k = 0, \dots, N_a$. Finally, $\widehat{\mathbf{U}}^{n,k}$ for $k \geq 1$ is computed using α_i^n , $\widehat{\mathbf{U}}^{n,k-1}$, $\widehat{\mathbf{U}}^{n-1,k}$ and $\widehat{\mathbf{U}}^{n-1,k-1}$ (blue arrows).

- Calculate α_i^n using the values of α_i^{n-1} and $(\widehat{\mathbf{U}}^{n-1,j})_{j=N_{a_1}-1}^{N_{a_2}}$:

$$\alpha_i^n = \frac{1}{1 + \sigma \Delta t} \left(\alpha_i^{n-1} - h \Delta t \sum_{k=N_{a_1}}^{N_{a_2}} \Pi(k \Delta a) (\boldsymbol{\eta}_i^k)^\mathbf{T} (\widehat{\mathbf{U}}^{n-1,k} - \widehat{\mathbf{U}}^{n-1,k-1}) \right. \\ \left. - \frac{\Delta a \Delta t}{h} \sum_{k=N_{a_1}}^{N_{a_2}} \Pi(k \Delta a) (\boldsymbol{\eta}_i^k)^\mathbf{T} \mathbb{K} \widehat{\mathbf{U}}^{n-1,k} + \sigma h \Delta a \Delta t \sum_{k=N_{a_1}}^{N_{a_2}} (\boldsymbol{\eta}_i^k)^\mathbf{T} \mathbf{Y}^{n,k} \right)$$

- $k = 0$: Initialization of $\widehat{\mathbf{U}}^{n,0}$ using the values of $(\widehat{\mathbf{U}}^{n-1,j})_{j=0}^{N_a}$:

$$\widehat{\mathbf{U}}^{n,0} = \sum_{k=0}^{N_a} \omega_k m(k \Delta a) \widehat{\mathbf{U}}^{n-1,k}$$

- For $k = 1, \dots, N_a$, solve the linear system

$$\mathbb{A} \widehat{\mathbf{U}}^{n,k} = \widehat{\mathbf{b}}^{n,k}$$

where

$$\mathbb{A} = \left(\Delta t + \frac{1}{2} \Delta a \right) \mathbb{I} + \frac{1}{2} \frac{\Delta t \Delta a}{h^2} \mathbb{K}$$

$$\widehat{\mathbf{b}}^{n,k} = \frac{1}{2} \Delta a \left(\widehat{\mathbf{U}}^{n-1,k} + \widehat{\mathbf{U}}^{n-1,k-1} \right) + \left[\left(\Delta t - \frac{1}{2} \Delta a \right) \mathbb{I} - \frac{1}{2} \frac{\Delta t \Delta a}{h^2} \mathbb{K} \right] \widehat{\mathbf{U}}^{n,k-1} \\ - \sigma \Delta a \Delta t \sum_{i=1}^N \alpha_i^n \mathbf{V}_i^k + \sigma h (\Delta a)^2 \Delta t \sum_{i=1}^N \left(\sum_{j=N_{a_1}}^{N_{a_2}} (\boldsymbol{\eta}_i^j)^\mathbf{T} \mathbf{Y}^{n,j} \right) \mathbf{V}_i^k$$

- $\widehat{\mathbf{P}}^{n,k} = \begin{pmatrix} \widehat{p}_1^{n,k} \\ \vdots \\ \widehat{p}_{N_x}^{n,k} \end{pmatrix}$ where $\widehat{p}_i^{n,k} = \Pi(a^k) \widehat{u}_i^{n,k}$.

6 Numerical results

We present in this section some numerical results illustrating the efficiency of our state reconstruction method. All the numerical tests are two-dimensional and run on Matlab.




Figure 5: The fertility and mortality functions.

Taking $a^* = 2$, we choose the fertility function to be (see Figure 5)

$$\beta(a) = 10 a(a^* - a) \exp \left\{ -20(a - a^*/3)^2 \right\},$$

while the mortality function is chosen as

$$\mu(a) = (a^* - a)^{-1}.$$

Note that for this particular choice of μ , the function $\Pi(a)$ can be computed explicitly:

$$\Pi(a) = \exp \left(- \int_0^a \mu(s) ds \right) = \frac{a^* - a}{a^*}.$$




Figure 6: Initial state corresponding to the first eigenfunction (3D and 2D representations).

6.1 Distributed observation in space and full observation in age

We take $\Omega = (0, \pi)$ (i.e. $\ell = \pi$), and we consider full observation in age (i.e. $(a_1, a_2) = (0, a^*)$) and partial observation in space : $(\ell_1, \ell_2) = (\ell/3, 2\ell/3)$. The time of observation is $T = 2a^*$ and the diffusion coefficient is $k = 1$. Under these assumptions, there is a unique unstable eigenvalue $\lambda_1 = \lambda_1^0 - 1$, where λ_1^0 is the unique real solution of the characteristic equation (2.2). Computing numerically this value, we obtain that $\lambda_1 = 0.239$.

We first choose as initial state an eigenfunction corresponding to the (unique) unstable eigenvalue λ_1 (see Figure 6), i.e. we take

$$p_0(a, x) = \varphi_1(a, x) = \varphi_1^0(a)\varphi_1^D(x) = \frac{a^* - a}{a^*} e^{-\lambda_1^0 a} \sin(x). \quad (6.1)$$

The solution of problem (5.1) is then simply given by

$$p(a, x, t) = e^{\lambda_1 t} p_0(a, x).$$




Figure 7: Estimated (left) and exact (right) solution at time $t = T$.

To validate our reconstruction method, we first generate numerically an output by solving problem (5.1) with p_0 given by (6.1). To avoid committing an “inverse crime” [16, p. 154], we use different space discretizations to generate the measurement and in the reconstruction step.

Using $N_x = 100$ points of discretization in space, $N_a = 120$ points in discretization in age and a time step $\Delta t = \Delta a/2$, we obtain an L^2 relative error of 4.07% (see Figure 7 for a representation of the estimated and exact solutions). In this numerical test, the gain coefficient is chosen as $\sigma = 2 + \lambda_1$.

From now on, we consider the more realistic situation of noisy observation and gaussian type initial data of the form (see Figure 8)

$$p_0(a, x) = \exp \left\{ - \left(30(a - a^*/4)^2 + 20(x - \ell/4)^2 \right) \right\}.$$

Using the same discretization parameters, we obtain an L^2 relative error of 2.99% with 5% of noise, 9.6% with 10% of noise and 16.3% with 15% of noise; see Figure 9 for a representation of the estimated and exact solutions at time $t = T$ in the case of 5% of noise.




Figure 8: Gaussian initial state (3D and 2D representations).




Figure 9: Estimated (left) and exact (right) solution at time $t = T$ (with 5% of noise).

We also plot on Figure 10 the estimated and exact* final total population : $P_T(x) = \int_0^{a^*} p(a, T, x) da$ and $\hat{P}_T(x) = \int_0^{a^*} \hat{p}(a, T, x) da$.

Remark 6.1. *One might wonder why the fact that the estimated final states obtained for two different initial data (the first eigenfunction and a initial state of gaussian type) look the same. In fact, this is not surprising as the asymptotic behavior (in time) of the solution of (1.1) is given by the first eigenfunction when there is only one unstable eigenvalue.*

Surprisingly enough, the reconstruction method is efficient even when the space interval of observation (ℓ_1, ℓ_2) is small. More precisely, for $N_x = 60$, $N_a = 80$ and 10% of noise, we obtain an L^2 -relative error of 19.9% when $\ell_2 - \ell_1 = 3h$, h being the mesh size (in space). The same experiment run for $\ell_2 - \ell_1 = \ell/3$ leads to a relative error of 9.1%.

6.2 Distributed observation in space and age

We consider now the case of local observation in age by taking successively $(a_1, a_2) = (0, a^*/20)$ and $(a_1, a_2) = (a^*/2, a^*)$. The observation interval in space is chosen in both cases to be $(\ell/3, 2\ell/3)$ and we use noisy data with 5% of noise. The relative errors on the

*Here, what we refer to as the “exact” solution is a numerical solution, computed assuming the initial data to be known.




Figure 10: Estimated (dashed line) and exact total population at time $t = T$ with 5% of noise (left) and 15% of noise (right).

final state are then respectively of 0.94% and 38.8% (see Figure 11). This shows that the quality of the reconstruction is not influenced by the size of the age observation interval but mainly by its location. One can obtain a very good reconstruction provided this interval is located near the origin. This shows the crucial role played by the observation of births in the quality of the observer.




Figure 11: Estimated (dashed line) and exact total population at time $t = T$ with age observation near $a = 0$ (left) and far from $a = 0$ (right).

6.3 Influence of the observation time

In this last experiment, we consider a configuration where there are two unstable eigenvalues ($\lambda_1 = 0.959$ and $\lambda_2 = 0.119$). This can be done by reducing the value of the diffusion coefficient $k = 0.28$, and taking the same parameters as in the previous numerical test. The observation interval in space is chosen to be $(\ell/3, 2\ell/3)$ and $(a_1, a_2) = (0, a^*/20)$.

For $T = a^*$, the observer yields a good approximation of the final state (the relative error is 3.83%), and an even better approximation of the total population (the relative error is 1.81%). For an observation time $T = 0.5a^*$, the estimate provided by the observer is not very accurate, the relative error being 27%. However, we still obtain a reasonable approximation of the total population, as it can be seen from Figure 12.




Figure 12: Estimated (dashed line) and exact total population at time $t = T$, for $T = a^*$ (left) and far from $T = 0.5a^*$ (right).

References

- [1] B. AINSEBA, *Exact and approximate controllability of the age and space population dynamics structured model*, J. Math. Anal. Appl., 275 (2002), pp. 562–574.
- [2] B. AINSEBA AND M. IANNELLI, *Exact controllability of a nonlinear population-dynamics problem*, Differential Integral Equations, 16 (2003), pp. 1369–1384.
- [3] B. AINSEBA AND M. LANGLAIS, *On a population dynamics control problem with age dependence and spatial structure*, J. Math. Anal. Appl., 248 (2000), pp. 455–474.
- [4] D. AUROUX, P. BANSART, AND J. BLUM, *An evolution of the back and forth nudging for geophysical data assimilation: application to Burgers equation and comparisons*, Inverse Probl. Sci. Eng., 21 (2013), pp. 399–419.
- [5] D. AUROUX AND J. BLUM, *Back and forth nudging algorithm for data assimilation problems*, C. R. Acad. Sci. Paris Sér. I Math., 340 (2005), pp. 873–878.
- [6] D. AUROUX AND M. NODET, *The back and forth nudging algorithm for data assimilation problems: theoretical results on transport equations*, ESAIM Control Optim. Calc. Var., 18 (2012), pp. 318–342.
- [7] B. P. AYATI, *A variable time step method for an age-dependent population model with nonlinear diffusion*, SIAM J. Numer. Anal., 37 (2000), pp. 1571–1589.
- [8] B. P. AYATI AND T. F. DUPONT, *Galerkin methods in age and space for a population model with nonlinear diffusion*, SIAM J. Numer. Anal., 40 (2002), pp. 1064–1076.
- [9] M. BADRA AND T. TAKAHASHI, *Stabilization of parabolic nonlinear systems with finite dimensional feedback or dynamical controllers: application to the Navier-Stokes system*, SIAM J. Control Optim., 49 (2011), pp. 420–463.
- [10] ———, *On the Fattorini criterion for approximate controllability and stabilizability of parabolic systems*, ESAIM Control Optim. Calc. Var., 20 (2014), pp. 924–956.

- [11] J. S. BARAS, A. BENSOUSSAN, AND M. R. JAMES, *Dynamic observers as asymptotic limits of recursive filters: special cases*, SIAM J. Appl. Math., 48 (1988), pp. 1147–1158.
- [12] V. BARBU AND R. TRIGGIANI, *Internal stabilization of Navier-Stokes equations with finite-dimensional controllers*, Indiana Univ. Math. J., 53 (2004), pp. 1443–1494.
- [13] W. L. CHAN AND B. Z. GUO, *On the semigroups of age-size dependent population dynamics with spatial diffusion*, Manuscripta Math., 66 (1989), pp. 161–181.
- [14] D. CHAPELLE, N. CÎNDEA, AND P. MOIREAU, *Improving convergence in numerical analysis using observers—the wave-like equation case*, Math. Models Methods Appl. Sci., 22 (2012), pp. 1250040, 35.
- [15] M. J. CHAPMAN AND A. J. PRITCHARD, *Finite-dimensional compensators for non-linear infinite-dimensional systems*, in Control theory for distributed parameter systems and applications (Vorau, 1982), vol. 54 of Lecture Notes in Control and Inform. Sci., Springer, Berlin, 1983, pp. 60–76.
- [16] D. COLTON AND R. KRESS, *Inverse acoustic and electromagnetic scattering theory*, vol. 93 of Applied Mathematical Sciences, Springer-Verlag, Berlin, second ed., 1998.
- [17] J.-M. CORON AND E. TRÉLAT, *Global steady-state controllability of one-dimensional semilinear heat equations*, SIAM J. Control Optim., 43 (2004), pp. 549–569 (electronic).
- [18] R. CURTAIN, M. DEMETRIOU, AND K. ITO, *Adaptive observers for structurally perturbed infinite dimensional systems*, in Decision and Control, 1997., Proceedings of the 36th IEEE Conference on, vol. 1, 1997, pp. 509–514 vol.1.
- [19] R. F. CURTAIN AND H. ZWART, *An introduction to infinite-dimensional linear systems theory*, vol. 21 of Texts in Applied Mathematics, Springer-Verlag, New York, 1995.
- [20] C. CUSULIN AND L. GERARDO-GIORDA, *A numerical method for spatial diffusion in age-structured populations*, Numer. Methods Partial Differential Equations, 26 (2010), pp. 253–273.
- [21] G. DI BLASIO AND A. LORENZI, *Direct and inverse problems in age-structured population diffusion*, Discrete Contin. Dyn. Syst. Ser. S, 4 (2011), pp. 539–563.
- [22] ———, *An identification problem in age-dependent population diffusion*, Numer. Funct. Anal. Optim., 34 (2013), pp. 36–73.
- [23] H. W. ENGL, W. RUNDELL, AND O. SCHERZER, *A regularization scheme for an inverse problem in age-structured populations*, J. Math. Anal. Appl., 182 (1994), pp. 658–679.
- [24] E. FRIDMAN, *Observers and initial state recovering for a class of hyperbolic systems via Lyapunov method*, Automatica, 49 (2013), pp. 2250 – 2260.

- [25] L. GERARDO-GIORDA, *Numerical approximation of density dependent diffusion in age-structured population dynamics*, in Conference Applications of Mathematics 2013 in honor of the 70th birthday of Karel Segeth, J. Brandts, S. Korotov, M. Kryzek, J. Sístek, and T. Vejchodský, eds., vol. 2013, Institute of Mathematics AS CR, Prague, 2013, pp. 88–97.
- [26] B. Z. GUO AND W. L. CHAN, *On the semigroup for age dependent population dynamics with spatial diffusion*, J. Math. Anal. Appl., 184 (1994), pp. 190–199.
- [27] M. E. GURTIN, *A system of equations for age-dependent population diffusion*, Journal of Theoretical Biology, 40 (1973).
- [28] M. E. GURTIN AND R. C. MACCAMY, *Diffusion models for age-structured populations*, Math. Biosci., 54 (1981), pp. 49–59.
- [29] M. GYLLENBERG, A. OSIPOV, AND L. PÄIVÄRINTA, *The inverse problem of linear age-structured population dynamics*, J. Evol. Equ., 2 (2002), pp. 223–239.
- [30] G. HAINE, *Recovering the observable part of the initial data of an infinite-dimensional linear system with skew-adjoint generator*, Mathematics of Control, Signals, and Systems, 26 (2014), pp. 435–462.
- [31] G. HAINE AND K. RAMDANI, *Reconstructing initial data using observers: error analysis of the semi-discrete and fully discrete approximations*, Numer. Math., 120 (2012), pp. 307–343.
- [32] Z. HIDAYAT, R. BABUSKA, B. DE SCHUTTER, AND A. NUNEZ, *Observers for linear distributed-parameter systems: A survey*, in Robotic and Sensors Environments (ROSE), 2011 IEEE International Symposium on, sept. 2011, pp. 166–171.
- [33] W. HUYER, *Semigroup formulation and approximation of a linear age-dependent population problem with spatial diffusion*, Semigroup Forum, 49 (1994), pp. 99–114.
- [34] ———, *Approximation of a linear age-dependent population model with spatial diffusion*, Commun. Appl. Anal., 8 (2004), pp. 87–108.
- [35] M. R. JAMES AND J. S. BARAS, *An observer design for nonlinear control systems*, in Analysis and optimization of systems (Antibes, 1988), vol. 111 of Lecture Notes in Control and Inform. Sci., Springer, Berlin, 1988, pp. 170–180.
- [36] T. KATO, *Perturbation theory for linear operators*, Classics in Mathematics, Springer-Verlag, Berlin, 1995.
- [37] O. KAVIAN AND O. TRAORÉ, *Approximate controllability by birth control for a nonlinear population dynamics model*, ESAIM Control Optim. Calc. Var., 17 (2011), pp. 1198–1213.
- [38] M.-Y. KIM, *Galerkin methods for a model of population dynamics with nonlinear diffusion*, Numerical Methods for Partial Differential Equations, 12 (1996).
- [39] M.-Y. KIM AND E.-J. PARK, *Mixed approximation of a population diffusion equation*, Comput. Math. Appl., 30 (1995), pp. 23–33.

- [40] M. KRSTIC, L. MAGNIS, AND R. VAZQUEZ, *Nonlinear control of the viscous Burgers equation: Trajectory generation, tracking, and observer design*, J. Dyn. Sys., Meas., Control, 131 (2009).
- [41] M. LANGLAIS, *A nonlinear problem in age-dependent population diffusion*, SIAM J. Math. Anal., 16 (1985), pp. 510–529.
- [42] ———, *Large time behavior in a nonlinear age-dependent population dynamics problem with spatial diffusion*, J. Math. Biol., 26 (1988), pp. 319–346.
- [43] L. LOPEZ AND D. TRIGIANTE, *A finite difference scheme for a stiff problem arising in the numerical solution of a population dynamic model with spatial diffusion*, Nonlinear Anal., 9 (1985), pp. 1–12.
- [44] D. LUENBERGER, *Observing the state of a linear system*, IEEE Trans. Mil. Electron., MIL-8 (1964), pp. 74–80.
- [45] F. A. MILNER, *A numerical method for a model of population dynamics with spatial diffusion*, Comput. Math. Appl., 19 (1990), pp. 31–43.
- [46] P. MOIREAU, D. CHAPELLE, AND P. LE TALLEC, *Joint state and parameter estimation for distributed mechanical systems*, Computer Methods in Applied Mechanics and Engineering, 197 (2008), pp. 659–677.
- [47] P. MOIREAU, D. CHAPELLE, AND P. LE TALLEC, *Filtering for distributed mechanical systems using position measurements: perspectives in medical imaging*, Inverse Problems, 25 (2009), pp. 035010, 25.
- [48] G. G. PELOVSKA, *Numerical investigations in the field of age-structured population dynamics*, PhD thesis, University of Trento, 2007.
- [49] K. D. PHUNG AND G. WANG, *An observability estimate for parabolic equations from a measurable set in time and its applications*, J. Eur. Math. Soc. (JEMS), 15 (2013), pp. 681–703.
- [50] M. PILANT AND W. RUNDELL, *Determining a coefficient in a first-order hyperbolic equation*, SIAM J. Appl. Math., 51 (1991), pp. 494–506.
- [51] K. RAMDANI, M. TUCSNAK, AND G. WEISS, *Recovering the initial state of an infinite-dimensional system using observers*, Automatica, 46 (2010), pp. 1616–1625.
- [52] J.-P. RAYMOND AND L. THEVENET, *Boundary feedback stabilization of the two dimensional Navier-Stokes equations with finite dimensional controllers*, Discrete Contin. Dyn. Syst., 27 (2010), pp. 1159–1187.
- [53] A. RHANDI, *Positivity and stability for a population equation with diffusion on L^1* , Positivity, 2 (1998), pp. 101–113.
- [54] W. RUNDELL, *Determining the death rate for an age-structured population from census data*, SIAM J. Appl. Math., 53 (1993), pp. 1731–1746.
- [55] A. SMYSHLYAEV AND M. KRSTIC, *Backstepping observers for a class of parabolic PDEs*, Systems Control Lett., 54 (2005), pp. 613–625.

- [56] J. SONG, J. Y. YU, X. Z. WANG, S. J. HU, Z. X. ZHAO, J. Q. LIU, D. X. FENG, AND G. T. ZHU, *Spectral properties of population operator and asymptotic behaviour of population semigroup*, Acta Math. Sci. (English Ed.), 2 (1982), pp. 139–148.
- [57] O. TRAORE, *Null controllability of a nonlinear population dynamics problem*, Int. J. Math. Math. Sci., (2006), pp. Art. ID 49279, 20.
- [58] ———, *Null controllability and application to data assimilation problem for a linear model of population dynamics*, Ann. Math. Blaise Pascal, 17 (2010), pp. 375–399.
- [59] R. TRIGGIANI, *On the stabilizability problem in Banach space*, J. Math. Anal. Appl., 52 (1975), pp. 383–403.
- [60] ———, *Addendum: “On the stabilizability problem in Banach space” (J. Math. Anal. Appl. 52 (1975), no. 3, 383–403)*, J. Math. Anal. Appl., 56 (1976), pp. 492–493.
- [61] ———, *Boundary feedback stabilizability of parabolic equations*, Appl. Math. Optim., 6 (1980), pp. 201–220.
- [62] C. WALKER, *Some remarks on the asymptotic behavior of the semigroup associated with age-structured diffusive populations*, Monatsh. Math., 170 (2013), pp. 481–501.
- [63] G. F. WEBB, *Theory of nonlinear age-dependent population dynamics*, vol. 89 of Monographs and Textbooks in Pure and Applied Mathematics, Marcel Dekker, Inc., New York, 1985.
- [64] ———, *Population models structured by age, size, and spatial position*, in Structured population models in biology and epidemiology, vol. 1936 of Lecture Notes in Math., Springer, Berlin, 2008, pp. 1–49.
- [65] J. ZABCZYK, *Remarks on the algebraic Riccati equation in Hilbert space*, Appl. Math. Optim., 2 (1975/76), pp. 251–258.