

Enumeration in algebra

Emmanuel Jeandel

▶ To cite this version:

Emmanuel Jeandel. Enumeration in algebra. 2015. hal-01146744v1

HAL Id: hal-01146744 https://inria.hal.science/hal-01146744v1

Preprint submitted on 29 Apr 2015 (v1), last revised 4 Aug 2017 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Enumeration in algebra

Emmanuel Jeandel

28th May 2015

Abstract

We show how the concept of enumeration reducibility from computability theory may be applied to obtain computability conditions on descriptions of substructures of algebras, more prominently on the language of forbidden words of a subshift, and on the word problem for finitely generated groups. Many of the results here are already known for recursively presented groups and effectively closed subshifts, but using enumeration reducibility we are able to generalize them to arbitrary objects. The proof is based on a topological framework similar to universal algebra, and generalizes in particular work by Kuznetsov and others.

There exist finitely presented groups with an uncomputable word problem [Nov55, Boo57a, Boo57b]. However finitely presented *simple* groups have a computable word problem [BH74].

Some subshifts of finite type have an uncomputable language [Rob71]. However *minimal* subshifts of finite type have a computable language [Hoc09, BJ08].

There is a pattern here. In fact, this pattern was already observed by Kuznetsov [Kuz58]: in the vocabulary of Maltsev [Mal61, Theorem 4.2.2]), every simple finitely presented algebra is constructive.

The goal of this paper is to reprove this statement using the vocabulary of topology rather than algebras. We will in particular be more general and drop the requirement that the algebra is finitely presented, or even recursively presented. We wil prove for example that, for a (finitely generated) simple group, the complement of the word problem is enumeration reducible to the word problem. This implies the previous result: a finitely presented group has a recursively enumerable word problem, thus the complement of the word problem of a simple f.p. group, being enumeration reducible to a recursively enumerable set, is recursively enumerable, and thus the word problem is recursive.

As evidenced in the previous paragraph, the key concept in this article is the notion of enumeration reducibility [FR59, Odi99]. Roughly speaking, a set A is enumeration reducible to a set B, if an enumeration of A can be obtained from any enumeration of B. We will see that this concept is of great interest in algebra for objects given by presentations, like groups (note that the notion of presentation we use here is the notion from group theory, which is different from the notion of presentation in computable algebra). We know for example that for every enumeration degree d, there exists a finitely generated group for which the word problem has enumeration degree d (the reduction is actually stronger, Dobritsa [Bel74, Theorem 2.4], see also [Zie76]) or that enumeration reducibility characterizes when a group G can be embedded into a group that is finitely presentend over a group H (C.F. Miller, see [HS88, Chapter 6]). We note in passing that various other reductions have been used in conjunction with algebraic objects, in particular quasi-reducibility [DLN98] and Ziegler-reducibility [Zie80]. However many of our theorems have converses (see in particular Theorem 4) which suggest enumeration-reducibility is indeed the right notion in our context.

The article is organized as follows: The set of all substructures of a given structure can be seen as a topological space, which is also a complete semilattice. In the case of finitely generated groups, this space is known as the space of marked groups [Gri85]. We call this space a quasivariety, by similarity with the corresponding notion from universal algebra. For groups, the points of the quasivariety V are (marked) groups, seen as normal subgroups of the free group. For shift spaces, the points of the quasivariety V correspond to subshifts, seen as factorial extensible languages of finite words.

A presentation of a point X is any set R so that X is the smallest point of V that contains R. The next section introduces this concept and show computability relations between R and X.

The next three sections introduce various way of limiting the power of a point X in a quasivariety, and the consequence in terms of computability. The first one correspond to the concept of simple group, or minimal subshifts, and is coined "maximal points". The next concept is borrowed from group theory [dCGP07] and provides a complete characterization of which objects are computable.

It is quite clear from the previous paragraphs that groups and subshifts are central examples of the theory. They will be running examples in all sections below. A third example from commutative algebra is also given, but less developed.

1 Definitions

Let I be a computable countable set, that we identify with the set of integers. In applications, I will be the set of words over a given finite alphabet, or the set of words in a free group.

In this article, we will always identify a subset $X \subseteq I$ and a point $x \in \{0, 1\}^I$.

We are interested in subsets $X \subseteq I$ that can be defined by some Horn formulas, i.e. by axioms of the type:

$$a \in X \land b \in X \land .. \land c \in X \implies z \in X$$

In the vocabulary of Higman, these are called *identical implications*.

In what follows, we will be given such a collection of formulas, and we will look at the set of all X that satisfies such a collection.

Definition 1.1. Let S be a recursive enumeration of finite sequences of elements of I.

A word $x \in \{0,1\}^I$ satisfies S if for all $(n_0, n_1, \dots n_k) \in S$,

$$x_{n_1} = 1 \land x_{n_2} = 1 \land \dots \land x_{n_k} = 1 \implies x_{n_0} = 1$$

Equivalently, a set $X \subseteq I$ satisfies S if for all $(n_0, n_1, \ldots n_k) \in S$

$$n_1 \in X \land n_2 \in X \land n_k \in X \implies n_0 \in X$$

The quasivariety V defined by S is the set of all words x (or all subsets $X \subseteq I$) that satisfy S.

The term quasivariety is used due to the similary between this notion and quasivariety in universal algebra.

The fact that the sequence S can be recursively enumerated is not mandatory, but happens in all interesting examples. This assumption can be dropped in almost all theorems, to obtain relativized versions of the theorems, by replacing all statements of the form $X \leq_e Y$ by $X \leq_e Y \oplus S$.

Example 1.1. Let Σ be a finite alphabet. A subset Y of $\Sigma^{\mathbb{Z}}$ is called a subshift [LM95] if it is topologically closed and invariant under translation. Y is entirely characterized by the set X of all finite words that do not appear in any word of Y.

In this way, the set of all subshifts over Σ is a quasivariety: A set X of words over Σ is (the forbidden language of) a subshift if it is extensible and factorial, that is:

- For any letter a, if $w \in X$ then $aw \in X$
- For any letter a, if $w \in X$ then $wa \in X$
- If $wa \in X$ for all letters a, then $w \in X$.
- If $aw \in X$ for all letters a, then $w \in X$.

The quasivariety V of subshifts contains two particular points: the point $X = \emptyset$ (which corresponds to the subshift $Y = \Sigma^{\mathbb{Z}}$) and the point $X = \Sigma^{\star}$ (which corresponds to the subshift $Y = \emptyset$)

Example 1.2. Let n be an integer. The set of all groups with n generators may be seen as a quasivariety. Indeed, such a group G can be seen (up to isomorphism) as a quotient of the free group \mathbb{F}_n , or equivalently as a normal subgroup R of \mathbb{F}_n (the subgroup R corresponds to the word problem of G, i.e. all combinations of generators of G that are equal to the identity).

Indeed, a set $X \subseteq \mathbb{F}_n$ is a normal subgroup of \mathbb{F}_n (i.e. codes a group) if:

- If (nothing) then $1 \in X$
- If $g \in X$ then $g^{-1} \in X$
- If $g \in X$, $h \in X$ then $gh \in X$

• For any h, if $g \in X$ then $hgh^{-1} \in X$

For example, for $\mathbb{Z}^2 = \{a, b | ab = ba\}$, we have $a \notin X$, $b \notin X$ but $aba^{-1}b^{-1} \in X$, $aaba^{-1}b^{-1}a^{-1} \in X$, and more generally, X is exactly the set of words of the free group for which the number of occurences of a is equal to the number of occurences of a^{-1} , and the same for b and b^{-1} .

The quasivariety V of groups contains two particular points: the point $X = \{1\}$ (which corresponds to the group $G = \mathbb{F}_n$) and the point $X = \mathbb{F}_n$ (which corresponds to the one-element group).

This particular quasivariety is usually called the space of marked groups, see Grigorchuck [Gri85].

Example 1.3. The set of all ideals of $\mathbb{Q}[t]$ is a quasivariety (once some recursive coding of $\mathbb{Q}[t]$ into I is given)

Indeed, a set $X \subseteq \mathbb{Q}[t]$ is an ideal if

- If (nothing) then $0 \in X$
- If $a, b \in X$ then $a + b \in X$
- If $a \in X$ then $-a \in X$
- For all f, if $a \in X$ then $fa \in X$

The same can be done for any commutative ring as long as it is computable in some sense.

It is easy to see from the definitions that V can be given the structure of a topological space, by inheriting the natural (product/Tychonoff) topology on $\{0,1\}^I$. As S is chosen to be computably enumerated, V is what is called a Π_1^0 set in computability theory [CR98, CR15]

It has also the structure of a complete semi-lattice, as evidenced by the following easy facts:

Fact 1.2. Let V be a quasivariety. Any intersection of elements of V is again in V In particular:

- V contains a minimal element, the intersection of all elements of V
- V contains a maximal element, the set I itself.
- For any set $Y \subseteq I$ there exists a smallest element X of V that contains Y.

2 Presentations

Definition 2.1. A presentation of $X \in V$ is a set Y so that X is the smallest element of V containing Y. X is finitely presented if it admits a finite presentation. X is recursively presented if it admits a recursively enumerable presentation.

Example 2.1. A finitely(resp. recursively) presented group is exactly what is usually called a finitely(resp. recursively) presented group.

A finitely presented subshift is usually called a subshift of finite type. A recursively presented subshift is usually called an effectively closed subshift.

A finitely presented ideal is usually called a finitely generated ideal.

Presentations, and in fact everything below, is related to the notion of enumeration reducibility, that we define now:

Definition 2.2. Let A, B two subsets of I.

We say that A is enumeration reducible to B, written $A \leq_e B$, if there exists a computable function $f: I \times \mathbb{N} \to P_f(I) \cup \{\{\bot\}\}$, where $P_f(I)$ is (a recursive encoding of) the finite sets of I and $\bot \notin I$, so that

$$x \in A \iff \exists n, f(x, n) \subseteq B$$

When we want to emphasize the computable function f, we will say that A is enumeration-reducible via f, written $A \leq_e^f B$. (Note that A is uniquely defined given f and B).

The following statement says that a point X in a quasivariety is enumerationreducible to any of its presentations. Thus any point has a smallest presentation (in terms of enumeration reducibility), which is X itself.

Proposition 2.3. Let V be a quasivariety.

Let Y be any presentation of X, then $X \leq_e Y$.

More precisely, there exists a computable function f (depending only on V) so that $X \leq_e^f Y$ iff Y is a presentation of X.

In particular a finitely/recursively presented point is recursively enumerable (as a subset of I).

(Finitely/recursively presented groups have a computably enumerable word problem. Subshifts of finite type and effectively closed subshifts have a computably enumerable set of forbidden words)

Proof. Let V be defined by a set S, and S' be the *closure* of S, i.e. all statements of the form

$$a \in X \land b \in X \land .. \land c \in X \implies z \in X$$

that are logical consequences of statements of S.

It is easy to see that S' is also computably enumerable, and of course define the same variety V. Note that S' contains in particular all implications of the form " $a \in X \implies a \in X$ ".¹

On input a, consider all finite sets F s.t. " $\bigwedge_{i \in F} i \in X \implies a \in X$ " is a statement of S'. All such finite sets can be enumerated, thus there is a computable function $f: I \times \mathbb{N} \to P_f(I)$ such that $f(a, \mathbb{N})$ is exactly the set of all F.

Then it is clear that $A \leq_e^f B$ iff B is a presentation of A.

It is interesting to note that most of the statements we obtain have a converse, and this is the case here: A is enumeration reducible to B, iff B is a presentation of A in a suitable quasivariety.

Proposition 2.4. Let $B \subseteq A$ so that $A \leq_e B$. Then there exists a quasivariety V such that B is a presentation of A.

Proof. Let $A \leq_e^f B$.

Let V be the quasivariety defined by :

• If $f(n, x) \subseteq X$ then $x \in X$.

V is by definition a quasivariety. And it is clear by definition that the smallest point of V that contains B is A.

3 Maximal elements

Definition 3.1. An element X of a quasivariety V is maximal if $X \subseteq Y$, with $Y \in V$, implies Y = X or Y = I.

Example 3.1. In the language of symbolic dynamics, this corresponds to minimal subshifts. In the language of groups, this corresponds to simple groups. In the language of ideals, this corresponds to a maximal ideal. In all three examples, X is implicitly supposed to be different from I.

Our first (easy) theorem generalizes the theorem of Kuznetsov, without any hypothesis on the computability of the structure:

Theorem 1. Let V be a quasivariety. If X is maximal then $\overline{X} \leq_e X$.

 $(\overline{X} \text{ is the complement of } X).$

¹It is easy to see that (if we drop the hypothesis for the base set S to be computably enumerated) the set of all quasivarieties V may be given itself the structure of a quasivariety VV, where each quasivariety $V \in VV$ is identified with the set S(V) of all implications true in any point of the quasivariety. In this identification, S(V) = S', and S above is just a particular presentation of V.

Proof. If X = I it is clear. Otherwise, let $a \notin X$.

Then $x \in \overline{X}$ iff the smallest point containing both X and x contains a. Recall there is a function f so that $A \leq_e^f B$ iff B is a presentation of A. Thus $x \in \overline{X}$ iff there exists n so that $f(n, a) \subseteq X \cup \{x\}$. Let $g(n, x) = f(n, a) \setminus \{x\}$. Then $x \in \overline{X}$ iff there exists n so that $g(n, x) \subseteq X$.

Corollary 3.2. Let V be a quasivariety. If X is finitely (or recursively) presentend and maximal, then X is computable.

Indeed, X is recursively enumerable, as X is finitely presented, and \overline{X} is recursively enumerable, as it is enumeration reducible to X.

Theorem 2 (Uniform version). Let V be a quasivariety so that I (the whole set) is finitely presented. Then there exists a computable function g so that if X is maximal, $X \neq I$, then $\overline{X} \leq_e^g X$.

Proof. By definition, there exists a finite set A s.t. any point containing all of A is equal to I. Say $A = \{a_1 \dots a_k\}$.

Thus $x \in \overline{X}$ iff for all $a \in A$, there exists n so that $f(n, a) \subseteq X \cup \{x\}$ Let $g(n_1, n_2 \dots n_k, x) = \bigcup_{i \leq k} f(n_i, a_i) \setminus \{x\}$. Then $x \in \overline{X}$ iff there exists $n_1 \dots n_k$ so that $g(n_1, n_2 \dots n_k, x) \subseteq X$.

Corollary 3.3. Let G be a finitely generated simple group. Then the complement of the word problem of G is enumeration reducible to the word problem of G. In particular [BH74], if G is a finitely generated, finitely presented, simple group, then G has a computable word problem.

Let S be a minimal subshift. Then the set of words that appear in S is enumeration-reducible to the set of words that do not appear in S. In particular [Hoc09, BJ08], if S is a minimal subshift of finite type, then S has a computable set of forbidden words.

In both cases, the problem is uniformly solvable: there exists an algorithm that, given a finite presentation of G (resp. S) decides the word problem of G (resp. the set of admissible words of S).

Theorem 3. Let $(S_n)_{n \in \mathbb{N}}$ be a computable collection of finite sets, and $(Y_n)_{n \in \mathbb{N}}$ be the points presented by S_n . Let $\mathcal{Y} = \{Y_n, n \in \mathbb{N}\}$

We say that X is maximal below \mathcal{Y} if $X \notin \mathcal{Y}$, but every point larger than X is in \mathcal{Y} .

Then $\overline{X} \leq_e X$, uniformly.

An example of such a collection S_n is the set of all finite groups with k generators: A group G with generators $a_1, \ldots a_k$ is finite iff there exists a size p s.t. all words of length p over the alphabet $\{a_1^{\pm 1} \ldots a_k^{\pm 1}\}$ are equal to a word of smaller length.

We can use this to obtain a recursive list of finite presentations that contain only finite groups, and all presentations of finite groups arise this way. Thus the theorem apply to all groups for which all normal subgroups are of finite index (these groups are called *just infinite groups*).

More generally, every finitely generated and finitely presented algebra on which all congruences have finite index is computable [Mal61].

Proof. Straightforward generalization of the previous theorem.

Let $(S_n)_n$ be the computable collection of finite sets, and write $S_n = \{a_1^n, \ldots, a_{h(n)}^n\}$, where h(n) is the (computable) size of S_n .

Recall there is a function f so that $A \leq_e^f B$ iff B is a presentation of A.

Then $x \in \overline{X}$ iff the smallest point (or any point) containing $\overline{X} \cup \{x\}$ is one of the sets of \mathcal{Y} .

Thus $x \in \overline{X}$ iff $\exists p, \exists n_1 \dots n_{h(p)}, \cup_{i \leq h(p)} f(n_i, a_i^p) \setminus \{x\} \subseteq X$

As before, it is interesting to note that the main theorem has a converse:

Theorem 4. Let $\overline{A} \leq_e A$. Then there exists a quasivariety V s.t. A is maximal.

Proof. Let $\overline{A} \leq_e^f A$.

Thus $x \in \overline{A}$ iff there exists n s.t. $f(n, x) \subseteq A$. Let V be the quasivariety defined by

• For all n, for all $x \in X$, for all $y \in I$, if $f(n, x) \subseteq X$ then $y \in X$.

A is in this variety: Indeed, there is no $x \in A$ and n so that $f(n, x) \subseteq A$, thus all premises are false.

It is clearly maximal: Let $A \subseteq A'$ and $a \in A' \setminus A$. Then $a \in \overline{A}$ thus there exists n s.t. $f(n, x) \subseteq A \subseteq A'$ thus for all $y, y \in A'$, thus A' = I.

4 Discriminable points

Discrimination is a generalization of maximal elements. The concept and the vocabulary comes from group theory, in particular [dCGP07].

Definition 4.1. Let V be a quasivariety.

A set Y is a discriminator for a point X if $Y \cap X = \emptyset$ and for every point X', $X \subseteq X'$, $X' \cap Y \neq \emptyset$.

X is finitely discriminable if it admits a finite discriminator. X is recursively discriminable if it admits a recursively enumerable discriminator.

Theorem 5. If X is recursively discriminable, then $\overline{X} \leq_e X$.

Proof. Let Y be the discriminator.

Recall again that there exists a computable function f s.t. $A \leq_e^f B$ iff B is a presentation of A

Now $x \in \overline{X}$ iff the point presented by $X \cup \{x\}$ contains some element of Y. Thus $x \in \overline{X} \iff \exists y \in Y, \exists n \in N, f(n, y) \subseteq X \cup \{x\}.$ Let $Y = (y_m)_{m \in \mathbb{N}}$ a computable enumeration of Y. We now define $g(n, m, x) = f(n, y_m) \setminus \{x\}$. Then $x \in \overline{X} \iff \exists n, m, g(n, m, x) \subseteq X$.

Corollary 4.2. Let V be a quasivariety and X a point of V.

X is computable iff it is recursively presented and recursively discriminated.

See [dCGP07] for the result for groups.

Proof. If X is computable, then X is a presentation of X which is recursive, and \overline{X} is a discriminator for X which is recursive.

Conversely, if X is recursively presented by Y, then X is recursively enumerable, as $X \leq_e Y$. As X is recursively discriminated, $\overline{X} \leq_e X$, thus \overline{X} is recursively enumerable, and X is computable.

5 Difference between X and \overline{X}

Many of the previous results are of the same form: from an enumeration of X we can compute an enumeration of \overline{X} . It is easy to provide examples where the converse is not true. Hence X gives more information that \overline{X} . Here we try to quantify what is this information.

Definition 5.1. Let $\overline{X} \leq_e^f X$. Let $g(x) = \min\{n \in \mathbb{N} | f(n, x) \subseteq X\}$. g is a partial map. We identify g with $G = \{(x, g(x)) | g(x) \text{ is defined}\}$

Proposition 5.2. Let $\overline{X} \leq_e^f X$. Then $G \leq_e X$.

Proof. Suppose we are given an enumeration of X, and at some point we conclude that $x \in \overline{X}$ because $f(n, x) \subseteq X$ for some n, and we have enumerated all elements of f(n, x). Then we know that $g(x) \leq n$. We then look at all sets f(i, x) for i < n. At some point in our enumeration, we will know the status of all points in $\bigcup_{i < n} f(i, x)$, either because they were enumerated in X, or we were able to prove that they are in \overline{X} . Thus we will be able to determine the exact value of g(x).

Proposition 5.3. Let h be a total function that is greater than $g: h(x) \ge g(x)$ whenever x is defined.

We identify h with the total set $H = \{(x, h(x)), x \in I\}$ Then $X \leq_e H \oplus \overline{X}$

Proof. $x \in X \iff \forall n \le h(x), f(n, x) \cap \overline{X} \ne \emptyset.$

The propositions are particularly interesting when we start from a quasivariety where I is finitely presented, so that for all maximal elements $X, \overline{X} \leq_e^f X$ for a given computable function f independent from X. In that case, it is possible to interpret the map g. We start with subshifts. A minimal subshift S has a quasiperiodicity function (also called uniform recurrence function): There exists a function g s.t. every word w of size n that appear in S is contained in every word of size g(n)that appear in S. For minimal subshifts, the previous theorems may thus be interpreted this way:

- The set of words that appear can be obtained from an enumeration of the set of words that do not appear
- The quasiperiodicity function can be obtained from an enumeration of the set of words that do not appear
- The set of words that do not appear can be obtained from an enumeration of both the quasiperiodicity function and the set of words that appear

We now introduce a similar concept for groups. Let S be a finitely generated, simple group, with generators $a_1 \ldots a_k$. For $w \in S, w \neq 1$, let g(w) be the smallest n s.t. all generators a_i can be written as products of less than n elements of the form hwh^{-1} or $hw^{-1}h^{-1}$ for $h \in S$.

g is well defined: Indeed the set of all elements that can be written this way is a normal subgroup of S that is nontrivial (it contains w), and thus is equal to S.

Note that g depends on the choices of generators of S, but it is easy to see that different choices of G only changes the function g up to a linear factor.

We now introduce another related function. Let S be a finitely generated, simple group, with generators $a_1 \ldots a_k$. For a word w over the generators $a_1 \ldots a_k$, let $g_1(w)$ be the smallest p s.t. all generators a_i can be written as products of less than p elements of the form hwh^{-1} or $hw^{-1}h^{-1}$ for $h \in S$, and each h is a product of less than p generators. $g_1(w)$ is defined only when w is not the identity element on S.

We now define $g'_1(n) = \max\{g_1(w)|w \in B_n, w \neq 1\}$, where B_n is the set of elements of G that can be written as a product of less than n generators.

Then, in S:

- The complement of the word problem on S can be enumerated from an enumeration of the word problem on S.
- The function g_1 can be computed from an enumeration of the word problem on S.
- The word problem on S can be enumerated from both the complement of the word problem on S and any bound t on g_1 . Indeed w = 1 iff there exists a generator a s.t. all products of less than t(w) terms of the form hwh^{-1} , where each h is the product of less than t(w) generators, are different from a.

Open Questions

This article presents how a few results in group theory and symbolic dynamics may be related once seen in the concept of universal algebra, and how they can be generalized for structures that are not recursively presented.

There exist other theorems which offer a striking similarity, but for which a general statement is not known, most proeminently Higman's embedding theorem and Boone-Higman's theorem. We concentrate here the discussion on the former theorem (The author claims he has a proof of an equivalent of the Boone-Higman theorem for subshifts, which will be found in a later paper).

Theorem 6 ([Hig61]). A finitely generated group can be embedded in a finitely presented groups iff it has a recursively enumerable set of defining relations.

Theorem 7 ([Kle52, CV58]). An arbitrary theory (with identity) is finitely axiomatisable using additional predicates iff it is recursively axiomatisable.

Theorem 8 ([Hoc09], see also [AS13, DRS10]). A symbolic system is isomorphic to the subaction of a sofic shift iff it is effectively closed.

The note by [Kuz58] also suggests an analog for universal algebras, and a similar theorem for semigroups also exist. Note also that the Relative Higman Embedding Theorem [HS88] also has an equivalent in the domain of subshifts [AS09].

Proofs of these theorems are tremendously combinatorial, as each proof needs to embed a Turing machine (or another computational device) into an algebraic system, and the methods to do this are quite different. However the fact remains that all these theorems have similar hypotheses and conclusions, so that either it is a striking coincidence, or something deep can be found here.

Acknowledgements

The author thanks Dmitry Sokolov for helping him with the translation of [Kuz58], and Yves de Cornulier for valuable comments on a first draft of this article.

References

- [AS09] Nathalie Aubrun and Mathieu Sablik. An order on sets of tilings corresponding to an order on languages. In 26th International Symposium on Theoretical Aspects of Computer Science, STACS 2009, February 26-28, 2009, Freiburg, Germany, Proceedings, pages 99– 110, 2009.
- [AS13] Nathalie Aubrun and Mathieu Sablik. Simulation of effective subshifts by two-dimensional subshifts of finite type. Acta Applicandae Mathematicae, 2013.

- [Bel74] O.V. Belegradek. Algebraically Closed Groups. Algebra and Logic, 13(3):135–143, May 1974.
- [BH74] W.W. Boone and G. Higman. An algebraic characterization of groups with soluble word problem. *Journal of the Australian Mathematical Society*, 18(1):41–53, August 1974.
- [BJ08] Alexis Ballier and Emmanuel Jeandel. Tilings and Model Theory. In Symposium on Cellular Automata Journées Automates Cellulaires (JAC), pages 29–39, Moscow, 2008. MCCME Publishing House.
- [Boo57a] William W. Boone. Certain Simple, Unsolvable Problems of Group Theory V. Indagationes Mathematicae, 60:22–27, 1957.
- [Boo57b] William W. Boone. Certain Simple, Unsolvable Problems of Group Theory VI. Indagationes Mathematicae, 60:227–232, 1957.
- [CR98] Douglas Cenzer and Jeffrey B. Remmel. Π_1^0 classes in mathematics. In Handbook of Recursive Mathematics - Volume 2: Recursive Algebra, Analysis and Combinatorics, volume 139 of Studies in Logic and the Foundations of Mathematics, chapter 13, pages 623–821. Elsevier, 1998.
- [CR15] Douglas Cenzer and Jeffrey B. Remmel. Effectively Closed Sets. ASL Lecture Notes in Logic, 2015. in preparation.
- [CV58] W. Craig and R. L. Vaught. Finite Axiomatizability Using Additional Predicates. *The Journal of Symbolic Logic*, 23(3):289–308, September 1958.
- [dCGP07] Yves de Cornulier, Luc Guyot, and Wolfgang Pitsch. On the isolated points in the space of groups. *Journal of Algebra*, 307:254–277, 2007.
- [DLN98] R. Downey, G. LaForte, and A. Nies. Computably enumerable sets and quasi-reducibility. Annals of Pure and Applied Logic, 95(1-3):1– 35, 1998.
- [DRS10] Bruno Durand, Andrei Romashchenko, and Alexander Shen. Effective Closed Subshifts in 1D Can Be Implemented in 2D. In *Fields of Logic and Computation*, number 6300 in Lecture Notes in Computer Science, pages 208–226. Springer, 2010.
- [FR59] Richard M. Friedberg and Hartley Rogers. Reducibility and Completeness for Sets of Integers. Zeitschrift für mathematische Logik und Grundlagen der Mathematik, 5:117–125, 1959.
- [Gri85] R. I. Grigorchuk. Degrees of Growth of Finitely Generated Groups, and the Theory of Invariant Means. *Mathematics of the USSR-Izvestiya*, 25(2):259–300, 1985.

- [Hig61] G. Higman. Subgroups of Finitely Presented Groups. Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences, 262(1311):455–475, August 1961.
- [Hoc09] Michael Hochman. On the dynamics and recursive properties of multidimensional symbolic systems. *Inventiones Mathematicae*, 176(1):2009, April 2009.
- [HS88] Graham Higman and Elizabeth Scott. *Existentially Closed Groups*. Oxford University Press, 1988.
- [Kle52] S.C. Kleene. Two Papers on the Predicate Calculus., chapter Finite Axiomatizability of Theories in the Predicate Calculus Using Additional Predicate Symbols. Number 10 in Memoirs of the American Mathematical Society. 1952.
- [Kuz58] A.V. Kuznetsov. Algorithms as operations in algebraic systems. Uspekhi Matematicheskikh Nauk, 13(3(81)):240–241, 1958. (in russian).
- [LM95] Douglas A. Lind and Brian Marcus. An Introduction to Symbolic Dynamics and Coding. Cambridge University Press, New York, NY, USA, 1995.
- [Mal61] A.I. Mal'tsev. Constructive Algebras I. Russian Mathematical Surveys, 16(3):77–129, 1961.
- [Nov55] P.S. Novikov. On the algorithmic unsolvability of the word problem in group theory. *Trudy Mat. Inst. Steklov*, 44:3–143, 1955.
- [Odi99] P.G. Odifreddi. Classical Recursion Theory Volume II, volume 143 of Studies in Logic and The Foundations of Mathematics. North Holland, 1999.
- [Rob71] Raphael M. Robinson. Undecidability and Nonperiodicity for Tilings of the Plane. *Inventiones Mathematicae*, 12(3):177–209, 1971.
- [Zie76] Martin Ziegler. Gruppen mit vorgeschriebenem Wortproblem. Mathematische Annalen, 219(1):43–51, 1976.
- [Zie80] Martin Ziegler. Algebraisch abgeschlossene gruppen. In Sergei I. Adian, William W. Boone, and Graham Higman, editors, Word Problems II, volume 95 of Studies in Logic and the Foundations of Mathematics, pages 449–576. North Holland, 1980.