



HAL
open science

Quadratic LYM-type inequalities for intersecting Sperner families

Christian Bey

► **To cite this version:**

Christian Bey. Quadratic LYM-type inequalities for intersecting Sperner families. 2005 European Conference on Combinatorics, Graph Theory and Applications (EuroComb '05), 2005, Berlin, Germany. pp.37-40, 10.46298/dmtcs.3418 . hal-01184375

HAL Id: hal-01184375

<https://inria.hal.science/hal-01184375v1>

Submitted on 17 Aug 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Quadratic LYM-type inequalities for intersecting Sperner families

Christian Bey

Fakultät für Mathematik, Otto-von-Guericke-Universität, Universitätsplatz 2, 39106 Magdeburg, Germany
email: christian.bey@mathematik.uni-magdeburg.de

Let $\mathcal{F} \subseteq 2^{[n]}$ be an intersecting Sperner family (i.e. $A \not\subseteq B$, $A \cap B \neq \emptyset$ for all $A, B \in \mathcal{F}$) with profile vector $(f_i)_{i=0 \dots n}$ (i.e. $f_i = |\mathcal{F} \cap \binom{[n]}{i}|$). We present quadratic inequalities in the f_i 's which sharpen the previously known linear LYM-type inequalities.

Keywords: Sperner family, antichain, LYM inequality

Let $[n]$ denote the set $\{1, 2, \dots, n\}$, $2^{[n]}$ the powerset of $[n]$, and $\binom{[n]}{i}$ the set of all i -element subsets of $[n]$. Given a set system $\mathcal{F} \subseteq 2^{[n]}$, put $f_i = |\mathcal{F} \cap \binom{[n]}{i}|$ and call $(f_i)_{i=0 \dots n}$ the *profile vector* of \mathcal{F} .

Recall that a set $\mathcal{F} \subseteq 2^{[n]}$ is called a *Sperner family* (or *antichain*) if there are no inclusions among the members of \mathcal{F} : $A \not\subseteq B$ for all $A, B \in \mathcal{F}$. The well-known theorem by Sperner (cf. [2]) says that the maximum size of an antichain in $2^{[n]}$ is $\binom{n}{\lfloor n/2 \rfloor}$, the only families achieving this cardinality being $\binom{[n]}{\lfloor n/2 \rfloor}$ and $\binom{[n]}{\lceil n/2 \rceil}$.

A generalization of Sperner's Theorem is the well-known *LYM inequality* (cf. [2]): If $\mathcal{F} \subseteq 2^{[n]}$ is a Sperner family with profile vector $(f_i)_{i=0 \dots n}$ then

$$\sum_{i=0}^n \frac{f_i}{\binom{n}{i}} \leq 1.$$

In [1] we have sharpened the LYM inequality by adding to its LHS all possible products of fractions $f_i / \binom{n}{i}$ with suitable coefficients. A special case of this sharpened version is the following *quadratic LYM inequality (QLYM)*:

Theorem 1 ([1]) *Let $\mathcal{F} \subseteq 2^{[n]}$ be a Sperner family with profile vector $(f_i)_{i=0 \dots n}$. Then*

$$\sum_{0 \leq i \leq n} \frac{f_i}{\binom{n}{i}} + \sum_{0 < i < j < n} \frac{n(j-i)}{i(n-j)} \frac{f_i}{\binom{n}{i}} \frac{f_j}{\binom{n}{j}} \leq 1.$$

The proof of Theorem 1 was based on the well-known Kruskal-Katona Theorem (cf. [2]) and induction on n .

Here we continue these investigations by presenting such sharpened LYM-type inequalities for the class of *intersecting Sperner families*. Recall that a set system $\mathcal{F} \subseteq 2^{[n]}$ is called *intersecting* if $A \cap B \neq \emptyset$ for all $A, B \in \mathcal{F}$.

The following LYM-type inequality for intersecting Sperner families $\mathcal{F} \subseteq 2^{[n]}$ was found in [3]:

$$\sum_{i \leq n/2} \frac{f_i}{\binom{n}{i-1}} + \sum_{j > n/2} \frac{f_j}{\binom{n}{j}} \leq 1. \quad (1)$$

We have the following sharpening:

Theorem 2 *Let $\mathcal{F} \subseteq 2^{[n]}$ be an intersecting Sperner family with profile vector $(f_i)_{i=0..n}$. Put*

$$c_{i,j} = \begin{cases} \frac{n(j-i)}{(i-1)j} & \text{if } i < j \leq n/2 \\ \frac{n(j-i)}{(n-i+1)(n-j)} & \text{if } n/2 < i < j \\ \frac{n(n+1-i-j)}{(i-1)j} & \text{if } i \leq n/2 < j, i+j \leq n+1 \\ \frac{n(i+j-n-1)}{(n-i+1)(n-j)} & \text{if } i \leq n/2 < j, i+j \geq n+1. \end{cases}$$

Then

$$\sum_{i \leq n/2} \frac{f_i}{\binom{n}{i-1}} + \sum_{j > n/2} \frac{f_j}{\binom{n}{j}} + \sum_{i < j} c_{i,j} \frac{f_i}{\binom{n}{i-1}} \frac{f_j}{\binom{n}{j}} \leq 1.$$

The proof of Theorem 2 is based on the previous one and the following connection between profile vectors of Sperner - and intersecting Sperner families.

Lemma 1 *Let $(f_i)_{i=0..n}$ be the profile vector of an intersecting Sperner family in $2^{[n]}$. Then $(g_i)_{i=0..n}$ defined by*

$$g_i = \begin{cases} f_{i+1} + f_{n-i} & \text{if } i < \frac{n-1}{2} \\ f_{i+1} & \text{if } i = \frac{n-1}{2} \\ 0 & \text{if } i > \frac{n-1}{2} \end{cases}$$

is profile vector of a Sperner family in $2^{[n]}$.

Besides (1), other LYM-type inequalities for intersecting Sperner families are known. In fact, in [4, 5] the convex hull of all profile vectors of such set systems was determined. In our talk we will also address the question on sharpening all facet inequalities of this convex hull.

References

- [1] C. Bey, Polynomial LYM inequalities. *Combinatorica* 25 (2005), 19-38.
- [2] B. Bollobás, *Combinatorics*. Cambridge Univ. Press, Cambridge 1986.
- [3] C. Greene, Gy.O.H. Katona and D.J. Kleitman, Extensions of the Erdős-Ko-Rado theorem. *SIAM Stud. Appl. Math* 55 (1976), 1-8.
- [4] P.L. Erdős, P. Frankl and Gy.O.H. Katona, Intersecting Sperner families and their convex hulls. *Combinatorica* 4 (1984), 21-34.

- [5] Gy.O.H. Katona and G. Schild, Linear inequalities describing the class of intersecting Sperner families of subsets, in: R. Bodendiek and R. Henn, eds, *Topics in Combinatorics and Graph Theory* (Physica-Verlag, Heidelberg, 1990), 413-420.

