



**HAL**  
open science

## Trees with product-form random weights

Konstantin Borovkov, Vladimir Vatutin

► **To cite this version:**

Konstantin Borovkov, Vladimir Vatutin. Trees with product-form random weights. Fourth Colloquium on Mathematics and Computer Science Algorithms, Trees, Combinatorics and Probabilities, 2006, Nancy, France. pp.423-426, 10.46298/dmtcs.3491 . hal-01184695

**HAL Id: hal-01184695**

**<https://inria.hal.science/hal-01184695v1>**

Submitted on 17 Aug 2015

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Recursive random trees with product-form random weights

Konstantin Borovkov<sup>1†</sup> and Vladimir Vatutin<sup>2‡</sup>

<sup>1</sup>Department of Mathematics and Statistics, The University of Melbourne, Parkville 3010, Australia

<sup>2</sup>Department of Discrete Mathematics, Steklov Mathematical Institute, Gubkin St. 8, 119991 Moscow, Russia

We consider growing random recursive trees in random environment, in which at each step a new vertex is attached according to a probability distribution that assigns the tree vertices masses proportional to their random weights. The main aim of the paper is to study the asymptotic behavior of the mean numbers of outgoing vertices as the number of steps tends to infinity, under the assumption that the random weights have a product form with independent identically distributed factors.

**Keywords:** random recursive trees, random environment, Sptizer's condition, distance to the root, outdegrees

## Introduction

We consider random recursive trees which are constructed incrementally as follows. Starting with a single vertex  $v(0)$  with weight  $w(0) = 1$  and label 0, at the first step a new vertex  $v(1)$  is added to the tree as a child of the initial vertex. It is labelled by 1 and assigned a random weight  $w(1) > 0$ . At step  $j > 1$ , given all the weights  $w(0), w(1), \dots, w(j-1)$ , first a node  $v(j^*)$  is chosen at random from the existing nodes  $v(0), v(1), \dots, v(j-1)$  with probabilities proportional to the nodes' weights, and then a new vertex  $v(j)$  is added to the tree as a child of  $v(j^*)$ . The new vertex has label  $j$  and a random weight  $w(j) > 0$ . As at the initial step (where we put  $1^* = 0$ ), the edge is directed from  $v(j^*)$  to its child vertex  $v(j)$ .

If  $w(j) = 1$  for all  $j$ , then we get the standard recursive tree. If  $w(j) = a^j$ , where  $a \neq 1$  is positive, we get the recursive tree considered in Borovkov and Motyer (3).

In this paper we consider the case when  $w(j) = a_1 \cdots a_j$ , where  $a_k$  are i.i.d. random variables. Our aim is to study the asymptotic behavior of the mean value of the outdegree  $N_n(j)$  of the vertex  $v(j)$ ,  $j \leq n$ , (i.e. the number of edges coming out of  $v(j)$  in the tree having  $n$  nonrooted vertices) as  $n \rightarrow \infty$ .

Denote by  $\mathcal{T}_n$ ,  $n = 0, 1, 2, \dots$ , the set of all rooted recursive trees with  $n$  nonrooted vertices. That is,  $\mathcal{T}_n$  consists of the rooted trees whose root is labelled by 0 and whose nonrooted vertices are labelled by numbers  $1, 2, \dots, n$  in such a way that for any  $j \leq n$  the shortest path leading from the vertex with label  $j$  to the root traverses only the vertices with labels  $k \leq j$ . For a  $t_n \in \mathcal{T}_n$ , let  $t_n(j) \in \mathcal{T}_{n+1}$  be the recursive tree which is obtained from  $t_n$  by adding a vertex labelled by  $n+1$  as a child of the vertex with the label  $j \in \{0, 1, \dots, n\}$ .

Now we can describe our construction of random recursive trees. Let  $\theta_j$ ,  $j = 1, 2, \dots, n$ , be i.i.d. r.v.'s. First, we run a random walk

$$S_0 = 0, \quad S_j = \theta_1 + \cdots + \theta_j, \quad j \geq 1. \quad (1)$$

Second, given  $S_j$ ,  $j = 0, 1, \dots, n$ , we construct a (conditional) Markov chain  $T_0, T_1, \dots, T_n$  with  $T_k \in \mathcal{T}_k$ ,  $k = 0, 1, \dots, n$ , by assigning the weight  $w(j) := e^{-S_j}$  to the vertex labelled by  $j \geq 0$  (so that  $w(j) = a_1 \cdots a_j$ ,  $j \geq 1$ , with  $a_j := e^{-\theta_j}$  being i.i.d. r.v.'s), setting, for  $r = 0, 1, \dots, n$ ,

$$W_r := \sum_{q=0}^r w(q) = \sum_{q=0}^r e^{-S_q}, \quad p_r(j) := e^{-S_j} W_r^{-1} = e^{-S_j} \left( \sum_{q=0}^r e^{-S_q} \right)^{-1}, \quad j = 0, 1, \dots, r,$$

<sup>†</sup>Supported by the ARC Centre of Excellence for Mathematics and Statistics of Complex Systems.

<sup>‡</sup>Supported by RFBR grant 05-01-00035 and by the program "Contemporary Problems of Theoretical Mathematics" of the Russian Academy of Sciences.

and letting, for any  $t_r \in \mathcal{T}_r$  and  $j = 0, 1, \dots, r$ ,

$$\mathbf{P}_w(T_{r+1} = t_r(j) | T_r = t_r) \equiv \mathbf{P}(T_{r+1} = t_r(j) | T_r = t_r; w(1), \dots, w(r)) : p_r(j).$$

Recalling that  $v(k^*)$  is the parent vertex for  $v(k)$  and denoting by  $I\{\mathcal{A}\}$  the indicator of the event  $\mathcal{A}$ , we can represent the outdegree  $N_n(j)$  of the vertex  $v(j)$ ,  $j \leq n$ , as

$$N_n(j) = \sum_{k=j+1}^n I\{v(k^*) = v(j)\}.$$

Therefore,

$$\begin{aligned} \mathbf{E}_w N_n(j) &:= \mathbf{E}[N_n(j) | w(1), \dots, w(n-1)] \\ &= \sum_{k=j+1}^n \mathbf{E}_w I\{v(k^*) = v(j)\} \sum_{k=j+1}^n p_{k-1}(j) = e^{-S_j} \sum_{k=j}^{n-1} W_k^{-1} \end{aligned}$$

and

$$\mathbf{E} N_n(j) = \sum_{k=j}^{n-1} \mathbf{E} e^{-S_j} W_k^{-1}.$$

In this note we formulate two theorems describing the asymptotic behavior (as  $n \rightarrow \infty$ ) of the expectations  $\mathbf{E} N_n(j)$  and the distributions of the random variables  $\mathbf{E}_w N_n(j)$  in different ranges of the parameter  $j$  values when the random walk (1) satisfies the following *Spitzer* condition: there exists a  $\rho \in (0, 1)$  such that

$$\frac{1}{n} \sum_{k=1}^n \mathbf{P}(S_k > 0) \rightarrow \rho \quad \text{as } n \rightarrow \infty. \tag{2}$$

Observe that if  $\mathbf{E}\theta_1 = 0$  and the distribution of  $\theta_1$  belongs to the domain of attraction of a stable law of order  $\alpha \in (1, 2]$  then (see e.g. Bingham *et al.* (2)) there exists  $\beta \in [-1, 1]$  such that

$$\rho = \frac{1}{2} + \frac{1}{\pi\alpha} \arctan\left(-\beta \tan \frac{\pi\alpha}{2}\right).$$

### Statement of results

Let

$$\gamma_0 := 0, \quad \gamma_{j+1} := \min\{n > \gamma_j : S_n < S_{\gamma_j}\}, \quad j \geq 0,$$

and

$$\Gamma_0 := 0, \quad \Gamma_{j+1} := \min\{n > \Gamma_j : S_n > S_{\Gamma_j}\}, \quad j \geq 0,$$

be the strict descending and ascending ladder epoches of  $\{S_n\}_{n \geq 0}$ , resp. Introduce the two renewal functions

$$\begin{aligned} U(x) &:= 1 + \sum_{j=1}^{\infty} \mathbf{P}(S_{\Gamma_j} < x), \quad x > 0; \quad U(0) = 1, \quad U(x) = 0, \quad x < 0, \\ V(x) &:= \sum_{j=0}^{\infty} \mathbf{P}(S_{\gamma_j} \geq -x), \quad x > 0; \quad V(0) = 1, \quad V(x) = 0, \quad x < 0. \end{aligned}$$

By means of  $V(x)$  and  $U(x)$  and the constant  $\phi := \sum_{j=1}^{\infty} j^{-1} \mathbf{P}(S_j = 0)$  one can specify two sequences of probability measures  $\{\mathbf{P}_n^-\}_{n \geq 1}$  and  $\{\mathbf{P}_n^+\}_{n \geq 1}$  (defined on the  $\sigma$ -algebras  $\Sigma_n = \sigma(S_1, \dots, S_n)$ , resp.), by setting for each  $\mathcal{A} \in \Sigma_n$

$$\mathbf{P}_n^-(\mathcal{A}) := e^\phi \int_{\mathcal{A}} U(-S_n) I\{\tilde{M}_n < 0\} d\mathbf{P}, \quad \mathbf{P}_n^+(\mathcal{A}) := \int_{\mathcal{A}} V(S_n) I\{L_n \geq 0\} d\mathbf{P}.$$

It is known (see e.g. Vatutin and Dyakonova (4)) that the measures are consistent, and therefore there exist measures  $\mathbf{P}^-$  and  $\mathbf{P}^+$  on the  $\sigma$ -algebra  $\sigma(S_1, S_2, \dots)$  such that their restrictions  $\mathbf{P}^\pm|_{\Sigma_n}$  to  $\Sigma_n$  coincide with  $\mathbf{P}_n^\pm$ ,  $n = 1, 2, \dots$

One can show (see Lemma 2.7 in Afanasyev *et al.* (1)) that under the condition (2)

$$\eta_1 := \sum_{k=1}^{\infty} e^{S_k} < \infty \quad \mathbf{P}^- \text{- a.s.}, \quad \eta_2 := \sum_{k=0}^{\infty} e^{-S_k} < \infty \quad \mathbf{P}^+ \text{- a.s.} \quad (3)$$

Set

$$L_n := \min_{0 \leq k \leq n} S_k, \quad \tilde{M}_n := \max_{1 \leq k \leq n} S_k.$$

It is known (see e.g. Bingham *et al.* (2)) that under the condition (2) there exist functions  $l_1(n)$  and  $l_2(n)$  slowly varying at infinity,  $l_1(n)l_2(n) \sim \pi^{-1} \sin \pi\rho$ ,  $n \rightarrow \infty$ , such that

$$\mathbf{P}(L_n \geq 0) \sim n^{\rho-1}l_1(n), \quad \mathbf{P}(\tilde{M}_n < 0) \sim n^{-\rho}l_2(n) \quad \text{as } n \rightarrow \infty. \quad (4)$$

**Theorem 1** *Let condition (2) hold. Then there exist positive sequences  $\{C_j\}_{j \geq 0}$  and  $\{D_j\}_{j \geq 0}$  such that for any fixed  $j \geq 0$*

$$\lim_{n \rightarrow \infty} \frac{\mathbf{E}N_n(j)}{n\mathbf{P}(L_n \geq 0)} = C_j \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{\mathbf{E}N_n(n-j)}{\mathbf{P}(\tilde{M}_n < 0)} = D_j.$$

In addition,

$$\lim_{j, n-j \rightarrow \infty} \frac{\mathbf{E}N_n(j)}{(n-j)\mathbf{P}(\tilde{M}_j < 0)\mathbf{P}(L_{n-j} \geq 0)} = \frac{1}{\rho}.$$

**Remark 2** *In view of (4), the assertions of Theorem 1 can be rewritten as*

$$\begin{aligned} \mathbf{E}N_n(j) &\sim C_j n^\rho l_1(n), \quad \mathbf{E}N_n(n-j) \sim D_j n^{-\rho} l_2(n) \quad \text{as } n \rightarrow \infty, \\ \mathbf{E}N_n(j) &\sim \rho^{-1} j^{-\rho} l_2(j) (n-j)^\rho l_1(n-j) \quad \text{as } j, n-j \rightarrow \infty. \end{aligned}$$

Our description of the asymptotic behavior of the distribution of the r.v.  $\mathbf{E}_w N_n(j)$  will only be limited to the case when  $j$  is located either to the right of the random epoch  $\tau(n)$  or in its small left vicinity.

Let  $\tau(n) := \min \{k \geq 0 : S_k \leq S_l, l \in [0, n]\}$  be the left-most point at which the minimal value of the random walk  $\{S_k\}_{0 \leq k \leq n}$  is attained.

**Theorem 3** *Let condition (2) hold and  $j = j(n)$  be an arbitrary (possibly random) sequence such that  $(\tau(n) - j)_+ = o(n)$  in probability as  $n \rightarrow \infty$ . Then*

$$\frac{e^{S_j - S_{\tau(n)}}}{n-j} \mathbf{E}_w N_n(j) \implies \frac{1}{\eta_1^- + \eta_2^+},$$

where  $\eta_1^-$  and  $\eta_2^+$  are independent copies of the r.v.'s  $\eta_1$  and  $\eta_2$  from (3), resp.

## References

- [1] Afanasyev, V.I., Geiger, J., Kersting, G. and Vatutin, V.A. *Criticality for branching processes in random environment.* Ann. Probab., 33 (2005), 2, 645–673.
- [2] Bingham, N.H., Goldie, C.M. and Teugels, J.L. *Regular variation.* Cambridge University Press, Cambridge (1987).
- [3] Borovkov, K.A. and Motyer, A. *On the asymptotic behavior of a simple growing point process model.* Statistics and Probability Letters, 72 (2005), 265–275.
- [4] Vatutin, V.A. and Dyakonova, E.E. *Galton-Watson branching processes in random environment. II: Finite-dimensional distributions.* Theory Probab. Appl., 49 (2004), 275–308.

