



HAL
open science

A combinatorial realization of Schur-Weyl duality via crystal graphs and dual equivalence graphs

S. Assaf

► **To cite this version:**

S. Assaf. A combinatorial realization of Schur-Weyl duality via crystal graphs and dual equivalence graphs. 20th Annual International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2008), 2008, Viña del Mar, Chile. pp.141-152, 10.46298/dmtcs.3626 . hal-01185161

HAL Id: hal-01185161

<https://inria.hal.science/hal-01185161v1>

Submitted on 19 Aug 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A combinatorial realization of Schur-Weyl duality via crystal graphs and dual equivalence graphs

Sami H. Assaf[†]

*Department of Mathematics
University of Pennsylvania
Philadelphia, PA 19104-6395*

Abstract. For any polynomial representation of the special linear group, the nodes of the corresponding crystal may be indexed by semi-standard Young tableaux. Under certain conditions, the standard Young tableaux occur, and do so with weight 0. Standard Young tableaux also parametrize the vertices of dual equivalence graphs. Motivated by the underlying representation theory, in this paper, we explain this connection by giving a combinatorial manifestation of Schur-Weyl duality. In particular, we put a dual equivalence graph structure on the 0-weight space of certain crystal graphs, producing edges combinatorially from the crystal edges. The construction can be expressed in terms of the local characterizations given by Stembridge for crystal graphs and the author for dual equivalence graphs.

Keywords: Schur-Weyl duality, zero weight spaces, crystal graphs, dual equivalence graphs

1 Introduction

Schur-Weyl duality [Wey39] is a powerful tool in the study of irreducible representations of the classical groups. The set-up is as follows. Let G be a complex, reductive Lie group, and let V be an irreducible finite-dimensional complex representation of G . Let V^0 be the space of vectors of weight 0, i.e. vectors which are fixed by a maximal torus. Then the Weyl group W acts naturally on V^0 . The case when $G = \mathrm{SL}_n$, and so then the Weyl group is isomorphic to \mathcal{S}_n , was studied by Gutkin [Gut73] who showed that if V is an irreducible component of the n th tensor power of the standard n -dimensional representation of SL_n , then the \mathcal{S}_n -module V^0 is irreducible and all irreducible \mathcal{S}_n -modules occur in this way. This was also observed by Kostant [Kos76]. Though there are descriptions of the Weyl group action on 0-weight spaces for SL_n , there is no general rule for how to decompose the action into irreducible representations of \mathcal{S}_n . This paper proposes a new approach to this problem by combinatorializing the representations using crystal graphs [Kas90, Kas91] for representations of SL_n and dual equivalence graphs [Ass07] for representations of \mathcal{S}_n .

[†]Work supported in part by NSF Mathematical Sciences Postdoctoral Research Fellowship DMS-0703567.

Kashiwara introduced in [Kas90, Kas91] the notion of crystal bases in his study of the representation theory of quantized universal enveloping algebras at $q = 0$. The theory of canonical bases, developed independently by Lusztig [Lus90a, Lus90b], studies the same problem from a different viewpoint, though for the purposes of this paper, we focus solely on the crystal approach. A crystal graph is a directed, colored graph with vertex set given by the crystal basis and directed edges given by deformations of the Chevalley generators e_i and f_i . The combinatorial structure of crystal graphs encodes important information for studying the representations; for instance, knowing the crystal immediately gives a formula for the character as well as tensor product and branching rules for the corresponding representation. For the quantum group $U_q(\mathfrak{sl}_n)$, the crystal basis can be indexed by semi-standard Young tableaux, and there is an explicit combinatorial construction of the crystal graph on tableaux [KN94, Lit95].

Dual equivalence graphs were first introduced in [Ass07] as a combinatorial tool for studying functions expressed in terms of quasi-symmetric functions. In particular, in [Ass07] they are used to give a purely combinatorial proof of symmetry and Schur positivity of Lascoux-Leclerc-Thibon polynomials and Macdonald polynomials. The original motivation for these graphs, based on ideas of Haiman [Hai05], was to mimic existing crystal-theoretic proofs used to establish positivity results for functions expressed in terms of monomials. Given this, a natural problem which this paper addresses is to explore the connections between crystal graphs and dual equivalence graphs. Combinatorially, the vertices of a dual equivalence graph are standard Young tableaux of a given shape, which also index a basis for irreducible representations of \mathcal{S}_n . When λ is a partition of n , these also index the 0-weight nodes in the crystal graph of the irreducible SL_n module with highest weight λ . From this observation we develop a correspondence between crystals of irreducible representations of SL_n of degree n and dual equivalence graphs.

This paper is organized as follows. Section 2 contains an exposition of type A crystal graphs. We begin by defining the combinatorial structure on semi-standard Young tableaux, and then present Stembridge's local characterization for crystals [Ste03] which will be essential for establishing our main result. Section 3 gives the analogous exposition for dual equivalence graphs, beginning with the combinatorial description in terms of standard Young tableaux before presenting the local characterization from [Ass07]. The correspondence is laid out in Section 4, where we give two separate proofs of the main result using the tableaux descriptions (Theorem 4.2) and the local characterizations (Theorem 4.3). Finally, in Section 5, we lay out how these methods may be extended to any irreducible representation of SL_n with a nontrivial 0-weight space, and we also suggest how this approach may be used to define dual equivalence graphs for other types.

2 Crystal graphs

In order to construct the crystal graph of an irreducible representation of SL_n , we must define Kashiwara's crystal operators in this setting. We depart somewhat from the original descriptions developed independently by Kashiwara and Nakashima [KN94] and Littelmann [Lit95] in favor of the presentation of the Littlewood-Richardson rule in [Mac95].

For a word w of length n , i.e. $w \in \mathbb{N}^n$, define $m_i(w, r)$ to be the number of i 's occurring in $w_r w_{r+1} \cdots w_n$ minus the number of $i-1$'s occurring in $w_r w_{r+1} \cdots w_n$. Then let $m_i(w)$ be the maximum $m_i(w, r)$ which occurs for any $r = 1, \dots, n$. Observe that if $m_i(w) > 0$ and w_p is the rightmost occurrence of this maximum, i.e. $m_i(w, p) = m_i(w)$ and $m_i(w, q) < m_i(w)$ for $q > p$, then $w_p = i$. The result of applying the crystal operator \tilde{e}_i to a semi-standard tableau T may be described as follows. Let w_T be the reading word for T . Then if $m_i(w_T) \leq 0$, $\tilde{e}_i(T) = 0$; otherwise $\tilde{e}_i(T)$ is the result of

changing the i which is the rightmost occurrence of $m_i(w_T)$ to an $i-1$. As \tilde{e}_i is invertible when nonzero, define \tilde{f}_i by requiring \tilde{e}_i and \tilde{f}_i to be inverses of one another whenever the image is not 0.




Fig. 1: The crystal graph $\mathcal{X}_{(2,2)}^4$, with edges $\tilde{f}_1 \nearrow$, $\tilde{f}_2 \rightarrow$, $\tilde{f}_3 \searrow$.

For a partition λ with at most n rows, the crystal graph \mathcal{X}_λ^n is the directed, colored graph on semi-standard tableaux of shape λ with entries from $[n]$ defined as follows. For $i = 1, \dots, n-1$, if $m_i(w_T) > 0$, define a directed i -edge from $\tilde{e}_i(T)$ to T . For example, Figure 1 gives the crystal graph $\mathcal{X}_{(2,2)}^4$.

In [Ste03], Stembridge gives a local characterization of crystal graphs which arise from representations for simply-laced types. In order to define Stembridge's axioms, we must first introduce some notation associated with a directed, colored graph. Say that \mathcal{X} has *degree* n if the largest color for an edge is $n-1$.

We have simplified things from the presentation in [Ste03] since we are considering only the type A case. The notations below may not be well-defined in general, but will be in the contexts in which they are used, i.e. for graphs satisfying axioms P1 and P2 below.

If $x \xleftarrow{i} y$ (resp. $x \xrightarrow{i} z$), write $y = E_i x$ (resp. $z = F_i x$). The i -string through x is the maximal path

$$F_i^{-d} x \xrightarrow{i} \dots \xrightarrow{i} F_i^{-1} x \xrightarrow{i} x \xrightarrow{i} F_i x \xrightarrow{i} \dots \xrightarrow{i} F_i^r x.$$

In this case we write $\delta(x, i) = -d$ and $\varepsilon(x, i) = r$. Finally, we have the following differences whenever E_i, F_i is defined at x .

$$\begin{aligned} \Delta_i \delta(x, j) &= \delta(E_i x, j) - \delta(x, j), & \nabla_i \delta(x, j) &= \delta(x, j) - \delta(F_i x, j), \\ \Delta_i \varepsilon(x, j) &= \varepsilon(E_i x, j) - \varepsilon(x, j), & \nabla_i \varepsilon(x, j) &= \varepsilon(x, j) - \varepsilon(F_i x, j). \end{aligned}$$

Definition 2.1 ([Ste03]) A directed, colored graph \mathcal{X} is regular if the following hold:

(P1) all monochromatic directed paths have finite length;

(P2) for every vertex x , there is at most one edge $x \xleftarrow{i} y$ and at most one edge $x \xrightarrow{i} z$;

(P3) assuming $E_i x$ is defined, $\Delta_i \delta(x, j) + \Delta_i \varepsilon(x, j) = \begin{cases} 2 & \text{if } j = i \\ -1 & \text{if } j = i \pm 1 \\ 0 & \text{if } |i - j| \geq 2 \end{cases}$;

(P4) assuming $E_i x$ is defined, $\Delta_i \delta(x, j), \Delta_i \varepsilon(x, j) \leq 0$ for $j \neq i$;

(P5) $\Delta_i \delta(x, j) = 0 \Rightarrow E_i E_j x = E_j E_i x = y$ and $\nabla_j \varepsilon(y, i) = 0$;
 $\nabla_i \varepsilon(x, j) = 0 \Rightarrow F_i F_j x = F_j F_i x = y$ and $\Delta_j \delta(y, i) = 0$;

(P6) $\Delta_i \delta(x, j) = \Delta_j \delta(x, i) = -1 \Rightarrow E_i E_j^2 E_i x = E_j E_i^2 E_j x = y$ and $\nabla_i \varepsilon(y, j) = \nabla_j \varepsilon(y, i) = -1$;
 $\nabla_i \varepsilon(x, j) = \nabla_j \varepsilon(x, i) = -1 \Rightarrow F_i F_j^2 F_i x = F_j F_i^2 F_j x = y$ and $\Delta_i \delta(y, j) = \Delta_j \delta(y, i) = -1$.




Fig. 2: An illustration of axioms P3 and P4, where $F_j \swarrow, F_i \searrow$.

Axioms P3 and P4 dictate how the maximal length of a j -string differs between x and $E_i x$; see Figure 2. When $|i - j| \geq 2$, there is no change, but when $j = i \pm 1$, the length must change by 1. Axioms P5 and P6 give information about how edges with different labels interact; see Figure 3. When $|i - j| \geq 2$, the conditions of P5 will always be satisfied, though when $j = i \pm 1$, either P5 or P6 could hold.

The following theorem shows that regular graphs corresponds precisely to the crystal graphs of representations in type A. This result allows us to combinatorialize the problem of studying representations of SL_n by instead studying regular graphs of degree n .




Fig. 3: An illustration of axioms P5 and P6, where $F_j \swarrow, F_i \searrow$.

Theorem 2.2 ([Ste03]) *Each \mathcal{X}_λ^n is a regular graph, and every connected component of a regular graph is isomorphic to \mathcal{X}_λ^n for some λ, n .*

The theorem is proved by showing that Littelmann’s path operators [Lit94] generate regular graphs, since the Path Model is known to generate the \mathcal{X}_λ^n . As remarked in [Ste03], it would be nice to have a graph-theoretic proof of Theorem 2.2 which bypasses the Path Model.

3 Dual equivalence graphs

Since the characters of irreducible representations of SL_n are Schur polynomials which form a basis for symmetric polynomials, crystal graphs can be used to prove that certain polynomials expressed in terms of monomials are symmetric and Schur positive. Motivated by this approach, Haiman [Hai05] suggested defining a graph structure on standard tableaux using the dual equivalence relation as a means of establishing the symmetry and Schur positivity of polynomials expressed in terms of quasi-symmetric functions. The result of this idea is the theory of dual equivalence graphs developed in [Ass07].

Following Haiman’s idea, we use the dual equivalence relation to construct a graph whose vertices are given by standard tableaux and whose connected components are indexed by partitions.

Definition 3.1 ([Hai92]) *An elementary dual equivalence on three consecutive letters, say $i-1, i, i+1$, of a permutation is given by switching the outer two letters whenever the middle letter is not i :*

$$\dots i \dots i \pm 1 \dots i \mp 1 \dots \equiv^* \dots i \mp 1 \dots i \pm 1 \dots i \dots$$

The standard dual equivalence graph associated to λ , denoted \mathcal{G}_λ , is the vertex-signed, edge-colored graph constructed in the following way. The vertices are given by all standard Young tableaux of shape λ . Two tableaux T and U are connected by an i -colored edge whenever their reading words, w_T and w_U , differ by an elementary dual equivalence for $i-1, i, i+1$. To each vertex T we associate the signature $\sigma(T)$ which indicates the descent set of the tableau:

$$\sigma(T)_i = \begin{cases} +1 & \text{if } i \text{ appears to the left of } i+1 \text{ in } w_T \\ -1 & \text{if } i+1 \text{ appears to the left of } i \text{ in } w_T \end{cases} \quad (3.1)$$

As the original motivation for these graphs was to prove Schur positivity, the role of the signatures is to index which quasi-symmetric function should be associated to each vertex.




Fig. 4: The standard dual equivalence graphs $\mathcal{G}_{(4,1)}$, $\mathcal{G}_{(3,2)}$ and $\mathcal{G}_{(3,1,1)}$.

In contrast with \mathcal{X}_λ^n , the bound on the edge colors (equivalently, on the entries of the tableaux) for \mathcal{G}_λ is implicit. More precisely, if λ is a partition of m , then \mathcal{G}_λ has degree m , meaning that the largest edge color is $m - 1$.

It follows from results in [Hai92] that each \mathcal{G}_λ is connected. It is easy to see from the construction that $\mathcal{G}_{\lambda'}$ can be obtained from \mathcal{G}_λ by conjugating each vertex and multiplying each entry of the signature by -1 . Several other nice properties of these graphs are given [Ass07], including the fact that the \mathcal{G}_λ are pairwise non-isomorphic and have no nontrivial automorphisms.

Motivated by the local characterization of crystal graphs \mathcal{X}_λ^n , we next present a local characterization of the graphs \mathcal{G}_λ . We will abuse notation by simultaneously referring to D_i as the collection of i -colored edges as well as an involution on vertices which have an i -edge, i.e. we write $D_i(v) = w$ whenever $\{v, w\} \in D_i$. This is justified by the first axiom.

Definition 3.2 ([Ass07]) A signed, colored graph $\mathcal{G} = (V, \sigma, D)$ is a dual equivalence graph if the following hold:


- (ax1) For $v \in V$ and $i > 1$, $\sigma(v)_{i-1} = -\sigma(v)_i$ if and only if there exists $w \in V$ such that $\{v, w\} \in D_i$. Moreover, w is unique when it exists.
- (ax2) Whenever $\{v, w\} \in D_i$,

$$\begin{aligned} \sigma(v)_j &= -\sigma(w)_j && \text{for } j = i-1, i; \\ \sigma(v)_h &= \sigma(w)_h && \text{for } h < i-2 \text{ and } h > i+1. \end{aligned}$$


(ax3) Whenever $\{v, w\} \in D_i$,

if $\sigma(v)_{i-2} = -\sigma(w)_{i-2}$, then $\sigma(v)_{i-2} = -\sigma(v)_{i-1}$;
 if $\sigma(v)_{i+1} = -\sigma(w)_{i+1}$, then $\sigma(v)_{i+1} = -\sigma(v)_i$.


(ax4) For $i > 3$, every non-trivial connected component of $(V, \sigma, D_{i-2} \cup D_{i-1} \cup D_i)$ is depicted below and there is a path containing at most one D_i edge between any two vertices of $(V, \sigma, D_2 \cup \dots \cup D_i)$.



(ax5) Whenever $|i - j| \geq 3$, $\{v, u\} \in D_i$ and $\{u, w\} \in D_j$, there exists $x \in V$ such that $\{v, x\} \in D_j$ and $\{x, w\} \in D_i$.



When $n = 4$, we stipulate the following in place of axiom 4: For $i > 2$, every non-trivial connected component of the sub-graph $(V, \sigma, D_{i-1} \cup D_i)$ is one of the following:



Next we have the analog of Theorem 2.2 for dual equivalence graphs. The proof of Theorem 3.3 is by a purely combinatorial, graph-theoretic argument.

Theorem 3.3 ([Ass07]) Each \mathcal{G}_λ is a dual equivalence graph, and every connected component of a dual equivalence graph is isomorphic to \mathcal{G}_λ for a unique partition λ .

4 From crystal graphs to dual equivalence graphs

Since standard tableaux are special cases of semi-standard tableaux, and since the construction of \mathcal{G}_λ was motivated by \mathcal{X}_λ^n , we would like to explore the connection between regular graphs and dual equivalence graphs. Since semi-standard tableaux index a basis for an irreducible representation of SL_n so that the content of the tableau corresponds to the weight of the basis vector, those basis vectors corresponding

to standard tableaux index the 0-weight space of the representations of dimension n . Therefore we may think of dual equivalence graphs as living in the 0-weight space of crystal graphs.

In general, the Weyl group acts naturally on the 0-weight space of a representation. In the case of SL_n , the Weyl group is isomorphic to \mathcal{S}_n . Since standard tableaux index the basis for an irreducible representation of \mathcal{S}_n , we would like to use dual equivalence graphs to combinatorialize the study of representations of \mathcal{S}_n in much the same way that crystal graphs combinatorialize the study of representations of SL_n . This is the motivation behind the following correspondence. In the interest of space, the longer proofs have been condensed by omitting the more mechanical details.

Combinatorially, the 0-weight space of a regular graph \mathcal{X} consists of all vertices which lie at the center of each i -string; i.e. $\varepsilon(x, i) = -\delta(x, i)$ for each i . Given a directed, colored graph \mathcal{X} , define a signed, colored graph $\mathcal{G}(\mathcal{X}) = (V, \sigma, D)$ by

$$V = \{x \in \mathcal{X} \mid \varepsilon(x, i) = -\delta(x, i) = 0 \text{ or } 1 \ \forall i, \}, \tag{4.1}$$

$$\sigma(x)_i = \begin{cases} +1 & \text{if } \varepsilon(x, i) = 1 \\ -1 & \text{if } \varepsilon(x, i) = 0 \end{cases}, \tag{4.2}$$

$$D_i(x) = \begin{cases} F_{i-1}F_iE_{i-1}E_ix & \text{if } \varepsilon(x, i) = 1 \text{ and } \varepsilon(x, i-1) = 0 \\ F_iF_{i-1}E_iE_{i-1}x & \text{if } \varepsilon(x, i) = 0 \text{ and } \varepsilon(x, i-1) = 1 \end{cases}. \tag{4.3}$$

An i -edge from x , or equivalently $D_i(x)$, is not defined if neither condition of equation (4.3) holds.




Fig. 5: Constructing $\mathcal{G}(\mathcal{X})$ from \mathcal{X} ; here we have $F_{i-1} \swarrow, F_i \searrow$.

Proposition 4.1 *If \mathcal{X} is a regular graph, then $\mathcal{G}(\mathcal{X})$ is well-defined.*

Proof: It suffices to show that D_i is well-defined. Note that if $\delta(x, i) = -1$, then E_ix is nonzero. If $\delta(x, i-1) = 0$, then by axiom P3 we must have $\delta(E_ix, i-1) + \varepsilon(E_ix, i-1) = -1$. The constraints of axiom P4 force $\varepsilon(E_ix, i-1) = 0$, and so we conclude that $\delta(E_ix, i-1) = -1$. Therefore $E_{i-1}E_ix$ is nonzero. Continuing thus, with liberal use of axioms P3 and P4, we eventually conclude that $F_{i-1}F_iE_{i-1}E_ix$ is nonzero. The analogous argument holds with the reverse assumptions for $\delta(x, i-1), \delta(x, i)$. \square

Below we give two separate proofs that if \mathcal{X} is a regular graph with 0-weight space given by V , then $\mathcal{G}(\mathcal{X})$ is a dual equivalence graph. The first proof utilizes the graph structures on tableaux, and the second uses the local characterizations. Note that the condition that V give the 0-weight space is highly restrictive.

In order for \mathcal{X}_λ^n to have a 0-weight space, λ must be a partition of kn into at most n parts. We first treat the case for \mathcal{X}_λ^n when $k = 1$, i.e. λ is a partition of n .

Theorem 4.2 *For every partition λ of n , we have $\mathcal{G}(\mathcal{X}_\lambda^n) = \mathcal{G}_\lambda$.*

Proof: In this case we may identify V with $\text{SYT}(\lambda)$. Let w_T be the reading word for a tableau T . For $T \in \text{SYT}(\lambda)$, note that $\varepsilon(T, i) = 0$ if and only if $i+1$ occurs to the left of i in w_T . Therefore equations 3.1 and 4.2 correspond. In particular, T will have an i -edge in \mathcal{G}_λ precisely when equation (4.3) is defined.

Suppose $D_i(T)$ is defined, and so T admits an elementary dual equivalence for $i-1, i, i+1$. By symmetry, we may assume i lies to the left of $i \pm 1$ in w_T . If $\varepsilon(T, i) = 1$, then ignoring all letters other than $i-1, i, i+1$ and regarding \tilde{e}_j, \tilde{f}_j ($j = i-1, i$) as operators on reading words, we have

$$\begin{aligned} \tilde{f}_{i-1}\tilde{f}_i\tilde{e}_{i-1}\tilde{e}_i w_T &= \tilde{f}_{i-1}\tilde{f}_i\tilde{e}_{i-1}\tilde{e}_i \cdots i \cdots i-1 \cdots i+1 \cdots \\ &= \tilde{f}_{i-1}\tilde{f}_i\tilde{e}_{i-1} \cdots i \cdots i-1 \cdots i \cdots \\ &= \tilde{f}_{i-1}\tilde{f}_i \cdots i \cdots i-1 \cdots i-1 \cdots \\ &= \tilde{f}_{i-1} \cdots i+1 \cdots i-1 \cdots i-1 \cdots \\ &= \cdots i+1 \cdots i-1 \cdots i \cdots . \end{aligned}$$

Therefore equation (4.3) corresponds to the elementary dual equivalences. The case when $\varepsilon(T, i-1) = 1$ is similar. \square

Going back to the representation theory of SL_n and \mathcal{S}_n , Theorem 4.2 reflects the fact that for λ a partition of n , the action of \mathcal{S}_n on the 0-weight space of the irreducible representation of SL_n corresponding to λ is given by the Specht module S^λ .

The following result generalizes Theorem 4.2. Here we present a self-contained proof based solely on the local characterizations of crystal graphs and dual equivalence graphs without appealing to Theorems 2.2 and 3.3 and without reference to tableaux.

Theorem 4.3 *If \mathcal{X} is a regular with 0-weight space given by V (see equation (4.1)), then $\mathcal{G}(\mathcal{X})$ is a dual equivalence graph.*

Proof: Comparing equation (4.2) with the conditions in equation (4.3) dictating when D_i is defined shows that $\sigma(x)_{i-1} = -\sigma(x)_i$ if and only if $D_i(x)$ is defined. Figure 5 makes evident the fact that $\sigma(x)_j = -\sigma(D_i(x))_j$ for $j = i-1, i$, though this can also be seen directly from the proof that D_i is well-defined. Therefore we have established dual equivalence axioms 1 and 2.

Axiom 3 can be demonstrated by showing that if $\varepsilon(x, i-2) \neq \varepsilon(D_i(x), i-2)$, then $D_i(x) = D_{i-1}(x)$. To do this, we show that the crystal graph operators going from x to $D_i(x)$ exactly mirror the situation in $\mathcal{X}_{(2,2)}^4$ for the two standard tableaux of shape $(2, 2)$; see Figure 1.

By crystal axioms P3, P4 and P5, whenever $|a - b| \geq 2$, we have $E_a E_b = E_b E_a$, $F_a E_b = E_b F_a$ and $F_a F_b = F_b F_a$. Dual equivalence axiom 5 is equivalent to showing that if D_i and D_j are both defined at x for some $|i - j| \geq 3$, then $D_i D_j(x) = D_j D_i(x)$. Therefore the identities above for crystal operators establish dual equivalence axiom 5.




Fig. 6: An illustration of the commutativity which establishes dual equivalence axiom 4; the 0-weight vertices are o's and crystal operators are $F_{i-3} \searrow, F_{i-2} \rightarrow, F_{i-1} \swarrow, F_i \downarrow$.

The most difficult step is to establish axiom 4. For this, note that it is sufficient to show that if D_{i-2}, D_{i-1}, D_i are all defined at x and are all distinct (i.e. x does not have a double edge), then $D_{i-2}D_i(x) = D_iD_{i-2}(x)$. By symmetry we may assume $\varepsilon(x, i-3) = 1 = \varepsilon(x, i-1)$ and $\varepsilon(x, i-2) = 0 = \varepsilon(x, i)$. Then the first part of axiom 4 is equivalent to establishing the following identity, illustrated in Figure 6.

$$(F_i F_{i-1} E_i E_{i-1})(F_{i-2} F_{i-3} E_{i-2} E_{i-3})x = (F_{i-2} F_{i-3} E_{i-2} E_{i-3})(F_i F_{i-1} E_i E_{i-1})x$$

The key to establishing this relation is to use the commutativity relations for crystal operators used to prove dual equivalence axiom 5, as well as the additional relation that both E_{i-1} and F_{i-1} commute with the sequence $F_{i-2} F_{i-3} E_{i-2}$.

For the second part of axiom 4, we first show that if \mathcal{X} is connected, then so is $\mathcal{G}(\mathcal{X})$. In particular, we show that any two vertices of V may be connected by a path consisting of sequences as in equation (4.3). Furthermore, for any two vertices we may choose a path containing at most one sequence using the maximal edge color. Now axiom 4 follows. \square

Corollary 4.4 *Let $\lambda = (\mu_1 + k, \mu_2 + k, \dots, \mu_n + k)$ for some partition μ of n and some non-negative integer k . Then $\mathcal{G}(\mathcal{X}_\lambda^n) = \mathcal{G}_\mu$.*

The partition λ is the result of adding k columns of height n to μ , and clearly has 0-weight space given by V . Such a graph corresponds to the irreducible representation of SL_n indexed by μ , since SL_n cannot discriminate between the irreducible representations indexed by λ and μ . Therefore Corollary 4.4 shows

that the corresponding dual equivalence graph decomposes (albeit trivially) the action of the Weyl group on the 0-weight space of the representation.

Remark 4.5 *By introducing weights to the regular graphs, we can recover the dimension of a regular graph, which in the context of Theorem 2.2 gives the size of the indexing partition. Letting k denote the ratio of the dimension of \mathcal{X} to the degree of \mathcal{X} , modify the definition of σ in equation (4.2) to be*

$$\sigma(x)_i = (-1)^{\varepsilon(x,i)+k}. \quad (4.4)$$

Since multiplying the signatures of a dual equivalence graph entry-wise by -1 has the effect of conjugating the indexing partition, under this definition Corollary 4.4 must read

$$\mathcal{G}(\mathcal{X}_\lambda^n) = \begin{cases} \mathcal{G}_\mu & \text{if } k \text{ is odd,} \\ \mathcal{G}_{\mu'} & \text{if } k \text{ is even.} \end{cases} \quad (4.5)$$

Representation theoretically, this corresponds to tensoring with the determinant representation in GL_n , which has the effect of conjugating the shape. So indeed, the dual equivalence graph on the 0-weight space of \mathcal{X}_λ^n decomposes the action of \mathcal{S}_n on the 0-weight space of the corresponding representation of GL_n with this minor modification of signatures.

5 Extensions

The ultimate goal is to extend the construction of $\mathcal{G}(\mathcal{X})$ for any regular graph \mathcal{X} which has a 0-weight space. The characterization of V in equation (4.1) would become

$$V = \{x \in \mathcal{X} \mid \varepsilon(x, i) = -\delta(x, i) \forall i\}. \quad (5.1)$$

In many cases, the 0-weight graphs obtained from regular graphs in a similar manner can be viewed as D graphs, which are generalizations of dual equivalence graphs; see [Ass07]. Each D graph can be transformed into a disjoint union of dual equivalence graphs, so the hope is that each 0-weight graph may be realized as a D graph which decomposes into dual equivalence graphs in the correct way. If successful, this would provide a combinatorial way to study the representations of Weyl groups on 0-weight spaces, at least in type A.

A further direction is to use the correspondence between type A crystal graphs and dual equivalence graphs to define dual equivalence graphs in other types. For the simply-laced types at least, this can be done by the same constructions using Stembridge's axioms. Using the combinatorial description of crystal bases in terms of generalized Young tableaux [KN94, Lit95], we may define the standard graphs which are the analogs of \mathcal{G}_λ , and then use the local characterization of crystal graphs to obtain a local characterization for dual equivalence graphs.

Acknowledgements

The author is grateful to Mark Haiman for the idea to construct the standard dual equivalence graph \mathcal{G}_λ and for suggesting investigating the connections with crystal graphs. The author also thanks Ian Grojnowski, Mark Haiman and Monica Vazirani for helpful conversations about crystal graphs and 0-weight spaces.

References

- [Ass07] S. Assaf, *A combinatorial proof of LLT and Macdonald positivity*, preprint, 2007.
- [Gut73] E. A. Gutkin, *Representations of the Weyl group in the space of vectors of zero weight*, *Uspehi Mat. Nauk* **28** (1973), no. 5 (173), 237–238.
- [Hai92] Mark D. Haiman, *Dual equivalence with applications, including a conjecture of Proctor*, *Discrete Math.*, **99** (1992), no. 1-3, 79–113.
- [Hai05] Mark Haiman, personal communication, 2005.
- [Kas90] Masaki Kashiwara, *Crystalizing the q -analogue of universal enveloping algebras*, *Comm. Math. Phys.* **133** (1990), no. 2, 249–260.
- [Kas91] M. Kashiwara, *On crystal bases of the Q -analogue of universal enveloping algebras*, *Duke Math. J.* **63** (1991), no. 2, 465–516.
- [KN94] Masaki Kashiwara and Toshiki Nakashima, *Crystal graphs for representations of the q -analogue of classical Lie algebras*, *J. Algebra* **165** (1994), no. 2, 295–345.
- [Kos76] Bertram Kostant, *On Macdonald's η -function formula, the Laplacian and generalized exponents*, *Advances in Math.* **20** (1976), no. 2, 179–212.
- [Lit94] Peter Littelmann, *A Littlewood-Richardson rule for symmetrizable Kac-Moody algebras*, *Invent. Math.* **116** (1994), no. 1-3, 329–346.
- [Lit95] Peter Littelmann, *Crystal graphs and Young tableaux*, *J. Algebra* **175** (1995), no. 1, 65–87.
- [Lus90a] G. Lusztig, *Canonical bases arising from quantized enveloping algebras*, *J. Amer. Math. Soc.* **3** (1990), no. 2, 447–498.
- [Lus90b] G. Lusztig, *Canonical bases arising from quantized enveloping algebras. II*, *Progr. Theoret. Phys. Suppl.* (1990), no. 102, 175–201 (1991), *Common trends in mathematics and quantum field theories* (Kyoto, 1990).
- [Mac95] I. G. Macdonald, *Symmetric functions and Hall polynomials*, second ed., *Oxford Mathematical Monographs*, The Clarendon Press Oxford University Press, New York, 1995, With contributions by A. Zelevinsky, Oxford Science Publications.
- [Ste03] John R. Stembridge, *A local characterization of simply-laced crystals*, *Trans. Amer. Math. Soc.* **355** (2003), no. 12, 4807–4823 (electronic).
- [Wey39] Hermann Weyl, *The Classical Groups. Their Invariants and Representations*, Princeton University Press, Princeton, N.J., 1939.