

Performance comparison between hybridizable DG and classical DG methods for elastic waves simulation in harmonic domain

 $\underline{\mathsf{M}}.\ \underline{\mathsf{Bonnasse-Gahot}}^{1,2},\ \underline{\mathsf{H}}.\ \mathsf{Calandra}^3,\ J.\ \mathsf{Diaz}^1\ \text{and}\ S.\ Lanteri^2$

¹ INRIA Bordeaux-Sud-Ouest, team-project Magique 3D
² INRIA Sophia-Antipolis-Méditerranée, team-project Nachos

³ TOTAL Exploration-Production





Examples of seismic applications

- -





M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 2/28

Imaging method : the full wave inversion

- Iterative procedure
- Inverse problem requiring to solve a lot of forward problems



Imaging method : the full wave inversion

- Iterative procedure
- Inverse problem requiring to solve a lot of forward problems

Seismic imaging : time-domain or harmonic-domain?

- Time-domain : imaging condition complicated but low computational cost
- Harmonic-domain : imaging condition simple but huge computational cost



Imaging method : the full wave inversion

- Iterative procedure
- Inverse problem requiring to solve a lot of forward problems

Seismic imaging : time-domain or harmonic-domain?

- Time-domain : imaging condition complicated but low computational cost
- Harmonic-domain : imaging condition simple but huge computational cost

Forward problem of the inversion process

- Elastic wave propagation in harmonic domain : Helmholtz equation
- Reduction of the size of the linear system

Seismic imaging in heterogeneous complex media

- Complex topography
- High heterogeneities



Seismic imaging in heterogeneous complex media

- Complex topography
- High heterogeneities

Use of unstructured meshes with FE methods



Seismic imaging in heterogeneous complex media

- Complex topography
- High heterogeneities

Use of unstructured meshes with FE methods

DG method

- ► Flexible choice of interpolation orders (*p* − *adaptativity*)
- Highly parallelizable method
- Increased computational cost as compared to classical FEM



Seismic imaging in heterogeneous complex media

- Complex topography
- High heterogeneities

Use of unstructured meshes with FE methods

DG method

- ▶ Flexible choice of interpolation orders (*p* − *adaptativity*)
- Highly parallelizable method
- Increased computational cost as compared to classical FEM











Objective of this work

- Development of an hybridizable DG (HDG) method
- Comparison with a reference method : a standard nodal DG method



$$\label{eq:Figure} \begin{split} &Figure: \text{Degrees of} \\ &freedom \ of \ DGM \end{split}$$



FIGURE : Degrees of freedom of HDGM



M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 5/28

HDG methods

HDG methods

- B. Cockburn, J. Gopalakrishnan, R. Lazarov Unified hybridization of discontinuous Galerkin, mixed and continuous Galerkin methods for second order elliptic problems, SIAM Journal on Numerical Analysis, Vol. 47 (2009)
- S. Lanteri, L. Li, R. Perrussel, Numerical investigation of a high order hybridizable discontinuous Galerkin method for 2d time-harmonic Maxwell's equations, COMPEL, Vol. 32 (2013) (time-harmonic domain)
- N.C. Nguyen, J. Peraire, B. Cockburn, High-order implicit hybridizable discontinuous Galerkin methods for acoustics and elastodynamics, J. of Comput. Physics, Vol. 230 (2011) (time domain for seismic applications)



2D Helmholtz equations

Contents

2D Helmholtz elastic equations

Notations and definitions

Hybridizable Discontinuous Galerkin method

Numerical results

Conclusions-Perspectives



M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 6/28

2D Helmholtz equations

2D Helmholtz elastic equations

First order formulation of Helmholtz wave equations $\mathbf{x} = (x, y) \in \Omega \subset \mathbb{R}^2$,

$$\begin{cases} i\omega\rho(\mathbf{x})\mathbf{v}(\mathbf{x}) = \nabla \cdot \underline{\sigma}(\mathbf{x}) + f_{s}(\mathbf{x}) \\ i\omega\underline{\sigma}(\mathbf{x}) = \underline{\underline{C}}(\mathbf{x}) \underline{\varepsilon}(\mathbf{v}(\mathbf{x})) \end{cases}$$

- Free surface condition : $\underline{\sigma}\mathbf{n} = 0$ on Γ_I
- Absorbing boundary condition : $\underline{\sigma}\mathbf{n} = v_{\rho}(\mathbf{v} \cdot \mathbf{n})\mathbf{n} + v_{s}(\mathbf{v} \cdot \mathbf{t})\mathbf{t}$ on Γ_{a}
- v : velocity vector
- <u>o</u> : stress tensor
- ► <u>ε</u> : strain tensor



2D Helmholtz equations

2D Helmholtz elastic equations

First order formulation of Helmholtz wave equations $\mathbf{x} = (x, y) \in \Omega \subset \mathbb{R}^2$,

$$\begin{cases} i\omega\rho(\mathbf{x})\mathbf{v}(\mathbf{x}) = \nabla \cdot \underline{\underline{\sigma}}(\mathbf{x}) + f_{s}(\mathbf{x}) \\ i\omega\underline{\underline{\sigma}}(\mathbf{x}) = \underline{\underline{C}}(\mathbf{x}) \underline{\underline{\varepsilon}}(\mathbf{v}(\mathbf{x})) \end{cases}$$

- Free surface condition : $\underline{\sigma}\mathbf{n} = 0$ on Γ_I
- Absorbing boundary condition : $\underline{\sigma}\mathbf{n} = v_p(\mathbf{v} \cdot \mathbf{n})\mathbf{n} + v_s(\mathbf{v} \cdot \mathbf{t})\mathbf{t}$ on Γ_a
- ρ : mass density
- <u>C</u>: tensor of elasticity coefficients

- v_p : P-wave velocity
- v_s : S-wave velocity
- f_s : source term, $f_s \in L^2(\Omega)$



Definitions

Contents

2D Helmholtz elastic equations

Notations and definitions

Hybridizable Discontinuous Galerkin method

Numerical results

Conclusions-Perspectives



M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 7/28

Notations

• \mathcal{T}_h mesh of Ω composed of triangles K





M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 8/28

Notations

- \mathcal{T}_h mesh of Ω composed of triangles K
- \mathcal{F}_h set of all faces F of \mathcal{T}_h





M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 8/28

Notations

- \mathcal{T}_h mesh of Ω composed of triangles K
- \mathcal{F}_h set of all faces F of \mathcal{T}_h
- **n** the normal outward vector of an element K





M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 8/28

Approximations spaces

• $P_p(K)$ set of polynomials of degree at most p on K



Approximations spaces

• $P_p(K)$ set of polynomials of degree at most p on K

►
$$\mathbf{V}_h^p = \{\mathbf{v} \in (L^2(\Omega))^2 : \mathbf{v}|_K \in \mathbf{V}^p(K) = (P_p(K))^2, \forall K \in \mathcal{T}_h\}$$



M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 9/28

Approximations spaces

▶ P_p(K) set of polynomials of degree at most p on K

►
$$\mathbf{V}_h^p = \{\mathbf{v} \in (L^2(\Omega))^2 : \mathbf{v}|_K \in \mathbf{V}^p(K) = (P_p(K))^2, \forall K \in \mathcal{T}_h\}$$

 $\blacktriangleright \Sigma_h^{p} = \{ \underline{\underline{\sigma}} \in \left(L^2(\Omega) \right)^3 : \underline{\underline{\sigma}} |_{\mathcal{K}} \in \Sigma^{p}(\mathcal{K}) = \left(P_{p}(\mathcal{K}) \right)^3, \forall \mathcal{K} \in \mathcal{T}_h \}$



M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 9/28

Approximations spaces

• $P_p(K)$ set of polynomials of degree at most p on K



M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 9/28

Definitions

• Jump $\llbracket \cdot \rrbracket$ of a vector **v** through F :

$$\llbracket \mathbf{v} \rrbracket = \mathbf{v}^+ \cdot \mathbf{n}^+ + \mathbf{v}^- \cdot \mathbf{n}^- = \mathbf{v}^+ \cdot \mathbf{n}^+ - \mathbf{v}^- \cdot \mathbf{n}^+$$

• Jump of a tensor $\underline{\sigma}$ through F :

$$\llbracket \underline{\underline{\sigma}} \rrbracket = \underline{\underline{\sigma}}^+ \mathbf{n}^+ + \underline{\underline{\sigma}}^- \mathbf{n}^- = \underline{\underline{\sigma}}^+ \mathbf{n}^+ - \underline{\underline{\sigma}}^- \mathbf{n}^+$$





M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 10/28

Contents

2D Helmholtz elastic equations

Notations and definitions

Hybridizable Discontinuous Galerkin method Formulation Algorithm

Numerical results

Conclusions-Perspectives



M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 10/28

Local HDG formulation

$$\begin{cases} i\omega\rho\mathbf{v}-\nabla\cdot\underline{\sigma} = 0\\ i\omega\underline{\sigma}-\underline{C}\underline{\varepsilon}(\mathbf{v}) = 0 \end{cases}$$

Innia

M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 11/28

Local HDG formulation

$$\begin{cases} \int_{K} i\omega\rho^{K} \mathbf{v}^{K} \cdot \mathbf{w} + \int_{K} \underline{\underline{\sigma}}^{K} : \nabla \mathbf{w} - \int_{\partial K} \underline{\widehat{\sigma}}^{\partial K} \cdot \mathbf{n} \cdot \mathbf{w} = 0\\ \int_{K} i\omega\underline{\underline{\sigma}}^{K} : \underline{\underline{\xi}} + \int_{K} \mathbf{v}^{K} \cdot \nabla \cdot \left(\underline{\underline{\zeta}}^{K} \underline{\underline{\xi}}\right) - \int_{\partial K} \mathbf{\widehat{v}}^{\partial K} \cdot \underline{\underline{\zeta}}^{K} \underline{\underline{\xi}} \cdot \mathbf{n} = 0\end{cases}$$

 $\underline{\widehat{g}}^{K}$ and $\mathbf{\widehat{v}}^{K}$ are numerical traces of $\underline{\underline{g}}^{K}$ and \mathbf{v}^{K} respectively on ∂K

Local HDG formulation

$$\begin{cases} \int_{K} i\omega\rho^{K} \mathbf{v}^{K} \cdot \mathbf{w} + \int_{K} \underline{\underline{\sigma}}^{K} : \nabla \mathbf{w} - \int_{\partial K} \underline{\underline{\hat{\sigma}}}^{\partial K} \cdot \mathbf{n} \cdot \mathbf{w} = 0\\ \int_{K} i\omega\underline{\underline{\sigma}}^{K} : \underline{\underline{\xi}} + \int_{K} \mathbf{v}^{K} \cdot \nabla \cdot \left(\underline{\underline{\zeta}}^{K} \underline{\underline{\xi}}\right) - \int_{\partial K} \mathbf{\hat{v}}^{\partial K} \cdot \underline{\underline{\zeta}}^{K} \underline{\underline{\xi}} \cdot \mathbf{n} = 0\end{cases}$$

We define :

$$\widehat{\mathbf{v}}^{F} = \lambda^{F}, \qquad \forall F \in \mathcal{F}_{h}, \\ \underline{\widehat{\sigma}}^{\partial K} \cdot \mathbf{n} = \underline{\sigma}^{K} \cdot \mathbf{n} - \tau \mathbf{I} \left(\mathbf{v}^{K} - \lambda^{\partial K} \right), \quad \text{on } \partial K$$

where au is the stabilization parameter (au > 0)



Local HDG formulation

We replace $\widehat{\mathbf{v}}^K$ and $\left(\underline{\widehat{c}}^K\cdot\mathbf{n}\right)$ by their definitions into the local equations

$$\begin{cases} \int_{K} i\omega \rho^{K} \mathbf{v}^{K} \cdot \mathbf{w} + \int_{K} \underline{\underline{\sigma}}^{K} : \nabla \mathbf{w} - \int_{\partial K} \underline{\underline{\sigma}}^{K} \cdot \mathbf{n} \cdot \mathbf{w} \\ + \int_{\partial K} \tau \mathbf{I} \left(\mathbf{v}^{K} - \lambda^{\partial K} \right) \cdot \mathbf{w} = 0 \\ \int_{K} i\omega \underline{\underline{\sigma}}^{K} : \underline{\underline{\xi}} + \int_{K} \mathbf{v}^{K} \cdot \nabla \cdot \left(\underline{\underline{\underline{\zeta}}}^{K} \underline{\underline{\xi}} \right) - \int_{\partial K} \lambda^{\partial K} \cdot \underline{\underline{\underline{\zeta}}}^{K} \underline{\underline{\xi}} \cdot \mathbf{n} = 0 \end{cases}$$



M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 12/28

Local HDG formulation

$$\begin{cases} \int_{K} i\omega \rho^{K} \mathbf{v}^{K} \cdot \mathbf{w} - \int_{K} \left(\nabla \cdot \underline{\underline{g}}^{K} \right) \cdot \mathbf{w} + \int_{\partial K} \tau \mathbf{I} \left(\mathbf{v}^{K} - \lambda^{\partial K} \right) \cdot \mathbf{w} = 0 \\ \int_{K} i\omega \underline{\underline{g}}^{K} : \underline{\underline{\xi}} + \int_{K} \mathbf{v}^{K} \cdot \nabla \cdot \left(\underline{\underline{\zeta}}^{K} \underline{\underline{\xi}} \right) - \int_{\partial K} \lambda^{\partial K} \cdot \underline{\underline{\zeta}}^{K} \underline{\underline{\xi}} \cdot \mathbf{n} = 0 \end{cases}$$



M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 13/28

Local HDG formulation

Innia

$$\begin{cases} \int_{K} i\omega \rho^{K} \mathbf{v}^{K} \cdot \mathbf{w} - \int_{K} \left(\nabla \cdot \underline{\underline{\sigma}}^{K} \right) \cdot \mathbf{w} + \int_{\partial K} \tau \mathbf{I} \left(\mathbf{v}^{K} - \lambda^{\partial K} \right) \cdot \mathbf{w} = 0 \\ \int_{K} i\omega \underline{\underline{\sigma}}^{K} : \underline{\underline{\xi}} + \int_{K} \mathbf{v}^{K} \cdot \nabla \cdot \left(\underline{\underline{\zeta}}^{K} \underline{\underline{\xi}} \right) - \int_{\partial K} \lambda^{\partial K} \cdot \underline{\underline{\zeta}}^{K} \underline{\underline{\xi}} \cdot \mathbf{n} = 0 \end{cases}$$

We define :

$$\underline{W}^{K} = \left(\underline{v_{x}}^{K}, \ \underline{v_{z}}^{K}, \ \underline{\sigma_{xx}}^{K}, \ \underline{\sigma_{zz}}^{K}, \ \underline{\sigma_{xz}}^{K}\right)^{T}$$

$$\underline{\Lambda} = \left(\underline{\Lambda}^{F_{1}}, \ \underline{\Lambda}^{F_{2}}, ..., \ \underline{\Lambda}^{F_{n_{f}}}\right)^{T}, \text{ where } n_{f} = \text{card}(\mathcal{F}_{h})$$

Discretization of the local HDG formulation

$$\mathbb{A}^{K}\underline{W}^{K} + \mathbb{C}^{K}\underline{\Lambda} = 0$$



March 5, 2015 - 13/28

Transmission condition

In order to determine λ^{K} , the continuity of the normal component of $\underline{\underline{\hat{\sigma}}}^{K}$ is weakly enforced, rendering this numerical trace conservative :

$$\int_{F} \llbracket \underline{\widehat{\sigma}}^{K} \cdot \mathbf{n} \rrbracket \cdot \eta = 0$$



Transmission condition

In order to determine λ^{K} , the continuity of the normal component of $\underline{\underline{\hat{\sigma}}}^{K}$ is weakly enforced, rendering this numerical trace conservative :

$$\int_{F} \llbracket \underline{\widehat{\sigma}}^{K} \cdot \mathbf{n} \rrbracket \cdot \eta = 0$$

Replacing $(\underline{\widehat{\sigma}}^K \cdot \mathbf{n})$ and summing over all faces, the transmission condition becomes :

$$\sum_{K\in\mathcal{T}_h}\int_{\partial K} \left(\underline{\underline{\sigma}}^K\cdot\mathbf{n}\right)\cdot\eta - \sum_{K\in\mathcal{T}_h}\int_{\partial K}\tau\mathbf{l}\left(\mathbf{v}^K-\lambda^{\partial K}\right)\cdot\eta = 0$$



M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 14/28

Transmission condition

In order to determine λ^{K} , the continuity of the normal component of $\underline{\underline{\hat{\sigma}}}^{K}$ is weakly enforced, rendering this numerical trace conservative :

$$\int_{F} \llbracket \underline{\widehat{\sigma}}^{K} \cdot \mathbf{n} \rrbracket \cdot \eta = 0$$

Replacing $(\underline{\widehat{\sigma}}^K \cdot \mathbf{n})$ and summing over all faces, the transmission condition becomes :

$$\sum_{\mathbf{K}\in\mathcal{T}_{h}}\int_{\partial K}\left(\underline{\underline{\sigma}}^{\mathbf{K}}\cdot\mathbf{n}\right)\cdot\eta-\sum_{\mathbf{K}\in\mathcal{T}_{h}}\int_{\partial K}\tau\mathbf{I}\left(\mathbf{v}^{\mathbf{K}}-\boldsymbol{\lambda}^{\partial \mathbf{K}}\right)\cdot\eta=0$$

Discretization of the transmission condition

$$\sum_{K\in\mathcal{T}_{h}}\left[\mathbb{B}^{K}\underline{W}^{K}+\mathbb{L}^{K}\underline{\Lambda}\right]=0$$

Global HDG formulation

$$\begin{cases} \int_{K} i\omega\rho^{K} \mathbf{v}^{K} \cdot \mathbf{w} - \int_{K} (\nabla \cdot \underline{\underline{\sigma}}^{K}) \cdot \mathbf{w} + \int_{\partial K} \tau \mathbf{I} (\mathbf{v}^{K} - \lambda^{\partial K}) \cdot \mathbf{w} = 0 \\ \int_{K} i\omega\underline{\underline{\sigma}}^{K} : \underline{\xi} + \int_{K} \mathbf{v}^{K} \cdot \nabla \cdot \left(\underline{\underline{C}}^{K} \underline{\xi}\right) - \int_{\partial K} \lambda^{\partial K} \cdot \underline{\underline{C}}_{K} \underline{\xi} \cdot \mathbf{n} = 0 \\ \sum_{K \in \mathcal{T}_{h}} \int_{\partial K} \left(\underline{\underline{\sigma}}^{K} \cdot \mathbf{n}\right) \cdot \eta - \sum_{K \in \mathcal{T}_{h}} \int_{\partial K} \tau \mathbf{I} (\mathbf{v}^{K} - \lambda^{\partial K}) \cdot \eta = 0 \end{cases}$$



Global HDG formulation

$$\begin{cases} \int_{K} i\omega \rho^{K} \mathbf{v}^{K} \cdot \mathbf{w} - \int_{K} (\nabla \cdot \underline{\underline{\sigma}}^{K}) \cdot \mathbf{w} + \int_{\partial K} \tau \mathbf{I} (\mathbf{v}^{K} - \lambda^{\partial K}) \cdot \mathbf{w} = 0\\ \int_{K} i\omega \underline{\underline{\sigma}}^{K} : \underline{\underline{\xi}} + \int_{K} \mathbf{v}^{K} \cdot \nabla \cdot \left(\underline{\underline{\zeta}}^{K} \underline{\underline{\xi}}\right) - \int_{\partial K} \lambda^{\partial K} \cdot \underline{\underline{\zeta}}_{K} \underline{\underline{\xi}} \cdot \mathbf{n} = 0\\ \sum_{K \in \mathcal{T}_{h}} \int_{\partial K} \left(\underline{\underline{\sigma}}^{K} \cdot \mathbf{n}\right) \cdot \eta - \sum_{K \in \mathcal{T}_{h}} \int_{\partial K} \tau \mathbf{I} (\mathbf{v}^{K} - \lambda^{\partial K}) \cdot \eta = 0\end{cases}$$

Global HDG discretization

$$\int_{K\in\mathcal{T}_{h}}^{K} \frac{W}{K}^{K} + \mathbb{C}^{K}\underline{\Lambda} = 0$$
$$\sum_{K\in\mathcal{T}_{h}} \left[\mathbb{B}^{K}\underline{W}^{K} + \mathbb{L}^{K}\underline{\Lambda}\right] = 0$$

Global HDG formulation

$$\begin{cases} \int_{K} i\omega \rho^{K} \mathbf{v}^{K} \cdot \mathbf{w} - \int_{K} (\nabla \cdot \underline{\underline{\sigma}}^{K}) \cdot \mathbf{w} + \int_{\partial K} \tau \mathbf{I} (\mathbf{v}^{K} - \lambda^{\partial K}) \cdot \mathbf{w} = 0\\ \int_{K} i\omega \underline{\underline{\sigma}}^{K} : \underline{\underline{\xi}} + \int_{K} \mathbf{v}^{K} \cdot \nabla \cdot \left(\underline{\underline{\zeta}}^{K} \underline{\underline{\xi}}\right) - \int_{\partial K} \lambda^{\partial K} \cdot \underline{\underline{\zeta}}_{K} \underline{\underline{\xi}} \cdot \mathbf{n} = 0\\ \sum_{K \in \mathcal{T}_{h}} \int_{\partial K} \left(\underline{\underline{\sigma}}^{K} \cdot \mathbf{n}\right) \cdot \eta - \sum_{K \in \mathcal{T}_{h}} \int_{\partial K} \tau \mathbf{I} (\mathbf{v}^{K} - \lambda^{\partial K}) \cdot \eta = 0\end{cases}$$

Global HDG discretization

$$\int \underline{W}^{\kappa} = -(\mathbb{A}^{\kappa})^{-1} \mathbb{C}^{\kappa} \underline{\Lambda}$$
$$\sum_{K \in \mathcal{T}_{h}} \left[\mathbb{B}^{\kappa} \underline{W}^{\kappa} + \mathbb{L}^{\kappa} \underline{\Lambda} \right] = 0$$

Global HDG formulation

$$\begin{cases} \int_{K} i\omega \rho^{K} \mathbf{v}^{K} \cdot \mathbf{w} - \int_{K} (\nabla \cdot \underline{\underline{\sigma}}^{K}) \cdot \mathbf{w} + \int_{\partial K} \tau \mathbf{I} (\mathbf{v}^{K} - \lambda^{\partial K}) \cdot \mathbf{w} = 0 \\ \int_{K} i\omega \underline{\underline{\sigma}}^{K} : \underline{\xi} + \int_{K} \mathbf{v}^{K} \cdot \nabla \cdot \left(\underline{\underline{C}}^{K} \underline{\underline{\xi}}\right) - \int_{\partial K} \lambda^{\partial K} \cdot \underline{\underline{C}}_{K} \underline{\underline{\xi}} \cdot \mathbf{n} = 0 \\ \sum_{K \in \mathcal{T}_{h}} \int_{\partial K} \left(\underline{\underline{\sigma}}^{K} \cdot \mathbf{n}\right) \cdot \eta - \sum_{K \in \mathcal{T}_{h}} \int_{\partial K} \tau \mathbf{I} (\mathbf{v}^{K} - \lambda^{\partial K}) \cdot \eta = 0 \end{cases}$$

Global HDG discretization

$$\sum_{K\in\mathcal{T}_h} \left[-\mathbb{B}^K (\mathbb{A}^K)^{-1} \mathbb{C}^K + \mathbb{L}^K \right] \underline{\Lambda} = 0$$

(nría_

M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 15/28

1. Construction of the linear system
$$\mathbb{M}\underline{\Lambda} = 0$$

with $\mathbb{M} = \sum_{K \in \mathcal{T}_h} \left[-\mathbb{B}^K (\mathbb{A}^K)^{-1} \mathbb{C}^K + \mathbb{L}^K \right]$
for $K = 1$ to Nb_{tri} do
Compute matrices \mathbb{B}^K , $(\mathbb{A}^K)^{-1}$, \mathbb{C}^K and \mathbb{L}^K
Construction of the corresponding section of
end for



M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 16/28

 \mathbb{K}

- 1. Construction of the linear system $\mathbb{M}\underline{\Lambda} = 0$
- 2. Construction of the right hand side ${\mathbb S}$



M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 16/28

- 1. Construction of the linear system $\mathbb{M}\underline{\Lambda} = 0$
- 2. Construction of the right hand side \mathbb{S}
- 3. Resolution $\mathbb{M}\underline{\Lambda} = \mathbb{S}$



- 1. Construction of the linear system $\mathbb{M}\underline{\Lambda} = 0$
- 2. Construction of the right hand side \mathbb{S}
- 3. Resolution $\mathbb{M}\underline{\Lambda} = \mathbb{S}$
- 4. Computation of the solutions of the initial problem

for
$$K = 1$$
 to Nb_{tri} do
Compute $\underline{W}^{K} = -(\mathbb{A}^{K})^{-1}\mathbb{C}^{K}\underline{\Lambda}$
end for



Contents

2D Helmholtz elastic equations

Notations and definitions

Hybridizable Discontinuous Galerkin method

Numerical results

Plane wave in an homogeneous medium Marmousi test-case

Conclusions-Perspectives



M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 16/28

Plane wave



 $\begin{array}{l} \text{Computational domain } \Omega \\ \text{setting} \end{array}$

Physical parameters :

•
$$\rho = 2000 kg.m^{-3}$$

•
$$\lambda = 16 GPa$$

Plane wave :

$$u = \nabla e^{i(k\cos\theta x + k\sin\theta y)}$$

where
$$k = \frac{\omega}{v_p}$$

- ► Three meshes :
 - 3000 elements
 - 10000 elements
 - 45000 elements



March 5, 2015 - 17/28

Plane wave



Convergence order of the HDG scheme



1. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 18/28

Marmousi test-case



Computational domain Ω composed of 235000 triangles



M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 19/28

Parallel results for the Marmousi test-case with the HDG-P3 scheme, f = 2Hz



Caracteristics of the computing processors used

• Plafrim platform

- Hardware specification : 16 nodes, 12 cores by nodes
- Caracteristics of computing nodes :
 - ▶ 2 Hexa-core Westmere Intel[®] Xeon[®] X5670
 - Frequency : 2,93 GHz
 - Cache L3 : 12 Mo
 - RAM : 96 Go
 - Infiniband DDR : 20Gb/s
 - Ethernet : 1Gb/s



Efficiency of the parallelism of the global matrix construction



Efficiency of the parallelism of the whole algorithm





M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 23/28

Speed up for the global matrix construction





M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 24/28

Speed up (Total simulation time)





M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 25/28

Numerical results Marmousi test-case

Memory required (GB) for the simulation





M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 26/28

Conclusion

Contents

2D Helmholtz elastic equations

Notations and definitions

Hybridizable Discontinuous Galerkin method

Numerical results

Conclusions-Perspectives



M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 26/28

Conclusions-Perspectives

Conclusions

On a same mesh, with the HDG method :

- Memory gain
- Computational time gain



Conclusions-Perspectives

Conclusions

On a same mesh, with the HDG method :

- Memory gain
- Computational time gain

Perspectives

- Develop 3D HDG formulation for Helmholtz equations
- Solution strategy for the HDG linear system



M. Bonnasse-Gahot - DG and HDG methods for Helmholtz wave equations

March 5, 2015 - 27/28

Conclusion

Thank you!

