



HAL
open science

Efficient Computation of Dual Space and Directional Multiplicity of an Isolated Point

Angelos Mantzaflaris, Hamid Rahkooy, Zafeirakis Zafeirakopoulos

► **To cite this version:**

Angelos Mantzaflaris, Hamid Rahkooy, Zafeirakis Zafeirakopoulos. Efficient Computation of Dual Space and Directional Multiplicity of an Isolated Point. [Technical Report] JKU. 2015. hal-01214195v1

HAL Id: hal-01214195

<https://inria.hal.science/hal-01214195v1>

Submitted on 10 Oct 2015 (v1), last revised 11 May 2016 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Efficient Computation of Dual Space and Directional Multiplicity of an Isolated Point

Angelos Mantzaflaris^a, Hamid Rahkooy^b, Zafeirakis Zafeirakopoulos^c

^a*Radon Institute for Computational and Applied Mathematics, Austrian Academy of Sciences, Linz, Austria*

^b*RISC & DK Computational Mathematics, Linz, Austria*

^c*Department of Mathematics, Galatasaray University, Istanbul, Turkey*

Abstract

Isolated singularities typically occur at self-intersection points of planar algebraic curves, curve offsets, intersections between spatial curves and surfaces, and so on. The information characterizing the singularity can be captured in a local dual basis, expressing combinations of vanishing derivatives at the singular point. Macaulay's algorithm is a classic algorithm for computing such a basis, for a point in an algebraic set. The integration method of Mourrain constructs much smaller matrices than Macaulay's approach, by performing integration on previously computed elements.

In this work we are interested in the efficiency of dual basis computation, as well as its relation to orthogonal projection. First, we introduce an easy to implement criterion that avoids redundant computations during the computation of the dual basis, by deleting certain columns from the matrices in the integration method. In doing so, we explore general (non-monomial) bases for the associated primal quotient ring. Experiments show the efficient behaviour of the improved method. Second, we introduce the notion of directional multiplicity, which expresses the multiplicity structure with respect to an axis, and is useful in understanding the geometry behind projection. We use this notion to shed light on the gap between the degree of the generator of the elimination ideal and the corresponding factor in the resultant.

Keywords: multiplicity structure, dual space, directional multiplicity, Gröbner basis, algebraic curves

2000 MSC: 13P10,

2000 MSC: 68W39

Email addresses: Angelos.Mantzaflaris@oeaw.ac.at (Angelos Mantzaflaris), rahkooy@risc.jku.at (Hamid Rahkooy), zafeirakopoulos@gmail.com (Zafeirakis Zafeirakopoulos)

Preprint submitted to Elsevier

1. Introduction

Multiple points correspond to singular intersections between geometric objects. Therefore special care has to be taken in geometric algorithms in the vicinity of such points [1, 13]. Singularities also arise while offsetting planar curves, and their robust computation is a long-standing problem. For example, in [22] the problem is treated using an algebraic approach. In implicit representations, one has access to evaluation of the level set function, rather than parametrized coordinates, therefore evaluation at arbitrary points in space is the basic tool at hand to be used for computations on this representation. For instance the topology or even the change of representation to implicit form is possible using solely the evaluation at arbitrary points [2, 6, 8]. Horner's scheme provides a stable and fast method to evaluate a polynomial in monomial representation. In the present paper we use value and derivative evaluations to identify singular solution via dual spaces.

The dual space of polynomial rings and algorithms for computing it, as well as the multiplicity structure of isolated points have been studied in different contexts in the literature, such as root deflation and certification, approximate root refinement, and so on [9, 12, 18, 19, 20, 21, 23, 31, 35] and has been recently studied for the positive dimensional case in [14, 15]. In the following we present a short literature review along with an introduction of notions, theorems and notation, which we will use in later sections.

Let $R = \mathbb{K}[x_1, \dots, x_n]$ and consider it as a vector space over \mathbb{K} . Denote by \hat{R} the dual of R . Let $\zeta = (\zeta_1, \dots, \zeta_n) \in \mathbb{K}^n$ and $\mathbf{a} = (a_1, \dots, a_n) \in \mathbb{N}^n$, and consider the differential operator $\partial_\zeta^{\mathbf{a}}$ that acts on $p \in R$ first by differentiation and then by evaluation at the point ζ . In [28], Proposition 2.2 states that there is an isomorphism of \mathbb{K} -vector spaces between \hat{R} and $\mathbb{K}[[\partial_\zeta]]$. This vector space, in general, is not finite dimensional. The orthogonal of an ideal I of R , i.e., $I^\perp = \left\{ \lambda \in \hat{R} : \lambda(f) = 0 \quad \forall f \in I \right\}$ can be seen as a linear subspace of $\mathbb{K}[[\partial_\zeta]]$, for every $\zeta \in \mathbb{K}^n$, as shown in Proposition 2.6 of [28]. For the rest of this work, unless otherwise stated, we assume that $\zeta = 0$ and we may omit the subscript, i.e., use $\partial^{\mathbf{a}}$ instead of $\partial_\zeta^{\mathbf{a}}$ and $\mathbb{K}[[\partial]]$ instead of $\mathbb{K}[[\partial_\zeta]]$.

In [27, 26], Marinari, Mora and Möller have shown that the \mathfrak{m}_ζ -primary ideals are in one-to-one correspondence with the non-null finite-dimensional vector spaces of $\mathbb{K}[[\partial]]$. Thus, the dual of an \mathfrak{m}_ζ -primary ideal can be treated computationally and its dimension is the multiplicity of the isolated point. From now on, given an ideal $I = \langle f_1, \dots, f_e \rangle \subseteq R$, we denote by Q_ζ , the \mathfrak{m}_ζ -primary component of I associated to an isolated point ζ in the variety of I . D will stand for a basis for Q_ζ^\perp . Therefore, according to Marinari, Mora and Möller $\langle D \rangle = Q_\zeta^\perp = I^\perp \cap \mathbb{K}[[\partial_\zeta]]$.

The maximum degree of the dual basis is also known as the depth. Together with the breadth, i.e. the co-rank of the Jacobian matrix, they are basic parameters that quantify the complexity of the multiplicity structure. The *Nil-index* of an \mathfrak{m}_ζ -primary ideal Q_ζ is the maximum integer $\mathcal{N} \in \mathbb{N}$ such that $\mathfrak{m}_\zeta^{\mathcal{N}} \not\subseteq Q_\zeta$. There is a tight connection between the dual space of \mathfrak{m}_ζ -primary ideals and

their Nil-index given by Mourrain as Lemma 3.3 in [28], namely, considering the differential operators as polynomials, the maximum degree of the elements of $I^\perp \cap \mathbb{K}[\partial_\zeta]$ is equal to the Nil-index of Q_ζ . Therefore, we can find the monomials of a basis D for Q^\perp by searching incrementally, i.e., degree by degree, among those monomials of I^\perp that have degree at most the Nil-index.

For a recent review on computing the dual space see [24]. The dual space of polynomial rings and its properties and applications have been thoroughly reviewed in [9]. Concerning literature on computations related to the dual bases, Wu and Zhi worked on a symbolic-numeric method for computing the primary components and the differential operators [34], which is based on an algorithm for determining the dual space appearing in [31] by Stetter. A specialized algorithm for the case of breadth (Jacobian cor-rank) one is devised in [17].

Below we present the main ideas behind the algorithms in the literature and in particular we explain Macaulay's algorithm [21] and the integration method [23, 28].

Due to the degree bound mentioned above, the algorithms that compute a basis D for Q^\perp proceed degree by degree. Let D_t be the subset of D that contains degree at most t elements of D (which is also a basis for the degree at most t part of Q^\perp , denoted by Q_t^\perp). Obviously $D_0 = \langle 1 \rangle$. The algorithms extend D_t into D_{t+1} , until $D_t = D_{t+1}$. Then we can conclude that $D = D_t$ and $t = \mathcal{N}$. We set $d_i := dx_i$ for presentation reasons in what follows.

A historical work conducted by Macaulay [21] shows how to construct matrices in order to compute a basis for the dual space Q^\perp . It is based on a simple condition that the coefficients of the elements of the dual space must fulfill. Let $\Lambda = \sum_{|\alpha| \leq \mathcal{N}} \lambda_\alpha d^\alpha$, where we use the multi-index notation $d = d_1 d_2 \cdots d_n$. Then

$\Lambda(f) = 0$ for all $f \in I$ if and only if $\Lambda(x^\beta f_i) = 0$ for all $\beta \in \mathbb{N}^n$ and $1 \leq i \leq e$. This observation, for $1 \leq |\beta| \leq \mathcal{N}$, reduces checking that $\Lambda(f) = 0$ for an infinite number of polynomials f into checking the finitely many conditions that are given in the right hand side. Namely, it suffices to impose conditions on λ_α 's, the coefficients of Λ . For $1 \leq |\beta| \leq \mathcal{N}$, we obtain a system of linear homogeneous equations and construct the corresponding matrix. The rows of this matrix are labeled by $x^\beta f_i$ and the columns are labeled by d^α . Every element in the kernel of this matrix is a coefficient vector, corresponding to an element of Q_ζ^\perp . Therefore, the problem is now reduced into the kernel computation of some specific matrices.

This algorithm is still used widely, although it is not efficient. The main obstacles are the large size of the matrices and that the new matrices built contain as submatrices previously constructed matrices, repeating some computations. In [35] Zeng used the ideas in Stetter's algorithm [31] and introduced his closedness property in order to make Macaulay's matrices smaller.

Mourrain, in [28], gave a new algorithm based on integration, which constructs matrices of smaller size than those in Macaulay's algorithm. Assume that we have computed $D_{t-1} = \{\Lambda_1, \dots, \Lambda_m\}$. The algorithm is based on *integrating* elements of D_{t-1} in order to ansatz the elements of D_t with symbolic

coefficients λ_{ij} . More precisely,

$$\Lambda = \sum_{i=1}^m \sum_{k=1}^n \lambda_{ik} \int_k \Lambda_i(d_1, \dots, d_k, 0, \dots, 0). \quad (1)$$

Then applying the following conditions on Λ guarantees that $\Lambda \in Q_t^\perp$.

$$\forall 1 \leq k \leq e, \quad \Lambda(f_k) = 0 \quad (2)$$

$$\forall 1 \leq k < l \leq n, \quad \sum_{i=1}^1 \lambda_{ik} d_l(\Lambda_i) - \sum_{i=1}^1 \lambda_{il} d_k(\Lambda_i) = 0. \quad (3)$$

From the above procedure one obtains a system of equations for λ_{ij} to which a matrix M_t is associated. Similar to Macaulay's algorithm, each vector in the kernel of M_t determines the coefficients of an element in Q_t^\perp . The columns of M_t are labeled by $\lambda_{ij} p_{ij}$ for $1 \leq i \leq e$ and $1 \leq j \leq n$, where $p_{ij} = \int_j \Lambda_i(d_1, \dots, d_j, 0, \dots, 0)$. By abuse of notation and for simplifying the presentation, we sometimes use λ_{ij} or p_{ij} instead of $\lambda_{ij} p_{ij}$ in order to label the columns of M_t .

Although the integration method constructs much smaller matrices than Macaulay's algorithm, however, it might recompute D_{t-1} in step t . In [23], the authors gave a method to overcome this problem. They use the tight connection between a dual basis for Q_{t-1}^\perp and a primal basis for the degree $t-1$ part of the quotient R/Q . More precisely, if $D_{t-1} = \{\Lambda_1, \dots, \Lambda_m\}$ is a basis of Q_{t-1}^\perp , where $\Lambda_i = d^{\beta_i} + \sum_{j=1}^{s-\mu} \lambda_{ij} d^{\gamma_j}$, then $\{x^{\beta_1}, \dots, x^{\beta_\mu}\}$ is a basis for the degree $t-1$ part of R/Q . Details of the above has been explained in [28] and [23]. The following theorem by the first author and Mourrain results in avoiding redundant computations.

Theorem 1 (Lemma 3.4 in [23]). *Let $D_t = \{\Lambda_1, \dots, \Lambda_m\}$ be a basis of Q_{t-1}^\perp , such that it yields the basis $\{x^{\beta_1}, \dots, x^{\beta_\mu}\}$ for the degree $t-1$ part of R/Q . An element $\Lambda \in \mathbb{K}[\partial]$ is not zero in $\Lambda \in D_t \setminus D_{t-1}$ if and only if in addition to conditions 2 and 3, we impose*

$$\Lambda(x^{\beta_i}) = 0, \quad \forall 1 \leq i \leq m. \quad (4)$$

Equation 4 adds new rows to the matrix M_t , which results in a larger matrix that we denote by \widetilde{M}_t . By Condition 4, we refer to Equation 4, indicating its complementary nature to the Conditions 2 and 3. However, by clever choice of a primal-dual basis, adding new rows can lead to removing some column.

Example 1. *Let*

$$\begin{aligned} f_1 &= x^2 + (y-1)^2 - 1 \\ f_2 &= y^2. \end{aligned}$$

Then for the root $(0, 0)$, we have

<i>Macaulay</i>	<i>Integration</i>
$M_1 = \begin{matrix} & 1 & d_1 & d_2 \\ f_1 & \begin{pmatrix} 0 & 0 & -2 \\ 0 & 0 & 0 \end{pmatrix} \\ f_2 & \end{matrix}$	$M_1 = \begin{matrix} & 1 & d_1 & d_2 \\ f_1 & \begin{pmatrix} 0 & 0 & -2 \\ 0 & 0 & 0 \end{pmatrix} \\ f_2 & \end{matrix}$

The kernel of M_1 yields the basis $D_1 = \{1, d_1\}$. In the second step, we have

<i>Macaulay</i>	<i>(Improved) Integration</i>
$M_2 = \begin{matrix} & 1 & d_1 & d_2 & d_1^2 & d_1 d_2 & d_2^2 \\ f_1 & \begin{pmatrix} 0 & 0 & -2 & 1 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & -2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -2 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \\ x_1 f_1 & \\ x_1 f_2 & \\ x_2 f_1 & \\ x_2 f_2 & \end{matrix}$	$\widetilde{M}_2 = \begin{matrix} & d_1 & d_2 & d_1^2 & d_1 d_2 \\ \text{Condition 4} & \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & -2 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix} \\ \text{Condition 3} & \\ \Lambda(f_1) & \\ \Lambda(f_2) & \end{matrix}$

from which we obtain $D_2 = \{1, d_1, 2d_1^2 + d_2\}$. Condition 4 implies that $\lambda_1 = 0$.

Therefore we can remove the first column from M_2 to obtain \widetilde{M}_2 .

The algorithm runs until step 4, during which we have $D_3 = D_4 = \{1, d_1, 2d_1^2 + d_2, 2d_1^3 + d_1 d_2\}$.

A detailed review of the integration method and its application to root deflation methods is given in [24]. In [3], Bates, Peterson and Sommese study the multiplicity of the primary components. Examining the multiplicity structure via deflation is exhibited in the work of Dayton and Zeng [7] and Leykin and Verschelde [16].

The present work was motivated (see [30]) and is closely connected to elimination theory. Marinari, Mora and Möller's work [26, 27] includes studying the behaviour of the dual space under projection, which is the base of our result related to the use of dual elements to study the elimination ideal in Section 3. There are different viewpoints on multiplicity structure. It can be studied via

- Gröbner bases: obtain global information on the elimination ideal and the multiplicity [4, 5].
- Resultants: obtain the multiplicity at a point, but not the elimination ideal [10, 11, 32, 33].
- Dual spaces: obtain the local structure of the multiplicity at that point as well as the elimination ideal and the Gröbner basis.

Apart from the application of Gröbner bases in computing a basis for the quotient ring and therefore the multiplicity, we will extensively use its elimination property [5] that allows computing elimination ideals. Mourrain in [28] proved that having a basis for the orthogonal of an ideal with an \mathfrak{m}_ζ -primary component, one can obtain the reduced Gröbner basis of this component.

Contributions. One contribution is improving the dual basis computation. Since the size of the matrices constructed in each step of the algorithms is the main obstacle in computations, we propose criteria that allow deleting some columns from the matrices in order to reduce the size of the matrices. Apart from those criteria, we introduce the notion of directional multiplicity, which gives more information about the singularity and its projections. An interesting interplay between the directional multiplicity and the degree of the elimination ideal is illustrated using intersections of algebraic curves.

For the integration method, the state of the art algorithm, in Proposition 5 we give an explicit generalization of the improvement in [23], as we detect and use a polynomial basis for the quotient rather than the monomial basis. The new primal bases are in accordance with [28, Prop. 3.7], which can be generalized for the case in question. Corollary 2 shows our criterion for deleting some columns such that the kernel of the new matrix only detects new members of a basis of the dual space, which avoids recomputing the lower degree basis elements that are obtained in the previous steps. The reduction of lower-degree elements has been employed in [23] using a different criterion; however, under certain circumstances this criterion can increase the row-size of the matrix.

Outline. Section 2 includes our main results. The main improvements for computing the dual space are presented and their advantages are discussed. isolated point. We introduce the notion of directional multiplicity and use the extended Buchberger diagram to illustrate it. We show bounds on the directional multiplicities with respect to Nil-index and the intersection multiplicity. Section 4 includes an empirical study of the algorithms presented. The theoretical expected behaviour is compared with the performance of the algorithms in practice. Finally, in Section 5 we summarize and conclude this work and provide future directions of research.

2. Efficient Algorithms for Dual Basis

Following the paradigm of [23], we provide modifications to the integration method and Macaulay's algorithms, which make computations more efficient. In particular, we give a more efficient criterion than Condition 4 for the integration method.

Modifications on Integration Method. Recall that, in the integration method, the columns of M_t are labeled by the $\lambda_{ij}p_{ij}$'s appearing in Λ , but we may refer to the label just by λ_{ij} or p_{ij} for simplicity. Fix one of the λ_{ij} 's and call it λ . We denote by v_λ the column of M_t that is indexed by λ and p_λ denotes the corresponding polynomial.

In step t of the algorithm, there exists a correspondence between D_t and K_t , a basis of $\text{Ker}(M_t)$. If there exists a vector $q \in K_t$ for which the coordinate corresponding to λ in this vector is nonzero, then we say that v_λ is active in D_t . In case we explicitly know such a vector q , i.e., a particular element of the kernel corresponding to an element E of D_t , then we say that v_λ is *active* in E .

Since M_{t-1} is a sub-matrix of M_t , if it is clear from the context, by a *column* of M_{t-1} we will refer to the corresponding column in M_t as well. If $\lambda_1, \dots, \lambda_m$ are the labels of some columns in M_t , then $M_t^{\lambda_1, \dots, \lambda_m}$ denotes the submatrix of M_t that is obtained by deleting columns $\lambda_1, \dots, \lambda_m$ from M_t . We work on M_t rather than \widetilde{M}_t in this section, although many of our arguments are correct for \widetilde{M}_t as well.

We start with a proposition that provides us with an improvement on the integration method, related to Condition 4.

Proposition 1. *Let $M_t, M_{t-1}, D_t, \Lambda_i$ ($1 \leq i \leq m$), λ, p_λ and v_λ be as above. Then the following hold.*

1. *If v_λ is a column of M_t , then v_λ is active in D_t if and only if v_λ can be reduced to zero by other columns of M_t .*
2. *For all $1 \leq i \leq m$, if v_{λ_i} is active in Λ_i , $K_t^{\lambda_i}$ is a basis for $\text{Ker}(M_t^{\lambda_i})$ and $D_t^{\lambda_i}$ is the set of its corresponding dual elements, then $\{\Lambda_i\} \cup D_t^{\lambda_i}$ is a basis for the degree t part of Q^\perp . Moreover, if v_{λ_i} is active in Λ_i , but is not active in Λ_j for $1 \leq j \neq i \leq m$, then there exists a basis $D_t^{\lambda_i}$ such that $\Lambda_j \in D_t^{\lambda_i}, j \neq i$.*
3. *Let $K_t^{\lambda_1 \dots \lambda_m}$ be a basis for $\text{Ker}(M_t^{\lambda_1 \dots \lambda_m})$ and $D_t^{\lambda_1 \dots \lambda_m}$ be the set of its corresponding dual elements. If for all $1 \leq i \leq m$, v_{λ_i} is active in Λ_i but not active in $\Lambda_1, \dots, \Lambda_{i-1}$, then $D_{t-1} \cup D_t^{\lambda_1 \dots \lambda_m}$ is a basis for the degree t part of Q^\perp .*

Proof. 1. Let $v_\lambda, v_1, \dots, v_k$ denote the columns of M_t and $p_\lambda, p_1, \dots, p_k$ be the polynomials labeling the columns of M_t . Then v_λ can be reduced to zero by v_1, \dots, v_k if and only if there exist $c_1, \dots, c_k \in \mathbb{K}$, such that $v_\lambda = c_1 v_1 + \dots + c_k v_k$, or equivalently $v_\lambda - c_1 v_1 - \dots - c_k v_k = 0$. This holds if and only if $q := (1, c_1, \dots, c_k) \in K_t$, which holds if and only if $\Lambda' := p_\lambda - c_1 p_1 - \dots - c_k p_k \in D_t$ (Note that this is exactly the fact that Λ' in D_t corresponds to $q \in K_t$). The latter is the case if and only if v_λ is active in Λ' , or equivalently v_λ is active in D_t .

2. Fix $1 \leq i \neq j \leq m$ and let q_i and q_j be the elements of K_t corresponding to Λ_i and Λ_j in D_t , respectively.

First we prove that for all $\Lambda' \in D_t$ if $\Lambda' \neq \Lambda_i$, then $\Lambda' \in \langle D_t^{\lambda_i} \cup \{\Lambda_i\} \rangle$.

Let q' be the corresponding element of Λ' in K_t . If v_{λ_i} is not active in Λ' , then by part 1 it cannot be reduced to zero by the active columns in Λ' . So the column v_{λ_i} is not involved in computing Λ' via column reducing in M_t . Thus, Λ' can be computed via column reducing in $M_t^{\lambda_i}$. Let q' be the corresponding element to Λ' in $\text{Ker}(M_t)$. Then $q' \in \text{Ker}(M_t^{\lambda_i})$. This means that $\Lambda' \in \langle D_t^{\lambda_i} \rangle$.

If v_{λ_i} is active in Λ' , then we prove that there exists a Λ'' in $D_t^{\lambda_i}$ such that $\Lambda' = \Lambda_i + \Lambda''$. Let $q' \in K_t$ be the element corresponding to $\Lambda' \in D_t$, such that that the first coordinate of q' corresponds to v_{λ_i} . Take $q' = (1, b_1, \dots, b_k)$. Then we have that $v_{\lambda_i} + b_1 v_1 + b_2 v_2 + \dots + b_k v_k = 0$, where

the columns v_1, \dots, v_k are as in the proof of part 1. Also, again as in the proof of the part 1, $v_{\lambda_i} = c_1 v_1 + \dots + c_k v_k$. Therefore $(b_1 - c_1)v_1 + \dots + (b_k - c_k)v_k = 0$, which means that $(0, b_1 - c_1, \dots, b_k - c_k) \in \text{Ker}(M_t)$, and therefore $q'' := (b_1 - c_1, \dots, b_k - c_k) \in \text{Ker}(M_t^\lambda)$. So one can construct a basis $K_t^{\lambda_i}$ in such a way that $q'' \in K_t^{\lambda_i}$. Let Λ'' be the member of $D_t^{\lambda_i}$ corresponding to q'' . Then $\Lambda' = \Lambda_i + \Lambda''$.

Secondly we note that if v_{λ_i} is not active in Λ_j , for $1 \leq j \neq i \leq m$, then by the above argument, one can compute a basis $K_t^{\lambda_i}$ (and respectively, $D_t^{\lambda_i}$) in such a way that $\Lambda_j \in D_t^{\lambda_i}$.

Thus, every element of D_t can be obtained from Λ_i and an element of $K_t^{\lambda_i}$ and therefore $\langle D_t \rangle \subseteq \langle \{\Lambda_i\} \cup D_t^{\lambda_i} \rangle$. Since the elements of $\{\Lambda_i\} \cup D_t^{\lambda_i}$ are linearly independent, we have $\langle \{\Lambda_i\} \cup D_t^{\lambda_i} \rangle = \langle D_t \rangle = Q^\perp$.

3. Let $K_t^{v_{\lambda_1} \dots v_{\lambda_m}}$ be a basis for $\text{Ker}(M_t^{\lambda_1 \dots \lambda_m})$ and $D_t^{\lambda_1 \dots \lambda_m}$ the corresponding dual elements. As in the proof of the previous parts, let K_t be a basis for $\text{Ker}(M_t)$ and $q_1, \dots, q_m \in K_t$ correspond to $\Lambda_1, \dots, \Lambda_m$ respectively. Then from the proof of part 2 we have that $\{q_1\} \cup K_t^{\lambda_1}$ is a basis for $\text{Ker}(M_t)$. Also by part 2 of the proposition, $q_2, \dots, q_m \in \langle K_t \rangle$ and correspondingly $\Lambda_2, \dots, \Lambda_m \in \langle D_t^{\lambda_1} \rangle$. Now consider the matrix $M_t^{\lambda_1}$ and the basis $D_t^{\lambda_1}$ obtained from it. Since v_{λ_2} is active in Λ_2 (which corresponds to q_2 in K_t), and it is not active in Λ_1 , then we can apply part 2 of the proposition to the matrix $M_t^{\lambda_1}$ and the basis $D_t^{\lambda_1}$ obtained by it. Then we will have that $\{q_2\} \cup K_t^{\lambda_1 \lambda_2}$ is a basis for $\text{Ker}(M_t^{\lambda_1})$ and $q_3, \dots, q_m \in \langle K_t^{\lambda_1 \lambda_2} \rangle$. Correspondingly, $\Lambda_3, \dots, \Lambda_m \in \langle D_t^{\lambda_1 \lambda_2} \rangle$. This implies that $\{q_1, q_2\} \cup K_t^{\lambda_1 \lambda_2}$ is a basis for $\text{Ker}(M_t)$. Continuing with v_{λ_i} , for $i \geq 3$, and considering the assumption that v_{λ_i} is not active in $\Lambda_1, \dots, \Lambda_{i-1}$, $j \neq i$, we finally get $\{q_1, \dots, q_m\} \cup K_t^{\lambda_1 \dots \lambda_m}$ as a basis for $\text{Ker}(M_t)$ and correspondingly $\{\Lambda_1, \dots, \Lambda_m\} \cup D_t^{\lambda_1 \dots \lambda_m}$ as a basis for the degree t part of Q^\perp .

□

The above proposition shows that deleting some columns from M_t helps to avoid re-computing the basis elements of degree at most $t-1$, which were already computed in the previous steps. Not every set of m active columns will give degree t elements of a basis. In fact, if we delete two columns that both are active in two different basis members of D_{t-1} , then we may not obtain some members of D_t . For instance, let $D_2 = \{\Lambda_1 = d_1 + d_2 + d_1^2 + d_2^2, \Lambda_2 = d_1 + d_2 + 2d_1^2 + d_1 d_2\}$ and $\Lambda' = d_1 + d_2^3 \in \text{Ker}(M_3)$. Then $\Lambda' \notin \text{Ker}(M_3^{d_1 d_2})$.

In the following we show how to detect columns $v_{\lambda_1}, \dots, v_{\lambda_m}$ that satisfy the assumption of part 3 of Proposition 1. This is done via changing the basis $\{\Lambda_1, \dots, \Lambda_m\}$ into a new *reduced* basis $\{\Lambda'_1, \dots, \Lambda'_m\}$, in which the leading terms satisfy the assumptions of part 3 of Proposition 1.

Let $D_{t-1} = \{\Lambda_1, \dots, \Lambda_m\}$ as above. In the improvement of the integration method given in [23], having D_{t-1} , one can construct a matrix D' in order

to obtain a basis for the degree t part of R/Q , so that Condition 4 can be applied (Equation 7 in [23]). Below we show constructing a similar, but smaller matrix which gives the desired set of active columns. Same as M_t , the columns of this matrix are labeled by the coefficients/polynomials that appear in Λ in Equation 1. Also same as D' , the rows of this matrix come from $\Lambda_1, \dots, \Lambda_m$. Let $v_{\lambda_1}, \dots, v_{\lambda_u}$ be the columns of M_t such that they are active in D_{t-1} . Construct the following matrix containing the columns $v_{\lambda_1}, \dots, v_{\lambda_u}$

$$M' = \begin{matrix} \Lambda_1 \\ \vdots \\ \Lambda_m \end{matrix} \begin{pmatrix} v_{\lambda_1} & \dots & v_{\lambda_u} \\ & & \\ & & \\ & & \end{pmatrix}. \quad (5)$$

Changing M' into a row echelon form matrix, after moving the pivot columns to the left hand, we will reach to a matrix of the form

$$G'^{-1}M' = \begin{matrix} \Lambda'_1 \\ \Lambda'_2 \\ \vdots \\ \Lambda'_m \end{matrix} \begin{pmatrix} * & & * & & & & & \\ 0 & \ddots & & & * & & & \\ \vdots & \vdots & \ddots & * & & & & \\ 0 & \dots & 0 & * & * & \dots & * & \end{pmatrix}, \quad (6)$$

where diagonal entries are nonzero and G' is the matrix that takes care of the operations done for the column swapping and the row echelon form. Note that we will not have any zero row. This is because otherwise, if we obtain a zero row in $G'^{-1}M'$, that row is linearly dependent to the other rows. But this is in contradiction with $\Lambda_1, \dots, \Lambda_m$ (and therefore $\Lambda'_1, \dots, \Lambda'_m$ as their linear combinations) being linearly independent. Then our basis will satisfy the conditions of part 3 of Proposition 1.

Now we are ready to prove the following, which provides us with an algorithmic improvement of the integration method, more efficient than Condition 4.

Corollary 2. (Criterion for Deleting Active Columns)

Let $D_{t-1} = \{\Lambda_1, \dots, \Lambda_m\}, M_t, D_t, v_{\lambda_1}, \dots, v_{\lambda_u}$ and $G'^{-1}M'$ be as above and (by abuse of notation) let $v_{\lambda_1}, \dots, v_{\lambda_m}$ be the columns of M_t corresponding to the first m columns in $G'^{-1}M'$. Also let $K_t^{\lambda_1 \dots \lambda_m}$ be a basis for $\text{Ker}(M_t^{\lambda_1 \dots \lambda_m})$ and $D_t^{\lambda_1 \dots \lambda_m}$ be the set of its corresponding dual elements. Then $D_{t-1} \cup D_t^{\lambda_1 \dots \lambda_m}$ is a basis for the degree t part of Q^\perp .

Proof. We only need to prove that the columns $v_{\lambda_1}, \dots, v_{\lambda_m}$ in $G'^{-1}M'$ satisfy the conditions of part 3 of Proposition 1. This is the case because for all $1 \leq i \leq m$, v_{λ_i} has zero in coordinates $i+1, \dots, m$ and has non-zero coordinate i , which is the row corresponding to Λ_i . This means that for all $1 \leq i \leq m$, v_{λ_i} is not active in $\Lambda_1, \dots, \Lambda_{i-1}$. The result follows directly from Proposition 1. \square

Corollary 2 provides us with an optimization in the integration method. Assume that the monomials $x^{\mathbf{a}_1}, \dots, x^{\mathbf{a}_m}$ form a basis for the degree $t-1$ part

of R/Q . If the monomial $dx^{\mathbf{a}_i}$ only appears once in Λ in Equation 1, then applying Condition 4, we have that

$$\Lambda(x^{\mathbf{a}_i}) = \lambda dx^{\mathbf{a}_i}(x^{\mathbf{a}_i}) = \lambda_i = 0.$$

This gives us an equation which adds a row to M_t . However, instead of adding the corresponding row to M_t , one can just plug in $\lambda_i = 0$ in the other equations obtained from Conditions 2 and 3. This will remove λ_i from the other equations, or equivalently will remove the column v_{λ_i} from M_t . If we let v_{λ_i} be the only column of M_t such that its label contains $dx^{\mathbf{a}_i}$, then v_{λ_i} is active in Λ_i and therefore according to Corollary 2, one can delete it from M_t in order to avoid re-computing D_{t-1} .

In what follows we show how our construction implies a basis for the quotient ring as well as a normal form algorithm and a Gröbner basis of the primary component corresponding to the isolated point in question.

Proposition 3. *Let $v_{\lambda_1}, \dots, v_{\lambda_m}$ be the columns in the criterion for deleting active columns, i.e., Corollary 2. Also assume that $p_1, \dots, p_m \in \mathbb{K}[x_1, \dots, x_n]$ are the polynomials corresponding to the coefficients $\lambda_1, \dots, \lambda_m$ in Λ in Equation 1. Consider p_1, \dots, p_m as polynomials in $\mathbb{K}[\partial]$, i.e., substitute $x_i = dx_i$ for $1 \leq i \leq m$. Then $\{p_1, \dots, p_m\}$ is a basis for the degree $t - 1$ part of R/Q .*

Proof. Let ℓ_1, \dots, ℓ_m be the leading terms of p_1, \dots, p_m . Then from the discussion in the integration method, we know that $\{\ell_1, \dots, \ell_m\}$ is a basis for the degree $t - 1$ part of R/Q . Since $p_1, \dots, p_m \in R/Q$ and also the cardinality of $\{\ell_1, \dots, \ell_m\}$ and $\{p_1, \dots, p_m\}$ is the same, then in order to prove that $\{p_1, \dots, p_m\}$ is a basis for R/Q , we just need to prove that p_1, \dots, p_m are linearly independent. Without loss of generality, we can assume that ℓ_i appears only in p_i for $1 \leq i \leq m$. Otherwise, we can reduce p_1, \dots, p_m with respect to each other so that we obtain polynomials p_1, \dots, p_m such that ℓ_1, \dots, ℓ_m are the leading terms of p_1, \dots, p_m and ℓ_i appears only in p_i , for $1 \leq i \leq m$ and also $\langle \ell_1, \dots, \ell_m \rangle = \langle p_1, \dots, p_m \rangle$. This shows that p_1, \dots, p_m are linearly independent, because each leading term only appears in one single polynomial and therefore no p_i can be in the span of the other p_j for $1 \leq j \neq i \leq m$. \square

Proposition 3 implies that the criterion for deleting active columns can be viewed as adding the equation $\Lambda(p_i) = 0$ for $1 \leq i \leq m$ to the integration method's conditions. Same as Condition 4, this equation leads to adding rows to M_t , however those rows are of the form $(0, \dots, 0, c, 0, \dots, 0)$, where c is a nonzero element in coordinate i for $1 \leq i \leq m$ and therefore they result in deleting the corresponding columns. We can say even more.

Proposition 4. *Let $\{p_1, \dots, p_m\} \subseteq \mathbb{K}[x_1, \dots, x_n]$ be a (not necessarily monomial) basis for the degree t part of R/Q such that no monomial of p_i is in Q and let $p_1, \dots, p_m \in \mathbb{K}[\partial]$ be the polynomials p_i after substituting $x_i = \partial_{x_i}$ for $1 \leq i \leq m$. For monomials $m_1, \dots, m_k \notin Q$ such that $m_1, \dots, m_k \notin \text{Supp}(p_1) \cup \dots \cup \text{Supp}(p_m)$, write $m_j = \sum_{i=1}^m \lambda_{ij} p_i'$. Then $\Lambda_i = p_i + \sum_{j=1}^k \lambda_{ij} m_j$*

for $1 \leq i \leq m$, is a basis for the degree t part of Q^\perp and the normal form of any $g \in \mathbb{K}[x_1, \dots, x_n]$ with respect to the basis $\{p_1, \dots, p_m\}$ is

$$NF(g) = \sum_{i=1}^m \Lambda_i(g)p_i.$$

Proof. $\Lambda_1, \dots, \Lambda_m$ are linearly independent because p_1, \dots, p_m are linearly independent in R/Q , due to Proposition 3. The rest of the proof is exactly the same as the proof of Lemma 2.4 in [23]. \square

If $\{p_1, \dots, p_m\} \subseteq \mathbb{K}[x_1, \dots, x_n]$ is an arbitrary basis of the quotient R/Q and $\{p_1, \dots, p_m\} \subseteq \mathbb{K}[\partial]$ are the corresponding differential polynomials, then removing from each p_i the monomials that are in Q , we will obtain a new basis for R/Q . Thus this assumption holds without loss of generality and we generalize Lemma 3.4 from [23].

Proposition 5. *Let $\{p_1, \dots, p_m\} \subseteq \mathbb{K}[x_1, \dots, x_n]$ be a basis for the degree t part of R/Q such that no monomial of p_i is in Q . An element $\Lambda \in \mathbb{K}[\partial]$ is not zero in $Q_t^\perp \setminus Q_{t-1}^\perp$ if and only if in addition to Equation 2 and Equation 3 it satisfies $\Lambda(p_i) = 0$ for $1 \leq i \leq m$.*

Constructing matrices M' and $G'^{-1}M'$ in order to choose particular active columns and deleting them is a special case of the above proposition. We have the following generalization of Proposition 3.7 in [28].

Proposition 6. *Let \prec be a term order and m_j, p_i, Λ_i , $1 \leq i \leq m$, $1 \leq j \leq k$ be as in Proposition 5. Also let w_1, \dots, w_s be the monomials different from $lt(p_i)$, $i = 1 \dots, m$. Write $w_i = \sum_{j=1}^m \gamma_{ij}p_j'$ and let*

$$W = \left\{ g_{w_i} := w_i + \sum_{j=1}^m \gamma_{ij}p_j' \mid 1 \leq i \leq s \right\}, G := \left\{ m_j + \sum_{i=1}^m \lambda_{ij}p_i \mid 1 \leq j \leq m \right\}$$

and $C := \{\mathbf{x}^c \mid c \in \mathbb{N}^n, |c| = \mathcal{N} + 1\}$. Then $G \cup W \cup C$ is a Gröbner basis for Q with respect to \prec .

Proof. Proof of Proposition 3.7 in [28] works here as well. We just need to note that for every $f \in Q$, $lt(f) \in \langle lt(G) \cup lt(W) \cup C \rangle$. \square

Note that unlike in Proposition 3.7 in [28] $G \cup C$ is not a Gröbner basis in this case as we don't necessarily have $\langle lt(Q) \rangle = \langle lt(G) \cup C \rangle$.

Example 2. *Let $I = \langle f_1, f_2 \rangle \subseteq \mathbb{K}[x, y]$, where $f_1 = x - y + x^2$ and $f_2 = x - y + y^2$. In step 2 of the algorithm we have that*

$$M_2 = \begin{array}{c} \text{Condition 3} \\ \Lambda(f_1) = 0 \\ \Lambda(f_2) = 0 \end{array} \begin{pmatrix} d_1 & d_2 & d_1^2 & d_1 d_2 + d_2^2 \\ 0 & 0 & 1 & -1 \\ 1 & -1 & 1 & 0 \\ 1 & -1 & 0 & 1 \end{pmatrix},$$

from which we obtain $D_2 = \{\Lambda_1 = 1, \Lambda_2 = d_1 + d_2, \Lambda_3 = d_2 + d_1^2 + d_1 d_2 + d_2^2\}$. The active columns in D_2 are v_1, v_2, v_3, v_4 , where v_i refers to column i and therefore matrix M' defined in Equation 5 (ignoring $\Lambda_1 = 1$) is

$$M' = \begin{matrix} & d_1 & d_2 & d_1^2 & d_1 d_2 + d_2^2 \\ \Lambda_2 & \left(\begin{array}{cccc} 1 & -1 & 0 & 0 \\ 0 & 1 & 1 & 1 \end{array} \right) \\ \Lambda_3 & & & & \end{matrix}.$$

Two instances of substituting some columns of M' and then computing its (reduced) echelon form are shown below. Matrix

$$G_1'^{-1}M' = \begin{matrix} & d_2 & d_1^2 & d_1 & d_1 d_2 + d_2^2 \\ \Lambda_2' & \left(\begin{array}{cccc} 1 & 0 & -1 & 0 \\ 0 & 1 & 1 & 1 \end{array} \right) \\ \Lambda_3' & & & & \end{matrix}$$

gives columns v_2 and v_3 and matrix

$$G_2'^{-1}M' = \begin{matrix} & d_2 & d_1 d_2 + d_2^2 & d_1 & d_1^2 \\ \Lambda_2' & \left(\begin{array}{cccc} 1 & 0 & -1 & 0 \\ 0 & 1 & 1 & 1 \end{array} \right) \\ \Lambda_3' & & & & \end{matrix}$$

gives columns v_2 and v_4 .

For instance, if we consider $G_2'^{-1}M'$, then $\Lambda_2' = d_2 + d_1 + d_1^2$ and $\Lambda_3' = d_1 d_2 + d_2^2 + d_1 + d_1^2$. Since d_2 only appears in λ_2' and $d_1 d_2 + d_2^2$ only appears in λ_3' , by deleting columns v_2 and v_4 from M_3 we will have the following matrix in step 3.

$$M_3^{\lambda_2, \lambda_4} = \begin{matrix} & d_1 & d_1^2 & d_1^3 - d_1^2 & d_2^3 + d_1 d_2^2 + d_1^3 d_2 - d_1 d_2 \\ \text{Condition 3} & \left(\begin{array}{cccc} 0 & 0 & 1 & 1 \\ 0 & 1 & 0 & 0 \\ \Lambda(f_1) = 0 & \left(\begin{array}{cccc} 1 & 1 & -1 & 0 \\ 1 & 0 & 0 & 0 \end{array} \right) \\ \Lambda(f_2) = 0 & & & \end{array} \right) \\ & & & & \end{matrix}.$$

Since $\text{Ker}(M_3^{\lambda_2, \lambda_4}) = 0$, the algorithm terminates. Using any of the pairs of columns obtained via other possible matrices the same basis would be obtained.

Modifications of Macaulay's Algorithm. The same idea can be used to delete some columns from the matrices obtained via Macaulay's algorithm. In this case, equivalent versions of Proposition 1 and Corollary 2 can be stated with slight modifications. For details see [25, 29].

The necessary modifications to Proposition 1 in order to adjust it to Macaulay's algorithm, are due to the different labeling of the matrices in the integration method and Macaulay's algorithm. In Macaulay's matrices, a column being active is reduced to having its label in the support of the dual basis elements in the previous step. In order to detect the columns to be deleted, one could construct matrix M' in Equation 5 with the labeling in Macaulay's matrices and change it into the echelon form. This will lead to the criterion for deleting active columns in Macaulay's algorithm, i.e., the modification of Corollary 2.

One can use the idea behind the integration method, that the monomials of the dual basis elements of degree t can be obtained by integrating *only* the elements of the dual basis of degree $t - 1$, in order to find some monomials labeling some columns of Macaulay's matrices in step t that cannot be obtained via integration and therefore cannot appear in the basis elements of degree t . This helps with deleting those columns in advance, which will reduce the size of the matrices in Macaulay's algorithm even further. After applying this trick, the monomials labeling the columns of the matrices in Macaulay's algorithm will all appear as monomials in the polynomials labeling the matrices of the integration method.

We explain the above improvement by redoing the calculations for step 3 of Example 1, using the above modifications on Macaulay's algorithm and comparing the computations.

Example 3. *After doing the computations in step 2, we have $D_2 = \{1, d_1, 2d_1^2 + d_2\}$. $d_2 \in \text{Supp}(D_2)$, but $d_2^2 \notin \text{Supp}(D_2)$. So, in step 3, by the above improvement, we can remove $v_{d_1 d_2^2}$ and $v_{d_2^3}$ from M_3 . Also we can remove the columns $v_1, v_{d_1}, v_{d_1^2}$ using the modifications induced by Proposition 1. So the new matrix has 5 columns, while the original matrix in Macaulay's method has 10 columns.*

Change of the Integration Order at Each Step. We conclude by another possible optimization strategy. One can change the order of the variables at each step of the integration method in order to gain some computational advantage. Suppose that we have computed $D_{t-1} = \{\Lambda_1, \dots, \Lambda_m\}$. Consider $n_i := \#\{dx_i^{\alpha_i} \in \bigcup_i \text{Supp}(\Lambda_i) \mid \alpha_i \in \mathbb{N}\}$. Now, re-order the variables in the following way: if $n_i \leq n_j$ then $x_i < x_j$. We call such an order a *good integrable* order. Assume that $x_{b_1} < x_{b_2} < \dots < x_{b_n}$ is a good integrable order, where $b_i \in \{1, \dots, n\}$. Now we consider $\Lambda_1, \dots, \Lambda_m$ as polynomials in $\mathbb{K}[dx_{b_1}, \dots, dx_{b_n}]$ and continue with the integration in the following order:

$$\begin{aligned} \Lambda &= \sum_i \lambda_{i1} \int_{b_1} \Lambda_i \Big|_{dx_{b_2}=\dots=dx_{b_n}=0} \\ &+ \dots \\ &+ \sum_i \lambda_{in-1} \int_{b_{n-1}} \Lambda_i \Big|_{dx_{b_2}=\dots=dx_{b_n}=0} \\ &+ \sum_i \lambda_{in} \int_{b_n} \Lambda_i. \end{aligned}$$

This way, least possible number of integrations will be performed. Note that the number of integrands and the number of basis elements of D_{t-1} are fixed and therefore there is no advantage in terms of the size of M_t . The following example illustrates this optimization.

Example 4. *Consider Example 1. In step two we have that*

$$D_2 = \langle \Lambda_1 = 1, \Lambda_2 = d_1 + d_2, \Lambda_3 = -d_1 + d_1^2 + d_1 d_2 + d_2^2 \rangle.$$

Then $n_1 = 3, n_2 = 2$. Therefore we change the order into $y < x$ and work on $\mathbb{K}[dy, dx]$. Then

$$\Lambda = \lambda_1 dy + \lambda_2 dx + \lambda_3 dy^2 + \lambda_4(dydx + dx^2) + \lambda_5(dy^3) + \lambda_6(dx^3 - dx^2 + dx^2 dy + dx dy^2).$$

There is only one monomial in the 5-th column of M_3 , while in the original ordering, there were two:

$$\begin{aligned} \Lambda &= \lambda_1 dx + \lambda_2 dy + \lambda_3 dx^2 + \lambda_4(dx dy + dy^2) \\ &\quad + \lambda_5(dx^3 - dx^2) + \lambda_6(dy^3 + dx dy^2 + dx^2 dy + dx dy). \end{aligned}$$

3. Directional Multiplicity

In this section, using the dual space of an ideal, we introduce the notion of *Directional Multiplicity*. Directional multiplicity provides a lot of information about the multiplicity structure at an isolated point. Using Lemma 9 we give a sound definition of directional multiplicity. Then, we show that directional multiplicities can be bounded and can bound other invariants of an ideal, namely the Nil-index and the intersection multiplicity.

Definition and Properties. The first step towards a sound definition of directional multiplicity is to observe that the set of monomials that appear in elements of Q^\perp is exactly the set of monomials $\partial^{\mathbf{a}}$ such that $x^{\mathbf{a}} \notin Q$, where $x^{\mathbf{a}} = x_1^{a_1} \cdots x_n^{a_n}$. Note that

$$\partial^{\mathbf{a}}(x^{\mathbf{b}}) = \prod \delta_{a_i, b_j} \quad (7)$$

where $\delta_{i,j}$ is the Kronecker delta. We prove the following result that first appeared without a proof in [23].

Proposition 7 (Characterization of Monomials in Q^\perp). *Let $Q = Q_\zeta$ be an \mathfrak{m}_ζ -primary ideal. Consider Q^\perp as a sub-vector space of $\mathbb{K}[\partial_\zeta]$ as above. Then*

$$\bigcup_{\Lambda \in Q^\perp} \text{supp}(\Lambda) = \left\{ \partial^{\mathbf{a}} \mid x^{\mathbf{a}} \notin Q \right\},$$

where $\text{supp}(\Lambda)$ is the set of monomials with nonzero coefficient in Λ .

Proof. By Theorem 3.2 in [28], we have that $f \in Q \Leftrightarrow (\lambda(f) = 0 \text{ for all } \lambda \in Q^\perp)$. This implies that $f \in Q \Leftrightarrow (\lambda(f) = 0 \text{ for all } \lambda \in D)$, for a basis $D \subset \mathbb{K}[\partial]$ of Q^\perp .

We are ready to prove the thesis:

- ” \subseteq “ If $\partial^{\mathbf{a}}$ is in $\text{supp}(\Lambda)$ then the monomial $x^{\mathbf{a}}$ is not annihilated by Λ (see Equation 7), which implies $x^{\mathbf{a}} \notin Q$.
- ” \supseteq “ If $x^{\mathbf{a}} \notin Q$, then there exists $\lambda \in D$ such that $\lambda(x^{\mathbf{a}}) \neq 0$. Let $\Lambda \in \mathbb{K}[\partial]$ be the differential operator corresponding to λ , so $\Lambda(x^{\mathbf{a}}) \neq 0$. By Equation 7, we know that $m(x^{\mathbf{a}}) = 0$ for all monomials m in $\text{supp}(\Lambda)$ which are different from $\partial^{\mathbf{a}}$. Hence $\partial^{\mathbf{a}}$ has to be in $\text{supp}(\Lambda)$.

□

After characterizing the monomials in Q^\perp , the next step is to study their behaviour under projection. Let I be an ideal in R and $I_{2,\dots,n} = I \cap \mathbb{K}[x_2, \dots, x_n]$ be its first elimination ideal. The following result shows that the objects introduced so far, behave well in the framework of elimination theory.

Proposition 8 ([9], Proposition 7.19). *Let π be the linear map*

$$\begin{array}{ccc} \pi : \mathbb{K}[[dx_1, \dots, dx_n]] & \longrightarrow & \mathbb{K}[[dx_2, \dots, dx_n]] \\ \Lambda & \longmapsto & \Lambda(0, dx_2, \dots, dx_n). \end{array}$$

Then $(I_{2,\dots,n})^\perp = \pi(I^\perp)$.

We use the above proposition in order to prove the Dual Projection Lemma, which shows how to get a basis of the dual space of the elimination ideal, having a basis for the dual space. Note that I in Proposition 8 can be any ideal, however the following lemma is valid only for the local case, i.e., when we are working on an \mathfrak{m}_ζ -primary ideal $Q = Q_\zeta$.

Lemma 9 (Dual Projection Lemma). *Let $Q = Q_\zeta$ be an \mathfrak{m}_ζ -primary ideal and $D = \{\Lambda_0, \Lambda_1, \dots, \Lambda_{\ell-1}\} \subset \mathbb{K}[\partial]$ be a basis of Q^\perp , seen as a sub-vector space of $\mathbb{K}[\partial_\zeta]$. Then*

$$Q_{2,\dots,n}^\perp = \langle \Lambda_0|_{dx_1=0}, \Lambda_1|_{dx_1=0}, \dots, \Lambda_{\ell-1}|_{dx_1=0} \rangle.$$

Proof. We prove the two inclusions:

(\supseteq) For $1 \leq i \leq \ell - 1$, since $\Lambda_i \in Q^\perp$, we have that $\Lambda_i|_{dx_1=0} \in Q^\perp|_{dx_1=0}$. Proposition 8 states that $Q^\perp|_{dx_1=0} = Q_{2,\dots,n}^\perp$ and thus $\Lambda_i|_{dx_1=0} \in Q_{2,\dots,n}^\perp$. This means $\langle \Lambda_0|_{dx_1=0}, \Lambda_1|_{dx_1=0}, \dots, \Lambda_{\ell-1}|_{dx_1=0} \rangle \subseteq Q_{2,\dots,n}^\perp$.

(\subseteq) Let $\Lambda' \in Q_{2,\dots,n}^\perp$. Since $Q^\perp|_{dx_1=0} = Q_{2,\dots,n}^\perp$, we have that $\Lambda' \in Q^\perp|_{dx_1=0}$. Therefore, there exists a $\Lambda \in Q^\perp$, such that $\Lambda' = \Lambda|_{dx_1=0}$. Since $Q^\perp = \langle \Lambda_0, \Lambda_1, \dots, \Lambda_{\ell-1} \rangle$, there exist $c_i \in \mathbb{K}$, such that $\Lambda = \sum_{i=0}^{\ell-1} c_i \Lambda_i$. Therefore

$$\Lambda|_{dx_1=0} = \sum_{i=0}^{\ell-1} c_i \Lambda_i|_{dx_1=0} \text{ and } \Lambda' = \sum_{i=0}^{\ell-1} c_i \Lambda_i|_{dx_1=0}, \text{ which implies that}$$

$$\Lambda' \in \langle \Lambda_0|_{dx_1=0}, \Lambda_1|_{dx_1=0}, \dots, \Lambda_{\ell-1}|_{dx_1=0} \rangle.$$

Thus, $Q_{2,\dots,n}^\perp \subseteq \langle \Lambda_0|_{dx_1=0}, \Lambda_1|_{dx_1=0}, \dots, \Lambda_{\ell-1}|_{dx_1=0} \rangle$.

□

The following immediate corollary deals projecting all but one dimension.

Corollary 10. Let $D = \{\Lambda_0, \Lambda_1, \dots, \Lambda_{l-1}\} \subset \mathbb{K}[\partial]$ be a basis of Q^\perp , and for $1 \leq i \leq n$ let $Q_i = Q \cap \mathbb{K}[x_i]$. Denote by $\Lambda|_{dx_i \neq 0}$ the polynomial obtained by substituting $dx_j = 0$ for $1 \leq i \neq j \leq n$ in Λ . Then

$$Q_i^\perp = \langle \Lambda_0|_{dx_i \neq 0}, \Lambda_1|_{dx_i \neq 0}, \dots, \Lambda_{l-1}|_{dx_i \neq 0} \rangle.$$

Moreover, there exists $\mu_i \in \mathbb{N}$ such that

$$Q_i^\perp = \langle 1, dx_i, \dots, dx_i^{\mu_i-1} \rangle.$$

Now we have the necessary tools to define the directional multiplicity.

Definition 1 (Directional Multiplicity). Let ζ be an isolated point in the variety of an ideal I and Q_ζ be the corresponding \mathfrak{m}_ζ -primary component. Using the notation of Corollary 10, for $1 \leq i \leq n$, we define the i -th directional multiplicity of ζ to be μ_i and denote it by $\mu_i(\zeta)$ or simply μ_i if ζ is clear from context.

In order to give an intuition of directional multiplicity, we take a look at the quotient R/Q_ζ . If we consider this quotient as a vector space, finding a basis for such a quotient was the task given to Buchberger for his PhD thesis by Gröbner, which led to the invention of Gröbner bases [4]. Let us recall that the multiplicity of ζ is defined as $\dim_{\mathbb{K}} R/Q_\zeta$. We will denote the multiplicity by $\mu(\zeta)$ or simply by μ if ζ is clear from the context.

As discussed earlier, we can find the monomials of D by searching among those monomials of I^\perp that have degree at most the Nil-index. These monomials are actually the monomials under the *Extended Buchberger Diagram* which is defined below.

Definition 2 (Extended Buchberger Diagram). The Extended Buchberger Diagram of an \mathfrak{m}_ζ -primary ideal Q_ζ is obtained by considering all the monomials that appear in a basis of the dual space of Q_ζ .

We can think of the Nil-index of Q_ζ as the largest degree of the monomials under the extended Buchberger diagram. Figure 1 shows the extended Buchberger diagram and all of its monomials for Example 1.

Note that the monomials under the Buchberger diagram with respect to an ordering form a vector space basis for R/Q . They include some monomials in a basis of Q^\perp , but they do not necessarily include all the monomials in D . In particular, they may not include the highest powers of dx_i , i.e., the monomials corresponding to the directional multiplicities. However in the extended Buchberger diagram, one can see all the possible monomials in D , which are all the monomials that do now appear in Q . These monomials include all the monomials in the Buchberger diagram of Q .

The above comments are illustrated in Figure 2. The black dots show a basis for R/Q , while the white dots are the rest of the monomials in the basis of Q^\perp , see also [24] for a similar diagram. Also Figures 3 and 4 show the quotient of the elimination ideal with respect to x and the quotient of the elimination ideal




Figure 1: Extended Buchberger Diagram for Example 1




Figure 2: Extended Buchberger diagram vs a basis for R/Q_ζ with respect to a degree ordering




Figure 3: Extended Buchberger diagram vs directional multiplicity with respect to x for Example 1




Figure 4: Extended Buchberger diagram vs directional multiplicity with respect to y for Example 1

with respect to y , respectively. In Figure 3, black dots are the basis for Q_2^\perp and the white dots are the rest of the monomials in the dual basis. In Figure 4, black dots are the basis for Q_1^\perp and the white dots are the rest of the monomials in the dual basis.

Considering the above figures, one can see that the extended Buchberger diagram includes the Buchberger diagram with respect to every order. \mathcal{N} is a bound for the degree of the members of a Gröbner basis with respect to every order. Directional multiplicity with respect to an axis is the largest intersection point of the extended Buchberger diagram with that axis. The Buchberger diagram does not necessarily have an intersection with the hyperplane $x_1 + \dots + x_n = \mathcal{N}$, but the extended Buchberger diagram does have at least a point in common with that hyperplane.

Example 5. Let $I = \langle f_1 = x^8 + y^5, f_2 = x^7 y^4 \rangle$. The origin is a root of the

system with multiplicity $\mu = 67$. We have that $\mathcal{N} = 18$, while $\mu_1 = 15, \mu_2 = 9$. The reduced Gröbner basis for I with respect to the lexicographic order ($x > y$) is $\{f_1 = x^8 + y^5, f_2 = x^7 y^4, g_y = y^9\}$, and with respect to lexicographic order ($y > x$) is $\{f_1 = y^5 + x^8, f_2 = y^4 x^7, g_x = x^{15}\}$, where g_y and g_x are the generators of the elimination ideal with respect to the lexicographic orders $x > y$ and $y > x$ respectively.

These observations give us the intuition that the directional multiplicities are at most as large as the Nil-index. Also their product gives us the volume of a cuboid which contains the Buchberger diagram. The following statements make the comments above more precise.

Let us first note that the Nil-index is at most as large as the multiplicity and the multiplicity is bounded by the number of lattice points in the n -simplex. Thus $\mathcal{N} \leq \mu \leq \text{Number of Lattice point in the } n\text{-simplex} = \binom{\mathcal{N}+n}{n}$.

Proposition 11. *Let μ be the multiplicity of an isolated point ζ . Then*

- $\mu_i \leq \mu$ for every $1 \leq i \leq n$.
- $\mu \leq \prod_{1 \leq i \leq n} \mu_i$.
- $\sum_{i=1}^n \mu_i - n + 1 \leq \mu$.

Proof. For the first part, recall that $\dim_{\mathbb{K}} Q_{\zeta}^{\perp} = \mu$ and that μ_i is the dimension of a vector subspace of Q_{ζ}^{\perp} . Thus $\mu_i \leq \mu$.

For the second part, for every $1 \leq i \leq n$, μ_i is the largest degree of the elements in $Q_{\zeta}^{\perp} \cap \mathbb{K}[d_i]$. This means that $\mu_i + 1$ is the largest possible degree of x_i in R/Q . Since $\mu = \dim_{\mathbb{K}} R/Q$, we conclude that $\mu \leq \prod_{1 \leq i \leq n} \mu_i$.

For the third statement, note that as argued above, $dx_i^{a_i} \in Q_{\zeta}^{\perp}$ if and only if $a_i < \mu_i$. This means that $x_i^{a_i} \notin Q_{\zeta}$ if and only if $a_i < \mu_i$. Now, for all $1 \leq i \leq n$, let $A_i := \{1, x_i, \dots, x_i^{\mu_i-1}\}$. Then, $\langle \bigcup A_i \rangle \subseteq R/Q_{\zeta}$ as vector spaces. Note that the elements of $\bigcup A_i$ are linearly independent. Then $\dim \langle \bigcup A_i \rangle = \sum \mu_i - n + 1 \leq \dim R/Q_{\zeta} = \mu$ and the result follows. \square

Proposition 12. *Let \mathcal{N} be the Nil-index of Q_{ζ} . Then*

- $\mathcal{N} \geq \mu_i$ for all $1 \leq i \leq n$,
- $\mathcal{N} \leq \sum_{1 \leq i \leq n} \mu_i - n$

Proof. According to the definition of the Nil-index we have $\mathfrak{m}_{\zeta}^{\mathcal{N}} \not\subseteq Q_{\zeta}$ and $\mathfrak{m}_{\zeta}^{\mathcal{N}+1} \subseteq Q_{\zeta}$. Since $\mathfrak{m}_{\zeta}^{\mathcal{N}} = \langle x_1 - \zeta_1, \dots, x_n - \zeta_n \rangle^{\mathcal{N}}$, we have that $(x_i - \zeta_i)^{\mathcal{N}} \notin Q_{\zeta}$ and $(x_i - \zeta_i)^{\mathcal{N}+1} \in Q_{\zeta}$. By the definition of μ_i and the Proposition 7, $dx_i^{\mu_i} (x_i - \zeta_i)^{\mathcal{N}} = 0$ and $dx_i^{\mu_i} (x_i - \zeta_i)^{\mathcal{N}-1} \neq 0$. Therefore $\mu_i \leq \mathcal{N}$.


For the second part, note that for all x_i , $d_{x_i}^{\mu_i-1} \in \text{supp}(Q_{\zeta}^{\perp})$ and $d_{x_i}^{\mu_i} \notin \text{supp}(Q_{\zeta}^{\perp})$. Therefore by Proposition 7, $x_i^{\mu_i-1} \notin Q_{\zeta}$ and $x_i^{\mu_i} \in Q_{\zeta}$. Consider

$A = \{\mathbf{a} \in \mathbb{N}^n \mid |\mathbf{a}| = \sum(\mu_i - 1) + 1\}$. By the Pigeonhole principle, there exists an i , $1 \leq i \leq n$, such that $x_i^{\mu_i} \mid x^{\mathbf{a}}$. Therefore $x^{\mathbf{a}} \in Q_\zeta$ for all $\mathbf{a} \in A$, which implies that $\mathfrak{m}_\zeta^{|\mathbf{a}|} \subseteq Q_\zeta$ and $\mathcal{N} < |\mathbf{a}| = 1 + \sum(\mu_i - 1)$. The result follows by minimality of \mathcal{N} . \square

The inequalities in the Propositions 11 and 12 are sharp. For example, in the univariate case, where $I = Q_\zeta \in \mathbb{K}[x]$, we have $\mathcal{N} = \mu_1 = \mu$.

A geometric interpretation of the i -th directional multiplicity at an intersection point is the number of instances of the intersection point that can be seen when we look at the intersection point in the direction parallel to the x_i axis. Some of the presented inequalities are direct consequences of the definitions, and reveal interesting properties of this new notion. In particular, knowing the directional multiplicities we can deduce information about the multiplicity or the Nil-index. Thus, the notion of directional multiplicity is, in this sense, a refinement of multiplicity and Nil-index. Moreover, in applications this refined information can be used. In particular, methods such the subdivision of [1], for the arrangement of several planar curves produces small boxes which are likely to contain isolated singularities. Taking the center of such a box as an approximation of a multiple root allows for an approximate dual computation that can reveal directions that lead to degenerate computations.

Geometry of the Elimination Ideal. Let $I \subseteq \mathbb{C}[x, y]$ be a zero dimensional ideal with no roots at infinity generated by two polynomials corresponding to two planar curves and $I_1 = I \cap \mathbb{C}[y] = \langle g \rangle$ be its elimination ideal. We illustrate the case of geometric degeneracy and how directional multiplicity can be used, in a concrete example. Let $f_1 = (y + 1)(y - x + 1)$ and $f_2 = x^2 + y^2 - 1$ as shown in the figure. The two curves intersect at two points, namely $(1, 0)$ and $(0, -1)$. Their Sylvester resultant is $2y(y + 1)^3$, which implies that the projection on the y -axis of the roots $(1, 0)$ and $(0, -1)$ have multiplicity 1 and 3 respectively. On the other hand, computing the Gröbner basis of the elimination ideal in $\mathbb{C}[y]$, we obtain the unique monic generator $g = y(y + 1)^2$.



$$\begin{array}{ll}
 f_1 & (y + 1)(y - x + 1) \\
 f_2 & x^2 + y^2 - 1 \\
 g & y(y + 1)^2 \\
 \text{resultant} & 2y(y + 1)^3
 \end{array}$$

Observing the difference in the multiplicities of the resultant and g , the questions “when does the multiplicity drop?” and “what does the multiplicity of a factor in g mean?” arise. Using the concept of directional multiplicity, we

are able to address these questions in the degenerate case, as the one in the example.

The exponent of the factor of g corresponding to an intersection point is the directional multiplicity at that point. The exponent of the corresponding factor of the resultant give us the multiplicity of the intersection points. However Gröbner basis did not say much about the geometry of the intersection. Now having the concept of directional multiplicity, we can explain the generator of the elimination ideal geometrically. In general given dense polynomials $f_1, \dots, f_n \in \mathbb{K}[x_1, \dots, x_n]$, let $I_1 = \langle g \rangle$ and R_1, \dots, R_k be the square-free factorization of the Macaulay resultant, with R_i corresponding to the isolated point ζ_i . Then $g = R_1^{\mu_1(\zeta_1)} \dots R_k^{\mu_k(\zeta_k)}$.

4. Benchmarks

We compare the methods described in the previous sections with respect to both the size of the matrices produced and the running time. The methods we compare are the following:

MACAULAY	The Macaulay algorithm in [21]
INTEGMM	The integration method of [23]
INTEGMMO	The integration method of [23] with optimizations
NEWINTEG	The integration method with the criterion of Corrolary 2
NEWINTEGO	The integration method with the criterion of Corrolary 2 and optimizations

The polynomial systems we use for the benchmarks are the following:

Name	Polynomials	Point
1	$x^3 - yz, y^3 - xz, z^3 - xy$	(0, 0, 0)
cbms2	$x^3 - 3x^2y + 3xy^2 - y^3 - z^2, z^3 - 3z^2x + 3zx^2 - x^3 - y^2, y^3 - 3y^2z + 3yz^2 - z^3 - x^2$	(0, 0, 0)
mth191	$x^3 + y^2 + z^2 - 1, x^2 + y^3 + z^2 - 1, x^2 + y^2 + z^3 - 1$	(0, 1, 0)
decker2	$x + y^3, x^2y - y^4$	(0, 0)
Ojika2	$x^2 + y + z - 1, x + y^2 + z - 1, x + y + z^2 - 1$	(1, 0, 0)
Ojika3	$x + y + z - 1, 2x^3 + 5y^2 - 10z + 5z^3 + 5, 2x + 2y + z^2 - 1$	(0, 0, 1)
KSS	$x^2 - 2x + x + y + z + w + t - 4, y^2 - 2y + x + y + z + w + t - 4, z^2 - 2z + x + y + z + w + t - 4, w^2 - 2w + x + y + z + w + t - 4, t^2 - 2t + x + y + z + w + t - 4$	(1, 1, 1, 1, 1)
Caprasse	$-x_1^3x_3 + 4x_1x_2^2x_3 + 4x_1^2x_2x_4 + 2x_2^3x_4 + 4x_1^2 - 10x_2^2 + 4x_1x_3 - 10x_2x_4 + 2, -x_1x_3^3 + 4x_2x_3^2x_4 + 4x_1x_3x_4^2 + 2x_2x_4^3 + 4x_1x_3 + 4x_3^2 - 10x_2x_4 - 10x_4^2 + 2, x_2^2x_3 + 2x_1x_2x_4 - 2x_1 - x_3, x_1x_4^2 + 2x_2x_3x_4 - x_1 - 2x_3$	$(2, -i\sqrt{3}, 2, i\sqrt{3})$
Cyclic 9	$\sum_{j=0}^8 \prod_{k=j}^{i+j-1} x_{k \bmod 9}, i = 1, \dots, 8$ and $1 - x_1x_2x_3x_4x_5x_6x_7x_8x_9$	$(-0.9383 - 0.3404i, \dots, 2.4602 + 0.89550i)$
DZ2	$x^4, x^2y + y^4, z + z^2 - 7x^3 - 8x^2$	(0, 0, -1)
DZ3	$14x + 33y - 3\sqrt{5}(x^2 + 4xy + 4y^2 + 2) + \sqrt{7} + x^3 + 6x^2y + 12xy^2 + 8y^3, 41x - 18y - \sqrt{5} + 8x^3 - 12x^2y + 6xy^2 - y^3 + 3\sqrt{7}(4xy - 4x^2 - y^2 - 2)$	(1.5055, 0.36528)

4.1. Optimizations

Pivoting. Below we show the role of pivoting within our improvement. Let us put an order on the monomials of D_{t-1} , e.g., degree lexicographic, and denote by $lt(\Lambda')$, the leading term of Λ' , for every $\Lambda' \in D_t$. Now one can reduce the members of a basis of D_t with respect to each other, so that $lt(\Lambda') \notin \text{Supp}(\Lambda'')$, for all $\Lambda' \neq \Lambda'' \in D_t$. We call such a basis a *reduced basis*. Then every leading term will be a monomial that uniquely appears in the reduced basis. If $\Lambda_1, \dots, \Lambda_m$ is a basis for D_{t-1} , then removing the columns corresponding to $lt(\Lambda_1), \dots, lt(\Lambda_m)$ from M_t is equivalent to part 3 of Proposition 1. Using part 1 of Proposition 1, one can efficiently check whether v_λ is active in D . This must be done with precise pivoting. For that, one must start with reducing v_λ with the *appropriate* columns, without doing the column reductions for the other columns, unless it is required. As a side remark, using row echelon form is also taking advantage of pivoting.

Reduction of the number of rows. Condition (2) expresses the fact that the dual is stable by derivation. The number of these polynomial conditions is $\binom{n}{2}$, and each of them implies m linear conditions on the coefficients. These conditions constitute the main part of the matrix. In the experiments we observe that many of these conditions are repeated for different combinations of i and k in (2), or are equal to null. Therefore, in the optimized versions of the algorithm we detect and purge repeating conditions before inserting them to the matrix. Moreover, several times a column appears with a single non-zero element. Such columns can be removed without affecting the kernel of the matrix. In particular, this is the reason why in the benchmarks the INTEGMMO method does not increase the number of columns as it ought to.

Finally, the element Λ_1 in (2) is always equal to the constant (evaluation functional in the dual). The result of the integration is always the same, independent of m , therefore m columns can be replaced by one; therefore the number of columns at the final step t becomes $m(n-1) + 1$.

4.2. Comparing Matrix Sizes

As size is the biggest obstacle in computations, an important metric for the comparison of the different methods is the size of the matrices constructed. The following matrix presents the sizes of the matrices at step t for each method, where e is the number of the polynomials in the given basis, n is the number of the variables in the polynomial ring and m is the dimension of Q_{t-1}^\perp :

method	rows	columns
MACAULAY	$\binom{t+n-1}{n}e$	$\binom{t+n}{n}$
INTEG	$\binom{n}{2}m + e$	nm
INTEGMM	$\binom{n}{2}m + e + m$	$(n-1)m + 1$
NEWINTEG	$\binom{n}{2}m + e$	$(n-1)m + 1$

Note that we denote by INTEG the original integration method in [28]. We observe that applying Condition 4 in the integration method, one gets m extra

rows, but it may lead to deleting at most those m rows and also at most m columns. However, this is exactly the size of the matrix obtained using the new modification to the integration method.

Using our implementation and applying the optimizations discussed above, at the final step of the algorithms, we obtain matrices of the following sizes:

System	MACAULAY	INTEGMM	INTEGMMO	NEWINTEG	NEWINTEGO
cbms1	105×56	46×33	27×23	36×23	27×23
cbms2	60×35	34×24	21×17	27×17	21×17
mth191	30×20	18×12	10×9	15×9	10×9
decker2	20×15	9×8	5×5	6×5	5×5
Ojika2	12×10	10×6	6×5	9×5	6×5
Ojika3	60×35	18×12	12×9	15×9	12×9
KSS	630×252	180×80	155×65	165×65	155×65
Caprasse	60×35	31×16	22×13	28×13	22×13
Cyclic 9	495×220	156×36	116×33	153×33	116×33
DZ2	360×165	66×48	44×33	51×33	44×33
DZ3	30×21	11×10	6×6	7×6	6×6

We observe that applying the optimizations described above in INTEGMMO, the extra rows added in INTEGMM are removed and the matrices obtained have exactly the same size as the ones obtained by the optimized version of the new method. Nevertheless, the non-optimized versions of the two methods produce matrices of different sizes, with the new method producing smaller ones. This has a direct effect in the time needed for the optimizations (as seen in the time comparisons). It is important to note that except for the extra rows, INTEGMM produces matrices with more columns as well, e.g., in cbms1 we have 33 versus 23 columns.

Using the new criterion for deleting columns and applying the optimizations, we obtain constantly the smallest matrix. In terms of columns, this is true even without the extra optimizations (note that in the last two columns of the comparison matrix, the number of columns does not change at all). In other words, the experiments confirm that the matrices produced using Corollary 2 are smaller than those produced by applying Condition 4.

4.3. Comparing Running Times

The benchmarks are performed in MAPLE 18 with 32 digits of precision. The hardware used is an 8 Intel(R) Xeon(R) CPU E5345 @ 2.33GHz core server with 16Gb or RAM, running Linux 3.16.0-4.

In the following table we provide the running times for each method:

System	MACAULAY	INTEGMM	INTEGMMO	NEWINTEG	NEWINTEGO
cbms1	3.080	0.492	0.291	0.363	0.275
cbms2	1.132	0.195	0.147	0.216	0.147
cbms1	0.259	0.064	0.056	0.112	0.056
decker2	0.195	0.052	0.043	0.044	0.043
Ojika2	0.060	0.027	0.024	0.075	0.024
Ojika3	0.863	0.112	0.103	0.088	0.083
KSS	85.684	7.095	5.404	5.287	4.783
Caprasse	1.284	1.020	1.388	0.775	0.775
Cyclic 9	37.867	6.192	6.139	6.504	5.747
DZ2	38.148	1.316	0.827	0.975	0.860
DZ3	0.456	0.168	0.087	0.079	0.080

As expected, the methods producing the smaller matrices are faster, since the kernel computation at the end of each step is the bottleneck of all algorithms tested. We notice that the new criterion is more efficient. It is simpler to implement and compute and thus, although the matrices produced in the optimized versions of the two algorithms are of equal size, the new method is faster.

5. Conclusions

There are two main algorithms for computing the orthogonal of an \mathfrak{m}_ζ -primary component of a given ideal. The first algorithm was given by Macaulay. Although it is still in use, and in spite of the improvements it saw since its inception, it is inefficient because of the construction of large matrices. The second algorithm, namely the integration method, reduces the size of the matrices drastically. In [23], the authors found a way to avoid redundant computations by computing a primal base for the quotient of the \mathfrak{m}_ζ -primary ideal along with its dual base. This condition adds new rows to the matrices, which in some cases may lead to deleting those rows and some columns.

Our work generalizes the method presented in [23] in two ways. On one hand by introducing a criterion to avoid redundant computations, on the other hand by computing a general basis. This is done via computing a general basis for the quotient, which does not necessarily consist of monomials. Our method leads to deleting some columns from the matrices in the integration method instead of adding new rows. As shown in the benchmarks, the method produces consistently the smallest matrix among all methods tested. Moreover, in terms of practical performance, the benchmarks show that the our implementation is consistently among the fastest of the methods tested.

Another aspect studied in the present work is that of local properties and geometry of an isolated point via its directional multiplicity. This gives the multiplicity structure with respect to an axis, and is useful in understanding the geometry behind projection. Moreover, it sheds light on the gap between the degree of the generator of the elimination ideal and the corresponding factor in the resultant.

Several research directions stem from the present work. A natural generalization would be to define the multiplicity in the direction of an arbitrary vector $v \in \mathbb{R}^n$, instead of only for the axis aligned directions. The directional multiplicities along these vectors can be used in studying singularities of curves. A second interesting aspect, arising naturally in almost all topics in elimination theory is sparsity. Exploiting the sparsity and structure of the matrices involved would improve the algorithms considerably. Last but not least, devising early termination criteria would be essential for the efficient computation of directional multiplicity.

Acknowledgments

The authors would like to express their sincere gratitude to Matteo Gallet for discussions and comments on the early versions of this work. The first author

is grateful to his thesis advisor, B. Mourrain for introducing him to dual bases of singular isolated points. Part of this work has been done during the second author's visit to the Centre de Recerca Matemàtica in Bellaterra, Spain. The third author was partially supported by the project 113F293 under the program 1001 of the Scientific and Technological Research Council of Turkey, and by the European Union (European Social Fund - ESF) and Greek national funds through the Operational Program "Education and Lifelong Learning" of the National Strategic Reference Framework (NSRF) - Research Funding Program: THALIS-UOA (MIS 375891).

References

- [1] L. Alberti, B. Mourrain, and J. Wintz. Topology and arrangement computation of semi-algebraic planar curves. *Computer Aided Geometric Design*, 25(8):631–651, 2008.
- [2] J. G. Alcázar and G. M. Díaz-Toca. Topology of 2d and 3d rational curves. *Computer Aided Geometric Design*, 27(7):483 – 502, 2010.
- [3] D. Bates, C. Peterson, and A. J. Sommese. A numerical-symbolic algorithm for computing the multiplicity of a component of an algebraic set. *Journal of Complexity*, 22(4):475–489, 2006.
- [4] B. Buchberger. *Ein Algorithmus zum Auffinden der Basiselemente des Restklassenringes nach einem Nulldimensionalen Polynomiideal (An Algorithm for Finding the Basis Elements in the Residue Class Ring Modulo a Zero Dimensional Polynomial Ideal)*. PhD thesis, Mathematical Institute, University of Innsbruck, Austria, 1965. (English translation in *Journal of Symbolic Computation*, Special Issue on Logic, Mathematics, and Computer Science: Interactions. Vol. 41, Number 3-4, Pages 475-511, 2006).
- [5] B. Buchberger. Ein Algorithmisches Kriterium für die Lösbarkeit eines Algebraischen Gleichungssystems (An Algorithmic Criterion for the Solvability of Algebraic Systems of Equations) . *Aequationes Mathematicae*, 3:374–383, 1970. (English translation in B. Buchberger, F. Winkler (eds.): *Gröbner Bases and Applications*, London Math. Society Lecture Note Series 251, Cambridge Univ. Press, 1998, Pages 535 -545).
- [6] R. M. Corless, G. M. Diaz-Toca, M. Fioravanti, L. Gonzalez-Vega, I. F. Rua, and A. Shakoori. Computing the topology of a real algebraic plane curve whose defining equations are available only "by values". *Computer Aided Geometric Design*, 30(7):675 – 706, 2013.
- [7] B. H. Dayton and Z. Zeng. Computing the multiplicity structure in solving polynomial systems. In *Proceedings of the 2005 International Symposium on Symbolic and Algebraic Computation*, ISSAC '05, pages 116–123, New York, NY, USA, 2005. ACM.

- [8] G. M. Diaz-Toca, M. Fioravanti, L. Gonzalez-Vega, and A. Shakoory. Using implicit equations of parametric curves and surfaces without computing them: Polynomial algebra by values. *Computer Aided Geometric Design*, 30(1):116 – 139, 2013. Recent Advances in Applied Geometry.
- [9] M. Elkadi and B. Mourrain. *Introduction à la Résolution des Systèmes Polynomiaux*, volume 59. Springer, 2007.
- [10] I. Emiris and B. Mourrain. Matrices in elimination theory. *Journal of Symbolic Computation*, 28(1-2):3–44, 1999.
- [11] I. M. Gelfand, M. M. Kapranov, and A. Zelevinski. *Discriminants, Resultants, and Multidimensional Determinants*. Birkhäuser, 1994.
- [12] J. D. Hauenstein, B. Mourrain, and A. Szanto. Certifying isolated singular points and their multiplicity structure. In *Proceedings of the 2015 ACM on International Symposium on Symbolic and Algebraic Computation, ISSAC '15*, pages 213–220, New York, NY, USA, 2015. ACM.
- [13] Y. Huang and D. Wang. Computing intersection and self-intersection loci of parametrized surfaces using regular systems and Gröbner bases. *Computer Aided Geometric Design*, 28(9):566 – 581, 2011.
- [14] R. Krone. Numerical algorithms for dual bases of positive-dimensional ideals. *Journal of Algebra and Its Applications*, 12(06):1350018, 2013.
- [15] R. Krone and A. Leykin. Eliminating dual spaces. *arXiv preprint arXiv:1503.02038*, 2015.
- [16] A. Leykin, J. Verschelde, and A. Zhao. Higher-order deflation for polynomial systems with isolated singular solutions. In *Algorithms in algebraic geometry*, pages 79–97. Springer, 2008.
- [17] N. Li and L. Zhi. Computing isolated singular solutions of polynomial systems: case of breadth one. *SIAM Journal on Numerical Analysis*, 50(1):354–372, 2012.
- [18] N. Li and L. Zhi. Computing the multiplicity structure of an isolated singular solution: Case of breadth one. *Journal of Symbolic Computation*, 47(6):700 – 710, 2012.
- [19] N. Li and L. Zhi. Verified error bounds for isolated singular solutions of polynomial systems. *SIAM Journal on Numerical Analysis*, 52(4):1623–1640, 2014.
- [20] Z. Li and L. Zhi. Computing the nearest singular univariate polynomials with given root multiplicities. *Theoretical Computer Science*, 479:150 – 162, 2013.

- [21] F. S. Macaulay. *The Algebraic Theory of Modular Systems*. Cambridge mathematical library. Cambridge University Press, Cambridge, New York, Melbourne, 1994.
- [22] T. Maekawa and N. M. Patrikalakis. Computation of singularities and intersections of offsets of planar curves. *Computer Aided Geometric Design*, 10(5):407 – 429, 1993.
- [23] A. Mantzaflaris and B. Mourrain. Deflation and certified isolation of singular zeros of polynomial systems. In *Proceedings of the 36th International Symposium on Symbolic and Algebraic Computation*, ISSAC '11, pages 249–256, New York, NY, USA, 2011. ACM.
- [24] A. Mantzaflaris and B. Mourrain. Singular zeros of polynomial systems. In T. Dokken and G. Muntingh, editors, *SAGA – Advances in ShApes, Geometry, and Algebra*, volume 10 of *Geometry and Computing*, pages 77–103. Springer International Publishing, 2014.
- [25] A. Mantzaflaris, H. Rahkooy, and Z. Zafeirakopoulos. On directional multiplicities of isolated points. Technical Report 15-08, RISC, Hagenberg, Austria, 2015.
- [26] M. Marinari, H. Möller, and T. Mora. On multiplicities in polynomial system solving. *Transactions of the American Mathematical Society*, 348(8):3283–3321, 1996.
- [27] M. G. Marinari, T. Mora, and H. M. Möller. Gröbner duality and multiplicities in polynomial system solving. In *Proceedings of the 1995 International Symposium on Symbolic and Algebraic Computation*, ISSAC '95, pages 167–179, New York, NY, USA, 1995. ACM.
- [28] B. Mourrain. Isolated points, duality and residues. *Journal of Pure and Applied Algebra*, 117:469–493, 1997.
- [29] H. Rahkooy. *Dual Space Algorithms for Computing Multiplicity Structure of Isolated Points*. PhD thesis, RISC, Hagenberg, Austria, 2015.
- [30] H. Rahkooy and Z. Zafeirakopoulos. Using resultants for inductive Gröbner bases computation. *ACM Communications in Computer Algebra*, 45(1), 2011.
- [31] H. J. Stetter. *Numerical Polynomial Algebra*. SIAM, 2004.
- [32] B. L. van der Waerden. *Algebra*, volume 1. Springer, New York, 7th edition, 1991. Based in part on lectures by E. Artin and E. Noether.
- [33] B. L. van der Waerden. *Algebra*, volume 2. Springer, New York, 5th edition, 1991. Based in part on lectures by E. Artin and E. Noether.

- [34] X. Wu and L. Zhi. Computing the multiplicity structure from geometric involutive form. In *Proceedings of the Twenty-first International Symposium on Symbolic and Algebraic Computation, ISSAC '08*, pages 325–332, New York, NY, USA, 2008. ACM.
- [35] Z. Zeng. The closedness subspace method for computing the multiplicity structure of a polynomial system. *Contemporary Mathematics*, 496:347, 2009.