



HAL
open science

0-Hecke algebra actions on coinvariants and flags

Jia Huang

► **To cite this version:**

Jia Huang. 0-Hecke algebra actions on coinvariants and flags. 23rd International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2011), 2011, Reykjavik, Iceland. pp.505-516, 10.46298/dmtcs.2929 . hal-01215048

HAL Id: hal-01215048

<https://inria.hal.science/hal-01215048>

Submitted on 13 Oct 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

0-Hecke algebra actions on coinvariants and flags

Jia Huang[†]

School of Mathematics, University of Minnesota, Minneapolis, MN, USA

Abstract. By investigating the action of the 0-Hecke algebra on the coinvariant algebra and the complete flag variety, we interpret generating functions counting the permutations with fixed inverse descent set by their inversion number and major index.

Résumé. En étudiant l'action de l'algèbre de 0-Hecke sur l'algèbre coinvariante et la variété de drapeaux complète, nous interprétons les fonctions génératrices qui comptent les permutations avec un ensemble inverse de descentes fixé, selon leur nombre d'inversions et leur "major index".

Keywords: 0-Hecke algebra, Ribbon number, Descent monomial, Demazure operator.

1 Introduction

A composition I of an integer n gives rise to a descent class of permutations in the symmetric group \mathfrak{S}_n ; the cardinality of this descent class is known as the *ribbon number* r_I and its inv-generating function is the q -*ribbon number* $r_I(q)$. Reiner and Stanton [15] defined a (q, t) -*ribbon number* $r_I(q, t)$, and gave an interpretation by representations of \mathfrak{S}_n and $GL(n, \mathbb{F}_q)$.

Our main object here is to obtain similar interpretations of various ribbon numbers by representations of the 0-Hecke algebra $H_n(0)$ of type A . Norton [14] decomposed $H_n(0)$ into a direct sum of 2^{n-1} distinct indecomposable $H_n(0)$ -submodules M_I indexed by compositions I of n . Consequently every indecomposable projective $H_n(0)$ -module is isomorphic to M_I for some I , and every simple $H_n(0)$ -module is isomorphic to $C_I = \text{top}(M_I) = M_I/\text{rad } M_I$ for some I .

1.1 Descent monomials and Demazure atoms

Our first result is related to the *descent monomials* within $\mathbb{Z}[\mathbf{x}] = \mathbb{Z}[x_1, \dots, x_n]$

$$x_w = \prod_{i \in D(w)} x_{w(i)}, \quad w \in \mathfrak{S}_n,$$

[†]Supported by NSF grant DMS-1001933.

introduced by Garsia [8] as a \mathbb{Z} -basis for the coinvariant algebra $\mathbb{Z}[\mathbf{x}]/(\mathbb{Z}[\mathbf{x}]_+^{\mathfrak{S}_n})$. The 0-Hecke algebra $H_n(0)$ acts on the coinvariant algebra $\mathbb{Z}[\mathbf{x}]/(\mathbb{Z}[\mathbf{x}]_+^{\mathfrak{S}_n})$ via the operators $\bar{\pi}_i = \pi_i - 1$, where π_i is the *Demazure operator* defined by

$$\pi_i f = \frac{x_i f - s_i(x_i f)}{x_i - x_{i+1}}.$$

Our first result shows that under this action:

- $\mathbb{Z}[\mathbf{x}]/(\mathbb{Z}[\mathbf{x}]_+^{\mathfrak{S}_n})$ is isomorphic to $H_n(0)$ as a (left) $H_n(0)$ -module;
- $\mathbb{Z}[\mathbf{x}]/(\mathbb{Z}[\mathbf{x}]_+^{\mathfrak{S}_n}) = \bigoplus_I N_I$, summed over all compositions of n , and $N_I \cong M_I$;
- $\mathbb{Z}[\mathbf{x}]/(\mathbb{Z}[\mathbf{x}]_+^{\mathfrak{S}_n})$ has a \mathbb{Z} -basis of certain *Demazure atoms* whose leading terms under some order are the descent monomials.

Since $\mathbb{Z}[\mathbf{x}]/(\mathbb{Z}[\mathbf{x}]_+^{\mathfrak{S}_n})$ has an extra grading by polynomial degree, it becomes a graded version of the left regular representation of $H_n(0)$.

1.2 A bigraded characteristic

Duchamp, Krob, Leclerc and Thibon [4] defined the *characteristic* of a finite dimensional $H_n(0)$ -module M with simple composition factors C_{I_1}, \dots, C_{I_k} to be

$$\mathcal{F}(M) = \sum_{i=1}^k F_{I_i}$$

where the F_I 's are *quasi-ribbon functions* which form a basis for the algebra of quasi-symmetric functions. Krob and Thibon [10] showed that $\mathcal{F}(M_I)$ is the *ribbon schur function* s_I , and thus $\mathcal{F}(M)$ is symmetric whenever M is projective.

If $M = H_n(0)v$ is cyclic then the *length filtration*

$$H_n(0)^{(\ell)} = \bigoplus_{\ell(w) \geq \ell} \mathbb{Z}T_w$$

induces a filtration of $H_n(0)$ -modules $M^{(\ell)} = H_n(0)^{(\ell)}v$, $k \geq 0$. This refines $\mathcal{F}(M)$ to a *graded characteristic*

$$\mathcal{F}_q(M) = \sum_{\ell \geq 0} q^\ell \mathcal{F}(M^{(\ell)}/M^{(\ell+1)}).$$

One has

$$\mathcal{F}_q(H_n(0)) = \sum_{w \in \mathfrak{S}_n} q^{\text{inv}(w)} F_{D(w^{-1})} = \sum_I r_I(q) F_I$$

and taking a limit as $q \rightarrow 1$ gives

$$\mathcal{F}(H_n(0)) = \sum_I r_I F_I.$$


If M has another filtration by $H_n(0)$ -modules M_d for $d \geq 0$, then one can look at the *bifiltration* by $H_n(0)$ -modules $M^{(\ell,d)} = M^{(\ell)} \cap M_d$ for $\ell, d \geq 0$, and define the *bigraded characteristic* to be

$$\mathcal{F}_{q,t}(M) = \sum_{\ell,d \geq 0} q^\ell t^d \mathcal{F} \left(M^{(\ell,d)} / (M^{(\ell+1,d)} + M^{(\ell,d+1)}) \right).$$

This happens to be the case for $M = \mathbb{Z}[\mathbf{x}] / (\mathbb{Z}[\mathbf{x}]_+^{\mathfrak{S}_n})$ with its length filtration (since $M \cong H_n(0)$) and its polynomial degree filtration by $M_d = \langle \bar{f} : \deg f \geq d \rangle_{\mathbb{Z}}$ for $d \geq 0$. Our next result is

- $\mathcal{F}_{q,t} \left(\mathbb{Z}[\mathbf{x}] / (\mathbb{Z}[\mathbf{x}]_+^{\mathfrak{S}_n}) \right) = \sum_{w \in \mathfrak{S}_n} t^{\text{maj}(w)} q^{\text{inv}(w)} F_{D(w^{-1})}$

which completes the following picture.



Here the left inverse descent set $D(w^{-1})$ is identified with its descent composition, and the equality $\mathcal{F}_q(H_n(0)) \xrightarrow{q \leftrightarrow t} \mathcal{F}_t(\mathbb{Z}[\mathbf{x}] / (\mathbb{Z}[\mathbf{x}]_+^{\mathfrak{S}_n}))$ comes from the equidistribution of inv and maj over inverse descent classes, proved by Foata and Schützenberger [7]. We shall see in Section 3 that r_I and $r_I(q)$ appear as coefficients of F_I in $\mathcal{F}(H_n(0))$ and $\mathcal{F}_q(H_n(0))$.

1.3 Complete flag variety


Consider the general linear group $G = GL(n, \mathbb{F}_q)$ over a finite field \mathbb{F}_q and its Borel subgroup B . The 0-Hecke algebra $H_n(0)$ acts on the *complete flag variety* $1_B^G = \mathbb{F}_q[G/B]$ by $T_w B = BwB$. This action induces an $H_n(0)$ -module structure on

$$\text{Hom}_{\mathbb{F}_q[G]} (1_B^G, \mathbb{F}_q[\mathbf{x}]) \cong \mathbb{F}_q[\mathbf{x}]^B$$

which is $\mathbb{F}_q[\mathbf{x}]^G$ -linear, hence inducing an $H_n(0)$ -module structure on $\mathbb{F}_q[\mathbf{x}]^B / (\mathbb{F}_q[\mathbf{x}]_+^G)$. Our next result is:

- $\mathcal{F}(1_B^G) = \sum_I r_I(q) F_I$;
- $\mathcal{F}_t(\mathbb{F}_q[\mathbf{x}]^B / (\mathbb{F}_q[\mathbf{x}]_+^G)) = \sum_I r_I(q, t) F_I$.

Therefore we have another picture as follows, which interprets all the ribbon numbers mentioned at the beginning.



1.4 Generalizations to other types

The $H_n(0)$ -actions on the coinvariants and complete flags can be generalized to the following setting. Let W be a Weyl group with weight lattice Λ . The 0-Hecke algebra \mathcal{H} of type W acts on the group ring $\mathbb{Z}[\Lambda]$ via the operators $\bar{\pi}_i = \pi_i - 1$ where π_i was originally considered by Demazure [2] in this setting. Garsia and Stanton [9] constructed the descent monomials in $\mathbb{Z}[\Lambda]$, which form a free basis over $\mathbb{Z}[\Lambda]^W$. We prove that, similarly to $\mathbb{Z}[\mathbf{x}] / (\mathbb{Z}[\mathbf{x}]_+^{\mathfrak{S}_n})$, taking a quotient of $\mathbb{Z}[\Lambda]$ by the ideal generated by “positive” W -invariants leads to an \mathcal{H} -module isomorphic to \mathcal{H} (without an extra grading). We also have an \mathcal{H} -action on 1_B^G by $T_w B = BwB$, where G is a finite group with split BN -pair of characteristic $p > 0$, whose Weyl group is W . We determine the characteristic $\mathcal{F}(1_B^G)$ in the same manner as for type A .

In Section 2 we review the definitions for the various ribbon numbers and their interpretation by representations of \mathfrak{S}_n and $GL(n, \mathbb{F}_q)$. In Section 3 we recall the representation theory of the 0-Hecke algebra. The result for the coinvariant algebra $\mathbb{Z}[\mathbf{x}] / (\mathbb{Z}[\mathbf{x}]_+^{\mathfrak{S}_n})$ is given in Section 4, and a similar result for $\mathbb{Z}[\Lambda]$ of the weight lattice Λ of a Weyl group is stated in Section 5. The \mathcal{H} -action on the complete flag variety 1_B^G and the coinvariant algebra of (G, B) is investigated in Section 6. Lastly some questions are asked in Section 7.

2 Ribbon numbers

We recall from Reiner and Stanton [15, §9, §10] the definitions and properties of the various ribbon numbers. Let $I = (i_1, \dots, i_k)$ be a composition of n , let $\sigma_j = i_1 + \dots + i_j$ for $j = 1, \dots, k$, and let the descent set of I be $D(I) = \{\sigma_1, \dots, \sigma_{k-1}\}$.

It is well known that compositions of n bijectively correspond to the subsets of $[n - 1]$ via their descent set; they also bijectively correspond to the ribbon diagrams, i.e. connected skew Young diagrams without 2×2 boxes, whose row sizes from bottom to top are i_1, \dots, i_n .

The *descent class* of I is the set of all permutations w in \mathfrak{S}_n with $D(w) = D(I)$, and the *inverse descent class* is the set of all w in \mathfrak{S}_n with $D(w^{-1}) = D(I)$. The *ribbon number* r_I is the cardinality of the descent class of I , and the *q-ribbon number* and *t-ribbon number* are

$$r_I(q) = \sum_{w \in \mathfrak{S}_n : D(w) = D(I)} q^{\text{inv}(w)} = [n]!_q \det \left(\frac{1}{[\sigma_j - \sigma_{i-1}]!_q} \right)_{i,j=1}^k,$$

$$r_I(t) = \sum_{w \in \mathfrak{S}_n : D(w) = D(I)} t^{\text{maj}(w^{-1})} = [n]!_t \det \left(\frac{1}{[\sigma_j - \sigma_{i-1}]!_t} \right)_{i,j=1}^k.$$

where $[n]!_q = [n]_q [n-1]_q \cdots [1]_q$ and $[n]_q = 1 + q + \cdots + q^{n-1}$. The notations make sense because $r_I(t) = r_I(q)|_{q=t}$, which is a consequence of the equidistribution of inv and maj on every inverse descent class, proved by Foata and Schützenberger [7]. A further generalization is the (q, t) -ribbon number

$$r_I(q, t) = n!_{q,t} \det \left(\varphi^{\sigma_{i-1}} \frac{1}{(\sigma_j - \sigma_{i-1})!_{q,t}} \right)_{i,j=1}^k$$


where $n!_{q,t} = (1 - t^{q^n-1})(1 - t^{q^n-q}) \cdots (1 - t^{q^n-q^{n-1}})$, and $\varphi : t \mapsto t^q$ is the *Frobenius operator*.

All these ribbon numbers can be calculated by similar determinantal formulae, and are interpreted by certain *homology representations* χ^I of \mathfrak{S}_n , and χ_q^I of $G = GL(n, \mathbb{F}_q)$, together with their intertwiners

$$M^I = \text{Hom}_{\mathbb{Z}\mathfrak{S}_n} (\chi^I, \mathbb{Z}[\mathbf{x}]), \text{ a } \mathbb{Z}[\mathbf{x}]^{\mathfrak{S}_n}\text{-module,}$$

$$M_q^I = \text{Hom}_{\mathbb{F}_q G} (\chi_q^I, \mathbb{F}_q[\mathbf{x}]), \text{ an } \mathbb{F}_q[\mathbf{x}]^G\text{-module.}$$

Here χ^I (χ_q^I resp.) is the top homology of the *rank-selected Coxeter complex* $\Delta(\mathfrak{S}_n)_I$ (*Tits building* $\Delta(G)_I$ resp.). Precisely, one has the following picture.



3 Representation theory of the 0-Hecke algebra

Recall from Norton [14] the representation theory of the 0-Hecke algebra. Let

$$W = \langle s_1, \dots, s_\ell : s_i^2 = 1, (s_i s_j s_i \cdots)_{m_{ij}} = (s_j s_i s_j \cdots)_{m_{ij}}, 1 \leq i \neq j \leq \ell \rangle$$

be a finite Coxeter group, where $(aba \cdots)_m$ denotes an alternating product of m terms. The 0-Hecke algebra \mathcal{H} of type W is an associative \mathbb{Z} -algebra generated by T_1, \dots, T_ℓ with relations

$$\begin{cases} T_i^2 = -T_i, & 1 \leq i \leq \ell, \\ (T_i T_j T_i \cdots)_{m_{ij}} = (T_j T_i T_j \cdots)_{m_{ij}}, & 1 \leq i \neq j \leq \ell. \end{cases}$$

Norton [14] decomposed \mathcal{H} into a direct sum of 2^ℓ distinct indecomposable submodules M_I indexed by compositions I of $\ell + 1$, with $C_I = \text{top}(M_I) = M_I/\text{rad } M_I$ being the (one-dimensional) simple module given by

$$\rho_I(T_i) = \begin{cases} -1, & \text{if } i \in D(I), \\ 0, & \text{if } i \notin D(I). \end{cases}$$

This gives a complete list of indecomposable projective \mathcal{H} -modules and simple \mathcal{H} -modules.

To explicitly construct M_I inside \mathcal{H} , let $T'_i = T_i + 1, 1 \leq i \leq \ell$. One can check that $(T'_i)^2 = T'_i$, i.e. $T_i T'_i = 0$, and $(T'_i T'_j T'_i \cdots)_{m_{ij}} = (T'_j T'_i T'_j \cdots)_{m_{ij}}, 1 \leq i \neq j \leq \ell$; see [5, Lemma 3.1] or [14, Lemma 4.3]. Thus $T_w = T_{i_1} \cdots T_{i_k}$ and $T'_w = T'_{i_1} \cdots T'_{i_k}$ are both well-defined if $w = s_{i_1} \cdots s_{i_k}$ is reduced.

Given a composition $I = (i_1, \dots, i_k)$, let $\bar{I} = (i_k, \dots, i_1)$ and let I^\sim be the conjugate composition of I obtained by reflecting the ribbon diagram of I across the diagonal. For example, if $I = (2, 1, 3)$ then $\bar{I} = (3, 1, 2)$ and $I^\sim = (1, 1, 3, 1)$. The descent class of I is an interval $[\alpha(I), \omega(I)]$ in the left weak order of W , where $\alpha(I)$ is the top element in the parabolic subgroup $W_{D(I)}$. One can write the module M_I in Norton's decomposition of \mathcal{H} as $M_I = \mathcal{H} \cdot T_{\alpha(I)} T'_{\alpha(I^\sim)}$, which has a \mathbb{Z} -basis given by

$$\left\{ T_w T'_{\alpha(I^\sim)} : w \in [\alpha(I), \omega(I)] \right\}.$$

4 Coinvariant algebra of \mathfrak{S}_n

The symmetric group \mathfrak{S}_n acts naturally on the polynomial ring $\mathbb{Z}[\mathbf{x}] = \mathbb{Z}[x_1, \dots, x_n]$ by permuting variables. The *Demazure operators* π_i are defined by

$$\pi_i f = \frac{x_i f - s_i(x_i f)}{x_i - x_{i+1}}$$

where s_i is the adjacent transposition $(i, i + 1)$. One checks that the operators $\bar{\pi}_i = \pi_i - 1$ satisfy the same relations as T_i , preserve the polynomial grading of $\mathbb{Z}[\mathbf{x}]$, and are $\mathbb{Z}[\mathbf{x}]^{\mathfrak{S}_n}$ -linear. Thus one has an $H_n(0)$ -action on the coinvariant algebra $\mathbb{Z}[\mathbf{x}]/(\mathbb{Z}[\mathbf{x}]_+^{\mathfrak{S}_n})$ by sending T_i to $\bar{\pi}_i$, or sending T_w to $\bar{\pi}_w = \bar{\pi}_{i_1} \cdots \bar{\pi}_{i_k}$ if $w = s_{i_1} \cdots s_{i_k}$ is reduced.

One sees from the definition that $\bar{\pi}_i$ fixes all variables x_j except x_i and x_{i+1} , and

$$\bar{\pi}_i(x_i^a x_{i+1}^b) = \begin{cases} x_i^{a-1} x_{i+1}^{b+1} + x_i^{a-2} x_{i+1}^{b+2} \cdots + x_i^b x_{i+1}^a, & \text{if } a > b, \\ 0, & \text{if } a = b, \\ -x_i^a x_{i+1}^b - x_i^{a+1} x_{i+1}^{b-1} - \cdots - x_i^{b-1} x_{i+1}^{a+1}, & \text{if } a < b. \end{cases} \tag{1}$$

The module $\mathbb{Z}[\mathbf{x}]$ is free over $\mathbb{Z}[\mathbf{x}]^{\mathfrak{S}_n}$, with a basis consisting of the *descent monomials*

$$x_w = \prod_{i \in D(w)} x_{w(i)}, \quad w \in \mathfrak{S}_n,$$

constructed by Garsia [8]. We shall obtain a \mathbb{Z} -basis for the $H_n(0)$ -module $\mathbb{Z}[\mathbf{x}]/(\mathbb{Z}[\mathbf{x}]_+^{\mathfrak{S}_n})$ from certain Demazure atoms, i.e. $\bar{\pi}_w x_I$ for certain permutations w and compositions I , where

$$x_I = \prod_{i \in D(I)} x_1 \cdots x_i.$$

See Mason [13] for more information on the Demazure atoms. The leading terms of these Demazure atoms are exactly the descent monomials, under any linear extension of the following partial order. Given two monomials x^d and x^e , say $x^d \prec x^e$ if $\lambda(d) <_L \lambda(e)$ in the lexicographic order, where $\lambda(d)$ is the unique partition obtained from rearranging the exponent vector d , and similarly for $\lambda(e)$.

Lemma 4.1 *Suppose that I is a composition of n and w is a permutation in \mathfrak{S}_n with $D(w) \subseteq D(I)$. Then*

$$\bar{\pi}_w x_I = wx_I + \sum_{d: x^d \prec x_I} c_d x^d$$

where $c_d \in \mathbb{Z}$; moreover, wx_I is a descent monomial if and only if $D(w) = D(I)$.

Theorem 4.2 *The coinvariant algebra $\mathbb{Z}[\mathbf{x}]/(\mathbb{Z}[\mathbf{x}]_+^{\mathfrak{S}_n})$ has a \mathbb{Z} -basis*

$$\{\bar{\pi}_w \bar{x}_I : w \in \mathfrak{S}_n, D(I) = D(w)\} \tag{2}$$

and decomposes into a direct sum of $H_n(0)$ -modules

$$N_I = H_n(0) \cdot \bar{\pi}_{\alpha(I)} \bar{x}_I$$

as I runs through all compositions of n ; moreover, N_I is isomorphic to $M_I \subseteq H_n(0)$ and has a basis $\{\bar{\pi}_w \bar{x}_I : w \in [\alpha(I), \omega(I)]\}$. Thus $\mathbb{Z}[\mathbf{x}]/(\mathbb{Z}[\mathbf{x}]_+^{\mathfrak{S}_n})$ is isomorphic to $H_n(0)$ as a left $H_n(0)$ -module.

Since each submodule N_I consists of homogeneous elements of degree $\text{maj}(w)$ for any $w \in [\alpha(I), \omega(I)]$, one has

$$\mathcal{F}_{q,t} \left(\mathbb{Z}[\mathbf{x}]/(\mathbb{Z}[\mathbf{x}]_+^{\mathfrak{S}_n}) \right) = \sum_{w \in \mathfrak{S}_n} t^{\text{maj}(w)} q^{\text{inv}(w)} F_{D(w^{-1})} \tag{3}$$

which relates inv to the equidistributed maj (on every inverse descent class). In fact, one can get

$$\mathcal{F}_t(\mathbb{Z}[\mathbf{x}]) = \frac{1}{(1-t)(1-t^2)\cdots(1-t^n)} \sum_{w \in \mathfrak{S}_n} t^{\text{maj}(w)} F_{D(w^{-1})} \tag{4}$$

directly from (1) using the filtration of $\mathbb{Z}[\mathbf{x}]$ induced from the following order (which appeared in Allen [1]): $x^d <_{ts} x^e$ if $\lambda(d) <_L \lambda(e)$, or if $\lambda(d) = \lambda(e)$ and $x^d <_L x^e$ in the lexicographic order. From (4) one can immediately deduce $\mathcal{F}_t \left(\mathbb{Z}[\mathbf{x}]/(\mathbb{Z}[\mathbf{x}]_+^{\mathfrak{S}_n}) \right)$, since $\bar{\pi}_i$ is $\mathbb{Z}[\mathbf{x}]^{\mathfrak{S}_n}$ -linear.

5 Coinvariant algebra of Weyl group

Demazure's character formula [2] expresses the character of highest weight modules over a semisimple algebra using the Demazure operators π_i on the group ring $\mathbb{Z}[\Lambda]$ of the weight lattice Λ . If e^λ is the

element in $\mathbb{Z}[\Lambda]$ corresponding to the weight $\lambda \in \Lambda$, and if $\lambda_1, \dots, \lambda_\ell$ denote the fundamental weights with $z_i = e^{\lambda_i}$, then

$$\mathbb{Z}[\Lambda] = \mathbb{Z}[z_1, \dots, z_\ell, z_1^{-1}, \dots, z_\ell^{-1}].$$

Suppose that $\alpha_1, \dots, \alpha_\ell$ are the simple roots which correspond to simple reflections s_1, \dots, s_ℓ . Then the Demazure operators are defined by

$$\pi_i = \frac{1 - e^{-\alpha_i s_i}}{1 - e^{-\alpha_i}}.$$

It follows easily that

$$\pi_i(e^\lambda) = \begin{cases} e^\lambda + e^{\lambda - \alpha_i} + \dots + e^{s_i \lambda}, & \text{if } \langle \alpha_i^\vee, \lambda \rangle \geq 0, \\ 0, & \text{if } \langle \alpha_i^\vee, \lambda \rangle = -1, \\ -e^{\lambda + \alpha_i} - \dots - e^{s_i \lambda - \alpha_i}, & \text{if } \langle \alpha_i^\vee, \lambda \rangle < -1. \end{cases} \tag{5}$$

See, for example, Kumar [12]. Demazure operators satisfy the braid relation [2, 5.5] and

$$s_i \pi_i = \pi_i, \quad \pi_i^2 = \pi_i.$$

Hence the 0-Hecke algebra \mathcal{H} of W acts on $\mathbb{Z}[\Lambda]$ by sending T_i to $\bar{\pi}_i = \pi_i - 1$.

Using the Stanley-Reisner ring of the Coxeter complex of W , Garsia and Stanton [9] showed that

$$\mathbb{Z}[\Lambda]^W = \mathbb{Z}[a_1, \dots, a_\ell]$$

where

$$a_i = \sum_{w \in W/W_{\{i\}^c}} e^{w\lambda_i}$$

and $\mathbb{Z}[\Lambda]$ has a free basis over $\mathbb{Z}[\Lambda]^W$, which consists of the *descent monomials*

$$z_w = \prod_{i \in D(w)} e^{w\lambda_i}, \quad w \in W$$

(see also Steinberg [17]). This basis induces a \mathbb{Z} -basis for $\mathbb{Z}[\Lambda]/(a_1, \dots, a_\ell)$.

The \mathcal{H} -action on $\mathbb{Z}[\Lambda]$ is $\mathbb{Z}[\Lambda]^W$ -linear, hence inducing an \mathcal{H} -action on $\mathbb{Z}[\Lambda]/(a_1, \dots, a_\ell)$.

For any weight λ in Λ , there exists a unique *dominant weight* (which is a nonnegative linear combination of $\lambda_1, \dots, \lambda_\ell$), denoted by $[\lambda]$, such that $\lambda = w[\lambda]$ for some w in W .

Lemma 5.1 *Given a composition I of $\ell + 1$, let $z_I = e^{\lambda_I}$ where*

$$\lambda_I = \sum_{i \in D(I)} \lambda_i.$$

Suppose that w lies in W with $D(w) \subseteq D(I)$. Then

$$\bar{\pi}_w z_I = e^{w\lambda_I} + \sum_{[\lambda] < \lambda_I} c_\lambda e^\lambda, \quad c_\lambda \in \mathbb{Z},$$

where $[\lambda] < \lambda_I$ means $\lambda_I - [\lambda]$ is a nonnegative linear combination of simple roots and $[\lambda] \neq \lambda_I$; moreover, $e^{w\lambda_I}$ is a descent monomial if and only if $D(w) = D(I)$.

Theorem 5.2 *The coinvariant algebra $\mathbb{Z}[\Lambda]/(a_1, \dots, a_\ell)$ has a \mathbb{Z} -basis*

$$\{\bar{\pi}_w \bar{z}_I : w \in W, D(I) = D(w)\}$$

and decomposes into a direct sum of \mathcal{H} -modules

$$N_I = \mathcal{H} \cdot \bar{\pi}_{\alpha(I)} \bar{z}_I$$

as I runs through all compositions of $\ell + 1$; moreover, N_I is isomorphic to $M_I \subseteq \mathcal{H}$ and has a basis $\{\bar{\pi}_w \bar{z}_I : w \in [\alpha(I), \omega(I)]\}$. Thus $\mathbb{Z}[\Lambda]/(a_1, \dots, a_\ell)$ is isomorphic to \mathcal{H} as an \mathcal{H} -module.

Remark 5.3 Garsia and Stanton [9] pointed out a way to reduce the descent monomials in $\mathbb{Z}[\Lambda]$ to the descent monomials in $\mathbb{Z}[\mathbf{x}]$ for type A . However, it does not give Theorem 4.2 directly from Theorem 5.2; instead, one should consider the Demazure operators on $\mathbb{Z}[X(T)]$ where $X(T)$ is the character group of the maximal torus T of $GL(n, \mathbb{C})$.

6 Complete flag variety 1_B^G and coinvariants of (G, B)

Let G be a finite group with split BN -pair of characteristic $p > 0$, whose Weyl group W is generated by ℓ simple reflections. Let $1_B^G = \mathbb{Z}[B \setminus G]$ be the induction of the right trivial representation of B to G , i.e. the permutation representation on the right cosets $B \setminus G$.

Given a subset $S \subseteq G$, let $\bar{S} = \sum_{s \in S} s$ in $\mathbb{Z}[G]$. Then $1_B^G \cong \bar{B} \cdot \mathbb{Z}[G]$ and $\text{End}_{\mathbb{Z}[G]}(1_B^G)$ has a basis $\{f_w : w \in \mathfrak{S}_n\}$, with f_w given by

$$f_w(\bar{B}) = \overline{BwB} = \overline{U_w w \bar{B}} \tag{6}$$

where U_w is the product of root subgroups U_α with $\alpha > 0, w^{-1}(\alpha) < 0$ [3, Proposition 1.7]. The endomorphism ring $\text{End}_{\mathbb{Z}[G]}(1_B^G)$ is isomorphic to the Hecke algebra of W with parameters $q_i = |U_{s_i}|$, since the relations satisfied by $\{f_w\}$ are the same as those satisfied by the standard basis $\{T_w\}$. It follows that the 0-Hecke algebra \mathcal{H} of type W acts on $1_B^G \otimes \mathbb{F}_p$ by (6). Dually, the left cosets of B give rise to a right \mathcal{H} -action. See Kuhn [11] for details.

Since we are mainly concerned with the 0-Hecke algebra, we shall write $1_B^G = 1_B^G \otimes \mathbb{F}_p$ for simplicity, and similar for $1_{P_I}^G$ where $P_I = BW_{D(I)}^c B$ is the *parabolic subgroup* of G indexed by the composition I of $\ell + 1$. For type A , one has $G = GL(n, \mathbb{F}_q)$ and P_I is the group of all upper triangular block matrices with invertible diagonal blocks of sizes given by the parts of I .

To determine the simple factors of an \mathcal{H} -module, we develop the following lemma, where the (graded) characteristic is a natural extension of that of type A , with the F_I 's simply being independent variables.

Lemma 6.1 *Given a finite dimensional graded \mathcal{H} -module Q , let Q_I be the submodule of elements that are annihilated by all T_j with $j \notin D(I)$, for any composition I of $\ell + 1$. Then*

$$\mathcal{F}_i(Q) = \sum_I c_I(Q) F_I,$$

summed over all compositions I of $\ell + 1$, where

$$c_I(Q) = \sum_{J: D(J) \subseteq D(I)} (-1)^{\ell(I, J)} \text{Hilb}(Q_J, t).$$

Define the q -ribbon number of type G to be

$$r_I(G) = \sum_{w \in W: D(w)=D(I)} |U_w|.$$

The characteristic of 1_B^G can be obtained by using Lemma 6.1 and the observations $(1_B^G)_I = 1_{P_I}^G$,

$$|G/P_I| = \sum_{w \in W: D(w) \subseteq D(I)} |U_w|.$$

Theorem 6.2 $\mathcal{F}(1_B^G) = \sum_I r_I(G)F_I$.

If G is a finite group of Lie type over a finite field \mathbb{F}_q , then $|U_w| = q^{\ell(w)}$. In particular, for type A , i.e. when $G = GL(n, \mathbb{F}_q)$, one has $r_I(G) = r_I(q)$. Furthermore, G acts on $\mathbb{F}_q[\mathbf{x}]$ and the $H_n(0)$ -action on 1_B^G induces an $H_n(0)$ -module structure on

$$\text{Hom}_{\mathbb{F}_q[G]}(1_B^G, \mathbb{F}_q[\mathbf{x}]) \cong \mathbb{F}_q[\mathbf{x}]^B$$

which is $\mathbb{F}_q[\mathbf{x}]^G$ -linear, hence inducing an $H_n(0)$ -module structure on $\mathbb{F}_q[\mathbf{x}]^B/(\mathbb{F}_q[\mathbf{x}]_+^G)$. Explicitly,

$$T_w(f) = \bar{U}_w wf, \text{ for all } f \text{ in } \mathbb{F}_q[\mathbf{x}]^B.$$

We observe that

$$(\mathbb{F}_q[\mathbf{x}]^B/(\mathbb{F}_q[\mathbf{x}]_+^G))_I = \{ \bar{f} : f \in \mathbb{F}_q[\mathbf{x}]^{P_I} \}.$$

It follows from Lemma 6.1 that

Theorem 6.3 *If $G = GL(n, \mathbb{F}_q)$ and B is the Borel subgroup of G , then*

$$\mathcal{F}_t(\mathbb{F}_q[\mathbf{x}]^B/(\mathbb{F}_q[\mathbf{x}]_+^G)) = \sum_I r_I(q, t)F_I.$$

We conclude this section with a description of the homology representation χ_q^I using the (G, \mathcal{H}) -bimodule structure on 1_B^G , which is now spanned by left B -cosets. Smith [16] showed the following left G -module decomposition

$$1_B^G = \bigoplus_J \chi_q^J. \tag{7}$$

On the other hand, the decomposition of the 0-Hecke algebra

$$\mathcal{H} = \bigoplus_I \mathcal{H} \cdot T_{\alpha(I)} T'_{\alpha(\bar{I} \sim)}$$

implies a unique way to write

$$1 = \sum_I h_I T_{\alpha(I)} T'_{\alpha(\bar{I} \sim)}, \quad h_I \in \mathcal{H}.$$

By the right action of \mathcal{H} on 1_B^G , one has

$$1_B^G = \sum_I 1_B^G h_I T_{\alpha(I)} T'_{\alpha(\bar{I} \sim)} \tag{8}$$

as a left G -module. We show that this is the same as the decomposition (7), that is,

Theorem 6.4

$$1_B^G h_I T_{\alpha(I)} T'_{\alpha(\tilde{I}^c)} = \chi_q^I.$$

One sees from the above equality that the left G -module χ_q^I is in general not a right \mathcal{H} -module; however, the trivial G -representation $\chi_q^{(n)}$ and the Steinberg representation $\chi_q^{(1^n)}$ are right (isotypic) \mathcal{H} -modules.

7 Remaining Questions

The equidistribution of inv and maj was first proved on permutations of multisets by P.A. MacMahon in the 1910s; applying an inclusion-exclusion would give their equidistribution on inverse descent classes of \mathfrak{S}_n . However, the first proof for the latter result appearing in the literature was by Foata and Schützenberger [7] in 1970, using a bijection constructed earlier by Foata [6]. Is there an algebraic proof from the (q, t) -bigraded characteristic (3) of $\mathbb{Z}[\mathbf{x}]/(\mathbb{Z}[\mathbf{x}]_+^{\mathfrak{S}_n})$, which involves inv , maj , and inverse descents?

The next question is on the \mathcal{H} -module 1_B^G ; its simple factors are given by Theorem 6.2, but its decomposition into indecomposable submodules is *not* obtained in general. Assume $G = GL(n, \mathbb{F}_q)$ below. Recall that

$$H_3(0) = M_3 \oplus M_{21} \oplus M_{12} \oplus M_{111}$$

where M_I is the indecomposable projective $H_3(0)$ -module indexed by the composition I . For $n = 3$ we have candidates for a q -analogous $H_3(0)$ -module decomposition

$$1_B^G \cong M_3 \oplus (M_{21} \oplus M_{12})^{\oplus \binom{q+1}{2}} \oplus (M_{111})^{\oplus q^3}$$

which has been checked correct for $q = 2, 3, 5, 7$, showing that 1_B^G is a projective $H_3(0)$ -modules in these cases. Is this true for all *primes* q ?

On the other hand, for $n = 3, q = 4, 8$, and $n = 4, q = 2, 3$, computations show that 1_B^G is *not* projective, although the characteristic of 1_B^G is always symmetric. In fact, using the RSK correspondence one can show that

$$\mathcal{F}(1_B^G) = \sum_{\lambda \vdash n} q^{b(\lambda)} \frac{[n]!_q}{\prod_{u \in \lambda} [h_u]_q} s_\lambda.$$

where h_u is the hook length of the box u in the Young diagram of λ and $b(\lambda) = \lambda_2 + 2\lambda_3 + 3\lambda_4 + \dots$.

Computations also show that $\mathbb{F}_q[\mathbf{x}]^B/(\mathbb{F}_q[\mathbf{x}]_+^G)$ is *not* projective for $n = 3, q = 2, 3$. Besides being curious to know its decomposition, we wonder if there is any q -analogue of the Demazure operators, which might give another $H_n(0)$ -action on $\mathbb{F}_q[\mathbf{x}]^B/(\mathbb{F}_q[\mathbf{x}]_+^G)$.

Acknowledgements

The author is grateful to Victor Reiner for asking questions which led to the research and for giving many constructive suggestions. He is indebted to Andrew Berget, Tom Denton, Anne Schilling, and Monica Vazirani for interesting conversations and hospitality during the visit to UC Davis. He thanks Marcelo Aguiar, Sami Assaf, Nantel Bergeron, Adriano Garsia, Patricia Hersh, Liping Li, Sarah Mason, Dennis Stanton, Peter Webb, and two anonymous referees for their helpful suggestions and comments.

References

- [1] E.E. Allen, The descent monomials and a basis for the diagonally symmetric polynomials, *J. Algebraic Combin.* 3 (1994) 5–16.
- [2] M. Demazure, Désingularisation des variétés de Schubert généralisées, *Ann. Sci. École Norm. Sup.* 7 (1974) 53–88.
- [3] F. Digne and J. Michel, Representations of Finite Groups of Lie Type, London Math. Soc. Student Texts 21, Cambridge University Press (1991).
- [4] G. Duchamp, D. Krob, B. Leclerc, and J.-Y. Thibon, Fonctions quasi-symétriques, fonctions symétriques non commutatives et algèbres de Hecke à $q = 0$. *C.R. Acad. Sci. Paris* 322 (1996) 107–112.
- [5] M. Fayers, 0-Hecke algebras of finite Coxeter groups, *J. Pure Appl. Algebra* 199 (2005) 27–41.
- [6] D. Foata, On the Netto inversion number of a sequence, *Proc. Amer. Math. Soc.* 19 (1968) 236–240.
- [7] D. Foata and M.-P. Schützenberger, Major index and inversion number of permutations, *Math. Nachr.* 83 (1970), 143–159.
- [8] A. Garsia, Combinatorial methods in the theory of Cohen-Macaulay rings, *Adv. Math.* 38 (1980) 229–266.
- [9] A. Garsia and D. Stanton, Group actions of Stanley-Reisner rings and invariant of permutation groups, *Adv. Math.* 51 (1984) 107–201.
- [10] D. Krob and J.-Y. Thibon, Noncommutative symmetric functions IV: Quantum linear groups and Hecke algebras at $q = 0$, *J. Algebraic Combin.* 6 (1997) 339–376.
- [11] N. Kuhn, The modular Hecke algebra and Steinberg representation of finite Chevalley groups, *J. Algebra* 91 (1984), 125–141.
- [12] S. Kumar, Demazure character formula in arbitrary Kac-Moody setting, *Invent. Math.* 89 (1987) 395–423.
- [13] S. Mason, An explicit construction of type A Demazure atoms. *J. Algebraic Combin.* 29 (2009), 295–313.
- [14] P.N. Norton, 0-Hecke algebras, *J. Austral. Math. Soc. A* 27 (1979) 337–357.
- [15] V. Reiner and D. Stanton, (q, t) -analogues and $GL_n(\mathbb{F}_q)$, *J. Algebr. Comb.* 31 (2010) 411–454.
- [16] S.D. Smith, On decomposition of modular representations from Cohen-Macaulay geometries, *J. Algebra* 131 (1990) 598–625.
- [17] R. Steinberg, On a theorem of Pittie, *Topology* 14 (1975) 173–177.