



HAL
open science

On the quasi-hydrostatic quasi-geostrophic model

Antoine Rousseau, James C. McWilliams

► **To cite this version:**

Antoine Rousseau, James C. McWilliams. On the quasi-hydrostatic quasi-geostrophic model. [Research Report] RR-7531, 2011, pp.12. hal-01232740v3

HAL Id: hal-01232740

<https://inria.hal.science/hal-01232740v3>

Submitted on 18 Apr 2011 (v3), last revised 27 Jan 2017 (v7)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



INSTITUT NATIONAL DE RECHERCHE EN INFORMATIQUE ET EN AUTOMATIQUE

*A propos du modèle quasi-geostrophique
quasi-hydrostatique*

Antoine Rousseau — James C. McWilliams

N° 7531

Mars 2011

Observation and Modeling for Environmental Sciences

A large, light gray, stylized letter 'R' is positioned to the left of the text. A horizontal gray bar is located below the text.

R
*apport
de recherche*

A propos du modèle quasi-geostrophique quasi-hydrostatique

Antoine Rousseau^{*}, James C. McWilliams[†]

Theme : Observation and Modeling for Environmental Sciences
Équipe-Projet Moise

Rapport de recherche n° 7531 — Mars 2011 — 12 pages

Résumé : Ce travail présente une dérivation rigoureuse des équations quasi-geostrophiques (QG) de l’océan grande échelle lorsque le nombre de Rossby tend vers 0. On suit les techniques classiques de dérivation de ces modèles (voir par exemple [BB94]), mais les équations d’origine sur lesquelles on s’appuie prennent en compte tous les termes de rotation, ainsi que dans [LPR10]. Le modèle ainsi obtenu est légèrement différent du modèle QG traditionnel, et présente une seconde direction verticale, légèrement inclinée. Ceci illustre des simulations numériques et des expérimentations antérieures (voir [She04, WB06, GZMvH08]).

Mots-clés : analyse multi-échelles, force de Coriolis, approximation traditionnelle, modèles de circulation générale océanique, limite quasi-geostrophique.

^{*} INRIA, Laboratoire Jean Kuntzmann, BP 53, 38041 Grenoble Cedex 9, France (Antoine.Rousseau@inria.fr).

[†] Dept. of Atmospheric and Oceanic Sciences University of California, Los Angeles (UCLA) Mathematical Sciences Building, Room 7983 Los Angeles, CA 90095-1565 (jcm@atmos.ucla.edu).

On the quasi-hydrostatic quasi-geostrophic model

Abstract: This paper introduces a rigorous derivation of the quasi-geostrophic (QG) equations of large scale ocean as the Rossby number goes to zero. We follow classical techniques for the derivation (as in [BB94]), but the primitive equations that we consider account for all the rotating terms, as in [LPR10]. We end up with a slightly different QG model with a tilted vertical direction, which has been illustrated in previous works (see [She04, WB06, GZMvH08]).

Key-words: multiscale analysis, Coriolis force, traditional approximation, ocean global circulation models, quasi-geostrophic limit.

Abstract

This paper introduces a rigorous derivation of the quasi-hydrostatic quasi-geostrophic (QHQG) equations of large scale ocean as the Rossby number goes to zero. We follow classical techniques for the derivation of the quasi-geostrophic (QG) equations (as in [BB94]), but the primitive equations that we consider account for all the rotating terms, as in [LPR10]. We end up with a slightly different QG model with a tilted vertical direction, which has been illustrated in previous works using the primitive equations (see [She04, WB06, GZMvH08, SKR02]).

1 Introduction

The quasi-geostrophic equations are very familiar to oceanographers and meteorologists for they have been extensively used for modeling oceanic and atmospheric circulations ([Cha71, BK75]). These equations are obtained from the 3D primitive (hydrostatic) equations thanks to an asymptotic expansion with respect to the Rossby number. The model is also very familiar to applied mathematicians, and several studies establishing the well-posedness of the corresponding boundary-value problem have been published (see *e.g.*, [BB94, GM97, Mas00]). The primitive equations from which the traditional QG equations are obtained rely on a few well-known hypotheses, among which the so-called *traditional approximation*, which consists in neglecting the rotating terms involving $2\Omega \cos \theta$ that appear in the zonal and vertical components (1a) and (1c) of the momentum equation. This approximation has also been widely discussed in the literature (see [Eck60] and the correspondence in [Phi66, Ver68, Phi68, Wan70]). One may choose not to do this approximation, as in [JMA97, WHRS05] : in this case, the primitive equations are called quasi-hydrostatic (see [LPR10] for a mathematical study).

Since the QG equations are obtained in the zero-limit of the Rossby number (large rotational effects), one could think of retaining all the rotating terms in the original equations, before performing the asymptotic analysis. This is the main purpose of this short paper. We will see in (20) that the modified QG equations, that we will call quasi-hydrostatic quasi-geostrophic (QHQG) equations, are very similar to the traditional QG ones, except that they raise a new vertical direction (denoted Z hereafter), which differ from the traditional vertical direction z . The tilt between z and Z is proportional to the nondimensional parameter λ introduced in (3), which measures the ratio between traditional and nontraditional Coriolis terms. Experimental and numerical evidences of this tilted vertical direction can be found in [She04, WB06, GZMvH08, SKR02] : hereafter we provide a first mathematical illustration.

The paper is organized as follows : in Section 2 we perform the rigorous derivation of the QHQG equations, starting from the QH primitive equations. Then, we present in Section 3 some simple physical properties of the QHQG model.

2 Derivation of the QHQG model

In this section we present the derivation of the quasi-hydrostatic quasi-geostrophic (QHQG) equations. The derivation follows classical principles (as in [BB94]) : scaling, asymptotic expansion with respect to a small parameter, equations satisfied at order zero and one. Here, the small parameter (denoted ε in the sequel) is the Rossby number, so that we underline the effect of rotating terms. In order to account for the complete Coriolis force (see *e.g.*, [LPR10] and references therein), we retain all the rotating terms in the original equations, including the terms that are usually neglected in the so-called *traditional approximation*¹ : see the discussion in [Phi66, Ver68, Phi68, Wan70]).

2.1 Scaling Parameters and Scaled Equations

We consider a three-dimensional domain with periodic boundary conditions in the horizontal directions, rigid lid and flat bottom in the vertical. The governing non-hydrostatic equations (sometimes called the incompressible Boussinesq equations), including the complete Coriolis force, read :

$$\frac{Du}{Dt} - fv + f^*w = -\frac{\partial\varphi}{\partial x}, \quad (1a)$$

$$\frac{Dv}{Dt} + fu = -\frac{\partial\varphi}{\partial y}, \quad (1b)$$

$$\frac{Dw}{Dt} - f^*u + \frac{g\rho}{\rho_0} = -\frac{\partial\varphi}{\partial z}, \quad (1c)$$

$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} = 0, \quad (1d)$$

$$\frac{D\rho}{Dt} = 0. \quad (1e)$$

Here (u, v, w) and ρ are respectively the three-dimensional velocity and density of the fluid, and φ is the renormalized pressure, $\varphi = p/\rho_0$. The scalars $f = 2\Omega \sin(\theta)$ and $f^* = 2\Omega \cos(\theta)$ are the Coriolis parameters where Ω stands for the angular velocity of the earth and θ is the latitude ; g is the universal gravity constant, and ρ_0 stands for the averaged density of the fluid.

Equations (1a) to (1c) describe the conservation of momentum, where D/Dt is the material derivative $D/Dt = \partial/\partial t + u\partial_x + v\partial_y + w\partial_z$, and Equation (1d) corresponds to the conservation of mass. Finally, Equation (1e) describes the advection of tracers (here the density ρ). The density and the pressure may be classically decomposed as

$$\rho(x, y, z, t) = \bar{\rho}(z) + \rho(x, y, z, t) \text{ and } \varphi(x, y, z, t) = \bar{\varphi}(z) + \phi(x, y, z, t), \quad (2)$$

where $\bar{\rho}$ and $\bar{\varphi}$ are the (known) background density and potential, depending only on the vertical variable. We also denote by $N^2(z) = -\bar{\rho}'(z)$ the buoyancy

1. The denomination of *traditional approximation* was introduced by Carl Eckart in his book [Eck60].

frequency, assuming that $\bar{\rho}'(z)$ is bounded away from zero.

Before going further in the derivation of the corresponding QG model, let us insist on the fact that we keep in equations (1a) and (1c) the Coriolis terms f^*w and f^*u . This is actually the novelty of this study and will finally lead to a slightly modified QG model (see (19)). We think that it is a relevant modification, since the QG approximation aims at underlying the earth's rotation effects : one should thus include every rotation term in the primitive equations prior to an asymptotic expansion with respect to the Rossby number. In the context of the β -plane approximation, we have², with θ_0 the average latitude :

$$f = f_0(1 + \varepsilon\beta_0 y) = 2\Omega \sin \theta_0(1 + \varepsilon\beta_0 y), \quad f^* = f_0^* = 2\Omega \cos \theta_0.$$

We now introduce the following dimensionless variables, as it is classically done in QG modeling :

$$\begin{aligned} (x, y) &= L(x', y'), & z &= H z', & t &= \frac{L}{U} t', \\ u &= U u', & v &= U v', & w &= \frac{UH}{L} w', \\ \bar{\rho} &= P \bar{\rho}', & \rho &= \frac{\varrho_0 f_0 UL}{gH} \rho', & \phi &= f_0 UL \phi'. \end{aligned}$$

The Rossby number $\varepsilon = U/f_0 L$ is the fundamental ordering parameter in the following asymptotic expansion. A secondary ordering parameter is the scale ratio of ρ to ϱ : this ratio is assumed to be ε , that is, we assume :

$$\frac{\varrho_0 f_0 UL}{gH} = P\varepsilon.$$

Finally, the density may be expressed in terms of nondimensional quantities :

$$\varrho = P \left(\bar{\rho}'(z) + \varepsilon \rho' \right).$$

Another usual nondimensional number is the aspect ratio $\delta = H/L$; δ also appears in the ratio between the two Coriolis terms in the zonal momentum equation (1a), which scales as

$$\lambda = \delta \cot[\theta_0]. \quad (3)$$

When we considered the scaling numbers introduced above, we have implicitly assumed that the leading term at the left-hand-side of Equation (1a) was $f v$, which means that $\delta \cot[\theta_0]$ should not be too large :

$$\lambda \lesssim 1. \quad (4)$$

Fortunately, because the aspect ratio $\delta = H/L$ is rather small in large ocean models, the condition (4) is easily satisfied. However, the objective of the present

². For simplicity we do not consider any Taylor expansion of f^* with respect to θ . In fact we only keep the 0^{th} order term, as suggested in [GZMvH08].

work is to draw the reader's attention on the fact that λ is not necessarily small, and that it may have some physical repercussions (see Section 3).

We end this section with the non-hydrostatic scaled equations (we naturally drop the primes) :

$$\varepsilon \frac{Du}{Dt} - (1 + \varepsilon\beta_0 y)v + \lambda w = -\frac{\partial\phi}{\partial x}, \quad (5a)$$

$$\varepsilon \frac{Dv}{Dt} + (1 + \varepsilon\beta_0 y)u = -\frac{\partial\phi}{\partial y}, \quad (5b)$$

$$\varepsilon\delta^2 \frac{Dw}{Dt} - \lambda u + \rho = -\frac{\partial\phi}{\partial z}, \quad (5c)$$

$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} = 0, \quad (5d)$$

$$\varepsilon \frac{D\rho}{Dt} + w\bar{\rho}_z = 0, \quad (5e)$$

where we recall that ε is the Rossby number (meant to go to zero), $\delta = H/L$ is the domain aspect ratio, and $\lambda = \delta \cot(\theta_0)$.

Remark 1 *Actually, the term $\varepsilon\delta^2 Dw/Dt$ could be set to zero in (5c) above with no modification in the sequel : indeed the reader will see below that $w^{(0)} = 0$, hence $\varepsilon\delta^2 Dw/Dt = O(\varepsilon^2)$ can be neglected prior to the QG approximation. The new model is thus called QHQG since the differences between the new QG model and the traditional one rely only on the terms related to the Coriolis force.*

2.2 Geostrophic Balance

We now consider an asymptotic expansion of all variables with respect to the Rossby number : for every unknown function ξ , we write the formal asymptotic expansion

$$\xi = \xi^{(0)} + \varepsilon\xi^{(1)} + \varepsilon^2\xi^{(2)} + \dots$$

where $(\xi^{(j)})_{j \geq 0}$ behave as $O(1)$ as ε goes to zero. Equations (5a)-(5c) give, keeping only the order zero terms in ε :

$$-v^{(0)} + \lambda w^{(0)} = -\phi_x^{(0)}, \quad (6)$$

$$u^{(0)} = -\phi_y^{(0)}, \quad (7)$$

$$-\lambda u^{(0)} + \rho^{(0)} = -\phi_z^{(0)}. \quad (8)$$

The incompressibility condition reads $w_z^{(0)} = -u_x^{(0)} - v_y^{(0)}$ and this *traditionally* leads to $w^{(0)} = 0$, thanks to equations (6), (7) and boundary conditions on w (see [BB94]). Here, the incompressibility condition does not provide $w_z^{(0)} = 0$, but we have, denoting $\partial_Z = \partial_z + \lambda\partial_y$:

$$\begin{aligned} w_Z^{(0)} &= w_z^{(0)} + \lambda w_y^{(0)} \\ &= -u_x^{(0)} - v_y^{(0)} + \lambda w_y^{(0)} \\ &= \text{curl}(\phi_y, \phi_x) \\ w_Z^{(0)} &= 0. \end{aligned} \quad (9)$$

Thanks to (9) and to homogeneous boundary conditions on $w^{(0)}$, we finally obtain³ that $w^{(0)} = 0$.

The geostrophic equations read :

$$-v^{(0)} = -\phi_x^{(0)}, \quad (10a)$$

$$u^{(0)} = -\phi_y^{(0)}, \quad (10b)$$

$$\rho^{(0)} = -\phi_z^{(0)} - \lambda\phi_y^{(0)} = -\phi_Z^0, \quad (10c)$$

$$w^{(0)} = 0. \quad (10d)$$

2.3 Quasi-Geostrophic Equations

Now we need the first order equations in order to determine the evolution of $\phi^{(0)}$. We denote by d_g the zero-order material derivative :

$$d_g = \partial_t + u^{(0)}\partial_x + v^{(0)}\partial_y.$$

The first order equations are :

$$d_g u^{(0)} - \beta_0 y v^{(0)} - v^{(1)} + \lambda w^{(1)} = -\phi_x^{(1)}, \quad (11a)$$

$$d_g v^{(0)} + \beta_0 y u^{(0)} + u^{(1)} = -\phi_y^{(1)}, \quad (11b)$$

$$-\lambda u^{(1)} + \rho^{(1)} = -\phi_z^{(1)}, \quad (11c)$$

$$u_x^{(1)} + v_y^{(1)} + w_z^{(1)} = 0, \quad (11d)$$

$$d_g \rho^{(0)} + w^{(1)} \bar{\rho}_z = 0. \quad (11e)$$

We now take the curl of equations (11a)-(11b) to obtain, thanks to Equation (11d)

$$d_g(v_x^{(0)} - u_y^{(0)}) - w_z^{(1)} - \lambda w_y^{(1)} + \beta_0 v^{(0)} = 0. \quad (12)$$

We notice, as for the traditional QG equations, that $\beta_0 v^{(0)} = d_g(\beta_0 y)$. We thus try to express $-w_z^{(1)} - \lambda w_y^{(1)} = w_Z^1$ as $d_g(\Gamma)$ where Γ is a function to be defined. To this aim, we will extensively make use of Equation (11e) that we reformulate :

$$w^{(1)} = N^{-2} d_g \rho^{(0)} = d_g(N^{-2} \rho^{(0)}). \quad (13)$$

Given (13), we may compute the required quantity

$$w_z^{(1)} + \lambda w_y^{(1)} = \left(d_g(N^{-2} \rho^{(0)}) \right)_z + \lambda \left(d_g(N^{-2} \rho^{(0)}) \right)_y. \quad (14)$$

We remark that for any function ξ and any variable $*$ we have the identity

$$\left(d_g(\xi) \right)_* = d_g(\xi_*) + u_*^{(0)} \partial_x \xi + v_*^{(0)} \partial_y \xi,$$

so that we can write

$$w_z^{(1)} + \lambda w_y^{(1)} = d_g \left((N^{-2} \rho^{(0)})_z \right) + \lambda d_g \left((N^{-2} \rho^{(0)})_y \right) + R, \quad (15)$$

3. Alternatively, we have equation (5e) which (written to the order zero and since $\bar{\rho}(z)$ never vanishes) leads to $w^{(0)} = 0$.

where the remainder R , according to the remark above, is

$$\begin{aligned} R &= u_z^{(0)}(N^{-2}\rho^{(0)})_x + v_z^{(0)}(N^{-2}\rho^{(0)})_y + \lambda u_y^{(0)}(N^{-2}\rho^{(0)})_x + \lambda v_y^{(0)}(N^{-2}\rho^{(0)})_y \\ &= (N^{-2}\rho^{(0)})_x (u_z^{(0)} + \lambda u_y^{(0)}) + (N^{-2}\rho^{(0)})_y (v_z^{(0)} + \lambda v_y^{(0)}). \end{aligned} \quad (16)$$

Using (10a)-(10c) again, we have

$$R = N^{-2}\rho_x^{(0)}\rho_y^{(0)} - N^{-2}\rho_y^{(0)}\rho_x^{(0)} = 0,$$

which simplifies Equation (15) as follows :

$$w_Z^{(1)} = w_z^{(1)} + \lambda w_y^{(1)} = d_g \left((N^{-2}\rho^{(0)})_z + \lambda (N^{-2}\rho^{(0)})_y \right). \quad (17)$$

Back to Equation (12), we obtain the quasi-hydrostatic quasi-geostrophic equation :

$$d_g \left(v_x^{(0)} - u_y^{(0)} - (N^{-2}\rho^{(0)})_z - \lambda (N^{-2}\rho^{(0)})_y + \beta_0 y \right) = 0. \quad (18)$$

The potential vorticity

$$v_x^{(0)} - u_y^{(0)} - (N^{-2}\rho^{(0)})_z - \lambda (N^{-2}\rho^{(0)})_y + \beta_0 y$$

is thus conserved along material paths.

Let us now rewrite Equation (18), expressing everything in terms of $\phi^{(0)}$. We have

$$\left(\partial_t - \phi_y^{(0)} \partial_x + \phi_x^{(0)} \partial_y \right) \left(\Delta \phi^{(0)} + N^{-2} (\partial_z + \lambda \partial_y)^2 \phi^{(0)} + N_z^{-2} (\partial_z + \lambda \partial_y) \phi^{(0)} + \beta_0 y \right) = 0, \quad (19)$$

where Δ is horizontal Laplacian operator. We could write this more compactly with $\partial_Z = \partial_z + \lambda \partial_y$,

$$\left(\partial_t - \phi_y^{(0)} \partial_x + \phi_x^{(0)} \partial_y \right) \left(\Delta \phi^{(0)} + (N^{-2} \phi_Z^{(0)})_Z + \beta_0 y \right) = 0. \quad (20)$$

One can thus easily recognize the traditional QG equation (see Equation (2.23) in [BB94]), except that the differential operator ∂_z is replaced by $\partial_Z = \partial_z + \lambda \partial_y$. We recall here that $\lambda = \delta \cot \theta_0$ is proportional to the domain aspect ratio. In particular, we recover the traditional QG equation when setting $\delta = 0$ in equations (18), (19) or (20).

3 Simple physical properties of the QH-QG model

We reformulate the QHQG equation as follows :

$$D[q + \beta y] = 0, \quad (21)$$

$$\left(\Delta + \partial_Z \left[\frac{1}{N^2} \partial_Z \right] \right) \phi = q, \quad (22)$$

where $D = \partial_t + J[\phi, \]$, $\Delta = \partial_x^2 + \partial_y^2$, $J[a, b] = a_x b_y - a_y b_x$, $\partial_Z = \partial_z + \lambda \partial_y$, and $\lambda = \frac{H}{L} \cot \theta_0$.

3.1 Coordinate transformation

We define a coordinate transform from (x, y, z, t) to (X, Y, Z, T) by

$$X = x, \quad Y = y - \lambda z, \quad Z = z, \quad T = t. \quad (23)$$

The transformed QHQG system is isomorphic to (22) because $D = \partial_T + J[\phi, \cdot]$, $\Delta = \partial_X^2 + \partial_Y^2$, and $J[a, b] = a_X b_Y - a_Y b_X$ have identical functional forms in the transformed coordinates; $N^2(z) = N^2(Z)$; and $\beta y = \beta(Y + \lambda Z)$ can be replaced by βY because $D[\lambda Z] = 0$. Thus, all QG solutions $\phi(\mathbf{x}, t)$ are also QHQG solutions $\phi(\mathbf{X}, T)$ if the initial and boundary conditions are consistent. Initial conditions are equivalently specified at $t = 0$ or $T = 0$. Planar vertical boundary surfaces at $z = z_0$ are also planar at $Z = z_0$. The same is true in x and X . Only in y is a planar surface in y no longer planar in Y ; therefore, the simple equivalence is for solutions that have an unbounded y domain, or are y -periodic, or decay away in y before the boundary.

3.2 Separable solutions and vertical modes

QHQG has vertically separable solutions when linearized by neglecting $J[\phi, q]$, even for general $N(z)$, $\phi = \Phi(X, Y, T)F(Z)$. (QG has an analogous property.) Such QHQG solutions have a vertically upward phase tilt in the (y, z) plane with a slope of $dz/dy = \lambda^{-1}$ relative to the z axis. This direction is aligned with the full rotation vector. Many phenomena influenced by the non-traditional approximation (*e.g.*, convection and centrifugal instability as reviewed in [GZMvH08]) are known to exhibit this type of phase tilt. If we assume constant density at the vertical boundaries, F is determined by the 1D eigenvalue problem,

$$(N^{-2}F_Z)_Z + \mathcal{R}^{-2}F = 0, \quad F_Z = 0 \text{ at } Z = Z_0, Z_1. \quad (24)$$

The eigenvalue \mathcal{R} is the horizontal radius of deformation. With N constant, the eigenmodes F are cosine functions in Z .

3.3 Thermal wind balance

The geostrophic equations (10a)-(10c) have a steady solution for a zonal flow. This can be expressed as $u = U(Y, Z)$, $\varrho = -\int^Z N^2(Z') dZ' + B(Y, Z)$ with $U_Z = -B_Y$, which is thermal wind balance.

3.4 Rossby wave modes

One class of simple solutions is Rossby wave modes. They satisfy a linearized PDE usually justified by an assumption of small amplitude flow,

$$\frac{\partial q}{\partial t} + \beta \frac{\partial \phi}{\partial x} = 0. \quad (25)$$

With constant N in either a vertically bounded or unbounded domain, eigenmodes are

$$\phi \propto e^{i(kx+ly+mz-\omega t)}, \quad (26)$$

with a dispersion relation,

$$\omega = \frac{-\beta k}{K^2}, \quad K^2 = k^2 + l^2 + \frac{1}{N^2}(m + \lambda l)^2. \quad (27)$$

We can equivalently write this as

$$\phi \propto e^{i(kX+lY+MZ-\omega T)}, \quad (28)$$

with $M = m + \lambda l$.

Thus, $\lambda \neq 0$ implies lower frequency and slower phase speed compared to QG Rossby waves with the same wavenumber (k, l, m) . We also consider the group velocity for wave energy propagation, $c_g = \partial_k \omega$. Its components are

$$\begin{aligned} \partial_k \omega &= \frac{\beta(2k^2 - K^2)}{K^4}, \\ \partial_l \omega &= \frac{2\beta k(l + \lambda(m + \lambda l)/N^2)}{K^4}, \\ \partial_m \omega &= \frac{2\beta k(m + \lambda l)/N^2}{K^4}. \end{aligned} \quad (29)$$

This usually has a more westward zonal propagation due to $\lambda \neq 0$, and its (y, z) propagation is altered as well. These effects also occur for the barotropic mode ($m = 0$) where the vertical propagation is nonzero because of the QH correction.

3.5 Vortex solutions

A simple vortex solution is a nonlinear stationary state when $\beta = 0$ and N is constant. In QG this occurs for any axisymmetric profile, $\phi(r, z)$, where $r = \sqrt{x^2 + y^2}$ is the radial coordinate, which is sufficient to make $J[\phi, q] = 0$ even when the arguments have large amplitude. Physical interest usually lies in profiles that are localized in r , *e.g.*, a monopole with $\phi(r)$ decaying away from a central extremum. In QHQG nonlinear stationary solutions exist for any profile $\phi(R, Z)$, where $R^2 = X^2 + Y^2$. Thus, QH vortices are meridionally tilted rather than vertically aligned. This tilted structure was previously proposed for Meddy vortices by [SS03].

3.6 Fourier-space estimates for non-traditional effects

Assuming a wavenumber-space characterization of the solution, as commonly done for turbulent flows, in terms of vertical and horizontal wavenumbers, kv and kh , or equivalently local values of H and L , we can ask when the λ value is not small.

In QG the common view (sometimes called Charney's stretched isotropy) is that the Burger number, $Bu = NH/2\Omega L = kh/(2\Omega kv/N)$, is order one while $2\Omega/N$ is small. If $H/L \sim 2\Omega/N \ll 1$, then $\lambda \sim (2\Omega/N) \cot \theta_0$, which will be small

except when θ_0 is very close to the Equator.

Alternatively, if we consider flow patterns with $2\Omega kv/Nkh \sim r \ll 1$, then $\lambda \sim (2\Omega/Nr) \cot \theta_0$. For small enough r , λ need not be small, and the QH correction to QG will be important. This happens when the aspect ratio H/L is large, the stratification is weak, and/or θ_0 is small. Large aspect ratio can be described from the QG perspective as atypically “tall” flows.

4 Summary

The quasi-hydrostatic equations have been used to derive a new quasi-geostrophic model for the simulation of geophysical fluid dynamics. The new QHQG model is derived in Section 2, in the same spirit than the usual derivation of the traditional QG model, such as in [BB94]. The resulting QHQG model (20) is slightly different from the traditional one, in the sense that the traditional operator ∂z is changed in $\partial Z = \partial z + \lambda \partial y$ where λ measures the effect of nontraditional terms of the Coriolis force, $\lambda = (H/L) \cot \theta_0$.

In Section 3, we discuss some physical properties of the QHQG model, and point the similarities / differences with the traditional QG model. This discussion can be seen as a first theoretical illustration of well-known phenomena that have already been mentioned in the literature.

Références

- [BB94] A. J. Bourgeois and J. T. Beale. Validity of the quasigeostrophic model for large-scale flow in the atmosphere and ocean. *SIAM Journal on Mathematical Analysis*, 25(4) :1023–1068, 1994.
- [BK75] F. J. Bretherton and M. J. Karweit. Mid-ocean mesoscale modeling. In *Numerical Models of Ocean Circulation*, pages 237–. Ocean Affairs Board, National Research Council, National Academy of Sciences, Washington, DC, 1975. [Quasi-geostrophic layered model with doubly-periodic horizontal boundary conditions and double Fourier pseudospectral algorithm.].
- [Cha71] J. G. Charney. Geostrophic turbulence. *Journal of the Atmospheric Sciences*, 28(6) :1087–1095, 1971.
- [Eck60] C. Eckart. *Hydrodynamics of oceans and atmospheres*. Pergamon Press, New York, 1960.
- [GM97] E. Grenier and N. Masmoudi. Ekman layers of rotating fluids, the case of well prepared initial data. *Comm. Partial Diff. Equations*, 22 :953–975, 1997.
- [GZMvH08] T. Gerkema, J. T. F. Zimmerman, L. R. M. Maas, and H. van Haren. Geophysical and astrophysical fluid dynamics beyond the traditional approximation. *Rev. Geophys.*, 46(2), 05 2008.

- [JMA97] L. Perelman J. Marshall, C. Hill and A. Adcroft. Hydrostatic, quasi-hydrostatic, and nonhydrostatic ocean modeling. *J. Geophys. Research*, 102 :5733–5752, 1997.
- [LPR10] C. Lucas, M. Petcu, and A. Rousseau. Quasi-hydrostatic primitive equations for ocean global circulation models. *Chin. Ann. Math.*, 31B(6) :1–20, 2010.
- [Mas00] N. Masmoudi. Ekman layers of rotating fluids : The case of general initial data. *Comm. Pure Appl. Math.*, 53(4) :432–483, 2000.
- [Phi66] N. A. Phillips. The equations of motion for a shallow rotating atmosphere and the "traditional approximation". *Journal of the atmospheric sciences*, 23 :626–628, 1966.
- [Phi68] N. A. Phillips. Reply (to George Veronis). *Journal of the atmospheric sciences*, 25 :1155–1157, 1968.
- [She04] V. A. Sheremet. Laboratory experiments with tilted convective plumes on a centrifuge : a finite angle between the buoyancy force and the axis of rotation. *Journal of Fluid Mechanics*, 506 :217–244, 2004.
- [SKR02] F. Straneo, M. Kawase, and S. C. Riser. Idealized models of slantwise convection in a baroclinic flow. *Journal of Physical Oceanography*, 32(2) :558–572, 2002.
- [SS03] I. P. Semenova and L. N. Slezkin. Dynamically equilibrium shape of intrusive vortex formations in the ocean. *Fluid Dynamics*, 38 :663–669, 2003. 10.1023/B :FLUI.0000007828.46531.6b.
- [Ver68] G. Veronis. Comments on phillips proposed simplification of the equations of motion for a shallow rotating atmosphere. *Journal of the atmospheric sciences*, 25 :1154–1155, 1968.
- [Wan70] R. K. Wangsness. Comments on "the equations of motion for a shallow rotating atmosphere and the 'traditionnal approximation' ". *Journal of the atmospheric sciences*, 27 :504–506, 1970.
- [WB06] A. Wirth and B. Barnier. Tilted convective plumes in numerical experiments. *Ocean Modelling*, 12(1-2) :101–111, 2006.
- [WHRS05] A. A. White, B. J. Hoskins, I. Roulstone, and A. Staniforth. Consistent approximate models of the global atmosphere : shallow, deep, hydrostatic, quasi-hydrostatic and non-hydrostatic. *Q. J. R. Meteorol. Soc.*, 131 :2081–2107, 2005.



Centre de recherche INRIA Grenoble – Rhône-Alpes
655, avenue de l'Europe - 38334 Montbonnot Saint-Ismier (France)

Centre de recherche INRIA Bordeaux – Sud Ouest : Domaine Universitaire - 351, cours de la Libération - 33405 Talence Cedex
Centre de recherche INRIA Lille – Nord Europe : Parc Scientifique de la Haute Borne - 40, avenue Halley - 59650 Villeneuve d'Ascq
Centre de recherche INRIA Nancy – Grand Est : LORIA, Technopôle de Nancy-Brabois - Campus scientifique
615, rue du Jardin Botanique - BP 101 - 54602 Villers-lès-Nancy Cedex
Centre de recherche INRIA Paris – Rocquencourt : Domaine de Voluceau - Rocquencourt - BP 105 - 78153 Le Chesnay Cedex
Centre de recherche INRIA Rennes – Bretagne Atlantique : IRISA, Campus universitaire de Beaulieu - 35042 Rennes Cedex
Centre de recherche INRIA Saclay – Île-de-France : Parc Orsay Université - ZAC des Vignes : 4, rue Jacques Monod - 91893 Orsay Cedex
Centre de recherche INRIA Sophia Antipolis – Méditerranée : 2004, route des Lucioles - BP 93 - 06902 Sophia Antipolis Cedex

Éditeur
INRIA - Domaine de Voluceau - Rocquencourt, BP 105 - 78153 Le Chesnay Cedex (France)
<http://www.inria.fr>
ISSN 0249-6399