



HAL
open science

Hypergeometric Expressions for Generating Functions of Walks with Small Steps in the Quarter Plane

Alin Bostan, Frédéric Chyzak, Mark van Hoeij, Manuel Kauers, Lucien Pech

► **To cite this version:**

Alin Bostan, Frédéric Chyzak, Mark van Hoeij, Manuel Kauers, Lucien Pech. Hypergeometric Expressions for Generating Functions of Walks with Small Steps in the Quarter Plane. 2016. hal-01332175v1

HAL Id: hal-01332175

<https://inria.hal.science/hal-01332175v1>

Preprint submitted on 15 Jun 2016 (v1), last revised 15 Nov 2016 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Hypergeometric Expressions for Generating Functions of Walks with Small Steps in the Quarter Plane

Alin Bostan, Frédéric Chyzak, Mark van Hoeij*, Manuel Kauers† and Lucien Pech

Abstract

We study nearest-neighbors walks on the two-dimensional square lattice, that is, models of walks on \mathbb{Z}^2 defined by a fixed step set that is a subset of the non-zero vectors with coordinates 0, 1 or -1 . We concern ourselves with the enumeration of such walks starting at the origin and constrained to remain in the quarter plane \mathbb{N}^2 , counted by their length and by the position of their ending point. Bousquet-Mélou and Mishna [Contemp. Math., pp. 1–39, Amer. Math. Soc., 2010] identified 19 models of walks that possess a D -finite generating function; linear differential equations have then been guessed in these cases by Bostan and Kauers [FPSAC 2009, Discrete Math. Theor. Comput. Sci. Proc., pp. 201–215, 2009]. We give here the first proof that these equations are indeed satisfied by the corresponding generating functions. As a first corollary, we prove that all these 19 generating functions can be expressed in terms of Gauss’ hypergeometric functions that are intimately related to elliptic integrals. As a second corollary, we show that all the 19 generating functions are transcendental, and that among their 19×4 combinatorially meaningful specializations only four are algebraic functions.

Keywords: Walks in the quarter plane; Generating functions; Hypergeometric functions

MSC2010: Primary 05A15, 14N10, 33F10, 68W30; Secondary 33C05, 97N80, 11J89.

1 Introduction

Context. An important problem in enumerative combinatorics is the study of lattice walks in restricted lattices. Many efforts have been deployed in recent years for classifying them, see e.g. the surveys [29] and [32] and the references therein. The generating functions of lattice walks are not only intriguing for combinatorial reasons, but also from the perspective of computer algebra. For combinatorial reasons they are interesting because, depending on the choice of admissible steps, the generating functions may have quite different algebraic and analytic properties. For computational reasons they are interesting because their descriptions (whether by a polynomial or by a linear differential equation, as we will see below) are sometimes so large in size that it becomes difficult to handle them with a reasonable efficiency.

In the present article, we consider *small step walks restricted to the quarter plane*, defined as follows. Let $\mathcal{S} \neq \emptyset$ be a fixed subset of $\{-1, 0, 1\}^2 \setminus \{(0, 0)\}$, which will contain all steps allowed in the walks. An \mathcal{S} -walk of length n starts at the origin $(0, 0)$ and consists of n consecutive steps, where a step from a point A to a point B is admitted if $B - A \in \mathcal{S}$, and both A and B belong to the quarter plane \mathbb{N}^2 . These walks are called *restricted* to the quarter plane because they are not allowed to step out of it, and *with small steps* because a single step changes the position by no more than 1 in each coordinate. As an example, for $\mathcal{S} = \{(1, 1), (-1, 0), (0, -1)\}$ (*Kreweras walks*), a possible walk of length six is

$$(0, 0) \rightarrow (1, 1) \rightarrow (2, 2) \rightarrow (1, 2) \rightarrow (2, 3) \rightarrow (2, 2) \rightarrow (2, 1).$$

A brute-force enumeration with rejection shows that, altogether, there are 125 different walks of length six for this particular step set. In the general case of an arbitrary step set \mathcal{S} , with q_n denoting the number of different \mathcal{S} -walks of length n , we are interested in the generating function $Q(t) := \sum_{n=0}^{\infty} q_n t^n \in \mathbb{Q}[[t]]$.

The generating function $Q(t)$ corresponding to the example step set \mathcal{S} above is *algebraic* [10, 22, 33], i.e., it satisfies a polynomial equation $P(t, Q(t)) = 0$ for some $P \in \mathbb{Q}[t, T] \setminus \{0\}$. But this is not the case for all other step sets. Still, among those step sets that induce a *transcendental* (i.e., non-algebraic) generating function $Q(t)$, some have a $Q(t)$ that is *D-finite*, i.e., that satisfies a linear differential equation with polynomial coefficients. The step set $\mathcal{S} = \{(1, 1), (-1, 1), (0, -1)\}$ is an example for this case [9, 12].

*Supported by NSF grant 1319547.

†Supported by FWF grants Y464-N18, F5004 and W1214.

Finally, there are also step sets whose corresponding generating function is not even D-finite; Mishna and Rechnitzer [40] proved that this is the case for example when $\mathcal{S} = \{(-1, 1), (1, 1), (1, -1)\}$.

More generally, for a fixed step set \mathcal{S} , one is interested in the study of the trivariate power series

$$Q(x, y; t) = \sum_{n=0}^{\infty} \sum_{i,j=0}^{\infty} q_{i,j;n} x^i y^j t^n,$$

where $q_{i,j;n}$ denotes the number of \mathcal{S} -walks of length n starting at $(0, 0)$ and ending at (i, j) . The power series $Q(x, y; t)$ is called the *complete generating function* for \mathcal{S} -walks. Note that the counting series $Q(t)$ introduced before is nothing but the specialization $Q(1, 1; t)$ of $Q(x, y; t)$ at $(x, y) = (1, 1)$. Other combinatorially meaningful specializations are $Q(0, 0; t)$, the generating function of \mathcal{S} -walks returning to the origin (also called *excursions*), $Q(1, 0; t)$, the generating function of \mathcal{S} -walks ending on the horizontal axis, and $Q(0, 1; t)$, the generating function of \mathcal{S} -walks ending on the vertical axis.

Bousquet-Mélou and Mishna [11] have undertaken a systematic classification of the 256 step sets $\mathcal{S} \subseteq \{-1, 0, 1\}^2 \setminus \{(0, 0)\}$, from the viewpoint of structural properties of the generating function $Q(x, y; t)$. Again, the concerned properties are algebraicity and D-finiteness, yet applied to a multivariate setting¹. They found out that there are 79 inherently different and nontrivial models to consider, of which they recognized 22 models for which $Q(x, y; t)$ is D-finite; a 23rd model, of the so-called *Gessel walks*, was proved to be D-finite (and even algebraic) by Bostan and Kauers [7]. These 23 models share the feature that a certain group associated to \mathcal{S} is finite. In the remaining 56 cases, it has been proved that the groups are infinite [11], and that the complete generating functions $Q(x, y; t)$ are not D-finite [8, 34, 38, 40].

Of the 23 D-finite generating functions, 4 were recognized to be algebraic [7, 11]: one corresponds to the Kreweras model $\mathcal{S} = \{(1, 1), (-1, 0), (0, -1)\}$, one to its reverse $\mathcal{S}^{\text{rev}} = \{(-1, -1), (1, 0), (0, 1)\}$, one to $\mathcal{S} \cup \mathcal{S}^{\text{rev}}$, and one to Gessel's. All the other 19 generating functions were proved to be transcendental [17], although some of their specializations are algebraic, e.g., for $\mathcal{S} = \{(-1, 0), (0, 1), (1, -1)\}$ [11]. These 19 models form the main object of this article; they are depicted in the third column of Table 1.

The proofs of D-finiteness given by Bousquet-Mélou and Mishna are implicit (i.e., qualitative): the existence of differential equations satisfied by the generating functions $Q(x, y; t)$ was shown without obtaining the differential equations explicitly. On the other hand, Bostan and Kauers [5, 6] provided explicit differential equations for (specializations of) the 23 D-finite generating functions, but these equations were determined only experimentally and, in most of the transcendental cases, they still lack formal proofs (in the 4 algebraic cases, differential equations are easily proved, starting from algebraic equations). Therefore, the following problems were left unsolved by Bousquet-Mélou and Mishna:

- (i) prove differential equations satisfied by $Q(0, 0; t)$ and $Q(t) = Q(1, 1; t)$ [11, §7.5];
- (ii) find closed form expressions for them [11, §7.3];
- (iii) classify models with algebraic generating function $Q(t)$ [11, §7.3].

Contributions. The original goal of the present paper was to answer question (i). In this regard, we rigorously prove the differential equations for $Q(0, 0; t)$ and $Q(1, 1; t)$ guessed by Bostan and Kauers [5, 6] for the 19 models with D-finite transcendental complete generating function. We actually do more, that is we also find and prove differential equations for $Q(x, 0; t)$ and for $Q(0, y; t)$, that specialize to equations for $Q(1, 0; t)$ and $Q(0, 1; t)$.

By solving these differential equations, we answer (ii) in the following sense: for all the 19 models mentioned above we uniformly find closed form expressions for $Q(x, y; t)$ in terms of Gauss' hypergeometric series ${}_2F_1$ with parameters $a, b, c \in \mathbb{Q}$, with $-c \notin \mathbb{N}$, defined by

$${}_2F_1 \left(\begin{matrix} a & b \\ c \end{matrix} \middle| t \right) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n} \frac{t^n}{n!}, \quad (1)$$

where $(x)_n$ denotes the Pochhammer symbol $(x)_n = x(x+1) \cdots (x+n-1)$ for $n \in \mathbb{N}$.

More precisely, we obtain the following structure result, that has been conjectured in [6, §3.2]. Note that a similar expression also appears in a related combinatorial context [4] for rook paths on a three-dimensional chessboard.

¹A trivariate power series $S \in \mathbb{Q}[[x, y, t]]$ is *algebraic* if it is the root of a nonzero polynomial $P \in \mathbb{Q}[x, y, t, T]$. It is called *D-finite* if the set of all partial derivatives of S spans a finite-dimensional vector space over $\mathbb{Q}(x, y, t)$.

	OEIS [45]	\mathcal{S}	$N(x, y)$	$S(x, y)$
1	A005566			$y + (x + \bar{x}) + \bar{y}$
2	A018224		$(x - \bar{x})(y - \bar{y})$	$(x + \bar{x})y + (x + \bar{x})\bar{y}$
3	A151312			$(x + 1 + \bar{x})y + (x + 1 + \bar{x})\bar{y}$
4	A151331			$(x + 1 + \bar{x})y + (x + \bar{x}) + (x + 1 + \bar{x})\bar{y}$
5	A151266		$(x - \bar{x}) \left(y - \frac{1}{x + \bar{x}} \bar{y} \right)$	$(x + \bar{x})y + \bar{y}$
6	A151307			$(x + \bar{x})y + (x + \bar{x}) + \bar{y}$
7	A151291		$(x - \bar{x}) \left(y - \frac{1}{x + 1 + \bar{x}} \bar{y} \right)$	$(x + 1 + \bar{x})y + \bar{y}$
8	A151326			$(x + 1 + \bar{x})y + (x + \bar{x}) + \bar{y}$
9	A151302		$(x - \bar{x}) \left(y - \frac{x + \bar{x}}{x + 1 + \bar{x}} \bar{y} \right)$	$(x + 1 + \bar{x})y + (x + \bar{x})\bar{y}$
10	A151329			$(x + 1 + \bar{x})y + (x + \bar{x}) + (x + \bar{x})\bar{y}$
11	A151261		$(x - \bar{x})(y - (x + 1 + \bar{x})\bar{y})$	$y + (x + 1 + \bar{x})\bar{y}$
12	A151297			$y + (x + \bar{x}) + (x + 1 + \bar{x})\bar{y}$
13	A151275		$(x - \bar{x}) \left(y - \frac{x + 1 + \bar{x}}{x + \bar{x}} \bar{y} \right)$	$(x + \bar{x})y + (x + 1 + \bar{x})\bar{y}$
14	A151314			$(x + \bar{x})y + (x + \bar{x}) + (x + 1 + \bar{x})\bar{y}$
15	A151255		$(x - \bar{x})(y - (x + \bar{x})\bar{y})$	$y + (x + \bar{x})\bar{y}$
16	A151287			$y + (x + \bar{x}) + (x + \bar{x})\bar{y}$
17	A001006		$xy - \bar{x}y^2 + \bar{x}^2y - \bar{x}\bar{y} + x\bar{y}^2 - x^2\bar{y}$	$y + \bar{x} + x\bar{y}$
18	A129400			$(1 + \bar{x})y + (x + \bar{x}) + (1 + x)\bar{y}$
19	A005558		$xy - \bar{x}y^2 + \bar{x}^3y^2 - \bar{x}^3y + \bar{x}\bar{y} - x\bar{y}^2 + x^3\bar{y}^2 - x^3\bar{y}$	$\bar{x}y + (x + \bar{x}) + x\bar{y}$

Table 1: The 19 models ($\bar{x} = x^{-1}$, $\bar{y} = y^{-1}$) and the corresponding rational functions N and S in Eq. (3). Cases 1–16 are numbered as in [11, Table 1]. Cases 17 and 18 are #1 and #2 in [11, Table 2]. Case 19 is #1 in [11, Table 3]. The OEIS tags correspond to coefficient sequences of the length generating function $Q(t) = Q(1, 1; t)$.

Theorem 1. *Let \mathcal{S} be one of the 19 models of small step walks in the quarter plane (see Table 1). The complete generating function $Q(x, y; t)$ is expressible as a finite sum of iterated integrals of products of algebraic functions in x, y, t and of expressions of the form ${}_2F_1 \left(\begin{smallmatrix} a & b \\ c \end{smallmatrix} \middle| w(t) \right)$, where $c \in \mathbb{N}$ and $w(t) \in \mathbb{Q}(t)$.*

The parameters a, b, c of the occurring ${}_2F_1$'s as well as the rational functions $w(t)$ are explicitly given in Table 2. The full expressions of the generating functions $Q(0, 0; t)$, $Q(0, 1; t)$, $Q(1, 0; t)$, $Q(1, 1; t)$, $Q(x, 0; t)$, $Q(0, y; t)$ and $Q(x, y; t)$ are too large to be displayed in this paper, and are available on-line at http://specfun.inria.fr/chyzak/ssw/closed_forms.html. It turns out by inspection that the involved hypergeometric functions have a very particular form: they are intimately related to elliptic integrals, namely to the complete elliptic integrals of first and second kinds,

$$K(k) = \int_0^{\pi/2} (1 - k^2 \sin^2 \theta)^{-1/2} d\theta = \frac{\pi}{2} {}_2F_1 \left(\begin{smallmatrix} \frac{1}{2} & \frac{1}{2} \\ 1 \end{smallmatrix} \middle| k^2 \right),$$

$$E(k) = \int_0^{\pi/2} (1 - k^2 \sin^2 \theta)^{1/2} d\theta = \frac{\pi}{2} {}_2F_1 \left(\begin{smallmatrix} -\frac{1}{2} & \frac{1}{2} \\ 1 \end{smallmatrix} \middle| k^2 \right).$$

For instance, for the step set $\{(1, 1), (0, 1), (-1, 1), (-1, 0), (-1, -1), (0, -1), (1, -1), (1, 0)\}$ of the so-called *king walks* (case 4 in Table 1), we prove that

$$Q(t) = \frac{1}{t} \int_0^t \frac{1}{(1 + 4x)^3} \cdot {}_2F_1 \left(\begin{smallmatrix} \frac{3}{2} & \frac{3}{2} \\ 2 \end{smallmatrix} \middle| \frac{16x(1+x)}{(1+4x)^2} \right) dx. \quad (2)$$

See Section 2 for a detailed presentation of this example. Alternatively, an expression of $Q(t)$ in terms of elliptic integrals is

$$Q(t) = \frac{1}{t} \int_0^t \frac{1}{\pi(1+4x)^2 \sqrt{x(1+x)}} \cdot K' \left(\frac{4\sqrt{x(1+x)}}{1+4x} \right) dx.$$

S	occurring	${}_2F_1$	w	S	occurring	${}_2F_1$	w
1		${}_2F_1$	$\left(\begin{matrix} \frac{1}{2} & \frac{1}{2} \\ 1 & 1 \end{matrix} \middle w\right)$	11		${}_2F_1$	$\left(\begin{matrix} \frac{1}{2} & \frac{1}{2} \\ 1 & 1 \end{matrix} \middle w\right)$
2		${}_2F_1$	$\left(\begin{matrix} \frac{1}{2} & \frac{1}{2} \\ 1 & 1 \end{matrix} \middle w\right)$	12		${}_2F_1$	$\left(\begin{matrix} \frac{1}{4} & \frac{3}{4} \\ 1 & 1 \end{matrix} \middle w\right)$
3		${}_2F_1$	$\left(\begin{matrix} \frac{1}{4} & \frac{3}{4} \\ 1 & 1 \end{matrix} \middle w\right)$	13		${}_2F_1$	$\left(\begin{matrix} \frac{1}{4} & \frac{3}{4} \\ 1 & 1 \end{matrix} \middle w\right)$
4		${}_2F_1$	$\left(\begin{matrix} \frac{1}{2} & \frac{1}{2} \\ 1 & 1 \end{matrix} \middle w\right)$	14		${}_2F_1$	$\left(\begin{matrix} \frac{1}{4} & \frac{3}{4} \\ 1 & 1 \end{matrix} \middle w\right)$
5		${}_2F_1$	$\left(\begin{matrix} \frac{1}{4} & \frac{3}{4} \\ 1 & 1 \end{matrix} \middle w\right)$	15		${}_2F_1$	$\left(\begin{matrix} \frac{1}{4} & \frac{3}{4} \\ 1 & 1 \end{matrix} \middle w\right)$
6		${}_2F_1$	$\left(\begin{matrix} \frac{1}{4} & \frac{3}{4} \\ 1 & 1 \end{matrix} \middle w\right)$	16		${}_2F_1$	$\left(\begin{matrix} \frac{1}{4} & \frac{3}{4} \\ 1 & 1 \end{matrix} \middle w\right)$
7		${}_2F_1$	$\left(\begin{matrix} \frac{1}{2} & \frac{1}{2} \\ 1 & 1 \end{matrix} \middle w\right)$	17		${}_2F_1$	$\left(\begin{matrix} \frac{1}{3} & \frac{2}{3} \\ 1 & 1 \end{matrix} \middle w\right)$
8		${}_2F_1$	$\left(\begin{matrix} \frac{1}{4} & \frac{3}{4} \\ 1 & 1 \end{matrix} \middle w\right)$	18		${}_2F_1$	$\left(\begin{matrix} \frac{1}{3} & \frac{2}{3} \\ 1 & 1 \end{matrix} \middle w\right)$
9		${}_2F_1$	$\left(\begin{matrix} \frac{1}{4} & \frac{3}{4} \\ 1 & 1 \end{matrix} \middle w\right)$	19		${}_2F_1$	$\left(\begin{matrix} \frac{1}{2} & \frac{1}{2} \\ 1 & 1 \end{matrix} \middle w\right)$
10		${}_2F_1$	$\left(\begin{matrix} \frac{1}{4} & \frac{3}{4} \\ 1 & 1 \end{matrix} \middle w\right)$				

Table 2: Hypergeometric series occurring in explicit expressions for $Q(x, y; t)$. The ${}_2F_1$ are given up to contiguity and derivation, that is, up to integer shifts of the parameters.

The relationship to elliptic integrals appears to hold true in a far more general setting. Indeed, taking Theorem 1 as starting point, one of us (van Hoeij) has checked that for *many* (more than 100) integer sequences $(a_n)_{n \geq 0}$ in the OEIS whose generating function $A(t) = \sum_{n \geq 0} a_n t^n$ is both D-finite and convergent in a small neighborhood of $t = 0$, all second-order irreducible factors of the minimal-order linear differential operator annihilating $A(t)$ are solvable either in terms of algebraic functions, or in terms of complete elliptic integrals. This surprisingly general feature, reminiscent of Dwork's conjecture mentioned in [6, §3.2], begs for a combinatorial explanation. See also [28, Section 8] for a similar discussion.

Finally, concerning question (iii), we start from the explicit differential equations and we exhaustively classify the algebraic cases among all the specializations of the generating function $Q(x, y; t)$ at points $(x, y) \in \{(0, 0), (1, 0), (0, 1), (1, 1)\}$. As a corollary, we reprove the transcendence of $Q(x, y; t)$. More precisely, we prove:

Theorem 2. *Let S be one of the 19 models of small step walks in the quarter plane (see Table 1), with complete generating function $Q(x, y; t)$. For any $(\alpha, \beta) \in \{(0, 0), (1, 0), (0, 1), (1, 1)\}$, the power series $Q(\alpha, \beta; t)$ is transcendental, except in the following four cases:*

- Case 18 at $(\alpha, \beta) = (1, 0)$ and at $(\alpha, \beta) = (0, 1)$,
- Cases 17 and 18 at $(\alpha, \beta) = (1, 1)$.

As a consequence, the power series $Q(x, y; t)$, $Q(x, 0; t)$, and $Q(0, y; t)$ are transcendental for all the 19 models. Additionally, the generating functions of the four algebraic cases are equal to:

- $Q(1, 1) = \frac{1}{2t^2} \left(1 - t - \sqrt{(1+t)(1-3t)}\right)$ in case 17,
- $Q(1, 1) = \frac{1}{8t^2} \left(1 - 2t - \sqrt{(1+2t)(1-6t)}\right)$ in case 18,
- $Q(1, 0) = Q(0, 1) = \frac{1}{32t^3} \left((1-6t)^{3/2}(1+2t)^{1/2} - 4t^2 + 8t - 1\right)$ in case 18.

As an aside, starting from the explicit expressions in terms of hypergeometric functions, we use singularity analysis and transfer theorems that are classical in Analytic Combinatorics to get some asymptotic formulas for the n th coefficient of $Q(0, 0; t)$, $Q(1, 0; t)$, $Q(0, 1; t)$ and $Q(1, 1; t)$.

Methodology. Our proofs of Theorems 1 and 2 are computer-driven and crucially rely on the use of several modern computer algebra algorithms. The starting point is a result by Bousquet-Mélou and Mishna [11], stating that for the 19 models in Table 1 the complete generating function $Q(x, y; t)$ can be expressed as the *positive part* of a certain rational function in three variables. The notion of positive part is one of the key mathematical ingredients in what follows. In one variable, extraction of the positive part is an operator, denoted $[x^>]$, which acts on formal Laurent series by cutting away all the terms with zero or negative exponents, leaving a formal power series with no constant term as a result. For example,

$$[x^>] \frac{1}{x^2(1-x)^2} = [x^>] \sum_{n=-2}^{\infty} (n+3)x^n = \sum_{n=1}^{\infty} (n+3)x^n = \frac{x(4-3x)}{(1-x)^2}.$$

Note that interpreting the rational function as a formal Laurent series in x^{-1} instead of x would lead to a different extraction map. Indeed, for this other definition of positive-part extraction, we would have

$$[x^>] \frac{1}{x^2(1-x)^2} = [x^>] \sum_{n=0}^{\infty} n \frac{1}{x^{n+3}} = [x^>] \sum_{n=-\infty}^{-3} -(n+3)x^n = 0.$$

Things get more complicated in the multivariate setting. Using the kernel method, Bousquet-Mélou and Mishna showed in [11, Prop. 8] that the generating function $Q(x, y; t)$, can be written in the form

$$Q(x, y; t) = \frac{1}{xy} [x^>][y^>] \frac{N(x, y)}{1 - tS(x, y)} \quad (3)$$

where $N(x, y)$ and $S(x, y)$ are certain (structured) Laurent polynomials in y with coefficients that are rational functions in x . These quantities depend on \mathcal{S} and are listed in Table 1. Since there is no unique natural way of mapping rational functions in several variables to multivariate formal Laurent series, it is a priori not clear how the positive-part extraction is defined in this context. Here is the intended reading of (3): first interpret $N(x, y)/(1 - tS(x, y))$ as an element of $\mathbb{Q}(x)[y, 1/y][[t]]$, owing to particular properties of N and S (see Lemma 9 below); let $[y^>]$ act term by term, obtaining a series in $\mathbb{Q}(x)[y][[t]]$ that can be shown to actually belong to $\mathbb{Q}[x, 1/x][y][[t]]$ for all cases in Table 1; then let $[x^>]$ act term by term, finally obtaining an element of $\mathbb{Q}[x][y][[t]]$. In this reading, the composition $[x^>][y^>]$ of positive-part operators is only applied to Laurent polynomials, for which it is certainly well-defined, in a unique way.

As pointed out by Bousquet-Mélou and Mishna, Equation (3) already implies the D-finiteness of $Q(x, y; t)$, since positive parts can be encoded as diagonals, and diagonals of D-finite power series are again D-finite [36]. This argument also implies an algorithm for computing linear differential equations satisfied by $Q(x, y; t)$, since the D-finiteness proof in [36] is effective and basically amounts to linear algebra. Therefore, from (3) one could, in principle, determine differential equations for $Q(x, y; t)$. To be more specific, the positive part of a formal power series $R \in \mathbb{Q}[[x, y, t]]$ can be encoded as

$$[x^>][y^>]R(x, y; t) = \frac{x}{1-x} \frac{y}{1-y} \odot_{x,y} R(x, y; t) = \text{Diag}_{x,x'} \text{Diag}_{y,y'} \frac{x}{1-x} \frac{y}{1-y} R(x', y'; t), \quad (4)$$

where the Hadamard product denoted $\odot_{x,y}$ is the term-wise product of two series, while the diagonal operator $\text{Diag}_{x,x'}$ selects those terms with equal exponents of x and x' . However, the direct use of (4) in our context leads to infeasible computations; worse, the intermediate algebraic objects involved in the calculations would probably have too large sizes to be merely written and stored. This is really unfortunate, since our need is mere evaluations of the diagonals in (4) at specific values for x and y .

To bypass this computational obstacle, we use two ingredients.

The first one is our main theoretical innovation: we reformulate the generating function $Q(x, y; t)$ in terms of *formal residues*. This idea is classical (and in fact already used in Lipshitz' proof [36]): we encode diagonals as residues, with the added advantage that early specialization of the variables x and/or y becomes possible. Additionally, our derivation bases on a positive-part extraction that differs from Bousquet-Mélou and Mishna's iterated operator $[x^>][y^>]$: we use a theory [2] of series with exponents that may be arbitrarily large in negative directions, but are restricted to fixed cones, together with a different, direct positive-part operator, $[x^>y^>]$, to be defined in Section 3. The outcome of this reformulation is the ability to compute linear differential equations with polynomial coefficients for the specializations $Q(x, 0; t)$ and $Q(0, y; t)$.

To perform these computations, we use a second ingredient, *creative telescoping*, an efficient algorithmic technique for the symbolic integration of multivariate functions. Indeed, a direct application of Lipshitz' linear-algebra algorithm (even with specialized variables) still leads to too large systems,

while creative telescoping succeeds in our cases of application; see [4, §2.3] for a related discussion. By specialization and recombination, the equations thus obtained give rise to rigorously proved differential equations for $Q(0, 0; t)$, $Q(1, 0; t)$, $Q(0, 1; t)$ and $Q(1, 1; t)$, thus answering question (i). The analysis of these differential equations combined with Kovacic’s algorithm [31] allows to answer question (iii) and to prove Theorem 2. Moreover, these differential equations are solved in explicit terms using symbolic algorithms for ODE factorization and ODE solving, leading to the proof of Theorem 1 and to the answer of question (ii).

The remarkable property that the differential equations in the 19×4 could *all* be solved in terms of hypergeometric functions relies on the fact that these operators share a very peculiar factorization pattern: they factor into factors that all have order 1 with the exception of the left-most one that can have order 1 or 2. The origin of this common mathematical feature deserves to be better understood.

Previous work. We conclude this introduction by mentioning previous contributions on the main topics of the article: D-finiteness, transcendence and explicit expressions for $Q(x, y; t)$ and its specializations.

D-finiteness. For the simplest models, the *square walk* (case 1) and the *diagonal walk* (case 2), D-finiteness is classical (e.g., [3]). Some more involved models have been considered sporadically: for case 19, Gouyou-Beauchamps [25] proved bijectively that $Q(1, 0; t)$ is D-finite; for cases 5 and 15 Mishna [41, §2.4.1, §2.4.2] showed that $Q(x, y; t)$ is D-finite using the kernel method; and for case 17 she proved [41, §2.3.3] that $Q(x, y; t)$ is D-finite by using a bijection with Young tableaux of height at most 3.

Several methods have been proposed to capture D-finiteness in a uniform way. For models with a vertical symmetry (cases 1–16), Bousquet-Mélou and Petkovšek [12, §2] proved that $Q(t) = Q(1, 1; t)$ is D-finite by a combinatorial argument, and Bousquet-Mélou [9, §3] proved D-finiteness of $Q(x, y; t)$ by an algebraic argument (a variation of the kernel method). For cases 17–19, Gessel and Zeilberger [23] proved D-finiteness of $Q(x, y; t)$ by using an algebraic version of the reflection principle; their argument works more generally when the step set is left invariant by a Weyl group and the walks are confined to a corresponding Weyl chamber. Bousquet-Mélou and Mishna [11] reproved D-finiteness of $Q(x, y; t)$ in all 19 cases by using the kernel equation and the group of the walk borrowed from [18]; their method generalizes the previous ones from [23] and [9]. Raschel [43] uses boundary value problems to get integral representations for $Q(x, y; t)$ that imply its D-finiteness in all 19 cases. By using methods from the book [18], Fayolle and Raschel reproved in [17, Theorem 1.1] that $(x, y) \mapsto Q(x, y; t_0)$ is D-finite for each $t_0 \in (0, \#S^{-1}]$ and in all 19 cases.

Transcendence. Algebraicity/transcendence proofs were first considered in some isolated cases: in case 15, $Q(x, y; t)$ was proved transcendental by Mishna [41, Th. 2.5]; in case 17, Mishna [41, §2.3.3], then Bousquet-Mélou and Mishna [11, §5.2], showed that $Q(x, y; t)$ and $Q(0, 0; t)$ are transcendental and that $Q(1, 1; t)$ is algebraic; in case 18, $Q(1, 1; t)$ was proved algebraic by Bousquet-Mélou and Mishna [11, §5.2]; in case 19, Bousquet-Mélou and Mishna [11, §5.3] showed that $Q(0, 0; t)$, $Q(0, 1; t)$, $Q(1, 0; t)$ and $Q(1, 1; t)$ are transcendental. The first unified transcendence proof for $Q(x, y; t)$ applying to all 19 cases is by Fayolle and Raschel [17, Theorem 1.1], although they attribute that result to Bousquet-Mélou and Mishna [11]. They actually proved more, namely that $Q(x, y; t_0)$ is transcendental for each $t_0 \in (0, \#S^{-1}]$, using the approach in [18, Chap. 4]. However, this result does not provide any transcendence information about specializations at $x, y \in \{0, 1\}$.

Explicit equations and formulas. For the simplest models (cases 1–2), simple formulas exist, see e.g. [9]; some of them admit bijective proofs, see e.g. [26]. For models 5 and 15, Mishna [41, Th. 2.5 and 2.6] gives explicit expressions of $Q(x, y; t)$ in terms of some auxiliary series. For model 17, basing on earlier work by Regev [44] and using a bijection with Young tableaux of height at most 3, Mishna [41, §2.3.3] shows that $q_{i,j;n}$ has a nice hypergeometric expression and that $Q(1, 1; t)$ is the (algebraic) generating function of Motzkin numbers. In cases 17–18, Bousquet-Mélou and Mishna [11, §5.2] gave explicit expressions for $Q(1, 1; t)$ (and more generally for $Q(x, 1/x; t)$). For model 19, it was proved by Gouyou-Beauchamps [25] that the number of n -step walks ending on the x -axis is a product of Catalan numbers; hypergeometric expressions for the total number of walks are derived in [11, §5.3], and for those ending at an arbitrary point (i, j) . Bostan and Kauers [5, 6] empirically determined differential equations for $Q(1, 1; t)$ and $Q(0, 0; t)$ for all 19 models. For the “highly symmetric” models (cases 1–4), Melczer and Mishna [37] proved differential equations for $Q(1, 1; t)$ conjectured by Bostan and Kauers [6]. Raschel [43] provides explicit integral representations of the complete generating function using a uniform analytic approach.

Before we enter into the details, Section 2 goes through the whole process on one concrete example. From now on, we more simply write \bar{x} and \bar{y} , respectively, for x^{-1} and y^{-1} .

2 A Worked Example: King Walks in the Quarter Plane

We illustrate our approach on an example. We choose the so-called *king walks* in the quarter plane (case 4 in Table 1), with step set $\mathcal{S} = \{(1, 1), (0, 1), (-1, 1), (-1, 0), (-1, -1), (0, -1), (1, -1), (1, 0)\}$. The first terms of the generating function $Q(1, 1; t)$ of king walks with prescribed length and arbitrary end point read (see <http://oeis.org/A151331>)

$$Q(1, 1; t) = 1 + 3t + 18t^2 + 105t^3 + 684t^4 + 4550t^5 + 31340t^6 + 219555t^7 + 1564080t^8 + \dots,$$

and the methods of the present article allow to obtain the above-mentioned closed formula (2) for it.

Here are the main steps of our approach. First, the classical kernel equation [11, Lemma 4] relates $Q(x, y; t)$ to $Q(x, 0; t)$, $Q(0, y; t)$, and $Q(0, 0; t)$:

$$Q(x, y; t) = 1 + t \left(S(x, y)Q(x, y; t) - \bar{y}(x + 1 + \bar{x})Q(x, 0; t) - \bar{x}(y + 1 + \bar{y})Q(0, y; t) + \bar{x}\bar{y}Q(0, 0; t) \right), \quad (5)$$

where $S(x, y)$ is the generating polynomial of the step set:

$$S(x, y) = \sum_{(i,j) \in \mathcal{S}} x^i y^j = xy + y + \bar{x}y + x + \bar{x} + x\bar{y} + \bar{y} + \bar{x}\bar{y}.$$

A simple but important observation is that the kernel $K(x, y; t) = 1 - tS(x, y)$ remains unchanged under the change of variables $(x, y) \leftarrow (x, \bar{y})$, $(x, y) \leftarrow (\bar{x}, y)$ and $(x, y) \leftarrow (\bar{x}, \bar{y})$. (Other step sets could require different changes of variables to provide the same property.)

Applying these rational transformations to the kernel equation (5) yields the four relations:

$$\begin{aligned} xyK(x, y; t)Q(x, y; t) &= xy - tx(x + 1 + \bar{x})Q(x, 0; t) - ty(y + 1 + \bar{y})Q(0, y; t) + tQ(0, 0; t), \\ -\bar{x}yK(x, y; t)Q(\bar{x}, y; t) &= -\bar{x}y + t\bar{x}(x + 1 + \bar{x})Q(\bar{x}, 0; t) + ty(y + 1 + \bar{y})Q(0, y; t) - tQ(0, 0; t), \\ \bar{x}\bar{y}K(x, y; t)Q(\bar{x}, \bar{y}; t) &= \bar{x}\bar{y} - t\bar{x}(x + 1 + \bar{x})Q(\bar{x}, 0; t) - t\bar{y}(y + 1 + \bar{y})Q(0, \bar{y}; t) + tQ(0, 0; t), \\ -x\bar{y}K(x, y; t)Q(x, \bar{y}; t) &= -x\bar{y} + tx(x + 1 + \bar{x})Q(x, 0; t) + t\bar{y}(y + 1 + \bar{y})Q(0, \bar{y}; t) - tQ(0, 0; t). \end{aligned}$$

Upon adding up these equations, all terms in the right-hand side involving Q disappear, resulting in

$$xyQ(x, y; t) - \bar{x}yQ(\bar{x}, y; t) + \bar{x}\bar{y}Q(\bar{x}, \bar{y}; t) - x\bar{y}Q(x, \bar{y}; t) = K(x, y; t)^{-1} (xy - \bar{x}y + \bar{x}\bar{y} - x\bar{y}).$$

Now, the main observation is that on the left-hand side, all terms except the first one involve negative powers either of x or of y . Therefore, extracting positive parts expresses the generating series $xyQ(x, y; t)$ as the positive part (w.r.t. x and y) of a trivariate rational function:

$$xyQ(x, y; t) = [x^>][y^>] \left(\frac{xy - \bar{x}y + \bar{x}\bar{y} - x\bar{y}}{1 - t(xy + y + y\bar{x} + \bar{x} + \bar{x}\bar{y} + \bar{y} + x\bar{y} + x)} \right). \quad (6)$$

Up to this point, our reasoning is borrowed from Bousquet-Mélou's and Mishna's article [11]. Combined with Lipshitz's result [36] that positive parts of D-finite functions are D-finite, it already implies that $Q(x, y; t)$ is D-finite; in particular, $Q(1, 1; t)$ is also D-finite. Our aim is to refine this qualitative result, and explicitly obtain a linear differential equation satisfied by $Q(1, 1; t)$. Such a differential equation was algorithmically *guessed* in [6] starting from the first terms in the power series expansion of $Q(1, 1; t)$; however the methods in [6] do not provide a *rigorous proof* of the correctness of that equation.

Starting from (6) and following more closely Lipshitz' encoding [36], a first observation is that $Q(x, y; t)$ is equal to the iterated diagonal

$$\text{Diag}_{x_1, x_2} \text{Diag}_{y_1, y_2} \left(\frac{x_2 y_2 (x_1 y_1 - \bar{x}_1 y_1 + \bar{x}_1 \bar{y}_1 - x_1 \bar{y}_1)}{(1 - x_2)(1 - y_2)(1 - t(x_1 y_1 + y_1 + y_1 \bar{x}_1 + \bar{x}_1 + \bar{x}_1 \bar{y}_1 + \bar{y}_1 + x_1 \bar{y}_1 + x_1))} \right). \quad (7)$$

There exist algorithms that take as input a rational function and compute a system of partial differential equations satisfied by its diagonal [14, 30, 35, 51]. However, in our case, these computations are too difficult, and exceed by far the limits of the best existing algorithms. The reason is that differential equations w.r.t. t and with polynomial coefficients in x, y, t are really huge, so the main limitation of algorithms computing (7) already comes from the size of the output. Another weakness of the diagonal encoding (7) is that it does not provide direct access to the univariate series $Q(1, 1; t)$, since taking diagonals and specializing variables are operations that do not commute.

To circumvent these difficulties and to make the computation feasible, our key idea is to encode the positive part in (6) as a formal residue:

$$Q(\alpha, \beta; t) = \text{Res}_{x,y} \left(\frac{xy - \bar{x}y + \bar{x}\bar{y} - x\bar{y}}{(1 - \alpha x)(1 - \beta y)(1 - t(xy + y + y\bar{x} + \bar{x} + \bar{x}\bar{y} + \bar{y} + x\bar{y} + x))} \right). \quad (8)$$

The formal proof of this encoding is delicate and is the topic of Section 3. The advantage of (8) over (7) is twofold. On the one hand, the residue computation can be carried out by using a single call to the creative-telescoping algorithm for rational functions, while the diagonal computation (7) has two steps, the first for a rational function in five variables, the second for an algebraic function in four variables. On the other hand, and more importantly, taking residues commutes with specialization, contrarily to positive parts and diagonals. Therefore, the generating series for walks $Q(1, 1; t)$ is equal to

$$Q(1, 1; t) = \text{Res}_{x,y} \left(\frac{xy - \bar{x}y + \bar{x}\bar{y} - x\bar{y}}{(1 - x)(1 - y)(1 - t(xy + y + y\bar{x} + \bar{x} + \bar{x}\bar{y} + \bar{y} + x\bar{y} + x))} \right).$$

A nice property of residues is that pure derivatives in x and y have zero residue. In order to compute a linear differential equation satisfied by $Q(1, 1; t)$ it thus suffices to find $U, V \in \mathbb{Q}(x, y; t)$ and a differential operator P in $\mathbb{Q}(t)\langle \partial_t \rangle$ such that

$$P \left(\frac{xy - \bar{x}y + \bar{x}\bar{y} - x\bar{y}}{(1 - x)(1 - y)(1 - t(xy + y + y\bar{x} + \bar{x} + \bar{x}\bar{y} + \bar{y} + x\bar{y} + x))} \right) = \partial_x(U) + \partial_y(V), \quad (9)$$

where ∂_t , ∂_x , and ∂_y stand for the derivation operators d/dt , d/dx , and d/dy , respectively. Indeed, since P depends neither on x nor on y , it commutes with the extraction of the residue with respect to x and y . The fact that the residue of $\partial_x(U) + \partial_y(V)$ is zero then proves that $P(Q(1, 1; t)) = 0$.

However, this last implication is more subtle than it looks: Equation (9) really is a relation between rational functions; the conclusion about $Q(t)$ is a conclusion about series. Owing to the ambiguity inherent to series expansions of rational functions, it is not clear a priori that the rational function in the left-hand side of (9) and U and V in its right-hand side can be expanded consistently and in accordance with the combinatorial interpretation for $Q(t)$. Describing the right expansion is the topic of Section 3: Theorem 5 provides the wanted implication.

Now, the existence of a non trivial (P, U, V) satisfying (9) is guaranteed by the theory of holonomic D-modules, which was first used in this context by Zeilberger [50] (see also [13]). The explicit form of a solution (P, U, V) can be determined using *creative telescoping* [14, 30, 51]; a detailed description of the algorithm is presented in [4, §2.4.2].

In our case, we find

$$P = t^2(1 + 4t)(8t - 1)(2t - 1)(1 + t)\partial_t^3 + t(200t^3 + 576t^4 - 33t - 252t^2 + 5)\partial_t^2 + 4(22t^3 - 117t^2 - 12t + 288t^4 + 1)\partial_t + 384t^3 - 12 - 144t - 72t^2. \quad (10)$$

Note that this is precisely the differential operator *guessed* in [6].

Moreover, factorization algorithms for linear differential operators [27] can be used to prove that $P = P_2P_1$, where $P_1 = \partial_t + 1/t$ and

$$P_2 = t^2(1 + 4t)(1 - 8t)(1 - 2t)(1 + t)\partial_t^2 + 2t(256t^4 + 80t^3 - 111t^2 - 14t + 2)\partial_t + 768t^4 + 8t^3 - 306t^2 - 30t + 2. \quad (11)$$

It follows that the Laurent power series

$$f(t) = \frac{dQ}{dt}(1, 1; t) + \frac{Q(1, 1; t)}{t} = t^{-1} + 6 + 54t + 420t^2 + 3420t^3 + 27300t^4 + 219380t^5 + O(t^6)$$

is a solution of P_2 . Starting from the second order operator P_2 , algorithmic methods explained in [4, §2.6] (see also [16]) allow to express $f(t)$ in terms of a ${}_2F_1$ hypergeometric function:

$$f(t) = \frac{1}{t(1 + 4t)^3} \cdot {}_2F_1 \left(\frac{3}{2}, \frac{3}{2} \middle| \frac{16t(1 + t)}{(1 + 4t)^2} \right).$$

Finally, solving the equation $d/dt Q(1, 1; t) + Q(1, 1; t)/t = f(t)$ yields formula (2).

Similarly, by deriving variants of (9) parametrized by *indeterminate* α and β we obtain the representations

$$Q(\alpha, 0; t) = \operatorname{Res}_{x,y} \left(\frac{xy - \bar{x}y + \bar{x}\bar{y} - x\bar{y}}{(1 - \alpha x)(1 - t(xy + y + y\bar{x} + \bar{x} + \bar{x}\bar{y} + \bar{y} + x\bar{y} + x))} \right)$$

and

$$Q(0, \beta; t) = \operatorname{Res}_{x,y} \left(\frac{xy - \bar{x}y + \bar{x}\bar{y} - x\bar{y}}{(1 - \beta y)(1 - t(xy + y + y\bar{x} + \bar{x} + \bar{x}\bar{y} + \bar{y} + x\bar{y} + x))} \right).$$

Creative-telescoping techniques still allow the effective computation of differential operators in $\mathbb{Q}(\alpha; t)\langle\partial_t\rangle$ for $Q(\alpha, 0; t)$, resp. in $\mathbb{Q}(\beta; t)\langle\partial_t\rangle$ for $Q(0, \beta; t)$. Owing to the additional symbolic indeterminate, the computations are much harder than for $Q(1, 1; t)$, but still feasible. Each of the resulting differential operators factors again, this time as a product of an order-two operator and of three order-one operators. Moreover, the second-order operators are again solvable in terms of ${}_2F_1$ functions. Finally, a closed formula for $Q(\alpha, \beta; t)$ is obtained from the closed formulas for $Q(\alpha, 0; t)$ and $Q(0, \beta; t)$ via the kernel equation (5). This detour is computationally crucial, since performing creative telescoping directly on the five-variable rational function from (8) is not feasible even using today's best algorithms.

3 Computing Positive Parts as Formal Residues

We will now put the assertions made in the previous section for one particular case on solid algebraic grounds by clarifying to which series domains we map our rational functions, by introducing formal notions of residue, Hadamard product, and positive part for the objects in these domains, and by showing that the differential equations we obtain from creative telescoping indeed annihilate the positive parts.

To this end, we study vector spaces of “series” that are just bilateral infinite arrays in §3.1, before elaborating in §3.2 on a theory of rings and fields of series with exponents in cones [2]. We are then able to represent positive parts as residues of suitable products in those rings in §3.3. In §3.4, a notion of cones in opposition is introduced to refine this representation and allow the early specialization we need for efficiency in computations. The connection to our application is done in §3.5, before we justify the use of creative telescoping in §3.6.

As the present section is of a more general nature than the application to small-step walks, we develop the theory here over a general field \mathbb{K} of characteristic 0.

3.1 Linear operations on infinite arrays

The set $\mathbb{K}^{\mathbb{Z}^n}$ consisting of all the series

$$\sum_{i_1, \dots, i_n \in \mathbb{Z}} a_{i_1, \dots, i_n} x_1^{i_1} \cdots x_n^{i_n} \tag{12}$$

is a \mathbb{K} -vector space. Here, we use the word “series” and the notation \sum just for convenience, disregarding any question of convergence. The set $\mathbb{K}^{\mathbb{Z}^n}$ is not a ring, but we can endow it with well-defined operations of residue (from $\mathbb{K}^{\mathbb{Z}^n}$ to \mathbb{K}) and positive part (from $\mathbb{K}^{\mathbb{Z}^n}$ to $\mathbb{K}^{\mathbb{N}^n}$): for a given series f , the residue $\operatorname{Res}_{x_1, \dots, x_n} f$ and positive part $[x_1^> \cdots x_n^>]f$ are defined in the natural way as

$$a_{-1, \dots, -1} \quad \text{and} \quad \sum_{i_1, \dots, i_n > 0} a_{i_1, \dots, i_n} x_1^{i_1} \cdots x_n^{i_n},$$

respectively. More generally, we may consider residue and positive-part extractions with respect to some of the variables only, and this will be obvious from the notation. This way, positive-part extractions compose nicely, e.g., $[x_1^>][x_2^>] = [x_2^>][x_1^>] = [x_1^> x_2^>]$. The vector space $\mathbb{K}^{\mathbb{Z}^n}$ also possesses an internal law of Hadamard product, defined for two series f and g with respective coefficients a_{i_1, \dots, i_n} and b_{i_1, \dots, i_n} by

$$f \odot g = \sum_{i_1, \dots, i_n \in \mathbb{Z}} a_{i_1, \dots, i_n} b_{i_1, \dots, i_n} x_1^{i_1} \cdots x_n^{i_n}.$$

Finally, for $k \leq n$, we will need more operations for series in different variable sets:

- for $f \in \mathbb{K}^{\mathbb{Z}^k}$ and $g \in \mathbb{K}^{\mathbb{Z}^n}$, we define

$$f \odot_{k \leq n} g = \sum_{i_1, \dots, i_n \in \mathbb{Z}} a_{i_1, \dots, i_k} b_{i_1, \dots, i_n} x_1^{i_1} \cdots x_n^{i_n}; \quad (13)$$

- for $f \in \mathbb{K}^{\mathbb{Z}^n}$ and $g \in \mathbb{K}^{\mathbb{Z}^k}$, we define

$$f \odot_{n \geq k} g = \sum_{i_1, \dots, i_n \in \mathbb{Z}} a_{i_1, \dots, i_n} b_{i_1, \dots, i_k} x_1^{i_1} \cdots x_n^{i_n}. \quad (14)$$

Beware that, despite the natural embedding of $\mathbb{K}^{\mathbb{Z}^k}$ into $\mathbb{K}^{\mathbb{Z}^n}$ mapping $f \in \mathbb{K}^{\mathbb{Z}^k}$ to $f x_{k+1}^0 \cdots x_n^0 \in \mathbb{K}^{\mathbb{Z}^n}$, $f \odot_{k \leq n} g$ in (13) is generally not equal to $(f x_{k+1}^0 \cdots x_n^0) \odot g$ and similarly $f \odot_{n \geq k} g$ in (14) is generally not equal to $f \odot (g x_{k+1}^0 \cdots x_n^0)$.

The vector space $\mathbb{K}^{\mathbb{Z}^n}$ also has a (linear-)differential structure: it can be endowed with a derivation operator ∂_1 that maps a series (12) to

$$\sum_{i_1, \dots, i_n \in \mathbb{Z}} i_1 a_{i_1, \dots, i_n} x_1^{i_1-1} x_2^{i_2} \cdots x_n^{i_n},$$

and with operators $\partial_2, \dots, \partial_n$ defined similarly. The following result is then obvious.

Lemma 1. *For any series $f \in \mathbb{K}^{\mathbb{Z}^n}$ and any j satisfying $1 \leq j \leq n$, $\text{Res}_{x_1, \dots, x_n} \partial_j f = 0$.*

Additionally, as Laurent polynomials are finite sums, $\mathbb{K}^{\mathbb{Z}^n}$ admits well-defined operations of multiplication by a Laurent polynomial. Together with closure under derivation, this makes it meaningful to consider differential equations in series of $\mathbb{K}^{\mathbb{Z}^n}$.

3.2 Series expansions with respect to a cone

In what follows, we introduce a family of rings of series with support restricted to cones of \mathbb{Z}^n . This extends to rings of series whose support enjoys a certain compatibility condition with a fixed monomial ordering. This discussion is mostly borrowed from [2]. See the introduction in that article for the comparison to the theory of Hahn series (later developed by Mal'cev, Neumann, and recently Xin) and to the theory of MacDonal series (recently developed by Aroca, Cano, and Jung).

In order to highlight the names and number of indeterminates in the notation for various rings and fields of polynomials, series, etc, in what follows, we will compactly write U_m to denote m indeterminates u_1, \dots, u_m in notations like $\mathbb{K}[U_m]$, $\mathbb{K}(U_m)$, $\mathbb{K}[[U_m]]$, etc. To refer to only the last $m - p$ indeterminates u_{p+1}, \dots, u_m of U_m , we will write $U_{m \setminus p}$. We will use similar notation for inverses and combinations of names of indeterminates, e.g., $\mathbb{K}[X_n, Y_{n \setminus k}, Y_k^{-1}]$ denotes $\mathbb{K}[x_1, \dots, x_n, y_{k+1}, \dots, y_n, y_1^{-1}, \dots, y_k^{-1}]$.

Definition 1. A topologically closed set $C \subseteq \mathbb{R}^n$ is called a *cone* if for all $u, v \in C$ and $\lambda, \mu \geq 0$, $\lambda u + \mu v \in C$. A cone C is called *line-free* if $u, -u \in C$ implies $u = 0$.

If C is a line-free cone, then the set consisting of all the series of the form (12) with $a_{i_1, \dots, i_n} = 0$ whenever $(i_1, \dots, i_n) \notin C$ forms a ring together with the natural addition and multiplication. We denote it by $\mathbb{K}_C[[x_1, \dots, x_n]]$ or $\mathbb{K}_C[[X_n]]$. It is a \mathbb{K} -algebra. For the trivial cone $C = \{0\}$, this is simply \mathbb{K} . The set $\mathbb{K}_C\langle\langle X_n \rangle\rangle := \bigcup_{(i_1, \dots, i_n) \in \mathbb{Z}^n} x_1^{i_1} \cdots x_n^{i_n} \mathbb{K}_C[[X_n]]$, forms another ring, in fact just $\mathbb{K}_C[[X_n]][X_n, X_n^{-1}]$. Its elements are series whose supports are contained in a finite union of translated copies $(i_1, \dots, i_n) + C$ of C . For the trivial cone $C = \{0\}$, this is simply the ring of Laurent polynomials, $\mathbb{K}[X_n, X_n^{-1}]$. The ring $\mathbb{K}_C\langle\langle X_n \rangle\rangle$ inherits the operations of residue and positive part, the internal law of Hadamard product, and the derivation operations. Observe as well that the Hadamard product $f_1 \odot f_2$ of series $f_1 \in \mathbb{K}_{C_1}\langle\langle X_n \rangle\rangle$ and $f_2 \in \mathbb{K}_{C_2}\langle\langle X_n \rangle\rangle$ for different C_1 and C_2 belongs to both rings, because the support of $f \odot g$ is the intersection of the supports of f and g .

Observe that rational functions from $\mathbb{K}(x_1, \dots, x_n)$, henceforth denoted $\mathbb{K}(X_n)$, can often be expanded as elements of different rings $\mathbb{K}_{C_1}\langle\langle X_n \rangle\rangle$ and $\mathbb{K}_{C_2}\langle\langle X_n \rangle\rangle$, potentially with trivial intersection $C_1 \cap C_2 = \{0\}$. This should not be understood as implying that the rational function is part of the intersection of both rings. In fact, we should refrain from viewing rational functions as elements of series rings: they only have images in those rings. As a consequence, residue and positive-part extractions of a rational function only make sense with respect to a given cone expansion. On the other hand, given two line-free cones $C \subseteq C'$, we can safely identify $\mathbb{K}_C[[X_n]]$ as a subring of $\mathbb{K}_{C'}[[X_n]]$, each given series having the

same coefficient family with regard to both expansions. The situation is the same with $\mathbb{K}_C\langle\langle X_n \rangle\rangle$ that we identify as a subring of $\mathbb{K}_{C'}\langle\langle X_n \rangle\rangle$. Similar identifications occur when extending the variable set: for instance, we will freely identify $\mathbb{K}_C[[X_n]]$ and $\mathbb{K}_{C \times \{0\}^m}[[X_{n+m}]]$.

To avoid blind identification of a rational function with a series, as the latter must depend on the cone used, we introduce the following notation: for a rational function $P/Q \in \mathbb{K}(X_n)$ which admits an expansion with respect to a line-free cone C , we write $[P/Q]_C$ for this expansion in $\mathbb{K}_C\langle\langle X_n \rangle\rangle$. As a consequence of [2, Thm 12], such an expansion exists if and only if Q can be factored in the form $x_1^{i_1} \cdots x_n^{i_n} \tilde{Q}$ for $\tilde{Q} \in \mathbb{K}_C[[X_n]]$ with non-zero constant term.

There will be situations where we need to know the existence of a cone C with respect to which a certain rational function can be expanded without having to know the exact cone C . This will be the case in §3.6 below, when we link creative telescoping to series expansions: we will want to know that certificates of creative telescoping can be expanded over some cone C without having to observe them to be able to make the cone C explicit. To this end, we borrow from [2] the definition of more rings and fields, defined with respect to a monomial order.

We start with a fixed monomial order \preccurlyeq on the monomials in x_1, \dots, x_n (with exponents in \mathbb{Z}), that is, with a total order that is monotonic with respect to product: for all a, b , and c , $a \preccurlyeq b$ implies $ac \preccurlyeq bc$.

Definition 2. A cone C is *compatible* with the monomial order \preccurlyeq if the apex of C coincides with its minimal element with respect to \preccurlyeq .

Given \preccurlyeq , the union over all cones C compatible with \preccurlyeq of the rings $\mathbb{K}_C[[X_n]]$ is a ring, denoted $\mathbb{K}_{\preccurlyeq}[[X_n]]$ in what follows. The set $\mathbb{K}_{\preccurlyeq}((X_n)) := \bigcup_{(i_1, \dots, i_n) \in \mathbb{Z}^n} x_1^{i_1} \cdots x_n^{i_n} \mathbb{K}_{\preccurlyeq}[[X_n]]$ now forms another ring. This ring is even a field [2, Thm 15]. Now, for a fixed cone C compatible with the monomial order \preccurlyeq , the ring $\mathbb{K}_C[[X_n]]$ is a subring of both $\mathbb{K}_C\langle\langle X_n \rangle\rangle$ and $\mathbb{K}_{\preccurlyeq}[[X_n]]$, and both of these rings are subrings of the field $\mathbb{K}_{\preccurlyeq}((X_n))$. All those inclusions are canonical, in the sense that they preserve the coefficients. In particular, a rational function F of the field $\mathbb{K}(X_n)$ maps to a well-defined series of $\mathbb{K}_{\preccurlyeq}((X_n))$, which we will denote $[F]_{\preccurlyeq}$. When C is compatible with \preccurlyeq , we will do the identification $[F]_C = [F]_{\preccurlyeq}$. As a consequence, the field $[\mathbb{K}(X_n)]_{\preccurlyeq} = \{[F]_{\preccurlyeq} : F \in \mathbb{K}(X_n)\}$ is a subfield of $\mathbb{K}_{\preccurlyeq}((X_n))$.

3.3 Positive parts as residues

For a given series $f \in \mathbb{K}_C\langle\langle X_n \rangle\rangle$, the positive part $[x_1^> \cdots x_n^>]f$ can be obtained by taking the Hadamard product with the expansion as a geometric series in $\mathbb{K}_{\mathbb{R}_{\geq 0}^n}[[X_n]]$ of $\frac{x_1 \cdots x_n}{(1-x_1) \cdots (1-x_n)}$. We will argue that the usual residue formula for computing Hadamard products holds in this setting.

Lemma 2. Let $\pi_1: \mathbb{R}^n \rightarrow \mathbb{R}^k$ and $\pi_2: \mathbb{R}^n \rightarrow \mathbb{R}^{n-k}$ be the projections to the first k and last $n-k$ coordinates, respectively, so that we have $u = (\pi_1(u), \pi_2(u))$ for every $u \in \mathbb{R}^n$. Each of the following constructions is a line-free cone:

1. the set $C_1 \star C_2 := \{(u, v - u) : u \in C_1, v \in C_2\} \subseteq \mathbb{R}^{2n}$ for line-free cones $C_1, C_2 \subseteq \mathbb{R}^n$,
2. the set $C_1 \star_{k \leq n} C_2 := \{(u, \pi_2(v), \pi_1(v) - u) : u \in C_1, v \in C_2\} \subseteq \mathbb{R}^{k+n}$ for line-free cones $C_1 \subseteq \mathbb{R}^k$ and $C_2 \subseteq \mathbb{R}^n$,
3. the set $C_1 \star_{n \geq k} C_2 := \{(u, v - \pi_1(u)) : u \in C_1, v \in C_2\} \subseteq \mathbb{R}^{n+k}$ for line-free cones $C_1 \subseteq \mathbb{R}^n$ and $C_2 \subseteq \mathbb{R}^k$.

Proof. For the first construction, given $u_1, u_2 \in C_1$, $v_1, v_2 \in C_2$, and $\lambda, \mu \geq 0$, $u := \lambda u_1 + \mu u_2$ and $v := \lambda v_1 + \mu v_2$ are in C_1 and C_2 , respectively, because C_1, C_2 are cones. Consequently, $\lambda(u_1, v_1 - u_1) + \mu(u_2, v_2 - u_2) = (u, v - u)$ is in $C_1 \star C_2$, and the latter is a cone. Next, given $(u, v - u)$ in $C_1 \star C_2$, if $-(u, v - u)$ is in $C_1 \star C_2$, too, then it can be written $(u', v' - u')$. From $(u, v - u) = -(u', v' - u')$ follows $u = -u'$, which implies $u = u' = 0$ because C_1 is line-free. This, in turn, implies that $v = -v'$, which implies that $v = v' = 0$ because C_2 is line-free. Therefore, $C_1 \star C_2$ is line-free.

The same argument applies with minor changes to the two other constructions. \square

As a consequence, each of $\mathbb{K}_{C_1 \star C_2}\langle\langle X_n, Y_n \rangle\rangle$, $\mathbb{K}_{C_1 \star_{k \leq n} C_2}\langle\langle X_n, Y_k \rangle\rangle$, and $\mathbb{K}_{C_1 \star_{n \geq k} C_2}\langle\langle X_n, Y_k \rangle\rangle$ is a bonafide ring for any two line-free cones C_1, C_2 of \mathbb{R}^n or \mathbb{R}^k (as required). Remark the intended choice to use X_n in the case $C_1 \subseteq \mathbb{Z}^k$ as well, and not X_k , in accordance to Equation (16) below. As well, observe $C_1 \star C_2 = C_1 \star_{n \leq n} C_2 = C_1 \star_{n \geq n} C_2$, and similar identities on the level of rings.

Lemma 3. (i) Let $C_1, C_2 \subseteq \mathbb{R}^n$ be two line-free cones. Then, for any $f \in \mathbb{K}_{C_1} \langle \langle X_n \rangle \rangle$ and $g \in \mathbb{K}_{C_2} \langle \langle X_n \rangle \rangle$,

$$f \odot g = \text{Res}_{y_1, \dots, y_n} \frac{1}{y_1 \cdots y_n} f \left(\frac{x_1}{y_1}, \dots, \frac{x_n}{y_n} \right) g(y_1, \dots, y_n), \quad (15)$$

where the argument of the residue is understood as a product in $\mathbb{K}_{C_1 \star C_2} \langle \langle X_n, Y_n \rangle \rangle$.

(ii) For $k \leq n$, let $C_1 \subseteq \mathbb{R}^k$ and $C_2 \subseteq \mathbb{R}^n$ be two line-free cones. Then, for any $f \in \mathbb{K}_{C_1} \langle \langle X_k \rangle \rangle$ and $g \in \mathbb{K}_{C_2} \langle \langle X_n \rangle \rangle$,

$$f \underset{k \leq n}{\odot} g = \text{Res}_{y_1, \dots, y_k} \frac{1}{y_1 \cdots y_k} f \left(\frac{x_1}{y_1}, \dots, \frac{x_k}{y_k} \right) g(y_1, \dots, y_k, x_{k+1}, \dots, x_n), \quad (16)$$

where the argument of the residue is understood as a product in $\mathbb{K}_{C_1 \star_{k \leq n} C_2} \langle \langle X_n, Y_k \rangle \rangle$.

(iii) For $k \leq n$, let $C_1 \subseteq \mathbb{R}^n$ and $C_2 \subseteq \mathbb{R}^k$ be two line-free cones. Then, for any $f \in \mathbb{K}_{C_1} \langle \langle X_n \rangle \rangle$ and $g \in \mathbb{K}_{C_2} \langle \langle X_k \rangle \rangle$,

$$f \underset{n \geq k}{\odot} g = \text{Res}_{y_1, \dots, y_k} \frac{1}{y_1 \cdots y_k} f \left(\frac{x_1}{y_1}, \dots, \frac{x_k}{y_k}, x_{k+1}, \dots, x_n \right) g(y_1, \dots, y_k), \quad (17)$$

where the argument of the residue is understood as a product in $\mathbb{K}_{C_1 \star_{n \geq k} C_2} \langle \langle X_n, Y_k \rangle \rangle$.

Before the proof, observe that (16) and (17) cannot just be obtained as specializations of (15).

Proof. We start by proving (15). Set $C := C_1 \star C_2 \subseteq \mathbb{R}^{2n}$, which is a line-free cone by Lemma 2. Now, observe that $f(x_1/y_1, \dots, x_n/y_n)$ and $g(y_1, \dots, y_n)$ are in $\mathbb{K}_C \langle \langle X_n, Y_n \rangle \rangle$, respectively because $\{(u, -u) : u \in C_1\} \subseteq C$ and because $\{(0, v) : v \in C_2\} \subseteq C$. So, the argument of the residue in (15) is a well-defined product. Call it h . To see that $\text{Res}_{y_1, \dots, y_n} h = f \odot g$, calculate:

$$h = \frac{1}{y_1 \cdots y_n} \left(\sum_{u_1, \dots, u_n} f_{u_1, \dots, u_n} \frac{x_1^{u_1} \cdots x_n^{u_n}}{y_1^{u_1} \cdots y_n^{u_n}} \right) \left(\sum_{v_1, \dots, v_n} g_{v_1, \dots, v_n} y_1^{v_1} \cdots y_n^{v_n} \right) = \sum_{i_1, \dots, i_n} \left(\sum_{(v_1, \dots, v_n) - (u_1, \dots, u_n) = (i_1, \dots, i_n)} f_{u_1, \dots, u_n} g_{v_1, \dots, v_n} x_1^{u_1} \cdots x_n^{u_n} \right) y_1^{i_1-1} \cdots y_n^{i_n-1}, \quad (18)$$

where the sums are meant to extend over all integers with the understanding that f_{u_1, \dots, u_n} and g_{v_1, \dots, v_n} are zero outside of the respective finite unions of translated cones. Applying $\text{Res}_{y_1, \dots, y_n}$ to this identity selects the term for which $i_1 = \dots = i_n = 0$, forcing $(v_1, \dots, v_n) = (u_1, \dots, u_n)$. Hence we have:

$$\text{Res}_{y_1, \dots, y_n} h = \sum_{(u_1, \dots, u_n)} f_{u_1, \dots, u_n} g_{u_1, \dots, u_n} x_1^{u_1} \cdots x_n^{u_n} = f \odot g,$$

as was to be proved to justify (15).

To prove (16), set $C' := C_1 \star_{k \leq n} C_2 \subseteq \mathbb{R}^{k+n}$, which is a line-free cone by Lemma 2. Now, observe that $f(x_1/y_1, \dots, x_k/y_k)$ and $g(y_1, \dots, y_k, x_{k+1}, \dots, x_n)$ are in $\mathbb{K}_{C'} \langle \langle X_n, Y_k \rangle \rangle$, respectively because $\{(u, 0, -u) : u \in C_1\} \subseteq C'$ and because $\{(0, \pi_2(v), \pi_1(v)) : v \in C_2\} \subseteq C'$. So, the argument of the residue in (16) is a well-defined product. The proof of (16) then follows calculations analogous to (18).

To prove (17), set $C'' := C_1 \star_{n \geq k} C_2 \subseteq \mathbb{R}^{n+k}$, which is a line-free cone by Lemma 2. Now, observe that $f(x_1/y_1, \dots, x_k/y_k, x_{k+1}, \dots, x_n)$ and $g(y_1, \dots, y_k)$ are in $\mathbb{K}_{C''} \langle \langle X_n, Y_k \rangle \rangle$, respectively because $\{(u, -\pi_1(u)) : u \in C_1\} \subseteq C''$ and because $\{(0, v) : v \in C_2\} \subseteq C''$. So, the argument of the residue in (17) is a well-defined product. The proof of (17) then follows calculations analogous to (18). \square

Recall the notation $[P/Q]_C$ to denote the expansion of a rational function with respect to a cone.

Lemma 4. For every line-free cone $C \subseteq \mathbb{R}^n$, every $\phi \in \mathbb{K}_C \langle \langle X_n \rangle \rangle$, and every $k \in \{1, \dots, n\}$, we have:

$$[x_1^> \cdots x_k^>] \phi = \text{Res}_{y_1, \dots, y_k} \frac{1}{y_1 \cdots y_k} \left[\frac{x_1 \cdots x_k}{(1 - \frac{x_1}{y_1}) \cdots (1 - \frac{x_k}{y_k})} \right]_{\mathbb{R}_{\geq 0}^k \times \mathbb{R}_{\leq 0}^k} \phi(y_1, \dots, y_k, x_{k+1}, \dots, x_n) \quad (19)$$

$$= \text{Res}_{y_1, \dots, y_k} \frac{1}{y_1 \cdots y_k} \phi \left(\frac{x_1}{y_1}, \dots, \frac{x_k}{y_k}, x_{k+1}, \dots, x_n \right) \left[\frac{y_1 \cdots y_k}{(1 - y_1) \cdots (1 - y_k)} \right]_{\mathbb{R}_{\geq 0}^k}, \quad (20)$$

where the brackets around rational functions in (19) and (20) denote taking their expansions in, respectively, $\mathbb{K}_{\mathbb{R}_{\geq 0}^k \times \mathbb{R}_{\leq 0}^k} \langle \langle X_k, Y_k \rangle \rangle$ and $\mathbb{K}_{\mathbb{R}_{\geq 0}^k} \langle \langle Y_k \rangle \rangle$.

Proof. Fix C' to $\mathbb{R}_{\geq 0}^k$ and $\psi \in \mathbb{K}_{C'} \langle \langle X_k \rangle \rangle$ to the geometric-series expansion of $\frac{x_1 \cdots x_k}{(1-x_1) \cdots (1-x_k)}$. The desired positive part can then be represented in two ways as variants of Hadamard products: $[x_1^> \cdots x_k^>] \phi = \psi \odot \phi = \phi \odot \psi$. We then use Lemma 3 twice:

- Firstly, by (16), $[x_1^> \cdots x_k^>] \phi = \text{Res}_{y_1, \dots, y_k} \frac{1}{y_1 \cdots y_k} \psi\left(\frac{x_1}{y_1}, \dots, \frac{x_k}{y_k}\right) \phi(y_1, \dots, y_k, x_{k+1}, \dots, x_n)$, where the product is in $\mathbb{K}_{C'} \underset{k \leq n}{\star} C \langle \langle X_n, Y_k \rangle \rangle$. More specifically, $\psi\left(\frac{x_1}{y_1}, \dots, \frac{x_k}{y_k}\right)$ is in $\mathbb{K}_{C'} \underset{k \leq n}{\star} \{0\}^n \langle \langle X_n, Y_k \rangle \rangle$, and with the proper identification in $\mathbb{K}_{\mathbb{R}_{\geq 0}^k \star \{0\}^k} \langle \langle X_k, Y_k \rangle \rangle$, hence (19) and the announced expansion.
- Secondly, by (17), $[x_1^> \cdots x_k^>] \phi = \text{Res}_{y_1, \dots, y_k} \frac{1}{y_1 \cdots y_k} \phi\left(\frac{x_1}{y_1}, \dots, \frac{x_k}{y_k}, x_{k+1}, \dots, x_n\right) \psi(y_1, \dots, y_k)$, where the product is in $\mathbb{K}_{C'} \underset{n \geq k}{\star} C' \langle \langle X_n, Y_n \rangle \rangle$. More specifically, $\psi(y_1, \dots, y_k)$ is in $\mathbb{K}_{\{0\}^n \underset{n \geq k}{\star} C'} \langle \langle X_n, Y_k \rangle \rangle$, and with the proper identification in $\mathbb{K}_{C'} \langle \langle Y_k \rangle \rangle$, hence (20) and the announced expansion. \square

3.4 Specializations of positive parts

Next, we would like to justify a formula to represent $([x_1^> \cdots x_k^>] f) \Big|_{x_1=\alpha_1, \dots, x_k=\alpha_k}$ as a Hadamard product, where $\alpha_1, \dots, \alpha_k$ are fixed elements of \mathbb{K} . For the evaluation of the positive part at some arbitrary field elements to make sense, we need to impose a restriction on the cones, which we will express using the following notion.

Definition 3. Let $\pi_2: \mathbb{R}^n \rightarrow \mathbb{R}^{n-k}$ be the projection to the last $n-k$ coordinates. We say that two line-free cones $C_1, C_2 \subseteq \mathbb{R}^n$ are *in opposition with respect to the first k variables* if $\pi_2(C_1), \pi_2(C_2)$ are line-free cones and $C_1 \cap C_2 \cap (\mathbb{R}^k \times \{0\}^{n-k}) = \{0\}^n$.

Lemma 5. Let C_1 and C_2 in \mathbb{R}^n be in opposition with respect to the first $k \leq n$ variables, and let $u, v \in \mathbb{R}^n$. Then, the set $(C_1 + u) \cap (C_2 + v) \cap (\mathbb{R}^k \times \{0\}^{n-k})$ is bounded.

Proof. Given $w, d \in \mathbb{R}^n$, denote by $R_{w,d}$ the set $\{w + cd : c \geq 0\}$. We recall without proof that: (i) for any $R_{w,d}$ contained in a cone C , both w and d are in C ; (ii) any unbounded convex set \mathbb{K} contains some $R_{w,d}$ with nonzero d ; (iii) as a consequence, for any cone C , any time $R_{w,d} \subseteq w' + C$, we have $w - w', d \in C$.

Now, if the (convex) set $(C_1 + u) \cap (C_2 + v) \cap (\mathbb{R}^k \times \{0\}^{n-k})$ was unbounded, it would contain some $R_{w,d}$ for a nonzero d . We would have in particular that d would be in C_1, C_2 , and $\mathbb{R}^k \times \{0\}^{n-k}$, implying $d = 0$. This would contradict that C_1 and C_2 are in opposition with respect to the first k variables. \square

Lemma 6 below gives a sufficient condition that allows a Hadamard product to specialize at $x_1 = \alpha_1, \dots, x_k = \alpha_k$ in a well defined way. Two potential obstructions to this are: (i) situations in which $f \odot g$ involves either of x_1, \dots, x_k with a negative exponent; (ii) situations in which the supports of f and g make it possible that the Hadamard product should involve some monomial in x_{k+1}, \dots, x_n having as coefficient an infinite sum in x_1, \dots, x_k that does not converge as a series at $x_1 = \alpha_1, \dots, x_k = \alpha_k$. In the lemma, the cones being in opposition ensures the finiteness of all sums under consideration in (ii), while case (i) is excluded by imposing $\alpha_j \neq 0$ when necessary. To state conditions for the latter point, for $1 \leq j \leq k \leq n$, for a series s in x_1, \dots, x_k and a series t in x_1, \dots, x_n , we introduce two predicates $H_j^n(t)$ and $H_j^{k \leq n}(s, t)$ that express, respectively, that t cannot be evaluated at $x_j = 0$ and that the (generally asymmetric) Hadamard product of s and t cannot be evaluated at $x_j = 0$; they read:

$H_j^n(t)$: There exists a monomial in x_1, \dots, x_n with negative exponent of x_j that occurs with non-zero coefficient in the series t .

$H_j^{k \leq n}(s, t)$: There exists a monomial m in x_1, \dots, x_k with negative exponent of x_j and a monomial m' in x_{k+1}, \dots, x_n , such that m occurs with non-zero coefficient in the series s and mm' occurs with non-zero coefficient in the series t .

Note that $H_j^{k \leq n}(s, t)$ implies $H_j^k(s)$ and $H_j^n(t)$.

Lemma 6. (i) Let $C_1, C_2 \subseteq \mathbb{R}^n$ be in opposition with respect to the first k variables. Further, let $f \in \mathbb{K}_{C_1} \langle \langle X_n \rangle \rangle$ and $g \in \mathbb{K}_{C_2} \langle \langle X_n \rangle \rangle$. Then, $(f \odot g) \Big|_{x_1=\alpha_1, \dots, x_k=\alpha_k}$ is well-defined for every $\alpha_1, \dots, \alpha_k \in \mathbb{K}$, provided that for each $1 \leq j \leq k$, $\alpha_j \neq 0$ if $H_j^{n \leq n}(f, g)$.

(ii) Let $C_1 \subseteq \mathbb{R}^k$ and $C_2 \subseteq \mathbb{R}^n$ be such that $C_1 \times \{0\}^{n-k}$ and C_2 are in opposition with respect to the first k variables. Further, let $f \in \mathbb{K}_{C_1} \langle \langle X_k \rangle \rangle$ and $g \in \mathbb{K}_{C_2} \langle \langle X_n \rangle \rangle$. Then, $(f \underset{k \leq n}{\odot} g)|_{x_1=\alpha_1, \dots, x_k=\alpha_k}$ is well-defined for every $\alpha_1, \dots, \alpha_k \in \mathbb{K}$, provided that for each $1 \leq j \leq k$, $\alpha_j \neq 0$ if $H_j^{k \leq n}(f, g)$.

(iii) Let $C_1 \subseteq \mathbb{R}^n$ and $C_2 \subseteq \mathbb{R}^k$ be such that C_1 and $C_2 \times \{0\}^{n-k}$ are in opposition with respect to the first k variables. Further, let $f \in \mathbb{K}_{C_1} \langle \langle X_n \rangle \rangle$ and $g \in \mathbb{K}_{C_2} \langle \langle X_k \rangle \rangle$. Then, $(f \underset{n \geq k}{\odot} g)|_{x_1=\alpha_1, \dots, x_k=\alpha_k}$ is well-defined for every $\alpha_1, \dots, \alpha_k \in \mathbb{K}$, provided that for each $1 \leq j \leq k$, $\alpha_j \neq 0$ if $H_j^{k \leq n}(g, f)$.

Proof. (i) Suppose first that there are vectors $e_1, e_2 \in \mathbb{R}^n$ such that the support of f is contained in $e_1 + C_1$ and the support of g is contained in $e_2 + C_2$. Fix $i = (i_{k+1}, \dots, i_n) \in \mathbb{Z}^{n-k}$ and introduce $i' = (0, \dots, 0, i_{k+1}, \dots, i_n) \in \mathbb{Z}^n$. From C_1 and C_2 being in opposition with respect to the first k variables, it follows by Lemma 5 that $(e_1 - i' + C_1) \cap (e_2 - i' + C_2) \cap (\mathbb{R}^k \times \{0\}^{n-k})$ is bounded. In other words, $(e_1 - i' + C_1) \cap (e_2 - i' + C_2) \cap \pi_2^{-1}(0)$ is bounded, and, after shifting by i' , so is the set $(e_1 + C_1) \cap (e_2 + C_2) \cap \pi_2^{-1}(i)$. Hence, there are at most finitely many vectors $(i_1, \dots, i_k) \in \mathbb{Z}^k$ such that the coefficient of $x_1^{i_1} \dots x_n^{i_n}$ in $f \odot g$ is nonzero.

In the general case, there exist finite families $e_1^{(1)}, \dots, e_1^{(r)}$ and $e_2^{(1)}, \dots, e_2^{(s)}$ such that the support of f is contained in $\bigcup_{i=1}^r e_1^{(i)} + C_1$ and such that the support of g is contained in $\bigcup_{i=1}^s e_2^{(i)} + C_2$. Given a fixed $i = (i_{k+1}, \dots, i_n) \in \mathbb{Z}^{n-k}$ again, the finiteness of the number of vectors $(i_1, \dots, i_k) \in \mathbb{Z}^k$ such that the coefficient of $x_1^{i_1} \dots x_n^{i_n}$ in $f \odot g$ is nonzero still holds, by distributivity and the previous argument.

The constraints on the α_i for a well-defined substitution follow in all cases.

(ii) The proof is like the case (i), with minimal changes: considering $e_1 \in \mathbb{R}^k$ instead of $e_1 \in \mathbb{R}^n$, replacing C_1 by $C_1 \times \{0\}^{n-k}$ in the use of Lemma 5, and replacing $f \odot g$ by $f \underset{k \leq n}{\odot} g$ in the conclusion.

(iii) The proof is like the case (i), with minimal changes: considering $e_2 \in \mathbb{R}^k$ instead of $e_2 \in \mathbb{R}^n$, replacing C_2 by $C_2 \times \{0\}^{n-k}$ in the use of Lemma 5, and replacing $f \odot g$ by $f \underset{n \geq k}{\odot} g$ in the conclusion. \square

The following lemma generalizes Lemma 2 to cones in opposition.

Lemma 7. For any $m \geq k \geq 0$, let $\tau: \mathbb{R}^m \rightarrow \mathbb{R}^{m-k}$ be the projection to the last $m - k$ coordinates.

(i) Given two cones $C_1, C_2 \subseteq \mathbb{R}^n$ that are in opposition with respect to the first k variables, the set $\tau(C_1 \star C_2) = \{(\pi_2(u), v - u) : u \in C_1, v \in C_2\} \subseteq \mathbb{R}^{(n-k)+n}$ is a line-free cone.

(ii) Given two cones $C_1 \subseteq \mathbb{R}^k$ and $C_2 \subseteq \mathbb{R}^n$ such that $C_1 \times \{0\}^{n-k}$ and C_2 are in opposition with respect to the first k variables, the set $\tau(C_1 \underset{k \leq n}{\star} C_2) = \{(\pi_2(v), \pi_1(v) - u) : u \in C_1, v \in C_2\} \subseteq \mathbb{R}^{k+n}$ is a line-free cone.

(iii) Given two cones $C_1 \subseteq \mathbb{R}^n$ and $C_2 \subseteq \mathbb{R}^k$ such that C_1 and $C_2 \times \{0\}^{n-k}$ are in opposition with respect to the first k variables, the set $\tau(C_1 \underset{n \geq k}{\star} C_2) = \{(\pi_2(u), v - \pi_1(u)) : u \in C_1, v \in C_2\} \subseteq \mathbb{R}^{n+k}$ is a line-free cone.

Proof. (i) The proof that $\tau(C_1 \star C_2)$ is a cone is similar to the proof for the case of $C_1 \star C_2$ in Lemma 2. We show that it is line-free: assume there is $w \in \tau(C_1 \star C_2)$ such that $-w \in \tau(C_1 \star C_2)$; we proceed to prove that $w = 0$. By assumption, there exist $u, u' \in C_1$ and $v, v' \in C_2$ such that $w = (\pi_2(u), v - u)$ and $-w = (\pi_2(u'), v' - u')$, and therefore such that $(\pi_2(u), v - u) = -(\pi_2(u'), v' - u')$. Then $\pi_2(u) = -\pi_2(u')$ implies $\pi_2(u) = \pi_2(u') = 0$, because $\pi_2(C_1)$ is line-free. Next, $v - u = -(v' - u')$ implies $\pi_2(v - u) = \pi_2(-(v' - u'))$, and therefore $\pi_2(v) = -\pi_2(v')$, because $\pi_2(u) = \pi_2(u') = 0$. It follows that $\pi_2(v) = \pi_2(v') = 0$ because $\pi_2(C_2)$ is line-free. Next, observe that $v + v' \in C_2 \cap (\mathbb{R}^k \times \{0\}^{n-k})$, because $\pi_2(v) = \pi_2(v') = 0$, and that $u + u' \in C_1 \cap (\mathbb{R}^k \times \{0\}^{n-k})$, because $\pi_2(u) = \pi_2(u') = 0$. From $v - u = -(v' - u')$ follows $v + v' = u + u' \in C_1 \cap C_2 \cap (\mathbb{R}^k \times \{0\}^{n-k})$. As C_1 and C_2 are in opposition with respect to the first k variables, both $v + v'$ and $u + u'$ are zero. Since C_1 , resp. C_2 , is line-free, this finally implies that $u = u' = 0$, resp. that $v = v' = 0$, thus that $w = 0$.

(ii) The proof that $\tau(C_1 \underset{k \leq n}{\star} C_2)$ is a cone is similar to the proof for the case of $C_1 \underset{k \leq n}{\star} C_2$ in Lemma 2.

We show that it is line-free: assume there is $w \in \tau(C_1 \underset{k \leq n}{\star} C_2)$ such that $-w \in \tau(C_1 \underset{k \leq n}{\star} C_2)$; we proceed to prove that $w = 0$. By assumption, there exist $u, u' \in C_1$ and $v, v' \in C_2$ such that $w = (\pi_2(v), \pi_1(v) - u)$ and $-w = (\pi_2(v'), \pi_1(v') - u')$, and therefore such that $(\pi_2(v), \pi_1(v) - u) = -(\pi_2(v'), \pi_1(v') - u')$. Then $\pi_2(v) = -\pi_2(v')$ implies $\pi_2(v) = \pi_2(v') = 0$, because $\pi_2(C_2)$ is line-free. Next, $\pi_1(v) - u = -(\pi_1(v') - u')$ implies $\pi_1(v + v') = u + u'$. Therefore, $v + v' = \pi_1(v + v') + \pi_2(v + v') = (u + u') + 0$. As $C_1 \times \{0\}^{n-k}$ and C_2 are in opposition with respect to the first k variables, both $v + v'$ and $u + u'$ are zero. Since C_1 , resp. C_2 , is line-free, this finally implies that $u = u' = 0$, resp. that $v = v' = 0$, thus that $w = 0$.

(iii) The proof that $\tau(C_1 \star_{n \geq k} C_2)$ is a cone is similar to the proof for the case of $C_1 \star_{n \geq k} C_2$ in Lemma 2.

We show that it is line-free: assume there is $w \in \tau(C_1 \star_{n \geq k} C_2)$ such that $-w \in \tau(C_1 \star_{n \geq k} C_2)$; we proceed to prove that $w = 0$. By assumption, there exist $u, u' \in C_1$ and $v, v' \in C_2$ such that $w = (\pi_2(u), v - \pi_1(u))$ and $-w = (\pi_2(u'), v' - \pi_1(u'))$, and therefore such that $(\pi_2(u), v - \pi_1(u)) = -(\pi_2(u'), v' - \pi_1(u'))$. Then $\pi_2(u) = -\pi_2(u')$ implies $\pi_2(u) = \pi_2(u') = 0$, because $\pi_2(C_1)$ is line-free. Next, $v - \pi_1(u) = -(v' - \pi_1(u'))$ implies $\pi_1(u + u') = v + v'$. Therefore, $u + u' = \pi_1(u + u') + \pi_2(u + u') = (v + v') + 0$. As C_1 and $C_2 \times \{0\}^{n-k}$ are in opposition with respect to the first k variables, both $u + u'$ and $v + v'$ are zero. Since C_1 , resp. C_2 , is line-free, this finally implies that $u = u' = 0$, resp. that $v = v' = 0$, thus that $w = 0$. \square

Hence, for any two line-free cones C_1, C_2 of \mathbb{R}^n or \mathbb{R}^k and with the relevant cones in opposition with respect to the first k variables (as required), each of $\mathbb{K}_{\tau(C_1 \star C_2)} \langle \langle X_{n \setminus k}, Y_n \rangle \rangle$, $\mathbb{K}_{\tau(C_1 \star_{k \leq n} C_2)} \langle \langle X_{n \setminus k}, Y_k \rangle \rangle$, and $\mathbb{K}_{\tau(C_1 \star_{n \geq k} C_2)} \langle \langle X_{n \setminus k}, Y_k \rangle \rangle$ is a well-defined ring, and a subring of, respectively, $\mathbb{K}_{C_1 \star C_2} \langle \langle X_n, Y_n \rangle \rangle$, $\mathbb{K}_{C_1 \star_{k \leq n} C_2} \langle \langle X_n, Y_k \rangle \rangle$, or $\mathbb{K}_{C_1 \star_{n \geq k} C_2} \langle \langle X_n, Y_k \rangle \rangle$.

The following lemma can now be viewed as a generalization of Lemma 3 above.

Lemma 8. (i) Let $C_1, C_2 \subseteq \mathbb{R}^n$ be two cones that are in opposition with respect to the first k variables. Let $f \in \mathbb{K}_{C_1} \langle \langle X_n \rangle \rangle$, $g \in \mathbb{K}_{C_2} \langle \langle X_n \rangle \rangle$, and $\alpha_1, \dots, \alpha_k, \beta_1, \dots, \beta_k \in \mathbb{K}$ satisfy: for each $1 \leq j \leq k$, $\alpha_j \neq 0$ if $H_j^n(f)$ and $\beta_j \neq 0$ if $H_j^n(g)$. Then,

$$(f \odot g) \Big|_{x_1 = \alpha_1 \beta_1, \dots, x_k = \alpha_k \beta_k} = \text{Res}_{y_1, \dots, y_n} \frac{1}{y_1 \cdots y_n} f \left(\frac{\alpha_1}{y_1}, \dots, \frac{\alpha_k}{y_k}, \frac{x_{k+1}}{y_{k+1}}, \dots, \frac{x_n}{y_n} \right) g(\beta_1 y_1, \dots, \beta_k y_k, y_{k+1}, \dots, y_n), \quad (21)$$

where the argument of the residue is understood as a product in $\mathbb{K}_{\tau(C_1 \star C_2)} \langle \langle X_{n \setminus k}, Y_n \rangle \rangle$.

(ii) For $k \leq n$, let $C_1 \subseteq \mathbb{R}^k$ and $C_2 \subseteq \mathbb{R}^n$ be two line-free cones, such that $C_1 \times \{0\}^{n-k}$ and C_2 are in opposition with respect to the first k variables. Let $f \in \mathbb{K}_{C_1} \langle \langle X_k \rangle \rangle$, $g \in \mathbb{K}_{C_2} \langle \langle X_n \rangle \rangle$, and $\alpha_1, \dots, \alpha_k, \beta_1, \dots, \beta_k \in \mathbb{K}$ satisfy: for each $1 \leq j \leq k$, $\alpha_j \neq 0$ if $H_j^k(f)$ and $\beta_j \neq 0$ if $H_j^n(g)$. Then,

$$(f \odot_{k \leq n} g) \Big|_{x_1 = \alpha_1 \beta_1, \dots, x_k = \alpha_k \beta_k} = \text{Res}_{y_1, \dots, y_k} \frac{1}{y_1 \cdots y_k} f \left(\frac{\alpha_1}{y_1}, \dots, \frac{\alpha_k}{y_k} \right) g(\beta_1 y_1, \dots, \beta_k y_k, x_{k+1}, \dots, x_n), \quad (22)$$

where the argument of the residue is understood as a product in $\mathbb{K}_{\tau(C_1 \star_{k \leq n} C_2)} \langle \langle X_{n \setminus k}, Y_k \rangle \rangle$.

(iii) For $k \leq n$, let $C_1 \subseteq \mathbb{R}^n$ and $C_2 \subseteq \mathbb{R}^k$ be two line-free cones, such that C_1 and $C_2 \times \{0\}^{n-k}$ are in opposition with respect to the first k variables. Let $f \in \mathbb{K}_{C_1} \langle \langle X_n \rangle \rangle$, $g \in \mathbb{K}_{C_2} \langle \langle X_k \rangle \rangle$, and $\alpha_1, \dots, \alpha_k, \beta_1, \dots, \beta_k \in \mathbb{K}$ satisfy: for each $1 \leq j \leq k$, $\alpha_j \neq 0$ if $H_j^n(f)$ and $\beta_j \neq 0$ if $H_j^k(g)$. Then,

$$(f \odot_{n \geq k} g) \Big|_{x_1 = \alpha_1 \beta_1, \dots, x_k = \alpha_k \beta_k} = \text{Res}_{y_1, \dots, y_k} \frac{1}{y_1 \cdots y_k} f \left(\frac{\alpha_1}{y_1}, \dots, \frac{\alpha_k}{y_k}, x_{k+1}, \dots, x_n \right) g(\beta_1 y_1, \dots, \beta_k y_k), \quad (23)$$

where the argument of the residue is understood as a product in $\mathbb{K}_{\tau(C_1 \star_{n \geq k} C_2)} \langle \langle X_{n \setminus k}, Y_k \rangle \rangle$.

Before the proof, observe that (22) and (23) cannot just be obtained as specializations of (21).

Proof. To prove (21), assume that $\alpha_1, \dots, \alpha_k, \beta_1, \dots, \beta_k \in \mathbb{K}$ satisfy: for each $1 \leq j \leq k$, $\alpha_j \neq 0$ if $H_j^n(f)$ and $\beta_j \neq 0$ if $H_j^n(g)$. As a first consequence, for each $1 \leq j \leq k$, $\alpha_j \beta_j \neq 0$ if $H_j^{n \leq n}(f, g)$, so that by Lemma 6, the left-hand side of (21) is well-defined. As a second consequence, $f \left(\frac{\alpha_1}{y_1}, \dots, \frac{\alpha_k}{y_k}, \frac{x_{k+1}}{y_{k+1}}, \dots, \frac{x_n}{y_n} \right)$ and $g(\beta_1 y_1, \dots, \beta_k y_k, y_{k+1}, \dots, y_n)$ are well defined, too, and are thus in $\mathbb{K}_{\tau(C_1 \star C_2)} \langle \langle X_{n \setminus k}, Y_n \rangle \rangle$. In a way similar to (18) in the proof of Lemma 3, we reformulate their product h in the form:

$$h = \frac{1}{y_1 \cdots y_n} \left(\sum_{u_1, \dots, u_n} f_{u_1, \dots, u_n} \frac{\alpha_1^{u_1} \cdots \alpha_k^{u_k} x_{k+1}^{u_{k+1}} \cdots x_n^{u_n}}{y_1^{u_1} \cdots y_n^{u_n}} \right) \left(\sum_{v_1, \dots, v_n} g_{v_1, \dots, v_n} \beta_1^{v_1} \cdots \beta_k^{v_k} y_1^{v_1} \cdots y_n^{v_n} \right) = \sum_{i_1, \dots, i_n} \left(\sum_{(v_1, \dots, v_n) - (u_1, \dots, u_n) = (i_1, \dots, i_n)} f_{u_1, \dots, u_n} g_{v_1, \dots, v_n} (\alpha_1^{u_1} \beta_1^{v_1}) \cdots (\alpha_k^{u_k} \beta_k^{v_k}) x_{k+1}^{u_{k+1}} \cdots x_n^{u_n} \right) y_1^{i_1-1} \cdots y_n^{i_n-1}.$$

Like in Lemma 3, extracting residues with respect to y_1, \dots, y_n selects the term for which $i_1 = \dots = i_n = 0$, forcing $(v_1, \dots, v_n) = (u_1, \dots, u_n)$, which yields (21).

The proofs of (22) and (23) are similar, basing on analogous calculations. \square

Likewise, the following theorem can now be viewed as a generalization of Lemma 4 above. For the proof, we proceed like for the proof of that lemma, invoking Lemma 8 in place of Lemma 3.

Theorem 3. *For every line-free cone $C \subseteq \mathbb{R}^n$ that is in opposition to $\mathbb{R}_{\geq 0}^n$ with respect to the first k coordinates, for every $\phi \in \mathbb{K}_C \langle \langle X_n \rangle \rangle$, and for every $\lambda_1, \dots, \lambda_k \in \mathbb{K}$, we have:*

$$\begin{aligned} & ([x_1^> \cdots x_k^>] \phi) \Big|_{x_1=\lambda_1, \dots, x_k=\lambda_k} \\ &= \text{Res}_{y_1, \dots, y_k} \frac{1}{y_1 \cdots y_k} \left[\frac{\frac{\lambda_1 \cdots \lambda_k}{y_1 \cdots y_k}}{\left(1 - \frac{\lambda_1}{y_1}\right) \cdots \left(1 - \frac{\lambda_k}{y_k}\right)} \right]_{\mathbb{R}_{\geq 0}^k} \phi(y_1, \dots, y_k, x_{k+1}, \dots, x_n) \end{aligned} \quad (24)$$

$$= \text{Res}_{y_1, \dots, y_k} \frac{1}{y_1 \cdots y_k} \phi \left(\frac{1}{y_1}, \dots, \frac{1}{y_k}, x_{k+1}, \dots, x_n \right) \left[\frac{\lambda_1 y_1 \cdots \lambda_k y_k}{(1 - \lambda_1 y_1) \cdots (1 - \lambda_k y_k)} \right]_{\mathbb{R}_{\geq 0}^k}, \quad (25)$$

where the brackets around rational functions in (24) and (25) denote taking their expansions in, respectively, $\mathbb{K}_{\mathbb{R}_{\geq 0}^k} \langle \langle Y_k \rangle \rangle$ and $\mathbb{K}_{\mathbb{R}_{\geq 0}^k} \langle \langle Y_k \rangle \rangle$.

Proof. Fix C' to $\mathbb{R}_{\geq 0}^k$ and $\psi \in \mathbb{K}_{C'} \langle \langle X_k \rangle \rangle$ to the geometric series expansion of $\frac{x_1 \cdots x_k}{(1-x_1) \cdots (1-x_k)}$. The desired positive part can then be represented in two ways as variants of Hadamard products:

$$([x_1^> \cdots x_k^>] \phi) \Big|_{x_1=\lambda_1, \dots, x_k=\lambda_k} = (\psi \underset{k \leq n}{\odot} \phi) \Big|_{x_1=\lambda_1, \dots, x_k=\lambda_k} = (\phi \underset{n \geq k}{\odot} \psi) \Big|_{x_1=\lambda_1, \dots, x_k=\lambda_k}.$$

We then use Lemma 8 twice, which applies as C is in opposition to $\mathbb{R}_{\geq 0}^n$ with respect to the first k coordinates, thus to $C' \times \{0\}^{n-k}$ as well:

- Firstly, Lemma 8 (ii) can be used with $C_1 = C'$, $C_2 = C$, $f = \psi$, $g = \phi$, $\alpha_i = \lambda_i$, $\beta_i = 1$, without restriction on $\lambda_i \in \mathbb{K}$ as ψ involves no monomial with negative exponent. Equation (22) then reads

$$([x_1^> \cdots x_k^>] \phi) \Big|_{x_1=\lambda_1, \dots, x_k=\lambda_k} = \text{Res}_{y_1, \dots, y_k} \frac{1}{y_1 \cdots y_k} \psi \left(\frac{\lambda_1}{y_1}, \dots, \frac{\lambda_k}{y_k} \right) \phi(y_1, \dots, y_k, x_{k+1}, \dots, x_n),$$

where $\psi \left(\frac{\lambda_1}{y_1}, \dots, \frac{\lambda_k}{y_k} \right)$ is in $\mathbb{K}_{\tau((C' \times \{0\}^{n-k}) \star_{k \leq n} \{0\}^n)} \langle \langle X_n \setminus k, Y_k \rangle \rangle$, and thus in $\mathbb{K}_{-C'} \langle \langle Y_k \rangle \rangle$, hence (24) and the announced expansion.

- Secondly, Lemma 8 (iii) can be used with $C_1 = C$, $C_2 = C'$, $f = \phi$, $g = \psi$, $\alpha_i = 1$, $\beta_i = \lambda_i$, without restriction on $\lambda_i \in \mathbb{K}$ as ψ involves no monomial with negative exponent. Equation (22) then reads

$$([x_1^> \cdots x_k^>] \phi) \Big|_{x_1=\lambda_1, \dots, x_k=\lambda_k} = \text{Res}_{y_1, \dots, y_k} \frac{1}{y_1 \cdots y_k} \phi \left(\frac{1}{y_1}, \dots, \frac{1}{y_k}, x_{k+1}, \dots, x_n \right) \psi(\lambda_1 y_1, \dots, \lambda_k y_k),$$

where $\psi(\lambda_1 y_1, \dots, \lambda_k y_k)$ is in $\mathbb{K}_{\tau(0 \star_{n \geq k} (C' \times \{0\}^{n-k}))} \langle \langle X_n \setminus k, Y_k \rangle \rangle$, and thus in $\mathbb{K}_{C'} \langle \langle Y_k \rangle \rangle$, hence (25) and the announced expansion. \square

The following variant formulation of Theorem 3 avoids to potentially get a tautology when some of the specialization point is zero.

Theorem 4. *For every line-free cone $C \subseteq \mathbb{R}^n$ that is in opposition to $\mathbb{R}_{\geq 0}^n$ with respect to the first k coordinates, for every $\phi \in \mathbb{K}_C \langle \langle X_n \rangle \rangle$, and for every $\lambda_1, \dots, \lambda_k \in \mathbb{K}$, we have:*

$$\begin{aligned} & \left(\frac{1}{x_1 \cdots x_k} [x_1^> \cdots x_k^>] \phi \right) \Big|_{x_1=\lambda_1, \dots, x_k=\lambda_k} \\ &= \text{Res}_{y_1, \dots, y_k} \frac{1}{y_1 \cdots y_k} \left[\frac{\frac{1}{y_1 \cdots y_k}}{\left(1 - \frac{\lambda_1}{y_1}\right) \cdots \left(1 - \frac{\lambda_k}{y_k}\right)} \right]_{\mathbb{R}_{\geq 0}^k} \phi(y_1, \dots, y_k, x_{k+1}, \dots, x_n) \end{aligned} \quad (26)$$

$$= \text{Res}_{y_1, \dots, y_k} \frac{1}{y_1 \cdots y_k} \phi \left(\frac{1}{y_1}, \dots, \frac{1}{y_k}, x_{k+1}, \dots, x_n \right) \left[\frac{y_1 \cdots y_k}{(1 - \lambda_1 y_1) \cdots (1 - \lambda_k y_k)} \right]_{\mathbb{R}_{\geq 0}^k}, \quad (27)$$

where the brackets around rational functions in (26) and (27) denote taking their expansions in, respectively, $\mathbb{K}_{\mathbb{R}_{\geq 0}^k} \langle \langle Y_k \rangle \rangle$ and $\mathbb{K}_{\mathbb{R}_{\geq 0}^k} \langle \langle Y_k \rangle \rangle$.

Proof. In a way very similar to the proof of Theorem 3, fix C' to $\mathbb{R}_{\geq 0}^k$ and $\tilde{\psi} \in \mathbb{K}_{C'} \langle \langle X_k \rangle \rangle$ to the geometric series expansion of $\frac{1}{(1-x_1)\dots(1-x_k)}$. Introduce as well $\tilde{\phi} = \phi/(x_1 \dots x_k)$. The desired positive part can then be represented in two ways as variants of Hadamard products:

$$\left(\frac{1}{x_1 \dots x_k} [x_1^> \dots x_k^>] \phi \right) \Big|_{x_1=\lambda_1, \dots, x_k=\lambda_k} = (\tilde{\psi} \underset{k \leq n}{\odot} \tilde{\phi}) \Big|_{x_1=\lambda_1, \dots, x_k=\lambda_k} = (\tilde{\phi} \underset{n \geq k}{\odot} \tilde{\psi}) \Big|_{x_1=\lambda_1, \dots, x_k=\lambda_k}.$$

The proof now follows the same lines as for Theorem 3, using Lemma 8 twice: Equations (26) and (27) follow after observing that all specializations of any λ_i (in particular, to 0) are well defined and that a factor $y_1 \dots y_k$ can move freely into or out of a bracket. \square

3.5 Generating series of walks as residues

Now, we want to apply Theorem 4 to find representations as residues of the specializations of a walk series $Q(x, y; t)$ at $x = \alpha$ and $y = \beta$ for numbers α and β : we need this in particular for $(\alpha, \beta) \in \{0, 1\}^2$. For technical reasons, below we will in fact need such representations for more general α and β , namely elements of a field \mathbb{K} of characteristic zero whose elements have zero derivative with respect to x and y .

We will obtain a formula of the form

$$Q(\alpha, \beta; t) = \left(\frac{1}{xy} [x^> y^>] \phi \right) \Big|_{x=\alpha, y=\beta} = \operatorname{Res}_{x,y} \frac{1}{xy} \left[\frac{\bar{x}\bar{y}}{(1-\alpha\bar{x})(1-\beta\bar{y})} \right]_{\mathbb{R}_{\geq 0}^2} \phi(x, y; t) = \operatorname{Res}_{x,y} \frac{1}{xy} \phi(\bar{x}, \bar{y}; t) \left[\frac{xy}{(1-\alpha x)(1-\beta y)} \right]_{\mathbb{R}_{\geq 0}^2}, \quad (28)$$

for some series ϕ to be determined and for residues that are residues of rational functions in $\mathbb{K}(x, y, t)$. Our first step is to express $Q(x, y; t)$ as a positive-part extraction. To this end, Bousquet-Mélou and Mishna provide an appealing formula [11, Prop. 8],

$$xyQ(x, y; t) = [x^>][y^>]R(x, y; t), \quad (29)$$

which represents the walk series as an extraction from the *rational function* $R(x, y; t) = N(x, y)/(1 - tS(x, y))$. Thus, their formula requires some care to be used in our context: we know that rational functions potentially admit several (distinct) positive parts in $\mathbb{Q}[[x, y, t]]$, depending on the cone used to expand it in $\mathbb{Q}^{\mathbb{Z}^3}$. So we need to determine a cone that results in the proper combinatorial representation of Q . Additionally, instead of our direct positive-part extraction $[x^> y^>]$ over $\mathbb{Q}^{\mathbb{Z}^3}$ (and more generally $\mathbb{K}^{\mathbb{Z}^3}$ after we specialize at $x = \alpha$ and $y = \beta$), they use iterated positive-part extractions: a first that operates coefficient-wise on series in t , namely $[y^>] : \mathbb{Q}(x)[y, \bar{y}][[t]] \rightarrow \mathbb{Q}(x)[y][[t]]$; a second that operates coefficient-wise on series in y and t , namely $[x^>] : \mathbb{Q}[x, \bar{x}][y][[t]] \rightarrow \mathbb{Q}[x, y][[t]]$. Then, they make the crucial observation that the (unambiguously defined) coefficients of $[y^>]R$ with respect to y and t are not only in $\mathbb{Q}(x)$, but in fact for the set of R under consideration more specifically in $\mathbb{Q}[x, \bar{x}]$, so that $[x^>]$ can be applied. At this point, it will be sufficient for us to determine a cone C such that

$$[x^> y^>] \phi = [x^>][y^>]R(x, y; t) \quad (30)$$

for the series expansion ϕ of R in $\mathbb{Q}_C \langle \langle x, y, t \rangle \rangle$.

The cone C that will do is the cone Γ generated by the vectors $(1, 1, 1)$, $(1, -1, 1)$, $(-1, 0, 0)$, so that elements of the ring $\mathbb{Q}_\Gamma[[x, y, t]]$ can only involve monomials $x^k y^m t^n$ for $k, m, n \in \mathbb{N}$ satisfying $-n \leq m \leq n$ and $k \leq n$. This cone is line-free, making $S_\Gamma := \mathbb{Q}_\Gamma \langle \langle x, y, t \rangle \rangle$ a well-defined ring. Now, for each step set in Table 1, the product $tS(x, y)$ involves only monomials $x^k y^m t$ satisfying $-1 \leq k, m \leq 1$, making the rational function $1/(1 - tS(x, y))$ admit the expansion $\sum_{n \geq 0} S(x, y)^n t^n$ in S_Γ . In addition, the coefficients $1/(x + \bar{x})$ and $1/(x + 1 + \bar{x})$ that occur in N for some of the step sets admit expansions in S_Γ , so that by ring operations, the rational function R admits an expansion in S_Γ , henceforth denoted $\phi = [R]_\Gamma$. Set $\Gamma' = \mathbb{R}_{\leq 0} \times \{0\}$, another line-free cone, so that we can introduce the ring $S_{\Gamma'} = \mathbb{Q}_{\Gamma'} \langle \langle x, y \rangle \rangle = \mathbb{Q}[[\bar{x}]]\langle \langle x, y, \bar{y} \rangle \rangle$. As vector spaces, we have $S_\Gamma \subset \sum_{j \in \mathbb{Z}} S_{\Gamma'} t^j$. With this observation, $\phi = \sum_{j \in \mathbb{Z}} [NS^j]_{\Gamma'} t^j$, where the brackets are taken in $S_{\Gamma'}$. Therefore, our $[x^> y^>] \phi$ equals Bousquet-Mélou and Mishna's $[x^>][y^>]R$ if and only if for any $j \geq 0$, our $[x^> y^>][NS^j]_{\Gamma'}$ equals their $[x^>][y^>](NS^j)$.

We now fix $j \geq 0$. There are two cases, depending on the step set in Table 1:

- *Cases 1 to 16.* The assumptions of Lemma 9 below are satisfied. The lemma proves that equality holds.
- *Cases 17 to 19.* Both N and S are finite sums in $\mathbb{Q}[x, \bar{x}, y, \bar{y}]$, so $[NS^j]_{\Gamma'}$ is just NS^j , and the equality $[x^>y^>][NS^j]_{\Gamma'} = [x^>][y^>](NS^j)$ holds.

Lemma 9. *Let $N(x, y) = N_1(x)y + N_{-1}(x)\bar{y}$ and $S(x, y) = A_1(x)y + A_0(x) + A_{-1}(x)\bar{y}$ be given by Laurent polynomials N_1, A_1, A_0 , and A_{-1} of $\mathbb{Q}[x, \bar{x}]$, and by a rational function N_{-1} of $\mathbb{Q}(x)$. Under the additional assumption that the denominator of N_{-1} divides A_1 in $\mathbb{Q}[x, \bar{x}]$, then for all $j \in \mathbb{N}$, $[y^>]NS^j \in \mathbb{Q}[x, \bar{x}, y]$ and $[x^>y^>][NS^j]_{\Gamma'} = [x^>][y^>](NS^j)$, where the brackets on the left-hand side are taken with respect to the cone $\Gamma' = \mathbb{R}_{\leq 0} \times \{0\}$ and the parentheses on the right-hand side are in $\mathbb{Q}(x)[y, \bar{y}]$.*

Proof. Write $S = A_1y + \tilde{A}$ where $\tilde{A} \in \mathbb{Q}[x, \bar{x}, \bar{y}]$ and set $P = N_{-1}A_1 \in \mathbb{Q}[x, \bar{x}]$. Writing brackets for expansions in $\mathbb{Q}_{\Gamma'}\langle\langle x, y \rangle\rangle$, we observe that $[N]_{\Gamma'} = N_1y + [N_{-1}]_{\Gamma'}\bar{y}$, $[S]_{\Gamma'} = S$, and $[N_{-1}]_{\Gamma'}A_1$ is in $\mathbb{Q}[x, \bar{x}]$, implying $[N_{-1}]_{\Gamma'}A_1 = N_{-1}A_1 = P$. Given $j \in \mathbb{N}$, a computation yields

$$NS^j = (N_1y + N_{-1}\bar{y})(A_1y + \tilde{A})^j = \left(N_1y(A_1y + \tilde{A})^j + P \sum_{\ell=1}^j \binom{j}{\ell} A_1^{\ell-1} y^{\ell-1} \tilde{A}^{j-\ell} \right) + N_{-1}\bar{y}\tilde{A}^j.$$

The large parenthesis in the right-hand side is in $\mathbb{Q}[x, \bar{x}, y, \bar{y}]$; we call it L . The last term $N_{-1}\bar{y}\tilde{A}^j$ is a polynomial in \bar{y} with no non-negative powers of y , and it is thus in $\mathbb{Q}(x)[\bar{y}]$. Consequently, $[x^>][y^>](NS^j) = [x^>][y^>]L$. A similar computation delivers $[NS^j]_{\Gamma'} = L + [N_{-1}]_{\Gamma'}\bar{y}\tilde{A}^j$, where the Laurent polynomial L is viewed in $\mathbb{Q}_{\Gamma'}\langle\langle x, t \rangle\rangle$. Applying our operator $[x^>y^>]$ kills the last term, returning $[x^>y^>][NS^j]_{\Gamma'} = [x^>y^>]L = [x^>][y^>]L$. The result is proved. \square

The following lemma summarizes our discussion.

Lemma 10. *For any field \mathbb{K} of characteristic zero whose elements have zero derivative with respect to x and y , for any $(\alpha, \beta) \in \mathbb{K}^2$, and for any step set in Table 1, Equation (28) above holds when ϕ is set to the expansion of $R = N/(1 - tS)$ in $\mathbb{Q}_{\Gamma'}\langle\langle x, y, t \rangle\rangle$, for the cone Γ generated by the vectors $(1, 1, 1)$, $(1, -1, 1)$, and $(-1, 0, 0)$.*

Proof. Set $\mathbb{K}, \alpha, \beta, \Gamma$, and ϕ as in the statement. The first identity in (28) follows from (29) and (30); the other identities are proven by Theorem 4, used with the field \mathbb{K} of the present lemma, provided we prove that Γ is in opposition to $\mathbb{R}_{\geq 0}^3$ with respect to the first two variables. To see this, note: (i) that both $\pi_2(\Gamma)$ and $\pi_2(\mathbb{R}_{\geq 0}^3)$ equal $\mathbb{R}_{\geq 0}$ and are thus line-free cones; (ii) that $\Gamma \cap \mathbb{R}_{\geq 0}^3$ is generated by $(1, 1, 1)$, $(1, 0, 1)$, $(0, 1, 1)$, and $(0, 0, 1)$, so that its intersection with $\mathbb{R}^2 \times \{0\}$ reduces to $\{(0, 0, 0)\}$. \square

3.6 Annihilating generating series of walks by creative telescoping

Our general goal is to justify via creative telescoping the computation of a differential equation in t satisfied by the residue with respect to x and y . The objects manipulated during the computation are rational functions, and it is usually taken for granted that results of these calculations can be ported without difficulties to other domains. Here, we want to formally prove this for the present situation. For this section again, \mathbb{K} is a field satisfying the conditions of Lemma 10.

Lemma 11. *Let $F \in \mathbb{K}(x, y, t)$. If there exist $U, V \in \mathbb{K}(x, y, t)$ such that $F = \partial_x U + \partial_y V$, then for any cone C for which F, U, V all admit an expansion, $\text{Res}_{x, y}[F]_C = 0$.*

Proof. From the hypothesis, it follows $[F]_C = \partial_x[U]_C + \partial_y[V]_C$. As the residue of a derivative is zero by Lemma 1, the result is proved by linearity. \square

As in our applications the creative-telescoping certificates U and V will be large and messy expressions, whose manipulation can be costly, it is comfortable to state a refinement of the previous lemma that avoids the need to inspect U and V so as to determine a cone C for which they admit expansions.

Lemma 12. *Let $F \in \mathbb{K}(x, y, t)$. If there exist $U, V \in \mathbb{K}(x, y, t)$ such that $F = \partial_x U + \partial_y V$, then for any line-free cone C for which F admits an expansion, $\text{Res}_{x, y}[F]_C = 0$.*

Proof. Since C is line free, there exists a monomial order \preceq such that C is compatible with \preceq . As F, U and V now admit expansions in the field $\mathbb{K}_{\preceq}\langle\langle x, y, t \rangle\rangle$, there is a cone C' containing C so that all three rational functions admit expansions in $\mathbb{K}_{C'}\langle\langle x, y, t \rangle\rangle$. Finally applying Lemma 11 proves the result. \square

Theorem 5. For some fixed step set \mathcal{S} , let $N(x, y)$ and $S(x, y)$ be as in Table 1, let $R(x, y; t)$ denote $N(x, y)/(1 - tS(x, y)) \in \mathbb{Q}(x, y; t)$, and let $Q(x, y; t) = \sum_{n,i,j=0}^{\infty} q_{i,j;n} x^i y^j t^n \in \mathbb{Q}[x, y][[t]]$ be the corresponding generating function of walks. Let Γ be the cone generated by the vectors $(1, 1, 1)$, $(1, -1, 1)$, and $(-1, 0, 0)$, so that the expansion $[R]_{\Gamma}$ exists in $\mathbb{Q}_{\Gamma}\langle\langle x, y; t \rangle\rangle$ and Q can be identified with its expansion there. For \mathbb{K} a field containing \mathbb{Q} , whose elements have zero derivative with respect to x and y , let $(\alpha, \beta) \in \mathbb{K}^2$ and set F to either of

$$F_1 = \frac{1}{xy} \frac{R(x, y; t)}{(\alpha - x)(\beta - y)} \quad \text{and} \quad F_2 = \frac{R(1/x, 1/y; t)}{(1 - \alpha x)(1 - \beta y)}.$$

Suppose $P \in \mathbb{K}[t]\langle\partial_t\rangle$ and $F, U, V \in \mathbb{K}(x, y; t)$ are such that $P(F) = \partial_x U + \partial_y V$. Then $P(Q(\alpha, \beta; t)) = 0$.

Proof. The line-free cones $C_1 = \tau((\mathbb{R}_{\geq 0}^2 \times \{0\}) \star_{2 \leq 3} \Gamma)$ and $C_2 = \tau(\Gamma \star_{3 \geq 2} (\mathbb{R}_{\geq 0}^2 \times \{0\}))$ are, by the construction of Lemma 10, the cones such that the product in the left-hand residue in (28) takes place in $\mathbb{K}_{C_1}\langle\langle t, x, y \rangle\rangle$ (with indeterminates in this order) while the product in the right-hand residue takes place in $\mathbb{K}_{C_2}\langle\langle t, x, y \rangle\rangle$ (same order). With this notation, Equation (28) rewrites

$$Q(\alpha, \beta; t) = \text{Res}_{x,y}[F_1]_{C_1} = \text{Res}_{x,y}[F_2]_{C_2}. \quad (31)$$

Set i to either of 1 and 2 so that $F = F_i$, then set C to C_i . By Lemma 11 and the expression of $P(F)$ in terms of U and V , we have $\text{Res}_{x,y} P([F]_C) = \text{Res}_{x,y} [P(F)]_C = 0$. Since P is free of x, y , we have $\text{Res}_{x,y} P([F]_C) = P(\text{Res}_{x,y} [F]_C)$. By (31), we have $P(Q(\alpha, \beta; t)) = P(\text{Res}_{x,y} [F]_C) = 0$, as claimed. \square

4 Results

4.1 Obtaining proved hypergeometric formulas

Let \mathbb{K} be a field of characteristic zero whose elements have zero derivative with respect to x and y . To obtain a proven expression for the specialization $Q(\alpha, \beta; t)$ of $Q(x, y; t)$ at $(x, y) = (\alpha, \beta) \in \mathbb{K}^2$, it is sufficient to find:

1. a nonzero operator $P \in \mathbb{K}[t]\langle\partial_t\rangle$ for which $P(Q(\alpha, \beta; t))$ is proved to be 0,
2. a closed-form solution $f(t)$ of P that agrees with $Q(\alpha, \beta; t)$ to a sufficiently high order.

Concerning point 1, Theorem 5 provides such a P by applying creative telescoping. The rational functions U and V that are also part of the output can be discarded, as they are not needed for our application to positive-part extraction. Taking $\mathbb{K} = \mathbb{Q}$, this is already sufficient to obtain formally proved differential equations for the generating functions $Q(t) = Q(1, 1; t)$, which was the initial goal of our work.

To obtain a proven equation for $Q(x, y; t)$, one could in principle follow the same strategy: introduce the bivariate function field $\mathbb{K} = \mathbb{Q}(\alpha, \beta)$, for indeterminates α and β ; use Theorem 5 for this \mathbb{K} ; then replace α and β by x and y at the end. However, this computation would be too slow to work in practice. One could instead consider computing operators $P_{\alpha,\beta} \in \mathbb{Q}[t]\langle\partial_t\rangle$ for sufficiently many points $(\alpha, \beta) \in \mathbb{Q}^2$ and then interpolate into $P \in \mathbb{Q}(x, y)[t]\langle\partial_t\rangle$. This would raise the delicate problem of determining how many interpolation points need to be taken to prove that P is correct.

What does work in practice is to refrain from computing an annihilator P for the complete generating function $Q(x, y; t)$. Instead, we obtain two formally proved differential equations: a first one for $Q(x, 0; t)$, by using creative telescoping over $\mathbb{K} = \mathbb{Q}(\alpha)$ and substituting x for α ; a second one for $Q(0, y; t)$, by using creative telescoping over $\mathbb{K} = \mathbb{Q}(\beta)$ and substituting y for β . In either case, the ground field \mathbb{K} to be used is a univariate rational-function field, which permits calculations to terminate in reasonable time. Although these operators do not lead to an annihilator for $Q(x, y; t)$, they are used below to get a closed form for it.

For all instances of creative telescoping, we effectively used Koutschan's package² `HolonomicFunctions` for Mathematica, then double-checked with Chyzak and Pech's package³ `Mgfun` for Maple.

Concerning point 2, the operators obtained by point 1 all factor miraculously into factors that have order 1 with the exception of the left-most one that can have order 1 or 2, as we could verify using Maple's `DEtools` [`Dfactor`]. At this point, we can capitalize on the work in [16] and obtain ${}_2F_1$ formulas

²<http://www.risc.jku.at/research/combinat/software/ergosum/RISC/HolonomicFunctions.html>

³<http://algo.inria.fr/chyzak/mgfun.html>


for $Q(1, 1; t)$, $Q(x, 0; t)$, and $Q(0, y; t)$ by the method already used in [4]. To this end, we appealed to Imamoglu's implementation⁴ for Maple.

Finally, the kernel equation (first equation in Lemma 4 of [11], which we reproduce here)

$$xy(1 - tS(x, y))Q(x, y; t) = xy - txA_{-1}(x)Q(x, 0; t) - tyB_{-1}(y)Q(0, y; t) + \epsilon tQ(0, 0; t),$$

where A_{-1} and B_{-1} are defined as in Lemma 9 and ϵ is 1 if and only if $(-1, -1) \in \mathcal{S}$ and 0 otherwise, expresses $Q(x, y; t)$ in terms of the obtained formulas for $Q(x, 0; t)$ and $Q(0, y; t)$. As observed in the introduction, all ${}_2F_1$ functions appearing in formulas can be expressed in terms of complete elliptic integrals. Furthermore, for each of the 19 step sets, those functions do not depend on the choice of specialization points α and β , so that $Q(x, y; t)$ involves a single hypergeometric function (up to variations under contiguity and derivatives).

We remark that the nature of the generating functions for our walk models is even more rigid: not only can each model be expressed in terms of a single ${}_2F_1$ function, but, if one can afford increasing degrees and/or algebraic extensions, all models can be expressed in terms of the same single ${}_2F_1$ function, namely ${}_2F_1\left(\begin{smallmatrix} \frac{1}{12} & \frac{5}{12} \\ 1 \end{smallmatrix} \middle| u\right)$. The emergence of this function is no surprise, owing to Takeuchi's classification [48]. This classification establishes connected components in the class of ${}_2F_1$ functions, under simple kinds of transformations. One of the connected components is well represented as the following diagram (see [49, Table 1, 'Classical transformations'] or [48, Section 4, Diagram (1)]):



This diagram should be read as follows: if $(a, b; c)$ and $(a', b'; c')$ are the endpoints of an edge labelled d , with the latter endpoint above the former in the diagram, then

$${}_2F_1\left(\begin{smallmatrix} a & b \\ c \end{smallmatrix} \middle| t\right) = \sqrt[m]{r(t)} \cdot {}_2F_1\left(\begin{smallmatrix} a' & b' \\ c' \end{smallmatrix} \middle| w(t)\right)$$

for some positive integer m , a rational function $w(t)$ of degree d , and another rational function $r(t)$.

Up to integer shifts, all entries in our table are written in terms of ${}_2F_1$ functions involving the boxed parameters of our diagram. If we do not mind increasing the degree of w by a factor 3, 4, or 6, they can all be rewritten in terms of ${}_2F_1\left(\begin{smallmatrix} \frac{1}{12} & \frac{5}{12} \\ 1 \end{smallmatrix} \middle| w\right)$, itself related to elliptic curves and modular forms (see, e.g., [46] or [47]). Alternatively, all of our formulas can be expressed up to algebraic extensions in terms of ${}_2F_1\left(\begin{smallmatrix} \frac{1}{2} & \frac{1}{2} \\ 1 \end{smallmatrix} \middle| w\right)$, itself related to the complete elliptic function $K(k)$.

The connections mentioned above can be made explicit by providing formulas for the main edges in Takeuchi's diagram:

- The hypergeometric series ${}_2F_1\left(\begin{smallmatrix} \frac{1}{2} & \frac{1}{2} \\ 1 \end{smallmatrix} \middle| w\right)$ and ${}_2F_1\left(\begin{smallmatrix} \frac{1}{4} & \frac{3}{4} \\ 1 \end{smallmatrix} \middle| w\right)$ are related by the well-known duplication formula [1, Eq. (3.1.7)]:

$$\left(1 - \frac{u}{2}\right)^{1/2} {}_2F_1\left(\begin{smallmatrix} \frac{1}{2} & \frac{1}{2} \\ 1 \end{smallmatrix} \middle| u\right) = {}_2F_1\left(\begin{smallmatrix} \frac{1}{4} & \frac{3}{4} \\ 1 \end{smallmatrix} \middle| \left(\frac{u}{2-u}\right)^2\right).$$

⁴<http://www.math.fsu.edu/~eimamogl/hypergeometricsols>

- Less well-known formulas [24, Eq. (119) & (126)] result, upon evaluating at $\eta = 1/12$, into

$$(1 + 3u)^{1/4} {}_2F_1\left(\frac{1}{4}, \frac{3}{4} \middle| u\right) = {}_2F_1\left(\frac{1}{12}, \frac{5}{12} \middle| \frac{27u(1-u)^2}{(1+3u)^3}\right),$$

$$\left(1 - \frac{8v}{9}\right)^{1/4} {}_2F_1\left(\frac{1}{3}, \frac{2}{3} \middle| v\right) = {}_2F_1\left(\frac{1}{12}, \frac{5}{12} \middle| \frac{64v^3(1-v)}{(9-8v)^3}\right).$$

This relates ${}_2F_1\left(\frac{1}{4}, \frac{3}{4} \middle| t\right)$ and ${}_2F_1\left(\frac{1}{3}, \frac{2}{3} \middle| t\right)$ via a degree-4 algebraic transformation.

Combining these formulas shows that ${}_2F_1\left(\frac{1}{3}, \frac{2}{3} \middle| t\right)$ can be expressed in terms of ${}_2F_1\left(\frac{1}{2}, \frac{1}{2} \middle| t\right)$ via a degree-4 function in a degree-6 algebraic extension, as indicated in the diagram.

4.2 Algebraic nature of the counting series

To prove the transcendence of one of our counting series $Q(x, y; t)$, it is sufficient to prove the transcendence of one of its evaluations for specific values α of x and β of y —and indeed, the complete generating functions for all cases listed in Table 1 are transcendental as a consequence of $Q(0, 0; t)$ being transcendental. The same holds true for $Q(x, 0; t)$ and $Q(0, y; t)$.

We now describe how we prove transcendence of a given enumerating series $Q(\alpha, \beta; t)$ at specific values for x and y , after we have computed an annihilating operator P for it. Remember from the discussion in Section 4.1 that P factors in the form LL' , where L has order 1 or 2. Define $\tilde{Q}(t) = L'(Q(\alpha, \beta; t))$, which has to be algebraic if $Q(\alpha, \beta; t)$ is algebraic. In each choice of a step set and of α, β in $\{0, 1\}$, the computation of a few terms of the series expansion of Q proves, upon applying L' , that \tilde{Q} is non-zero. Now, if L has order 2, Kovacic's algorithm [31] decides if it has (non-zero) algebraic solutions: if not, this proves that Q cannot be algebraic. The case when L has order 1 is similar, and simpler as solving a first-order linear ODE is very elementary. In our calculations, we used Maple's `DEtools[kovacicssols]` in the second-order case and `DEtools[expssols]` in the first-order case for this test of existence of algebraic solutions of the operators L . This proves the first part of Theorem 2, namely that among the 19×4 combinatorially meaningful specializations $Q(\alpha, \beta; t)$ ($\alpha, \beta \in \{0, 1\}$) of the complete generating function $Q(x, y; t)$, only four cases could possibly be algebraic functions: Case 17 for x and y specialized to 1, and Case 18 for (x, y) specialized to $(0, 1)$, $(1, 0)$, and $(1, 1)$.

In these four cases, we indeed prove the algebraicity of the corresponding counting series. The procedure we followed was simply to use specific algorithms when solving the relevant differential equation $Py = 0$, and note that it admits a basis of algebraic solutions. Explicit expressions in these four cases are then found using the initial terms of the counting series. For instance, in case 18 for $(\alpha, \beta) = (1, 0)$ and for $(\alpha, \beta) = (0, 1)$, the operator P proved to annihilate $Q(1, 0; t)$ and $Q(0, 1; t)$ is

$$P = t^3(2t+1)(6t-1)\partial_t^4 + 4t^2(48t^2+13t-3)\partial_t^3 + 36t(3t+1)(8t-1)\partial_t^2 + (1152t^2+168t-24)\partial_t + 288t + 24,$$

which admits the basis of solutions

$$\left\{ s_1 = \frac{1}{t}, s_2 = \frac{4t^2 - 8t + 1}{t^3}, s_3 = \frac{12t^2 - 1}{t^3}, s_4 = \frac{(2t+1)^{1/2}(1-6t)^{3/2}}{t^3} \right\}.$$

Therefore, $Q(1, 0; t)$ is equal to a linear combination $c_1s_1 + c_2s_2 + c_3s_3 + c_4s_4$ for some constants c_1, c_2, c_3, c_4 . Identification of initial terms using $Q(1, 0; t) = 1 + t + 4t^2 + 12t^3 + O(t^4)$ provides a linear system in the c_i 's, finally proving that $Q(1, 0; t)$ is equal to $s_4 - s_2$. This concludes the proof of Theorem 2.

Note that, for any of the 19 models, the excursions generating functions $Q(0, 0; t)$ could alternatively be proved transcendental by an argument based on asymptotics, similar to the one in [8]: using [15] the coefficient of t^{12n} in $Q(0, 0; t)$ grows like $\kappa\rho^n n^\alpha$ for $\alpha \in \{-3, -4, -5\}$ (see Table 3), and this implies transcendence of $Q(0, 0; t)$ by [19, Theorem D]. By contrast, note that this asymptotic argument is *not* sufficient to prove the transcendence of *all* the other transcendental specializations, as showed for instance by Case 7 at $(0, 1)$ ($\alpha = -3/2$) and at $(1, 1)$ ($\alpha = -1/2$), and by Case 17 at $(1, 0)$ ($\alpha = -5/2$), see Tables 4, 5 and 6.

4.3 Asymptotic formulas for coefficients

This section discusses how to develop asymptotic estimates on the counting coefficients from the closed forms for the generating series. This bases on asymptotics-transfer theorems (developed in Analytic

Combinatorics) to get the results. Not all asymptotic formulas could be proved by our approach, so we report here on a reduction of the proofs of asymptotic formulas to the proof of a number of integral representation of rational constants.

Consider the 19 walk models with prescribed length (and unprescribed end point). What is most difficult is proving the constants (κ below) in front of the pattern ρ^n/n^α , which were initially heuristically discovered by numerical calculations (number recognition) [6].

All of our generating series consist of hypergeometric functions, rational or algebraic factors, and iterated primitivations. In there, hypergeometric terms take the form ${}_2F_1\left(\begin{matrix} a & b \\ c \end{matrix} \middle| w\right)$, with good constraints on a , b , and c , so that enough of their asymptotic behavior (e.g., at $w = 1$) can be derived easily, and where w is a rational or algebraic term in t . Singularities of the generating series can occur at poles or branch point of the cofactors, or at values of t where $w = w(t)$ becomes one of the singularities of the ${}_2F_1$, namely 0, 1, and ∞ .

We first expected that the problem of proving all coefficient asymptotics would be just a problem of Analytic Combinatorics (in the sense of the theory well exposed by Flajolet and Sedgewick in [21, Chap. VI], largely basing on transfer theorems by Flajolet and Odlyzko [20]). It turned out that the part of these theories to extract asymptotics is minimal, most of the difficulties being in computing asymptotic expansions of non-standard representations of functions. (These expansions are then input to transfer theorems.)

What also made the problem more difficult is a broad spectrum of the possible asymptotic phenomena / asymptotic patterns (i.e., what ingredient of the function contributes to the asymptotics). In particular, one situation is recurrent: that when the dominant singularity is caused by an algebraic factor between two primitivation operators. In some cases, this is worsened by a polar part of the integrand in the inner primitivation: this part is necessary and disappears in the formal-power-series expansion after the alternation of primitivations and multiplications by algebraic factors, but to get the constants κ by analytic means, it is easier to remove this polar part in calculations of asymptotic estimates.

In view of this, we have in fact formally proved only some of the constants κ . However, in all the cases, those κ —as well as other constants that appear in the asymptotics—can be formulated as definite integrals of univariate functions obtained from $Q(x, y; t)$. Proving the asymptotic formulas up to the constants is equivalent to proving closed-form evaluations of those integrals. This is exemplified and described now in four typical cases that cover all difficulties that may occur.

- For Case 4, the generating series is:

$$\frac{1}{t} \int_0^t \frac{1}{(1+4u)^3} {}_2F_1\left(\begin{matrix} \frac{3}{2} & \frac{3}{2} \\ 2 \end{matrix} \middle| \frac{16u(1+u)}{(1+4u)^2}\right) du.$$

Its dominant singularity is at $1/8$, where $w = 1$. The asymptotic analysis of the ${}_2F_1$ at 1 (using normalization by (15.8.12) in [42] before using (15.8.10) in [42]⁵) delivers the asymptotic equivalent $\frac{8}{3\pi}8^n$, with proved constant, for the coefficient of the integrand, and $\frac{1}{t} \int$ is responsible for a factor $1/n$, leading to the final proved $\frac{8}{3\pi} \frac{8^n}{n}$.

- For Case 3, the generating series is:

$$\frac{1}{t} \int_0^t \frac{1-2u}{((1+2u)(1+6u))^{3/2}} {}_2F_1\left(\begin{matrix} \frac{3}{2} & \frac{3}{2} \\ 2 \end{matrix} \middle| \frac{16u}{(1+2u)(1+6u)}\right) du.$$

It has two dominant singularities at $+1/6$, where $w = 1$, and at $-1/6$, where $w = -\infty$. The same approach as above, replicated in parallel, delivers two contributions, respectively $\frac{\sqrt{6}}{\pi}6^n$ and $\frac{\sqrt{6}}{4\pi} \frac{6^n}{n}$. Only the first one remains in an asymptotic equivalent, and after $\frac{1}{t} \int$, the asymptotics of the coefficient in the generating series is $\frac{\sqrt{6}}{\pi} \frac{6^n}{n}$.

- For Case 7, the generating series is:

$$\frac{1}{t(t-1)} \int_0^t \frac{u}{(1-4u)^{3/2}} \left\{ 4 + \int_0^v \frac{(1-4v)^{1/2}(\frac{1}{2}+v)}{v^2} \left[1 + \frac{1}{2v(1+2v)(1+4v^2)^{1/2}} \times \right. \right. \\ \left. \left. \left((1-v) {}_2F_1\left(\begin{matrix} \frac{3}{2} & \frac{1}{2} \\ 1 \end{matrix} \middle| \frac{16v^2}{1+4v^2}\right) - (1+v)(1-4v+8v^2) {}_2F_1\left(\begin{matrix} \frac{1}{2} & \frac{1}{2} \\ 1 \end{matrix} \middle| \frac{16v^2}{1+4v^2}\right) \right) \right] dv \right\} du.$$

⁵Alternative references are <http://dlmf.nist.gov/15.8.E12> and <http://dlmf.nist.gov/15.8.E10>, respectively.

Here, we have a linear combination of hypergeometric functions, but this is no problem. The dominant singularity is at $t = 1/4$, and dominates $\frac{1}{2\sqrt{3}}$, where $w = 16t^2/(1 + 4t^2) = 1$. But $1/4$ and $\frac{1}{3\sqrt{3}}$ being very close, this makes numerical evaluations delicate (or, similarly, series expansions at $1/4$ seem to diverge). Call f the innermost integrand. It proved successful to remove from f the singular part $\frac{1}{t^2}$, which is the polar part at the origin, and whose contribution after double integration can be obtained by an easy use of the theory. Proceeding by linearity, we get that the announced asymptotic formula, providing $\frac{4}{3}\sqrt{\frac{1}{\pi}}\frac{4^n}{\sqrt{n}}$, will be proved if and only if we can prove the numerical integration

$$\int_0^{1/4} \frac{(1-4v)^{1/2}(\frac{1}{2}+v)}{v^2} \left[1 + \frac{1}{2v(1+2v)(1+4v^2)^{1/2}} \times \left((1-v)_2F_1\left(\frac{3}{2}, \frac{1}{2} \middle| \frac{16v^2}{1+4v^2}\right) - (1+v)(1-4v+8v^2)_2F_1\left(\frac{1}{2}, \frac{1}{2} \middle| \frac{16v^2}{1+4v^2}\right) \right) \right] - \frac{1}{v^2} dv = -2,$$

which one can check numerically.

- For Case 5, the generating series is:

$$\frac{1}{t(t-1)} \int_0^t \frac{u^2}{(1+u)^{1/2}(1-3u)^{3/2}} \left\{ -7 + \int_0^u \frac{(1-3v)^{1/2}}{v^3(1+v)^{1/2}} \times \left[1 + (1-10v^3)_2F_1\left(\frac{3}{4}, \frac{5}{4} \middle| 64v^4\right) + 6v^3(3-8v+14v^2)_2F_1\left(\frac{5}{4}, \frac{7}{4} \middle| 64v^4\right) \right] dv \right\} du.$$

The situation is similar to Case 7, starting with a dominant singularity at $t = 1/3$, very close to $\frac{1}{\sqrt{8}}$, where $w = 64t^4 = 1$. In this case, removing the polar part of the innermost integrand f at the origin, namely $\frac{2}{t^3} - \frac{4}{t^2}$, is enough to find an equivalent formulation of the result as a numerical integral.

Proceeding by linearity, we get that the announced asymptotic formula, providing $\frac{1}{2}\sqrt{\frac{3}{\pi}}\frac{3^n}{\sqrt{n}}$, will be proved if and only if we can prove the numerical integration

$$\int_0^{1/3} \frac{(1-3v)^{1/2}}{v^3(1+v)^{1/2}} \times \left[1 + (1-10v^3)_2F_1\left(\frac{3}{4}, \frac{5}{4} \middle| 64v^4\right) + 6v^3(3-8v+14v^2)_2F_1\left(\frac{5}{4}, \frac{7}{4} \middle| 64v^4\right) \right] - \frac{2}{v^3} + \frac{4}{v^2} dv = 1,$$

which one can again check numerically.

In all 19×4 cases, our hypergeometric expressions for the counting functions produce similar equalities of integrals, whose correctness implies the asymptotics in Tables 3, 4, 5, and 6 in the Appendix.

4.4 Additional tables

The results of our computations are described in this section, where the most characteristic features of the computed data are provided as tables. For the data in full, we refer our readers to the web page of this article <http://specfun.inria.fr/chyzak/ssw/>.

The hypergeometric series occurring in explicit expressions for $Q(1, 1; t)$ and for $Q(x, y; t)$ are respectively given in Table 2, together with the rational-function substitution in those series. The complete closed forms we obtained for $Q(0, 0; t)$, $Q(0, 1; t)$, $Q(1, 0; t)$, $Q(1, 1; t)$, $Q(x, 0; t)$, $Q(0, y; t)$, and $Q(x, y; t)$ are given on the web site.

Various degrees, the differential order with respect to t , and the maximal size of integers in the annihilating operator and certificates that we computed for each step set can be obtained from the web site as well. Thus, the annihilators—the same as in [6]—are finally proved. Annihilators and certificates can also be downloaded in full form from the web site, both as human-readable Maple sources and as a pre-digested Maple library.

For each of the step set in Table 1, the algebraic or transcendental nature of the series $Q(0, 0; t)$, $Q(0, 1; t)$, $Q(1, 0; t)$, and $Q(1, 1; t)$, as well as asymptotic equivalent for the coefficients of t^n in these series, are respectively given in Tables 3, 4, 5, and 6 in the Appendix. There, ‘N’ denotes a transcendental series and ‘Y’ an algebraic one, and the result has been proved by the method of Section 4.2. Regarding the

status of asymptotic equivalents, the constants in the tables (named κ in Section 4.3) refine (and correct) those provided in [5].

The possibility of several interlaced regimes with same parameters ρ and α but non-zero different κ was first observed and suggested to us by Melczer. By redoing the calculations as in [5], using convergence acceleration of (subsequences of) the sequences $(q_{x,y;n})$ and (q_n) and using the PSLQ algorithm, we numerically guessed those constants κ , confirming the known ones and obtaining the new ones. That [5] could overlook those constants is explained by the nature of the method of convergence acceleration, which in its native form considers values at powers of 2 only, and is thus not able to distinguish regimes for odd vs even n , let alone for various residues modulo 3 or 4. In their recent manuscript [39], Melczer and Wilson proved our guessed constants in all 19×4 cases via analytic combinatorics in several variables.

References

- [1] George E. Andrews, Richard Askey, and Ranjan Roy. *Special functions*, volume 71 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1999.
- [2] Ainhoa Aparicio Monforte and Manuel Kauers. Formal Laurent series in several variables. *Expo. Math.*, 31(4):350–367, 2013.
- [3] Didier Arquès. Dénombrements de chemins dans \mathbf{R}^2 soumis à contraintes. *RAIRO Inform. Théor. Appl.*, 20(4):473–482, 1986.
- [4] Alin Bostan, Frédéric Chyzak, Mark van Hoeij, and Lucien Pech. Explicit formula for the generating series of diagonal 3D rook paths. *Séminaire Lotharingien de Combinatoire*, B66a, 2011.
- [5] Alin Bostan and Manuel Kauers. Unpublished notes, 2008.
- [6] Alin Bostan and Manuel Kauers. Automatic classification of restricted lattice walks. In *21st International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2009)*, Discrete Math. Theor. Comput. Sci. Proc., AK, pages 201–215. Assoc. Discrete Math. Theor. Comput. Sci., Nancy, 2009.
- [7] Alin Bostan and Manuel Kauers. The complete generating function for Gessel walks is algebraic. *Proc. Amer. Math. Soc.*, 138(9):3063–3078, 2010. With an appendix by Mark van Hoeij.
- [8] Alin Bostan, Kilian Raschel, and Bruno Salvy. Non-D-finite excursions in the quarter plane. *J. Combin. Theory Ser. A*, 121:45–63, 2014.
- [9] Mireille Bousquet-Mélou. Counting walks in the quarter plane. In *Mathematics and computer science, II (Versailles, 2002)*, Trends Math., pages 49–67. Birkhäuser, Basel, 2002.
- [10] Mireille Bousquet-Mélou. Walks in the quarter plane: Kreweras’ algebraic model. *Ann. Appl. Probab.*, 15(2):1451–1491, 2005.
- [11] Mireille Bousquet-Mélou and Marni Mishna. Walks with small steps in the quarter plane. In *Algorithmic probability and combinatorics*, volume 520 of *Contemp. Math.*, pages 1–39. Amer. Math. Soc., Providence, RI, 2010.
- [12] Mireille Bousquet-Mélou and Marko Petkovšek. Walks confined in a quadrant are not always D-finite. *Theoret. Comput. Sci.*, 307(2):257–276, 2003. Random generation of combinatorial objects and bijective combinatorics.
- [13] Pierre Cartier. Démonstration “automatique” d’identités et fonctions hypergéométriques (d’après D. Zeilberger). *Astérisque*, (206):Exp. No. 746, 3, 41–91, 1992. Séminaire Bourbaki, Vol. 1991/92.
- [14] Frédéric Chyzak. An extension of Zeilberger’s fast algorithm to general holonomic functions. *Discrete Math.*, 217(1-3):115–134, 2000.
- [15] Denis Denisov and Vitali Wachtel. Random walks in cones. *Ann. Probab.*, 43(3):992–1044, 2015.
- [16] Tingting Fang and Mark van Hoeij. 2-descent for second order linear differential equations. In *ISSAC 2011—Proceedings of the 36th International Symposium on Symbolic and Algebraic Computation*, pages 107–114. ACM, New York, 2011.

- [17] G. Fayolle and K. Raschel. On the holonomy or algebraicity of generating functions counting lattice walks in the quarter-plane. *Markov Process. Related Fields*, 16(3):485–496, 2010.
- [18] Guy Fayolle, Roudolf Iasnogorodski, and Vadim Malyshev. *Random walks in the quarter-plane*, volume 40 of *Applications of Mathematics (New York)*. Springer-Verlag, Berlin, 1999. Algebraic methods, boundary value problems and applications.
- [19] Philippe Flajolet. Analytic models and ambiguity of context-free languages. *Theoret. Comput. Sci.*, 49(2-3):283–309, 1987. Twelfth international colloquium on automata, languages and programming (Nafplion, 1985).
- [20] Philippe Flajolet and Andrew Odlyzko. Singularity analysis of generating functions. *SIAM J. Discrete Math.*, 3(2):216–240, 1990.
- [21] Philippe Flajolet and Robert Sedgewick. *Analytic combinatorics*. Cambridge University Press, Cambridge, 2009.
- [22] Ira M. Gessel. A probabilistic method for lattice path enumeration. *J. Statist. Plann. Inference*, 14(1):49–58, 1986.
- [23] Ira M. Gessel and Doron Zeilberger. Random walk in a Weyl chamber. *Proc. Amer. Math. Soc.*, 115(1):27–31, 1992.
- [24] Édouard Goursat. Sur l'équation différentielle linéaire qui admet pour intégrale la série hypergéométrique. *Ann. Sci. École Norm. Sup. (2)*, 10:3–142, 1881.
- [25] Dominique Gouyou-Beauchamps. Chemins sous-diagonaux et tableaux de Young. In *Combinatoire énumérative (Montreal, Que., 1985/Quebec, Que., 1985)*, volume 1234 of *Lecture Notes in Math.*, pages 112–125. Springer, Berlin, 1986.
- [26] Richard K. Guy, C. Krattenthaler, and Bruce E. Sagan. Lattice paths, reflections, & dimension-changing bijections. *Ars Combin.*, 34:3–15, 1992.
- [27] Mark van Hoeij. Factorization of differential operators with rational functions coefficients. *J. Symbolic Comput.*, 24(5):537–561, 1997.
- [28] Mark van Hoeij and Vijay Jung Kunwar. Classifying (near)-Belyi maps with five exceptional points. Technical Report 1604.08158, arXiv, 2016. Available at <http://arxiv.org/abs/1604.08158>.
- [29] Katherine Humphreys. A history and a survey of lattice path enumeration. *Journal of Statistical Planning and Inference*, 140(8):2237–2254, 2010.
- [30] Christoph Koutschan. A fast approach to creative telescoping. *Math. Comput. Sci.*, 4(2-3):259–266, 2010.
- [31] Jerald J. Kovacic. An algorithm for solving second order linear homogeneous differential equations. *J. Symbolic Comput.*, 2(1):3–43, 1986.
- [32] Christian Krattenthaler. Lattice path enumeration. In *Handbook of enumerative combinatorics*, Discrete Math. Appl. (Boca Raton), pages 589–678. CRC Press, Boca Raton, FL, 2015.
- [33] Germain Kreweeras. Sur une classe de problèmes liés au treillis des partitions d'entiers. *Cahiers du B.U.R.O.*, 6:5–105, 1965.
- [34] Irina Kurkova and Kilian Raschel. On the functions counting walks with small steps in the quarter plane. *Publ. Math. Inst. Hautes Études Sci.*, 116:69–114, 2012.
- [35] Pierre Lairez. Computing periods of rational integrals. *Math. Comp.*, 85(300):1719–1752, 2016.
- [36] L. Lipshitz. The diagonal of a D -finite power series is D -finite. *J. Algebra*, 113(2):373–378, 1988.
- [37] Stephen Melczer and Marni Mishna. Enumerating lattice paths with symmetries through multivariate diagonals. In *AofA 2014, Paris, France*, Discrete Math. Theor. Comput. Sci. Proc., AK, pages 1–12. 2014.

- [38] Stephen Melczer and Marni Mishna. Singularity analysis via the iterated kernel method. *Combin. Probab. Comput.*, 23(5):861–888, 2014.
- [39] Stephen Melczer and Mark C. Wilson. Asymptotics of lattice walks via analytic combinatorics in several variables. Technical Report 1511.02527, arXiv, 2015. Available at <http://arxiv.org/abs/1511.02527>.
- [40] M. Mishna and A. Rechnitzer. Two non-holonomic lattice walks in the quarter plane. *Theoretical Computer Science*, 410(38–40):3616–3630, 2009.
- [41] Marni Mishna. Classifying lattice walks restricted to the quarter plane. *J. Combin. Theory Ser. A*, 116(2):460–477, 2009.
- [42] Frank W. J. Olver, Daniel W. Lozier, Ronald F. Boisvert, and Charles W. Clark, editors. *NIST handbook of mathematical functions*. U.S. Department of Commerce, National Institute of Standards and Technology, Washington, DC; Cambridge University Press, Cambridge, 2010.
- [43] Kilian Raschel. Counting walks in a quadrant: a unified approach via boundary value problems. *J. Eur. Math. Soc. (JEMS)*, 14(3):749–777, 2012.
- [44] Amitai Regev. Asymptotic values for degrees associated with strips of Young diagrams. *Adv. in Math.*, 41(2):115–136, 1981.
- [45] N. J. A. Sloane. *The On-Line Encyclopedia of Integer Sequences*. 2010. Published electronically at <https://oeis.org>.
- [46] Jan Stienstra and Frits Beukers. On the Picard-Fuchs equation and the formal Brauer group of certain elliptic $K3$ -surfaces. *Math. Ann.*, 271(2):269–304, 1985.
- [47] P. F. Stiller. Classical automorphic forms and hypergeometric functions. *J. Number Theory*, 28(2):219–232, 1988.
- [48] Kisao Takeuchi. Commensurability classes of arithmetic triangle groups. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.*, 24(1):201–212, 1977.
- [49] Raimundas Vidūnas. Transformations of some Gauss hypergeometric functions. *J. Comput. Appl. Math.*, 178(1-2):473–487, 2005.
- [50] Doron Zeilberger. A holonomic systems approach to special functions identities. *J. Comput. Appl. Math.*, 32(3):321–368, 1990.
- [51] Doron Zeilberger. The method of creative telescoping. *J. Symbolic Comput.*, 11(3):195–204, 1991.

Appendix

The tables on the next pages gather information described in Section 4.4.

	OEIS	S	alg	equiv
1	A005568		N	$\begin{cases} \frac{32}{\pi} \frac{4^n}{n^3} & (n = 2p) \\ 0 & (n = 2p + 1) \end{cases}$
2	A001246		N	$\begin{cases} \frac{8}{\pi} \frac{4^n}{n^3} & (n = 2p) \\ 0 & (n = 2p + 1) \end{cases}$
3	A151362		N	$\begin{cases} \frac{3\sqrt{6}}{\pi} \frac{6^n}{n^3} & (n = 2p) \\ 0 & (n = 2p + 1) \end{cases}$
4	A172361		N	$\frac{128}{27\pi} \frac{8^n}{n^3}$
5	A151332		N	$\begin{cases} \frac{16\sqrt{2}}{\pi} \frac{(2\sqrt{2})^n}{n^3} & (n = 4p) \\ 0 & (n = 4p + 1) \\ 0 & (n = 4p + 2) \\ 0 & (n = 4p + 3) \end{cases}$
6	A151357		N	$\frac{2A^{3/2}}{\pi} \frac{(2A)^n}{n^3}$
7	A151341		N	$\begin{cases} \frac{12\sqrt{3}}{\pi} \frac{(2\sqrt{3})^n}{n^3} & (n = 2p) \\ 0 & (n = 2p + 1) \end{cases}$
8	A151368		N	$\frac{2B^{3/2}}{\pi} \frac{(2B)^n}{n^3}$
9	A151345		N	$\begin{cases} \frac{24\sqrt{30}}{25\pi} \frac{(2\sqrt{6})^n}{n^3} & (n = 2p) \\ 0 & (n = 2p + 1) \end{cases}$
10	A151370		N	$\frac{2\mu^3 C^{3/2}}{\pi} \frac{(2C)^n}{n^3}$
11	A151341		N	$\begin{cases} \frac{12\sqrt{3}}{\pi} \frac{(2\sqrt{3})^n}{n^3} & (n = 2p) \\ 0 & (n = 2p + 1) \end{cases}$
12	A151368		N	$\frac{2B^{3/2}}{\pi} \frac{(2B)^n}{n^3}$
13	A151345		N	$\begin{cases} \frac{24\sqrt{30}}{25\pi} \frac{(2\sqrt{6})^n}{n^3} & (n = 2p) \\ 0 & (n = 2p + 1) \end{cases}$
14	A151370		N	$\frac{2\mu^3 C^{3/2}}{\pi} \frac{(2C)^n}{n^3}$
15	A151332		N	$\begin{cases} \frac{16\sqrt{2}}{\pi} \frac{(2\sqrt{2})^n}{n^3} & (n = 4p) \\ 0 & (n = 4p + 1) \\ 0 & (n = 4p + 2) \\ 0 & (n = 4p + 3) \end{cases}$
16	A151357		N	$\frac{2A^{3/2}}{\pi} \frac{(2A)^n}{n^3}$
17	A151334		N	$\begin{cases} \frac{81\sqrt{3}}{\pi} \frac{3^n}{n^4} & (n = 3p) \\ 0 & (n = 3p + 1) \\ 0 & (n = 3p + 2) \end{cases}$
18	A151366		N	$\frac{27\sqrt{3}}{\pi} \frac{6^n}{n^4}$
19	A138349		N	$\begin{cases} \frac{768}{\pi} \frac{4^n}{n^5} & (n = 2p) \\ 0 & (n = 2p + 1) \end{cases}$

Table 3: Nature of the generating series for $(x, y) = (0, 0)$ and coefficient asymptotics

$$A = 1 + \sqrt{2}, \quad B = 1 + \sqrt{3}, \quad C = 1 + \sqrt{6}, \quad \lambda = 7 + 3\sqrt{6}, \quad \mu = \sqrt{\frac{4\sqrt{6}-1}{19}}$$

	OEIS	S	alg	equiv
1	A005558		N	$\frac{8}{\pi} \frac{4^n}{n^2}$
2	A151392		N	$\begin{cases} \frac{4}{\pi} \frac{4^n}{n^2} & (n = 2p) \\ 0 & (n = 2p + 1) \end{cases}$
3	A151478		N	$\frac{3\sqrt{6}}{2\pi} \frac{6^n}{n^2}$
4	A151496		N	$\frac{32}{9\pi} \frac{8^n}{n^2}$
5	A151380		N	$\frac{3}{4} \sqrt{\frac{3}{\pi}} \frac{3^n}{n^{3/2}}$
6	A151450		N	$\frac{5}{16} \sqrt{\frac{10}{\pi}} \frac{5^n}{n^{3/2}}$
7	A148790		N	$\frac{8}{3\sqrt{\pi}} \frac{4^n}{n^{3/2}}$
8	A151485		N	$\sqrt{\frac{3}{\pi}} \frac{6^n}{n^{3/2}}$
9	A151440		N	$\frac{5}{24} \sqrt{\frac{10}{\pi}} \frac{5^n}{n^{3/2}}$
10	A151493		N	$\frac{7}{54} \sqrt{\frac{21}{\pi}} \frac{7^n}{n^{3/2}}$
11	A151394		N	$\begin{cases} \frac{36\sqrt{3}}{\pi} \frac{(2\sqrt{3})^n}{n^3} & (n = 2p) \\ \frac{54}{\pi} \frac{(2\sqrt{3})^n}{n^3} & (n = 2p + 1) \end{cases}$
12	A151472		N	$\frac{3B^{7/2}}{2\pi} \frac{(2B)^n}{n^3}$
13	A151437		N	$\begin{cases} \frac{72\sqrt{30}}{5\pi} \frac{(2\sqrt{6})^n}{n^3} & (n = 2p) \\ \frac{864\sqrt{5}}{25\pi} \frac{(2\sqrt{6})^n}{n^3} & (n = 2p + 1) \end{cases}$
14	A151492		N	$\frac{6\lambda\mu^3 C^{5/2}}{5\pi} \frac{(2C)^n}{n^3}$
15	A151375		N	$\begin{cases} \frac{448\sqrt{2}}{9\pi} \frac{(2\sqrt{2})^n}{n^3} & (n = 4p) \\ \frac{640}{9\pi} \frac{(2\sqrt{2})^n}{n^3} & (n = 4p + 1) \\ \frac{416\sqrt{2}}{9\pi} \frac{(2\sqrt{2})^n}{n^3} & (n = 4p + 2) \\ \frac{512}{9\pi} \frac{(2\sqrt{2})^n}{n^3} & (n = 4p + 3) \end{cases}$
16	A151430		N	$\frac{4A^{7/2}}{\pi} \frac{(2A)^n}{n^3}$
17	A151378		N	$\frac{27}{8} \sqrt{\frac{3}{\pi}} \frac{3^n}{n^{5/2}}$
18	A151483		Y	$\frac{27}{8} \sqrt{\frac{3}{\pi}} \frac{6^n}{n^{5/2}}$
19	A005568		N	$\begin{cases} \frac{32}{\pi} \frac{4^n}{n^3} & (n = 2p) \\ 0 & (n = 2p + 1) \end{cases}$

Table 4: Nature of the generating series for $(x, y) = (0, 1)$ and coefficient asymptotics

$$A = 1 + \sqrt{2}, \quad B = 1 + \sqrt{3}, \quad C = 1 + \sqrt{6}, \quad \lambda = 7 + 3\sqrt{6}, \quad \mu = \sqrt{\frac{4\sqrt{6}-1}{19}}$$

	OEIS	\mathcal{S}	alg	equiv
1	A005558		N	$\frac{8}{\pi} \frac{4^n}{n^2}$
2	A151392		N	$\begin{cases} \frac{4}{\pi} \frac{4^n}{n^2} & (n = 2p) \\ 0 & (n = 2p + 1) \end{cases}$
3	A151471		N	$\begin{cases} \frac{2\sqrt{6}}{\pi} \frac{6^n}{n^2} & (n = 2p) \\ 0 & (n = 2p + 1) \end{cases}$
4	A151496		N	$\frac{32}{9\pi} \frac{8^n}{n^2}$
5	A151379		N	$\begin{cases} \frac{4\sqrt{2}}{\pi} \frac{(2\sqrt{2})^n}{n^2} & (n = 2p) \\ 0 & (n = 2p + 1) \end{cases}$
6	A148934		N	$\frac{\sqrt{2}A^{3/2}}{\pi} \frac{(2A)^n}{n^2}$
7	A151410		N	$\begin{cases} \frac{4\sqrt{3}}{\pi} \frac{(2\sqrt{3})^n}{n^2} & (n = 2p) \\ 0 & (n = 2p + 1) \end{cases}$
8	A151464		N	$\frac{2B^{3/2}\sqrt{3}}{3\pi} \frac{(2B)^n}{n^2}$
9	A151423		N	$\begin{cases} \frac{4\sqrt{30}}{5\pi} \frac{(2\sqrt{6})^n}{n^2} & (n = 2p) \\ 0 & (n = 2p + 1) \end{cases}$
10	A151490		N	$\frac{\sqrt{6}\mu C^{3/2}}{3\pi} \frac{(2C)^n}{n^2}$
11	A151410		N	$\begin{cases} \frac{4\sqrt{3}}{\pi} \frac{(2\sqrt{3})^n}{n^2} & (n = 2p) \\ 0 & (n = 2p + 1) \end{cases}$
12	A151464		N	$\frac{2B^{3/2}\sqrt{3}}{3\pi} \frac{(2B)^n}{n^2}$
13	A151423		N	$\begin{cases} \frac{4\sqrt{30}}{5\pi} \frac{(2\sqrt{6})^n}{n^2} & (n = 2p) \\ 0 & (n = 2p + 1) \end{cases}$
14	A151490		N	$\frac{\sqrt{6}\mu C^{3/2}}{3\pi} \frac{(2C)^n}{n^2}$
15	A151379		N	$\begin{cases} \frac{4\sqrt{2}}{\pi} \frac{(2\sqrt{2})^n}{n^2} & (n = 2p) \\ 0 & (n = 2p + 1) \end{cases}$
16	A148934		N	$\frac{\sqrt{2}A^{3/2}}{\pi} \frac{(2A)^n}{n^2}$
17	A151497		N	$\frac{27}{8} \sqrt{\frac{3}{\pi}} \frac{3^n}{n^{5/2}}$
18	A151483		Y	$\frac{27}{8} \sqrt{\frac{3}{\pi}} \frac{6^n}{n^{5/2}}$
19	A005817		N	$\frac{32}{\pi} \frac{4^n}{n^3}$

Table 5: Nature of the generating series for $(x, y) = (1, 0)$ and coefficient asymptotics

$$A = 1 + \sqrt{2}, \quad B = 1 + \sqrt{3}, \quad C = 1 + \sqrt{6}, \quad \lambda = 7 + 3\sqrt{6}, \quad \mu = \sqrt{\frac{4\sqrt{6}-1}{19}}$$

	OEIS	\mathcal{S}	alg	equiv
1	A005566		N	$\frac{4}{\pi} \frac{4^n}{n}$
2	A018224		N	$\frac{2}{\pi} \frac{4^n}{n}$
3	A151312		N	$\frac{\sqrt{6}}{\pi} \frac{6^n}{n}$
4	A151331		N	$\frac{8}{\pi} \frac{8^n}{n}$
5	A151266		N	$\frac{1}{2} \sqrt{\frac{3}{\pi}} \frac{3^n}{n^{1/2}}$
6	A151307		N	$\frac{1}{4} \sqrt{\frac{10}{\pi}} \frac{5^n}{n^{1/2}}$
7	A151291		N	$\frac{4}{3\sqrt{\pi}} \frac{4^n}{n^{1/2}}$
8	A151326		N	$\frac{2}{3} \sqrt{\frac{3}{\pi}} \frac{6^n}{n^{1/2}}$
9	A151302		N	$\frac{1}{6} \sqrt{\frac{10}{\pi}} \frac{5^n}{n^{1/2}}$
10	A151329		N	$\frac{1}{9} \sqrt{\frac{21}{\pi}} \frac{7^n}{n^{1/2}}$
11	A151261		N	$\begin{cases} \frac{12\sqrt{3}}{\pi} \frac{(2\sqrt{3})^n}{n^2} & (n = 2p) \\ \frac{18}{\pi} \frac{(2\sqrt{3})^n}{n^2} & (n = 2p + 1) \end{cases}$
12	A151297		N	$\frac{\sqrt{3}B^{7/2}}{2\pi} \frac{(2B)^n}{n^2}$
13	A151275		N	$\begin{cases} \frac{12\sqrt{30}}{\pi} \frac{(2\sqrt{6})^n}{n^2} & (n = 2p) \\ \frac{144\sqrt{5}}{5\pi} \frac{(2\sqrt{6})^n}{n^2} & (n = 2p + 1) \end{cases}$
14	A151314		N	$\frac{\sqrt{6}\lambda\mu C^{5/2}}{5\pi} \frac{(2C)^n}{n^2}$
15	A151255		N	$\begin{cases} \frac{24\sqrt{2}}{\pi} \frac{(2\sqrt{2})^n}{n^2} & (n = 4p) \\ \frac{32}{\pi} \frac{(2\sqrt{2})^n}{n^2} & (n = 4p + 1) \\ \frac{24\sqrt{2}}{\pi} \frac{(2\sqrt{2})^n}{n^2} & (n = 4p + 2) \\ \frac{32}{\pi} \frac{(2\sqrt{2})^n}{n^2} & (n = 4p + 3) \end{cases}$
16	A151287		N	$\frac{2\sqrt{2}A^{7/2}}{\pi} \frac{(2A)^n}{n^2}$
17	A001006		Y	$\frac{3}{2} \sqrt{\frac{3}{\pi}} \frac{3^n}{n^{3/2}}$
18	A129400		Y	$\frac{3}{2} \sqrt{\frac{3}{\pi}} \frac{6^n}{n^{3/2}}$
19	A005558		N	$\frac{8}{\pi} \frac{4^n}{n^2}$

Table 6: Nature of the generating series for $(x, y) = (1, 1)$ and coefficient asymptotics

$$A = 1 + \sqrt{2}, \quad B = 1 + \sqrt{3}, \quad C = 1 + \sqrt{6}, \quad \lambda = 7 + 3\sqrt{6}, \quad \mu = \sqrt{\frac{4\sqrt{6}-1}{19}}$$