



HAL
open science

Iterative construction of replicated designs based on Sobol' sequences

Laurent Gilquin, Lluís Antoni Jiménez Rugama, Élise Arnaud, Fred J. Hickernell, Hervé Monod, Clémentine Prieur

► **To cite this version:**

Laurent Gilquin, Lluís Antoni Jiménez Rugama, Élise Arnaud, Fred J. Hickernell, Hervé Monod, et al.. Iterative construction of replicated designs based on Sobol' sequences. 2016. hal-01349444v2

HAL Id: hal-01349444

<https://inria.hal.science/hal-01349444v2>

Preprint submitted on 1 Dec 2016 (v2), last revised 3 Jan 2017 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Iterative construction of replicated designs based on Sobol' sequences

Laurent Gilquin^a, Lluís Antoni Jiménez Rugama^b, Elise Arnaud^c, Fred J. Hickernell^b,
Hervé Monod^d, Clémentine Prieur^c

^a*Inria Grenoble - Rhône-Alpes, Inovallée, 655 avenue de l'Europe, 38330 Montbonnot*

^b*Illinois Institute of Technology, Rettaliata Engineering Center, 10 W 32st, Chicago, IL 60616*

^c*Univ. Grenoble Alpes, Jean Kunzmann Laboratory, F-38000 Grenoble, France*

CNRS, LJK, F-38000 Grenoble, France, Inria

^d*MaIAGE, INRA, Université Paris-Saclay, 78350 Jouy-En-Josas, France*

Received *****, accepted after revision +++++

Presented by

Abstract

In the perspective of estimating main effects of model inputs, two approaches are studied to iteratively construct replicated designs based on Sobol' sequences. Space-filling properties of the resulting designs are studied based on two criteria. *To cite this article: L. Gilquin, Ll. A. Jiménez Rugama, E. Arnaud, F. J. Hickernell, H. Monod, C. Prieur, C. R. Acad. Sci. Paris, Ser. I 340 (2016).*

Résumé

Construction de plans répliqués à partir de séquences de Sobol'. Dans l'objectif d'estimer les effets principaux des paramètres d'un modèle, nous proposons d'étudier deux approches pour construire itérativement des plans répliqués à partir de séquences de Sobol'. Les propriétés de remplissage de l'espace des plans construits sont étudiées sur la base de deux critères. *Pour citer cet article : L. Gilquin, Ll. A. Jiménez Rugama, E. Arnaud, F. J. Hickernell, H. Monod, C. Prieur, C. R. Acad. Sci. Paris, Ser. I 340 (2016).*

1. Introduction

Mathematical models often involve a substantial number of poorly known parameters. The effect of these parameters on the output of the model can be assessed through sensitivity analysis. Global sensitivity

Email addresses: laurent.gilquin@inria.fr (Laurent Gilquin), ljimene1@hawk.iit.edu (Lluís Antoni Jiménez Rugama).

analysis methods are useful tools to identify the parameters having the most influence on the output. A well known approach is the variance based method introduced by Sobol' in [11]. This method estimates sensitivity indices called Sobol' indices that summarize the influence of each model input. Among all Sobol' indices, one can distinguish the first-order indices that estimate the main effect of each input.

The procedure to estimate first-order Sobol' indices proposed by Sobol' and its improvements (see Saltelli [9] for an exhaustive survey) all suffer from a prohibitive number of model evaluations that grows with respect to the input space dimension. An elegant solution to reduce this number relies on the construction of particular designs of experiments called replicated designs. The notion of replicated designs was first introduced by McKay through its definition of replicated Latin Hypercubes in [4]. Below we provide this definition in a more general framework:

Definition 1.1 Consider $\mathbf{x} \in [0, 1]^s$, and $\mathbf{x}_u \in [0, 1]^{|u|}$ the subset of elements of \mathbf{x} given by $u \subsetneq \{1, \dots, s\}$, where $|u|$ is the cardinality of u . Let $\mathcal{P} = \{\mathbf{x}_i\}_{i=0}^{n-1}$ and $\mathcal{P}' = \{\mathbf{x}'_i\}_{i=0}^{n-1}$ be two point sets in $[0, 1]^s$, and denote by $\mathcal{P}^u = \{\mathbf{x}_{i,u}\}_{i=0}^{n-1}$ (resp. \mathcal{P}'^u) the subset of elements of points in \mathcal{P} (resp. \mathcal{P}') indexed by u . We say that \mathcal{P} and \mathcal{P}' are two replicated designs of order $a = 1, \dots, s - 1$, if for any $u \subsetneq \{1, \dots, s\}$ with $|u| = a$, \mathcal{P}^u and \mathcal{P}'^u are the same point set in $[0, 1]^a$, perhaps in a different order.

The replication procedure described in [2,12] allows the estimation of all first-order Sobol' indices with only two replicated designs of order 1. This procedure has the major advantage of reducing the number of model evaluations, evaluating only on designs \mathcal{P} and \mathcal{P}' regardless of the input space dimension. However, Sobol' indices estimates may still not be accurate enough if designs \mathcal{P} and \mathcal{P}' do not explore the input space properly.

In this note, we propose two different constructions of replicated designs of order 1 based on Sobol' sequences. Both constructions ensure that the input space is properly explored and can be used within the replication procedure to estimate all first-order Sobol' indices. The definition of these constructions is recursive, therefore one can iteratively refine each replicated design by adding the corresponding new set of points. We first provide a brief introduction on digital sequences and then present two iterative approaches to construct the two replicated point sets. We end this note by analyzing the space-filling properties of the two designs constructed.

2. Digital sequences background

2.1. Preliminaries

Digital nets and sequences were first introduced by Niederreiter [6] in the numerical integration framework to define good uniformly distributed points in $[0, 1]^s$. They can also appear in the literature as digital (t, m, s) -nets and digital (t, s) -sequences, or simply (t, m, s) -nets and (t, s) -sequences. Sobol' and Niederreiter-Xing sequences are two examples of digital sequences detailed in [10] and [7].

Definition 2.1 Let \mathcal{A} be the set of all elementary intervals $A \subset [0, 1]^s$ where $A = \prod_{j=1}^s [\alpha_j b^{-\gamma_j}, (\alpha_j + 1)b^{-\gamma_j})$, with integers $s \geq 1$, $b \geq 2$, $\gamma_j \geq 0$, and $b^{\gamma_j} > \alpha_j \geq 0$. For $m \geq t \geq 0$, the point set $\mathcal{P} \in [0, 1]^s$ with b^m points is a (t, m, s) -net in base b if every A with volume b^{t-m} contains b^t points of \mathcal{P} .

Thus, a (t, m, s) -net is defined such that all elementary intervals of volume b^{t-m} will enclose the same proportion of points of \mathcal{P} , namely $b^{t-m}|\mathcal{P}|$ points. The most evenly spread nets are $(0, m, s)$ -nets, since each elementary interval of the smallest volume possible, b^{-m} , contains exactly one point. The quality of any (t, m, s) -net is therefore measured by the parameter t , called t -value.

By increasing m , we increase the number of points of the (t, m, s) -net. In the limiting case where $m \rightarrow \infty$, we can define the (t, s) -sequence as:

Definition 2.2 For integers $s \geq 1$, $b \geq 2$, and $t \geq 0$, the sequence $\{\mathbf{x}_i\}_{i \in \mathbb{N}_0}$ is a (t, s) -sequence in base b , if for every set $\mathcal{P}_{\ell, m} = \{\mathbf{x}_i\}_{i=\ell b^m}^{(\ell+1)b^m-1}$ with $\ell \geq 0$ and $m \geq t$, $\mathcal{P}_{\ell, m}$ is a (t, m, s) -net in base b .

The replicated design properties can also apply to digital sequences. Hence, we introduce the following definition,

Definition 2.3 Two digital sequences $\{\mathbf{x}_i\}_{i \in \mathbb{N}_0}$ and $\{\mathbf{x}'_i\}_{i \in \mathbb{N}_0}$ are digitally replicated of order a if for all $m \geq 0$, $\{\mathbf{x}_i\}_{i=0}^{b^m-1}$ and $\{\mathbf{x}'_i\}_{i=0}^{b^m-1}$ are two replicated designs of order a .

2.2. Sobol' sequences

Sobol' sequences in dimension s are digital sequences in base $b = 2$ that can be computed using the *generating matrices*, a set of s full rank infinite dimensional upper triangular matrices over the Galois field $\mathbb{F}_2 := \{0, 1\}$. These generating matrices are recursively constructed given some primitive polynomials and initial directional numbers. In [1], Kuo and Joe detail this construction and also suggest a particular choice for these matrices that optimize the 2 dimensional projection t -values.

Consider the generating matrices C_1, \dots, C_s , and C_1^m, \dots, C_s^m their upper left corner blocks of size $m \times m$. Although C_1, \dots, C_s are of infinite size, one only requires the knowledge of C_1^m, \dots, C_s^m to construct the first 2^m Sobol' points: for each $i = 0, \dots, 2^m - 1$, the point $\mathbf{x}_i = (x_{i,1}, \dots, x_{i,s})^\top$ of the sequence is obtained dimension-wise by:

$$(x_{i,j,1}, \dots, x_{i,j,m})^\top = C_j^m \mathbf{i}, \quad j = 1, \dots, s, \quad (1)$$

where $x_{i,j} = \sum_{k \geq 1} x_{i,j,k} 2^{-k}$ is the binary expansion of $x_{i,j}$ and $\mathbf{i} = (i_0, \dots, i_{m-1})^\top$ is the vector obtained from the binary expansion of $i = \sum_{k \geq 0} i_k 2^k$. All matrix operations are performed in \mathbb{F}_2 . Below we provide an example of how to compute $x_{11,1} = 0.8125$ and $x_{9,2} = 0.4375$,

$$\begin{pmatrix} x_{11,1,1} \\ x_{11,1,2} \\ x_{11,1,3} \\ x_{11,1,4} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}}_{C_1^4} \underbrace{\begin{pmatrix} 1 \\ 1 \\ 0 \\ 1 \end{pmatrix}}_{\mathbf{11}} = \begin{pmatrix} 1 \\ 1 \\ 0 \\ 1 \end{pmatrix}, \quad \begin{pmatrix} x_{9,2,1} \\ x_{9,2,2} \\ x_{9,2,3} \\ x_{9,2,4} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 & 1 & 1 \\ 0 & 1 & 0 & 1 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}}_{C_2^4} \underbrace{\begin{pmatrix} 1 \\ 0 \\ 0 \\ 1 \end{pmatrix}}_{\mathbf{9}} = \begin{pmatrix} 0 \\ 1 \\ 1 \\ 1 \end{pmatrix}$$

To compute the next 2^m points of the sequence, one can infer from (1) that

$$(x_{i+2^m,j,1}, \dots, x_{i+2^m,j,m+1})^\top = (x_{i,j,1}, \dots, x_{i,j,m}, 0)^\top \oplus (c_j^{m+1})^\top, \quad i = 0, \dots, 2^m - 1, \quad (2)$$

where c_j^{m+1} is the last column of C_j^{m+1} and \oplus is the addition in \mathbb{F}_2 .

In addition, Sobol' sequences have good group structure properties. For any $m \geq 1$, the first 2^m points of the sequence form an Abelian group under \oplus :

$$\begin{aligned} \mathbf{x} &= \left(\sum_{k=1}^m x_{1,k} 2^{-k}, \dots, \sum_{k=1}^m x_{s,k} 2^{-k} \right)^\top, & \mathbf{z} &= \left(\sum_{k=1}^m z_{1,k} 2^{-k}, \dots, \sum_{k=1}^m z_{s,k} 2^{-k} \right)^\top, \\ \mathbf{x} \oplus \mathbf{z} &:= \left(\sum_{k=1}^m (x_{1,k} + z_{1,k} \bmod 2) 2^{-k}, \dots, \sum_{k=1}^m (x_{s,k} + z_{s,k} \bmod 2) 2^{-k} \right)^\top. \end{aligned} \quad (3)$$

From equation (1) above, one may see that the first 2^m points of the sequence are elements of $(\mathbb{F}_2^m)^s$.

Lemma 2.4 All Sobol' sequences are digitally replicated of order 1.

Proof: Consider any two s -dimensional Sobol' sequences generated by C_1, \dots, C_s and C'_1, \dots, C'_s respectively. Since any two matrices C_j^m and $C'_j{}^m$ are square and full rank, the operations $\mathbf{i} \mapsto C_j^m \mathbf{i}$ and $\mathbf{i} \mapsto C'_j{}^m \mathbf{i}$ are one-to one and onto for all $i = 0, \dots, 2^m - 1$. Therefore, they generate the same point sets.

3. Iterative constructions of replicated point sets

In this section we propose two different approaches to iteratively construct two replicated point sets, \mathcal{P}_ℓ and \mathcal{P}'_ℓ , based on Sobol' sequences. These two constructions are carried out according to the following recursive scheme :

$$\begin{cases} \mathcal{P}_0 = B_0 \\ \mathcal{P}_\ell = \mathcal{P}_{\ell-1} \cup B_\ell \end{cases} \quad \begin{cases} \mathcal{P}'_0 = B'_0 \\ \mathcal{P}'_\ell = \mathcal{P}'_{\ell-1} \cup B'_\ell \end{cases}, \quad \ell \geq 1,$$

where B_ℓ and B'_ℓ are new sets of points added at step ℓ to refine $\mathcal{P}_{\ell-1}$ and $\mathcal{P}'_{\ell-1}$. For all $\ell \geq 0$, \mathcal{P}_ℓ and \mathcal{P}'_ℓ are two replicated designs of order 1.

The first approach is called multiplicative because $|\mathcal{P}_\ell| = 2^\ell$ while the second one is called additive and $|\mathcal{P}_\ell| = \ell|B_0|$. In the multiplicative case, we will directly use 2 s -dimensional sequences as a result from Lemma 2.4. However, for the additive case, we will consider an initial set of Sobol' points and apply different scramblings and digital shifts to extend the point sets. Additionally, in both cases one can randomize the points using Owen's scrambling [8] as long as same coordinates of \mathcal{P}_ℓ and \mathcal{P}'_ℓ share the same scrambling.

3.1. Multiplicative approach

Two replicated point sets of order 1, \mathcal{P}_ℓ and \mathcal{P}'_ℓ , can be constructed using two s -dimensional Sobol' sequences. We note C_1, \dots, C_s the generating matrices used to generate \mathcal{P}_ℓ and C'_1, \dots, C'_s those used to generate \mathcal{P}'_ℓ . To ensure that \mathcal{P}_ℓ and \mathcal{P}'_ℓ are as uniform as possible, these generating matrices need to be different from each other. We choose $C_1, \dots, C_s, C'_1, \dots, C'_s$ to be the first $2s$ generating matrices suggested by Joe and Kuo in [1], not necessarily in this order. In [1], Joe and Kuo minimized the t -value for all 2 dimensional projections.

In order to iteratively extend designs \mathcal{P}_ℓ and \mathcal{P}'_ℓ , one just needs to compute $B_\ell = \{\mathbf{x}_{2^{\ell-1}}, \dots, \mathbf{x}_{2^\ell-1}\}$ and $B'_\ell = \{\mathbf{x}'_{2^{\ell-1}}, \dots, \mathbf{x}'_{2^\ell-1}\}$, starting with $B_0 = B'_0 = \{\mathbf{0}\}$, with $\mathbf{0}$ the null vector. Each set B_ℓ and B'_ℓ can be constructed using (2) applied to $\mathcal{P}_{\ell-1}$ and $\mathcal{P}'_{\ell-1}$.

As a direct consequence of Lemma 2.4, at each step ℓ designs \mathcal{P}_ℓ and \mathcal{P}'_ℓ are two replicated designs of order 1. Furthermore, they both inherit the space-filling properties of (t, ℓ, s) -nets.

3.2. Additive Approach

With the multiplicative approach, the size of designs \mathcal{P}_ℓ and \mathcal{P}'_ℓ is multiplied by 2 each time ℓ is increased by one. This growth rate may be inadequate for some applications. The additive approach presented in this section is attractive due to a slower size growth. Given an initial choice of $r \geq 1$ that specifies the size of B_0 and B'_0 , only $|B_0| = 2^r$ points are added to both designs at each step. The main drawback of this approach is that \mathcal{P}_ℓ and \mathcal{P}'_ℓ do not inherit the structure of a Sobol' sequence when $\ell \geq 1$. Nevertheless, both designs display good space-filling properties as shown in the next section.

Analogously to the multiplicative case, the two replicated point sets \mathcal{P}_ℓ and \mathcal{P}'_ℓ constructed with the additive approach are iteratively refined with B_ℓ and B'_ℓ . First $\{\mathbf{x}_i\}_{i=0}^{2^r-1}$ and $\{\mathbf{x}'_i\}_{i=0}^{2^r-1}$ are set to be the first 2^r points of two s -dimensional Sobol' sequences. The generating matrices of these two sequences are selected as in the multiplicative approach. Then, B_ℓ (resp. B'_ℓ), for $\ell \geq 0$, is obtained from $\{\mathbf{x}_i\}_{i=0}^{2^r-1}$ (resp. $\{\mathbf{x}'_i\}_{i=0}^{2^r-1}$) by carrying out digital shifts and scrambling operations. Therefore, both B_ℓ and B'_ℓ inherit the (t, r, s) -net structure of these initial sets.

At step $\ell = 0$, $B_0 = \{\mathbf{x}_i^{(0)}\}_{i=0}^{2^r-1}$ and $B'_0 = \{\mathbf{x}'_i{}^{(0)}\}_{i=0}^{2^r-1}$ are generated as follows: for each $i = 0, \dots, 2^r - 1$, points $\mathbf{x}_i^{(0)} = (x_{i,1}^{(0)}, \dots, x_{i,s}^{(0)})^\top$ and $\mathbf{x}'_i{}^{(0)} = (x'_{i,1}{}^{(0)}, \dots, x'_{i,s}{}^{(0)})^\top$ are obtained by linearly transforming \mathbf{x}_i ,

$$\begin{aligned} (x_{i,j,1}^{(0)}, \dots, x_{i,j,r}^{(0)})^\top &= L(x_{i,j,1}, \dots, x_{i,j,r})^\top, & j = 1, \dots, s, \\ (x'_{i,j,1}{}^{(0)}, \dots, x'_{i,j,r}{}^{(0)})^\top &= L'(x_{i,j,1}, \dots, x_{i,j,r})^\top, & j = 1, \dots, s, \end{aligned} \quad (4)$$

where $x_{i,j}^{(0)} = \sum_{k=1}^r x_{i,j,k}^{(0)} 2^{-k}$, $x'_{i,j}{}^{(0)} = \sum_{k=1}^r x'_{i,j,k}{}^{(0)} 2^{-k}$ and L, L' are two distinct full rank lower triangular matrices of size $r \times r$ over \mathbb{F}_2 . Left multiplications by L and L' in (4) are called linear scramblings [3].

At step $\ell \geq 1$, using addition as defined in (3) we construct:

$$B_\ell = \{\mathbf{x}_i^{(0)} \oplus \mathbf{e}_\ell\}_{i=0}^{2^r-1}, \quad B'_\ell = \{\mathbf{x}'_i{}^{(0)} \oplus \mathbf{e}'_\ell\}_{i=0}^{2^r-1}. \quad (5)$$

where $\mathbf{e}_\ell \in (\mathbb{F}_2^r)^s \setminus \mathcal{P}_{\ell-1}$ and $\mathbf{e}'_\ell \in (\mathbb{F}_2^r)^s \setminus \mathcal{P}'_{\ell-1}$. Indeed, B_0 is a subgroup of $(\mathbb{F}_2^r)^s$. Therefore, we construct $B_\ell = B_0 \oplus \mathbf{e}_\ell$ as a coset of B_0 under the right choice of \mathbf{e}_ℓ . The same reasoning applies to B'_0 . Additions by vectors \mathbf{e}_j and \mathbf{e}'_j in (5) are referred as digital shifts.

There are exactly 2^{rs} different choices of \mathbf{e}_ℓ and \mathbf{e}'_ℓ in $(\mathbb{F}_2^r)^s$, and $2^{r(r-1)/2}$ choices of L and L' . The maximum number of iterations ℓ that can be performed equals the number of cosets, $2^{r(s-1)}$. If this maximum is reached, designs \mathcal{P}_ℓ and \mathcal{P}'_ℓ both correspond to a full grid design of 2^{rs} points with 2^r distinct values per input.

Lemma 3.1 *Under the additive construction \mathcal{P}_ℓ and \mathcal{P}'_ℓ are two replicated designs of order 1, $\ell \geq 0$.*

Proof: For each $\ell \geq 1$, digital shift and scrambling operations carried out in (4) and (5) are bijections from \mathbb{F}_2^r and $(\mathbb{F}_2^r)^s$. Hence, the same coordinates of B_ℓ and B'_ℓ will contain the same set of 2^r values.


4. Space-filling properties

To analyze the space-filling properties of both approaches we will use two criteria. The first one is the L_2 star discrepancy that measures the uniformity of a point set. This criterion can be easily computed through an analytical expression provided by [5]. The lower the criterion, the more uniform the point set is. Sobol' sequences were originally constructed to be low discrepancy sequences and therefore, behave well under this criterion. The second criterion is called maximin distance and measures the regularity of a point set. It returns the minimum of all Euclidean distances between pairs of points in the set. The higher the criterion, the more spread the point set is.

We name by design M (resp. A) one of the two replicated designs constructed with the multiplicative (resp. additive) approach, and set the input space dimension s to 6. For design M , we study the properties of $\mathcal{P}_8, \dots, \mathcal{P}_{12}$, and for design A , we set the initial value to $r = 8$ and study the properties of $\mathcal{P}_0, \dots, \mathcal{P}_{15}$. For both M and A we generate 100 designs whose average criteria are presented in Figure 1. While each of the 100 designs M is independently randomized applying Owen's scrambling [8], designs A are constructed by randomly selecting matrix L (uniformly among all $2^{r(r-1)/2}$ choices) and vectors \mathbf{e}_ℓ used in equation (5). In addition, we also show in Figure 1 the results obtained with an optimized Latin Hypercube design (LH design) according to each criterion for an equal number of points as in design A . The x-axis corresponds to the number of design points N and the y-axis to the value of the criterion. Although LH designs are the standard designs used in the replication procedure to estimate first-order indices, the replicated design of an optimized LH design is not guaranteed to be optimal.

As expected, design M performs the best and the estimated slope of the discrepancy curve (-0.72) falls within the expected range for Sobol' sequences [5]. Although design A does not perform as well as design M , it outperforms an optimized LH design for the discrepancy criterion. However for the maximin

Figure 1. Log-log graph of averaged L_2 star discrepancy (left) and maximin (right) for designs M and A over 100 repetitions.



criterion, design A shows slightly worse results. Concerning the number of steps, design A allows 15 refinement steps from 2^8 to 2^{12} points, instead of only 4 steps for design M .

Acknowledgements

This work is supported by the CITiES project funded by the Agence Nationale de la Recherche (grant ANR-12-MONU-0020) and by the United States National Science Foundation (grant DMS-1522687).

References

- [1] S. Joe, F. Y. Kuo, Constructing Sobol Sequences with Better Two-Dimensional Projections, *SIAM J. Sci. Comput.* 30(8), 2008, pp. 2635–2654.
- [2] T. A. Mara, O. R. Joseph, Comparison of some efficient methods to evaluate the main effect of computer model factors, *Journal of Statistical Computation and Simulation* 78(2), 2008, pp. 167–178.
- [3] J. Matoušek, On the L_2 -discrepancy for anchored boxes, *Journal of Complexity* 14, 1998, pp. 527–556.
- [4] M. D. McKay, W. J. Conover, R. J. Beckman, A comparison of three methods for selecting values of input variables in the analysis of output from a computer code, *Technometrics* 21, 1979, pp. 239–245.
- [5] W. J. Morokoff, R. E. Caflisch, Quasi-random sequences and their discrepancies, *SIAM Journal on Scientific Computing* 15, 1994, pp.1251?–1279.
- [6] H. Niederreiter, Random Number Generation and Quasi-Monte Carlo Methods, *CBMS-NSF Regional Conference Series in Applied Math.* 63, SIAM, Philadelphia, PA, 1992.
- [7] H. Niederreiter, C. Xing, Low-Discrepancy Sequences and Global Function Fields with Many Rational Places, *Finite Fields and Their Applications* 2(3), 1996, pp. 241–273.
- [8] A. B. Owen, Scrambling Sobol’ and Niederreiter-xing points, *J. Complexity*, 14(4), 1998, pp. 466–489.
- [9] A. Saltelli, Making best use of model evaluations to compute sensitivity indices, *Computer Physics Communications* 145(2), 2002, pp. 280–297.
- [10] I. M. Sobol’, On the distribution of points in a cube and the approximate evaluation of integrals, *USSR Computational Mathematics and Mathematical Physics* 7(4), 1967, pp. 86–112.
- [11] I. M. Sobol’, Sensitivity indices for nonlinear mathematical models, *Mathematical Modeling and Computational Experiment* 1, 1993, pp. 407–414.
- [12] J. Y. Tissot, C. Prieur, A randomized orthogonal array-based procedure for the estimation of first- and second-order Sobol’ indices, *J. Statist. Comput. Simulation* 85, 2014, pp. 1358–1381.