

# The Development of Decision Support System for Production of Layer

Jianhua Xiao, Hongbin Wang, Luyi Shi, Mingzhe Lv, Haikun Ma

## ▶ To cite this version:

Jianhua Xiao, Hongbin Wang, Luyi Shi, Mingzhe Lv, Haikun Ma. The Development of Decision Support System for Production of Layer. 5th Computer and Computing Technologies in Agriculture (CCTA), Oct 2011, Beijing, China. pp.161-168, 10.1007/978-3-642-27281-3\_21. hal-01351807

# HAL Id: hal-01351807 https://inria.hal.science/hal-01351807v1

Submitted on 4 Aug 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

## The Development of Decision Support System for Production of Layer

Jianhua Xiao, Hongbin Wang, Luyi Shi, Mingzhe Lv, Haikun Ma

College of Veterinary Medicine, Northeast Agricultural University, Harbin 150030, China

**Abstract.** The Decision Support System for Production of Commercial Layer was developed based on the demand analysis. VB .NET 2005 and SQL SERVER 2000 was used as main methods for system development. With this system, the data of alternation, growth, breeding, environment, immune for layer production can be recorded, edited, analyzed and be given by a report forms. With this system, the production efficiency can be raised.

Keywords: layer; DSS.

## 1 Introduction

For a large-scale layer enterprise, there must be many batches of chickens kept at the same time. And there must be many differences for observed indicators among those batches. Therefore, it was very difficult to ensure that each chicken goes according to plan production. This problem can be solved by developing one decision support system for production of commercial layer. In terms of computer software for layer production, one computer software for breeding was developed by Cai Juandeng, breeding data can be treated and various kinds of genetic parameters can be obtained by this soft [1]. One management system for chicken breeding data was developed by Xiao Fan, which has the pedigree matching, hatching management, determination of egg, report query etc[2]. One system of data acquisition and input for egg production was developed in 1993 by Yang Ning using bar code technology and handheld computer technology [3]. One digital breeding platform for breeding pig was developed by Zhao ruixue [4]. One management information system for layer farm was developed by Wu Xiaohong [5]. In addition, the computer software is also used in retrospective [6], disease diagnosis [7] and so on. The Decision Support System for Production of Commercial Layer was developed in this study aimed at recording and analyzing the data of growth, feeding, egg production, environment, immunization, and disease, changes during brood period, Incubation period, and laying period and out. By management and statistical analysis of production data, the overall situation and problems can be controlled by managers with this system, and the objective, scientific decision can be made by managers.

## 2 Principle and methods

#### 2.1 The production flow-sheet of layer

There are four stages during a production cycle of goods layers. They can be listed as follows: birth, incubation, egg, and out. The brood period is from its birth to six weeks. The incubation period is from the seventh to twentieth weeks. The egg period is from twenty to seventy-two weeks; then layer is out, different indicators need to be obtained in each period.

#### 2.2 The development methods and system framework

In general, one chicken farm is composed of several branches. moreover easy of use is required, and therefore the C/S framework was used in this system. One computer was set as server and the computer in every branch was installed client soft; to access the data in server by set the only IP address. The operating system for developed this soft was selected as windows XP, and Microsoft VB.NET 2005 was selected as development language[9], the excel component of Microsoft office was used to import and export data, the third party component Dotnet Charting was used to generate figure. Microsoft SQL Server 2000 was used as database.

## 3 Result

The system was divided into 6 modules based on the production processes of layer: the management of farm and house, data input, statistics and report forms, curve diagram, system management and help. The initial data of one batch of chickens was recorded after being introduced; the data generating in every production processes was record synchronously. All of data were done statistics in a statistics and report forms module, and also produced daily, weekly, periodic report. The function and structure of system can be found in figure 1.




Fig. 1. The function and structure of production and management system for layer

#### 3.1 The introduction of one batch of chicken

The basic data of one batch of chicken include information generated when they were introduced and information generated in every production stages. Those data mainly included serial number, species, source, approach date, born weight, quantity, stocking rate and survival rate after 24 weeks, date when increase light, egg date etc. because different species of chicken have different breeding standard, and those procedures such as increasing light and egg, mix chicken groups must be completed within the limited periods, therefore the standard of production was compiled in software, and the program will compute everyday according to this standard, the current status of chickens can be get, and the attention for change house, increase light, mix groups, etc. was gave to user automatically (fig. 2).

1000	× C 秘光亡 入	e wer	A THE	# a	<b>a</b> itt	_			_									
191722								鸡汁	畔信!	5								
815	19 B#				综合 存在公務		松正幸			入育諸會	入寬成會	入产蛋白	760%C	R.W	12.24	3		
2009		200	205 29 90 10			4300 4904	<ul> <li>原章</li> <li>波季</li> </ul>	<b>本語</b> 地形派		-72009-09-0 成入業入	2 J2009-10- 度入来入	14 度入宗入	Eastat	22*22	但况未况群 3	2.		
2009		200	135 19 290 40			4937	172 172	内要古 除学道		度入未入. 度入未入。	臣入未入 臣入未入	5日度入声 成入ま入		今日見重	应用米混料 应用米混料 5			
																1		
	存朽	É公鸡	音	Ę	顺道	李	*	源		代次	入育	雏舍	入育	成舍	٨ř	蛋舍	加光	见蛋
		122	410	DO .	. 顺	季	本	地	. Ī	前品鸡	√200	09-09-02	20	09-10-1	14 应入	.未入	应加未加光	应见未
	4	195	498	84	. 谜	季	Pi	小演	Ē	66品鸡	应入	未入	应入	未入				
	4	168	493	37	. 顺	季	Þ	「蒙古	7	祖代	应入	未入	应入	.未入	. 5日危	入产		今日.
		0	495	50	1165	季	D	尔滨	: 7	商品鸡	应λ	未入	应λ	未入	应λ	未入	应加未加光	应见未

**Fig.2**. The introduction of chickens and attentions in key production processes The data can be modified at any time and export into excel table. The production data need to be backed up and separated when production cycle of one batch of chicken has closed, then the data of this batch of chicken can be deleted from database.

#### 2.2 The management of change data for one batch of chicken

Death, eliminate, marketing, output and input of chickens are main changes of one batch of chicken in production. These data and data of herbs must be recorded in to computer every day. In order to facilitate data recording, few data (such as death and eliminate etc.) need to be input and closing stock will be computed by numbers of yesterday automatically. By this ways, the efficiency would be improved significantly. The changes of data can be statistics, daily, weekly, and period report and figure of one phase can be generated in system. By these results, more scientific measure can be made by managers. By contrasted to standard, the conclusion of production can be got(fig. 3).




Fig. 3. The changes data of one batch of chicken

#### 2.3 The management of growth data for one batch of chicken

The growth was the main observation for management during brood and incubation period. Sound skeletal structure is necessary for egg production. It is very important to monitor redords and analyzation of the chickens growth information. Only the weight of chicken at Pre-laying (from 21 to 28 week) need to be record weekly. Based on the general requirement of layer farm, body weight, uniformity,

average uniformity, change of feather, length of keel, length of tibia need to be managed in this system. The growth data can be statistics, daily report, weekly report, report of one phase can be generated in system (fig.4).

导航区	_		X	0 2 >		_		_	_	_	_	_	_				_
9 <b>Co</b> 19	6合管理 3 瑞场管理			1870 1872 80	96 BB				_								_
1.49	3合管理			🗌 生长日期	2010年 3月25日	至 2010	年3月25日 💌										
T G	647首位 引入 分會 美術 生時辨 学項 环境			地群洋细谋息						生长旗图							
	分合			斜编号	出生日期	存栏合计	存栏公略	存栏母19		生物	日期	体置(公)	<b>体重(母)</b>	龙骨长度(公)	龙骨长度(母)	<b>胫骨长度</b> (公)	<b>股骨</b> 核
Ğ	生长			200912001	2009-09-01	4522		4100			9109102	90	40	3.4	3.4	3.4	
12	「四方			200912002	2009-12-22	5479	495	4984			9-09-09	140	90	4.2	4.2	4.2	4
149	环境			200912003	2009-11-10	5380		4929		-	9-09-16	180	145	4.9	4.9	4.9	-
	F1.42.88			200912005	2009-06-18	4959	0	4959		200	9-09-23	270	220	5.8	5.8	5.8	
	■ 高齢生长編集 ○注約4人											- 🛛 🗖 -	360	7.4	0.0 T.4	7.4	
	38.8168.4												410	8	8	1.5	
		_											500		9	9	-
	8. P						体置(8)			标准值			590			9	-
	出生日期	8 200	9-09-0	1			俸重(¥)	40 E		标准值(	0至90		680	9	9	9	
	存栏公瑞(ま	) 422		R 109	5月 001		沈骨长度(8)	3.4 0		80#10 B	4至3.4		730	9	9	9	
	存栏母端(平			я в	29 1		龙骨长度(*)	3.4			4至3.4		820	9	9	9	
	存栏合计				新 (JR)天							_	910	9	9	9	
	(PCa)		-	1 1	av 004034	_	<b>胫骨长度(8)</b>				.4 至 3.4		960	9	9	9	
	地群分布						胫骨长度(*)	3.4 0		标准值 2	4至3.4		1040	9	9	9	
	渴舍名称	2	<b>亨杜公</b> 第	6 存栏母稿 在	F栏合计		均均度(8)	0.8		标准值	至1		1090	9	9	9	
0 <b>G</b>	1등		499	4992	5491		均均度(年)	0.8		标准值	至1		1140	9	9	9	
							全群均匀度	0.8		标准值							
8 <b>G</b>										ALC: NO.	_	_					
- 1																	
							备 注										
÷ 🖓																	
										(87		28日					
. L		_	_							140		20					
	<b>环境指标</b>																
- 18	建鸡品种 1间科种类																
										<							>
当前用户	:系统管理员	삶	司名称:														
1 7	187 B	18) F		· · · · · · · · · · · · · · · · · · ·		19,15	🔁 546	12		8理系统 -	(38	Ja Hicroso	ft Pore	<b>副 瑞羅主氏病</b>		10066	15:24

Fig.4. The management of growth data for one batch of chicken

#### 2.4 The management of feeding data for one batch of chicken

The feeding is associated with each period of production. The difference among those periods is the feed. Therefore feeding methods, the number of feed, feed consumption, water consumption, feeding time, difference between planning and the actual feed consumption, protein and metabolic energy intake and other related information need to be recorded in database. By comparing, the information whether the actual feed consumption has met the standards will be reminder to user. In addition, the feeding data can be statistics, daily report, weekly report, report of one phase and figure of changing can be generated in system. Thus managers can investigate the reason for insufficiency or excessive, and take effective measures timely (fig.5).




Fig.5. The management of feeding data for one batch of chicken

#### 2.5 The management of production data for one batch of chicken

Egg number, number of unqualified eggs, fertilized eggs, ratio of feed to egg need to managed. Those data can be input from three routes: ①input by manager directly, ②generated from system by computed, ③obtain from computer directly or generate from system daily. The number of hatching egg, number of goods egg, number of double-yolked egg, number of soft egg need to be recorded by manager daily. And laying rate, pass rate of hatching egg, ratio of commodity egg to total egg, double yellow egg ratio, ratio of soft eggs, broken egg rate, fertilization rate, hatching rate, etc generated from system by computed. Every data can be count(fig.6), and statistical graph can be generated(fig.7).

				_		_							_
场合管理	城市 截 副除 3	2 2 2											
18 地场管理 18 地合管理 18 地容管理	□严重日期 2010	年 3月25日 🖌	至 2010年 3	月25日 💌									
4 6 6 7 12 ( 1 3 1 2 5 1 2 1 2	观寻洋洋相信息						"面信息						
- Log 分音 - Log 支动		生日期	存胜合计	存胜公顷			产蛋日期	約(日本)	产派的	种滥款	商品運動	双黄蛋胶	\$1
重生长		009-09-01	4522	422	4100		2009-11-10	4990	250	100	150	2	
- Co P 面		009-12-22	5479	495	4984		2009-11-11	4989	251	120	131	4	-
ित मध्र		009-11-10	5389	450	4929		2009-11-12	4989	240	100	140	2	-
■ 江戸統计 ■ 水香产 承編編	200912005 5				4959		2009-11-13	4900	260	100	160	40	-
n and an			0.0						500	200	300	80	+
19新橋令 201212003									505	298	201	90	+
品种富万斯			入會瑞数 4990	8	产要率	E 01	标准値の至		2187	850	1337	90	+
出生日期 2009-05-10						40.00	転産値の至こ		2690	800	1890	600	+
EPE DW 2009-00-10			产蛋数 250	教	种重比率				3505	500	3005	700	-
			种蛋款 100	枚	商品重比率		标准値 0 至 :		3608	500	3108	2	+
存把公路(8) 0 5	(词参员 001		商品重数 150	枚	軟皮蛋比率		标准値 0 至 :		3527	500	3027	4	+
			软成蛋数 2	权	<b>双黄蛋比率</b>	0.90	标准値 0 至 :	1	3T00	500	3200	500	-
存栏每路(平) 4059 5			双黄蛋胶 2	35	破损重比率	3.20	标准値 0 至 :	1	3708	500	3208	344	
存栏合计 4959 5	8 周 龄 20周5天		N 15 20 20 0	82	合稿率	90.00	标准値 0 至 :	1	3709	500	3209	2	
地群分布 助会名称 存托公路	存托曼地 存托会计	- 10	第合結款 90	27	入射量比率	38.00	标准值 0 至 :	1	3941	500	3441	4	
지방권하 (FE23) 5등순 0			入解漏数 90	n	受機率	91.44	枢虐伯 0 至 :	1	3470	500	2970	500	
248 0	4990 4990		受機置数 85	17	(入耕酒)耕化率	88.89	标准值 0 至 :	1	4000	1000	3000	800	
				8	使始率	97.50	彩岸伯		4021	500	3521	400	
			出始数 80		受持限制化率				4101	500	3601	4	
			修建数 78	<b>R</b>			标准値 0 至 :		4202	500	3702	2	
			产置驾数 0.9		产重数/入会路		标准值 0 至 :		4282	500	3782	4	
					种蛋数/入舍鸡	0.02	都准値 0 至 :		4321	500	3621	2	
		全群	日产蛋重 20	815	蘆蘆/枚	80.00	€ 标准值 50 至	80	4482	500	3982	4	
		E.	产次营营 0.76	815	料漏比		标准値 0 至 :	1	4489	500	3989	4	_
					产蛋本增强	0.00		_	4467	500	3967	2	+
					上次产量率				4541	500	4031	4	-
									4507	500	4007	2	-
		1	塘 表 人 系统制	理想	备 注				4541 4567		4031	2	-
									4567	500	4067	2	+-
							保存	退出	4578	500	4010	300	-

Fig.6. The management of production data for one batch of chicken




Fig.7. The statistical graph of production data for one batch of chicken

### 4 Discussion

#### 4.1 Data inputting of management system

Too many items and large amounts of data are main factors that restrict the application of software in practices. Although some data can be recorded by auto technology, however, some data still be input manually. It is so important that reduce the input by hand and improve the efficiency. The reducing of input was considered in any parts of this system, besides the keying order of textbox was adjusted, many data can be computed by system automatically. Moreover, the association of items among different modules was enhanced in system, and user can find the data that he wanted in place where he wanted.

#### 4.2 The intelligence of management system

Generally, many batches of chicken will be feed simultaneously in one large-scale farm, and different batches of chicken will be produced at different stages. Different items need to be managed for each batch of chicken. In order to improve the intelligence of system, the production information will be retrieved and computed by production standard every day, and the works need to be completed can be displayed in a form. The software can be used to look over new task.. By this way, the fault in will be reduced significantly.

#### 4.3 decision support for production of management system

The main purpose that a manager using the software is to do statistics and analyzes production data, thereby finding out the problem during the production, so that making the work plans on the basis of current data. This quest was considered thoroughly when this system being designed and developed. The production data in every procedure can be statistics by day, week and period, and the statistical graph can be generated directly.

### reference

- 1. Cai J., Tong H.B.. The Application of Computer in Breed for Chicken. Shandong Poultry. 5,14--16(1997).
- 2. Xiao fan, The Design and Application of Data Management System for High Quality Fowl Breeding.China Poultry, 31,4—7(2009)
- 3. Yang Ning, Shan Chonghao, Li Jixiang. Research on Record System Without Paper for Fowl Egg Number. Journal of Chinese Animal Husbandry, 29,3—5(1993).
- 4. Zhao Ruixue, Zhao Pengjv, Gong Chengfeng. Research on Digital Cultivation Technique Platform for Goods Layer. Heilongjiang Animal Husbandry and Veterinary,12,47—48(2006).
- Wu Xiaohong, Bin Shuying. The Computer Network Management for Modern Fowl Fram. Journal of Zhongkai Agrotechnical College, 6,53—59(1993)
- 6. Zhao Jinshi. The Analysis for Application Current Situation of Trace System for Chicken Quality in China. Chinese Journal of Animal Science, 47, 45—48(2011)
- Xu Jianqin. The Diagnosis Expert System for Common Disease of Fowl. Chinese Journal of Veterinary Medicine, 18,40—41(1992).
- 8. Yin Haiqiang. Useful Course of Studies for Visual Basic.NET. Beijing: China Railway Publishing House. (2003)