



Irreversible Electroporation: Disappearance of Observable Changes at Imaging Does Not Always Imply Complete Reversibility of the Underlying Causal Tissue Changes

Olivier Seror, Clair Poignard, Olivier Gallinato, Rafik Belkacem-Ourabia,
Olivier Sutter

► To cite this version:

Olivier Seror, Clair Poignard, Olivier Gallinato, Rafik Belkacem-Ourabia, Olivier Sutter. Irreversible Electroporation: Disappearance of Observable Changes at Imaging Does Not Always Imply Complete Reversibility of the Underlying Causal Tissue Changes. *Radiology*, 2017, 282, pp.301 - 302. 10.1148/radiol.2017161809 . hal-01421863

HAL Id: hal-01421863

<https://inria.hal.science/hal-01421863>

Submitted on 23 Dec 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Irreversible Electroporation: Disappearance of observable changes at imaging does not always imply complete reversibility of the underlying causal tissue changes

Olivier Seror, MD, PhD; Clair Poignard, PhD; Olivier Gallinato, PhD; Rafik Belkacem-Ourabia, MD, Olivier Sutter, MD¹

¹ From the Service de Radiologie de l’Hôpital Jean Verdier, Hôpitaux universitaires Paris-Seine-Saint-Denis, Assistance publique Hôpitaux de Paris, Bondy, France (O.S., R.B-O., O.Su.), the Unité mixte de Recherche 1162, Génomique fonctionnelle des Tumeurs solides, Institut National de la Santé et de la Recherche médicale, Paris, France (O.S.), the Unité de Formation et de Recherche Santé Médecine et Biologie humaine, Université Paris 13, Communauté d’Universités et Etablissements Sorbonne Paris Cité, Bobigny, France (O.S., O.Su.), Institut National de Recherche en Informatique et en Automatique, Bordeaux-Sud-Ouest, F-33400, Talence, France (C.P., O.G.), Centre National de la Recherche Scientifique, Institut de Mathématiques de Bordeaux, Unité Mixte de Recherche 5251, F-33400, Talence, France (C .P., O.G.)

Letter to editor

Irreversible Electroporation: Disappearance of observable changes at imaging does not always imply complete reversibility of the underlying causal tissue changes

We read with great interest Padia et al. article emphasizing the potential role of early MR assessment of irreversible electroporation (IRE) ablation in hepatocellular carcinoma (HCC) patients (1). We especially appreciated that authors pointed out that IRE efficacy cannot be assessed like for other physical ablative methods because cell death is no longer mainly related to thermal coagulative necrosis (2). One singularity of the technique is that below 600V/cm the change in permeability of cell membrane are assumed to be reversible (3). Thus, Padia et al. suggested that the transient changes seen at the periphery of IRE ablation zone on the early MR examinations were related to reversible effect of the treatment.

From our point of view, this explanatory hypothesis calls some comments. The biological effects of IRE on tissues are still unclear. Cell membrane electroporation leads to a swelling of the affected cells as well as a release of cellular materials, which generate edema and inflammation in the tissue (4). Such inflammation, visible with MRI disappears within few days (4, 5). On the other hand, it has been reported that reversible electroporation lasts several tens of minutes and any way much shorter than several hours (6). Interestingly, peripheral delayed enhanced zones have been also described with perfusion CT performed immediately after IRE in livers of pigs (7).

Histopathological analysis of livers harvested one day after showed a strong correlation between these peripheral zones and red zones which contained hepatocytes involved in terminal apoptotic process. Preliminary numerical simulations of IRE on tissue model showed also that the ablation zone induced is wider than its central necrotic part (8). Indeed, IRE is a multiprobe ablative technique requiring the placement of multiple electrodes into the tumor margin rather than in the centre of the lesion(9). In this setting, we suggest that the disappearance of peripheral delayed enhanced zones could be related to fast remodeling of apoptotic component of IRE ablation combining resolution of inflammation, phagocytosis and replacement of dead cells, and not necessarily to the recovery of basal membrane permeability of cells that were submitted to amount of energy below irreversibility threshold.

References:

1. Padia SA, Johnson GE, Yeung RS, Park JO, Hippe DS, Kogut MJ. Irreversible Electroporation in Patients with Hepatocellular Carcinoma: Immediate versus Delayed Findings at MR Imaging. *Radiology*. 2016;278(1):285-94.
2. Edd JF, Horowitz L, Davalos RV, Mir LM, Rubinsky B. In vivo results of a new focal tissue ablation technique: irreversible electroporation. *IEEE Trans Biomed Eng*. 2006;53(7):1409-15.
3. Davalos RV, Mir IL, Rubinsky B. Tissue ablation with irreversible electroporation. *Ann Biomed Eng*. 2005;33(2):223-31.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

4. Mahmood F, Hansen RH, Agerholm-Larsen B, Gissel H, Ibsen P, Gehl J. Detection of electroporation-induced membrane permeabilization states in the brain using diffusion-weighted MRI. *Acta Oncol.* 2015;54(3):289-97.

5. Al-Sakere B, Andre F, Bernat C, Connault E, Opolon P, Davalos RV, et al. Tumor ablation with irreversible electroporation. *PLoS One.* 2007;2(11):e1135.

6. Marty M, Sersa G, Garbay JR, al. e. Electrochemotherapy—an easy, highly effective and safe treatment of cutaneous and subcutaneous metastases: results of ESOPE (European Standard Operating Procedures of Electrochemotherapy) study. *Eur J Cancer* 2006;Suppl(4):3–13.

7. Chung DJ, Sung K, Osuagwu FC, Wu HH, Lassman C, Lu DS. Contrast Enhancement Patterns after Irreversible Electroporation: Experimental Study of CT Perfusion Correlated to Histopathology in Normal Porcine Liver. *J Vasc Interv Radiol.* 2016;27(1):104-11.

8. Laguebe M, Silve A, Mir LM, Poignard C. Conducting and permeable states of cell membrane submitted to high voltage pulses: mathematical and numerical studies validated by the experiments. *J Theor Biol.* 2014;360:83-94.

9. Seror O. Percutaneous hepatic ablation: what needs to be known in 2014. *Diagn Interv Imaging.* 2014;95(7-8):665-75.