A Simple Model of Coevolution for Macroscopic and Microscopic Levels

Matteo Morini^{1,2}, Sarah de Nigris¹

¹Université de Lyon, Inria, UCB Lyon 1, Laboratoire de l'Informatique du Parallélisme UMR 5668, DANTE, CNRS, Institut Rhône-Alpin des Systèmes Complèxes, ENS Lyon

²Department of Economics and Statistics, University of Torino

IMT, Lucca, 22-23 February, 2017

Outline

Introduction

Segregation Metric

Palatability Map

Influence

Relocation rule

Preliminary Results

References

Future lines of research

Questions

The Schelling Model of Ethnic Residential Dynamics

- The Schelling Model of Ethnic Residential Dynamics
- Measured

- The Schelling Model of Ethnic Residential Dynamics
- Measured
 - Segregation metric (spatial autocorrelation)

- The Schelling Model of Ethnic Residential Dynamics
- Measured
 - Segregation metric (spatial autocorrelation)
 - Detection of Stationary States

- The Schelling Model of Ethnic Residential Dynamics
- Measured
 - Segregation metric (spatial autocorrelation)
 - Detection of Stationary States
 - Intrinsic tolerance

- The Schelling Model of Ethnic Residential Dynamics
- Measured
 - Segregation metric (spatial autocorrelation)
 - Detection of Stationary States
 - Intrinsic tolerance
 - "Social" tolerance

- The Schelling Model of Ethnic Residential Dynamics
- Measured
 - Segregation metric (spatial autocorrelation)
 - Detection of Stationary States
 - Intrinsic tolerance
 - "Social" tolerance
 - Expectation: segregation reinforcement

- The Schelling Model of Ethnic Residential Dynamics
- Measured
 - Segregation metric (spatial autocorrelation)
 - Detection of Stationary States
 - Intrinsic tolerance
 - "Social" tolerance
 - Expectation: segregation reinforcement
 - Apparent outcome: alter transition towards segregation in form of mixed patterns (not segregated)

- The Schelling Model of Ethnic Residential Dynamics
- Measured
 - Segregation metric (spatial autocorrelation)
 - Detection of Stationary States
 - Intrinsic tolerance
 - "Social" tolerance
 - Expectation: segregation reinforcement
 - Apparent outcome: alter transition towards segregation in form of mixed patterns (not segregated)

- The Schelling Model of Ethnic Residential Dynamics
- Measured
 - Segregation metric (spatial autocorrelation)
 - Detection of Stationary States
 - Intrinsic tolerance
 - "Social" tolerance
 - Expectation: segregation reinforcement
 - Apparent outcome: alter transition towards segregation in form of mixed patterns (not segregated)

Augmented

- The Schelling Model of Ethnic Residential Dynamics
- Measured
 - Segregation metric (spatial autocorrelation)
 - Detection of Stationary States
 - Intrinsic tolerance
 - "Social" tolerance
 - Expectation: segregation reinforcement
 - Apparent outcome: alter transition towards segregation in form of mixed patterns (not segregated)

- Palatability Map
- Networking

- The Schelling Model of Ethnic Residential Dynamics
- Measured
 - Segregation metric (spatial autocorrelation)
 - Detection of Stationary States
 - Intrinsic tolerance
 - "Social" tolerance
 - Expectation: segregation reinforcement
 - Apparent outcome: alter transition towards segregation in form of mixed patterns (not segregated)

- Palatability Map
- Networking
- Influence

- The Schelling Model of Ethnic Residential Dynamics
- Measured
 - Segregation metric (spatial autocorrelation)
 - Detection of Stationary States
 - Intrinsic tolerance
 - "Social" tolerance
 - Expectation: segregation reinforcement
 - Apparent outcome: alter transition towards segregation in form of mixed patterns (not segregated)

- Palatability Map
- Networking
- Influence
- Informed relocation decision

Segregation Metric

Spatial Autocorrelation Metric

Moran's Global I (Anselin, 1995)

$$I = n/S_0 \frac{\sum_i \sum_j z_i w_{i,j} z_j}{\sum_i z_i z_i}$$

$$z_i = y_i - \overline{y}$$

$$w_{i,j} =$$
spatial weight



Segregation Metric

Spatial Autocorrelation Metric

- Moran's Global I (Anselin, 1995)
 - "Queen" contiguity

$$I = n/S_0 \frac{\sum_i \sum_j z_i w_{i,j} z_j}{\sum_i z_i z_i}$$

$$z_i = y_i - \overline{y}$$

$$w_{i,j} =$$
spatial weight



Segregation Metric

Spatial Autocorrelation Metric

- Moran's Global I (Anselin, 1995)
 - "Queen" contiguity
 - Missing neighbors

$$I = n/S_0 \frac{\sum_i \sum_j z_i w_{i,j} z_j}{\sum_i z_i z_i}$$

$$z_i = y_i - \overline{y}$$

$$w_{i,j} =$$
 spatial weight



Palatability Map

▶ Individual "world" map $_{p}M_{i}$

- Individual "world" map pMi
- Tiles rating updated tick after tick

- Individual "world" map pMi
- Tiles rating updated tick after tick
- Time spent

- Individual "world" map pMi
- Tiles rating updated tick after tick
- Time spent
 - on tile: increase tile value (incrementally)

- Individual "world" map pMi
- Tiles rating updated tick after tick
- Time spent
 - on tile: increase tile value (incrementally)
 - moving away: decrease tile value (one-shot)

Networked Agents

- Networked Agents
 - Spatial Neighbors become Network Neighbors

- Networked Agents
 - Spatial Neighbors become Network Neighbors
 - Time Persistence

- Networked Agents
 - Spatial Neighbors become Network Neighbors
 - Time Persistence
 - "Type" Distribution according to **Tolerance Parameter** $\tau/2$

- Networked Agents
 - Spatial Neighbors become Network Neighbors
 - Time Persistence
 - "Type" Distribution according to **Tolerance Parameter** $\tau/2$
 - $\tau = 0$: same-type "friends" only

- Networked Agents
 - Spatial Neighbors become Network Neighbors
 - Time Persistence
 - "Type" Distribution according to **Tolerance Parameter** $\tau/2$
 - $\tau = 0$: same-type "friends" only
 - $\tau = 1$: equal share of types

- Networked Agents
 - Spatial Neighbors become Network Neighbors
 - Time Persistence
 - "Type" Distribution according to **Tolerance Parameter** $\tau/2$
 - $\tau = 0$: same-type "friends" only
 - $\tau = 1$: equal share of types

• Stubborness: $\sigma \in [0, 1]$

- Networked Agents
 - Spatial Neighbors become Network Neighbors
 - Time Persistence
 - "Type" Distribution according to **Tolerance Parameter** $\tau/2$
 - $\tau = 0$: same-type "friends" only
 - $\tau = 1$: equal share of types
- Stubborness: $\sigma \in [0, 1]$
 - Incorporating Palatability maps from networked agents

Relocation rule

Final decision

Final decision

_pM_i [Personal matrix, a.k.a. palatability map]

Final decision

- _pM_i [Personal matrix, a.k.a. palatability map]
- ▶ _pM_j(j = 1,...,n) [Connected agents matrixes]

Final decision

- _pM_i [Personal matrix, a.k.a. palatability map]
- ▶ _pM_j(j = 1,..., n) [Connected agents matrixes]

•
$$_f M_{i,j(u)} = \sigma_p M_i + (1-\sigma) \sum_{j=1p}^n M_j$$

Final decision

- pMi [Personal matrix, a.k.a. palatability map]
- ▶ _pM_j(j = 1,..., n) [Connected agents matrixes]

•
$$_f M_{i,j(u)} = \sigma_p M_i + (1-\sigma) \sum_{j=1p}^n M_j$$

Hop to best, empty tile

Preliminary Results

Segregation without influence



Figure 2 : Standard model.

Preliminary Results

Segregation with influence



Figure 3 : Augmented model.

Preliminary Results

Segregation with influence (Zoom on steps [400:410])



Figure 4 : Augmented model cyclic behaviour.

References

- Schelling T.C., Dynamic models of segregation, J Math Soc 1.2 (1971): 143-186
- 2. Grauwin S. et al., *Competition Between Collective and Individual Dynamics*, PNAS 106(49) (2009): 20622-20626
- 3. Erez H. and Benenson I., *The Schelling Model of Ethnic Residential Dynamics: Beyond the Integrated - Segregated Dichotomy of Patterns*, JASSS 15 (1) 6 (2012)
- Anselin L., Local Indicators of Spatial Association, Geogr An 27(2) (1995): 93-115
- 5. Lee S.-I., *Developing a bivariate spatial association measure: an integration of Pearson's r and Moran's I*, J Geogr Syst 3 (2001): 369-385.
- Gandica, Y., Gargiulo, F., & Carletti, T., Can topology reshape segregation patterns?, Chaos, Solitons & Fractals 90 (2016): 46-54.

Segregation Metric Refinements

Moran's Global I (Anselin, 1995)

Segregation Metric Refinements

- Moran's Global I (Anselin, 1995)
- Refined measure (Lee, 2001): L statistic integrating

Segregation Metric Refinements

- Moran's Global I (Anselin, 1995)
- ▶ Refined measure (Lee, 2001): L statistic integrating
 - Pearson's r as an aspatial bivariate association measure

Segregation Metric Refinements

- Moran's Global I (Anselin, 1995)
- ▶ Refined measure (Lee, 2001): *L* statistic integrating
 - Pearson's r as an aspatial bivariate association measure
 - Moran's I as a univariate spatial association measure

Questions

Questions

Questions