



**HAL**  
open science

# Rational invariants of ternary forms under the orthogonal group

Paul Görlach, Evelyne Hubert, Théo Papadopoulo

► **To cite this version:**

Paul Görlach, Evelyne Hubert, Théo Papadopoulo. Rational invariants of ternary forms under the orthogonal group. 2017. hal-01570853v2

**HAL Id: hal-01570853**

**<https://inria.hal.science/hal-01570853v2>**

Preprint submitted on 18 Aug 2017 (v2), last revised 12 Jul 2018 (v4)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# RATIONAL INVARIANTS OF TERNARY FORMS UNDER THE ORTHOGONAL GROUP

PAUL GÖRLACH, EVELYNE HUBERT, AND THÉO PAPADOPOULOU

ABSTRACT. In this article we determine a generating set of rational invariants of minimal cardinality for the action of the orthogonal group  $O_3$  on the space  $\mathbb{R}[x, y, z]_{2d}$  of ternary forms of even degree  $2d$ . The construction relies on two key ingredients: On one hand, the Slice Lemma allows us to reduce the problem to determining the invariants for the action on a subspace of the finite subgroup  $B_3$  of signed permutations. On the other hand, our construction relies in a fundamental way on specific bases of harmonic polynomials. These bases provide maps with prescribed  $B_3$ -equivariance properties. Our explicit construction of these bases should be relevant well beyond the scope of this paper. The expression of the  $B_3$ -invariants can then be given in a compact form as the composition of two equivariant maps. Instead of providing (cumbersome) explicit expressions for the  $O_3$ -invariants, we provide efficient algorithms for their evaluation and rewriting. We also use the constructed  $B_3$ -invariants to determine the  $O_3$ -orbit locus and provide an algorithm for the inverse problem of finding an element in  $\mathbb{R}[x, y, z]_{2d}$  with prescribed values for its invariants. These are the computational issues relevant in brain imaging.

## CONTENTS

1. Introduction	1
2. Preliminaries	6
3. The Slice Method	10
4. Invariants of ternary quartics	13
5. Harmonic bases with $B_3$ -symmetries and invariants for higher degree	18
6. Solving the main algorithmic challenges	28
References	32

## 1. INTRODUCTION

Invariants allow to classify objects up to the action of a group of transformations. In this article we determine a set of generating rational invariants of minimal cardinality for the action of the orthogonal group  $O_3$  on the space  $\mathbb{R}[x, y, z]_{2d}$  of ternary forms of even degree  $2d$ . We do not give the explicit expression for these invariants, but provide an algorithmic way of evaluating them for any ternary form.

Classical Invariant Theory [GY10] is centered around the action of the general linear group on homogeneous polynomials, with an emphasis on binary forms. Yet, the orthogonal group arises in applications as the relevant group of transformations, especially in three-dimensional space. Its relevance to brain imaging is the original motivation for the present article.

Computational Invariant Theory [DK02, GY10, Stu08] has long focused on polynomial invariants. In the case of the group  $O_3$ , any two real orbits are separated by a generating set of polynomial invariants. The generating set can nonetheless be very large. For instance a generating set of 64 polynomial invariants for the action of  $O_3$  on  $\mathbb{R}[x, y, z]_4$  is determined as a subset of a minimal generating set of polynomial invariants of the action of  $O_3$  on the *elasticity tensor* in [OKA17]. There, the problem is mapped to the joint action of

---

*Date:* August 18, 2017.

$\mathrm{SL}_2(\mathbb{C})$  on binary forms of different degrees and resolved by Gordan’s algorithm [GY10, Oli16] so that the invariants are given as transvectants.

A generating set of rational invariants separates general orbits – this remains true for any groups, even for non-reductive groups. Rational invariants can thus prove to be sufficient, and sometimes more relevant, in applications [HL12, HL13, HL16] and in connection with other mathematical disciplines [HK07b, Hub12]. A practical and very general algorithm to compute a generating set of these first appeared in [HK07a]; see also [DK15]. The case of the action of  $\mathrm{O}_3$  on  $\mathbb{R}[x, y, z]_4$ , a 15-dimensional space, is nonetheless not easily tractable by this algorithm. In the case of  $\mathbb{R}[x, y, z]_4$ , the 12 generating invariants we construct in this article are seen as being uniquely determined by their restrictions to a *slice*  $\Lambda_4$ , which is here a 12-dimensional subspace. The knowledge of these restrictions is proved to be sufficient to evaluate the invariants at any point in the space  $\mathbb{R}[x, y, z]_4$ . The systematic computational use of characterization of invariants in terms of their restrictions to a *slice* is a novel approach to Computational Invariant Theory that we shall introduce.

More generally, by virtue of the so-called *Slice Lemma*, the field of rational invariants of the action of  $\mathrm{O}_3$  on  $\mathbb{R}[x, y, z]_{2d}$  is isomorphic to the field of rational invariants for the action of the finite subgroup  $\mathrm{B}_3$  of signed permutations on a subspace  $\Lambda_{2d}$  of  $\mathbb{R}[x, y, z]_{2d}$ . Working with this isomorphism, we provide efficient algorithms for the evaluation of a (minimal) generating set of  $\mathrm{O}_3$  invariants based on a (minimal) generating set of  $\mathrm{B}_3$ -invariants. Finding an element of  $\mathbb{R}[x, y, z]_{2d}$  with prescribed values of  $\mathrm{O}_3$ -invariants and rewriting any  $\mathrm{O}_3$ -invariant in terms of the generating set are also made possible through the specific generating sets of  $\mathrm{B}_3$ -invariants we construct.

$\mathbb{R}[x, y, z]_{2d}$  decomposes into a direct sum of  $\mathrm{O}_3$ -invariant vector spaces determined by the harmonic polynomials of degree  $2d, 2d - 2, \dots, 2, 0$ . The construction of the  $\mathrm{B}_3$ -invariants relies in a fundamental way on specific bases of harmonic polynomials. These bases provide maps with prescribed  $\mathrm{B}_3$ -equivariance properties. Our explicit construction of these bases should be relevant well beyond the scope of this paper. The explicit expression of the  $\mathrm{B}_3$ -invariants can then be given in a compact form as the composition of two equivariant maps. Both the rewriting and the inverse problem can be made explicit for this well-structured set of invariants.

The whole construction is first made explicit for  $\mathbb{R}[x, y, z]_4$  and then extended to  $\mathbb{R}[x, y, z]_{2d}$ . It should then be clear how one can obtain the rational invariants of the joint action of  $\mathrm{O}_3$  on similar spaces, as for instance the integrity tensor.

In the rest of this section we first introduce the geometrical motivation and then the context of application for our constructions around the invariants of the action of  $\mathrm{O}_3$  on  $\mathbb{R}[x, y, z]_{2d}$ . Notations, preliminary material and a formal statement of the problems appear then in Section 2. In Section 3, we describe the technique for reducing the construction of invariants for the action of  $\mathrm{O}_3$  on  $\mathbb{R}[x, y, z]_{2d}$  to the one for the action of a finite subgroup, the signed symmetric group, on a subspace. Based on this, we construct a generating set of rational invariants for the case of ternary quartic forms in Section 4. We extend this approach in Section 5 to ternary forms of arbitrary even degree; central there is the construction of bases of vector spaces of harmonic polynomial with specific equivariant properties with respect to the signed symmetric group. Finally, we solve the principal algorithmic problems associated with rational invariants in Section 6.

**1.1. Motivation: spherical functions up to rotation.** In geometric terms, the results in this paper allow to determine when two general centrally symmetric closed surfaces in  $\mathbb{R}^3$  differ only by a rotation.

The surfaces we consider are given by a continuous function  $f: \mathbb{S}^2 \rightarrow \mathbb{R}$  on the unit sphere  $\mathbb{S}^2 := \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 = 1\}$ . The surface is then defined as the set of points

$$\mathcal{S} := \{(x f(x, y, z), y f(x, y, z), z f(x, y, z)) \in \mathbb{R}^3 \mid (x, y, z) \in \mathbb{S}^2\} \subset \mathbb{R}^3,$$

i.e. for each point on the unit sphere we rescale its distance to the origin according to the function  $f$ . For example, the surface described by a constant function  $f \equiv r$  (for some  $r \in \mathbb{R}$ ) is the sphere with radius  $|r|$  centered at the origin. As  $f$  takes on more general functions, a large variety of different surfaces can be described. For the surface to be symmetric with respect to the origin, one needs the function  $f$  to satisfy




FIGURE 1.1. Two centrally symmetric surfaces only differing by a rotation.

the property

$$(1.1) \quad f(p) = f(-p) \quad \forall p \in \mathbb{S}^2.$$

Since the unit sphere  $\mathbb{S}^2 \subset \mathbb{R}^3$  is a compact set, the Stone–Weierstraß Theorem implies that a given continuous function  $\mathbb{S}^2 \rightarrow \mathbb{R}$  can be approximated arbitrarily well by polynomial functions, i.e. by functions  $f: \mathbb{S}^2 \rightarrow \mathbb{R}$  of the form

$$(1.2) \quad f(x, y, z) = \sum_{i+j+k \leq d} a_{ijk} x^i y^j z^k.$$

The symmetry property (1.1) is then fulfilled if and only  $a_{ijk} = 0$  whenever  $i + j + k$  is odd, in other words,  $f$  must only consist of even-degree terms. We can then rewrite  $f$  in such a way that all its monomials are of the same degree, by multiplying monomials of small degree with a suitable power of  $x^2 + y^2 + z^2$  (which does not change the values of  $f$  since  $x^2 + y^2 + z^2 = 1$  for all points  $(x, y, z)$  on the sphere).

We therefore model closed surfaces that are centrally symmetric with respect to the origin by  $f: \mathbb{S}^2 \rightarrow \mathbb{R}$ , a polynomial function of degree  $2d$ :

$$f(x, y, z) = \sum_{i+j+k=2d} a_{ijk} x^i y^j z^k.$$

The modeled surface can then be encoded *exactly* by simply storing the  $\frac{1}{2}(2d+2)(2d+1)$  numbers  $a_{ijk} \in \mathbb{R}$ . However, from such an encoding of a surface via the coefficients  $a_{ijk}$  of its defining polynomial, it is not immediately apparent when the defined surfaces have the same geometric shape, only differing by a rotation. As an example, Figure 1.1 depicts the two surfaces  $\mathcal{S}_1$  and  $\mathcal{S}_2$  described by

$$\begin{aligned} f_1(x, y, z) &:= 18x^2 - 27y^2 + 18z^2 \quad \text{and} \\ f_2(x, y, z) &:= 13x^2 + 20xy - 20xz - 2y^2 + 40yz - 2z^2, \end{aligned}$$

respectively, whose numerical description in terms of coefficients look very distinct, but whose shapes look very similar. Indeed  $\mathcal{S}_2$  arises from  $\mathcal{S}_1$  by applying the rotation determined by the matrix

$$\frac{1}{3} \begin{pmatrix} 2 & -1 & -2 \\ 2 & 2 & 1 \\ 1 & -2 & 2 \end{pmatrix},$$

so that

$$f_2(x, y, z) = f_1\left(\frac{2x + 2y + z}{3}, \frac{-x + 2y - 2z}{3}, \frac{-2x + y - 2z}{3}\right).$$

In general, we want to consider surfaces to be of the same shape if they differ by a rotation or, equivalently, by an orthogonal transformation: Since we only consider surfaces symmetric with respect to the origin, two surfaces differing by an orthogonal transformation also differ by a rotation.

The question which arises is: *How can we (algorithmically) decide whether two surfaces only differ by an orthogonal transformation, only by examining the coefficients of their defining polynomial?*

Regarding this question, we should to be aware that if  $f: S^2 \rightarrow \mathbb{R}$  describes  $\mathcal{S} \subset \mathbb{R}^3$ , then its negative  $-f$  describes the same set  $\mathcal{S} \subset \mathbb{R}^3$ . With the intuition of describing the surface by deforming the unit sphere, this ambiguity corresponds to turning the closed surface  $\mathcal{S}$  inside out. We typically want to think of the surfaces defined by  $f$  and by  $-f$  as two distinct objects (even though they are equal as subsets of  $\mathbb{R}^2$ ). For example, the surface defined by the constant function  $f \equiv 1$  is the unit sphere, while the surface defined by  $f \equiv -1$  should be considered as the unit sphere turned inside out.

Then the question above corresponds to the following algebraic question:

**Question 1.1.** *Given two polynomials  $f = \sum_{i+j+k=2d} a_{ijk} x^i y^j z^k$  and  $g = \sum_{i+j+k=2d} b_{ijk} x^i y^j z^k$ , how can we decide in terms of their coefficients  $a_{ijk}$  and  $b_{ijk}$  whether or not there exists an orthogonal transformation  $\varphi: \mathbb{R}^3 \rightarrow \mathbb{R}^3$  such that  $f = g \circ \varphi$  as functions  $\mathbb{R}^3 \rightarrow \mathbb{R}$ ?*

Once we can decide whether two polynomials define equally shaped surfaces, a natural question is how to uniquely encode the shape of a surface, i.e. an equivalence class of surfaces up to orthogonal transformations. The corresponding question for polynomials is:

**Question 1.2.** *How can we encode in a unique way equivalence classes of polynomials up to orthogonal transformations?*

For the purpose of illustration we discuss next the case of homogeneous polynomials of degree two, i.e. *quadratic forms*. In the following section, we shall describe the mathematical setup for addressing Questions 1.1 and 1.2 with the rational invariants of the action of the orthogonal group on ternary forms of even degree.

*Illustration for the case of quadratic surfaces.* Quadratic forms are in one-to-one correspondence with symmetric  $3 \times 3$ -matrices, as indeed we can write (1.2) as:

$$(1.3) \quad f(x, y, z) = \underbrace{\begin{pmatrix} x & y & z \end{pmatrix} \begin{pmatrix} a_{200} & \frac{1}{2}a_{110} & \frac{1}{2}a_{101} \\ \frac{1}{2}a_{110} & a_{020} & \frac{1}{2}a_{011} \\ \frac{1}{2}a_{101} & \frac{1}{2}a_{011} & a_{002} \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix}}_{=:A}$$

When we compose  $f$  with a rotation, given by a matrix  $R$ , we obtain another homogeneous polynomial of degree 2. The defining symmetric matrix (as in 1.3) of the obtained polynomial is then  $RAR^T$ . For two symmetric matrices  $A$  and  $B$ , Linear Algebra then tells us that there is an orthogonal matrix  $R$  such that  $B = RAR^T$  if and only if  $A$  and  $B$  have the same eigenvalues. Yet, eigenvalues are algebraic functions of the entries of a matrix and thus cannot be easily expressed symbolically. It is thus easier to compare the coefficients of the characteristic polynomials. Up to scalar factors, these are:

$$(1.4) \quad \begin{aligned} e_1 &:= a_{200} + a_{020} + a_{002}, \\ e_2 &:= 4a_{020}a_{200} + 4a_{002}a_{020} + 4a_{002}a_{200} - a_{110}^2 - a_{101}^2 - a_{011}^2, \\ e_3 &:= 4a_{002}a_{020}a_{200} + a_{101}a_{110}a_{011} - a_{101}^2a_{020} - a_{011}^2a_{200} - a_{002}a_{110}^2. \end{aligned}$$

The result is that two homogeneous polynomial of degree 2 are obtained from one another by an orthogonal transformation if and only if the functions  $e_1, e_2, e_3$  of their coefficients take the same values. This answers Question 1.1 in this case. The functions  $e_1, e_2, e_3$  also provide a solution to Question 1.2. To each polynomial of degree 2 we can associate a point in  $\mathbb{R}^3$  whose components are the values of the functions  $e_1, e_2, e_3$  for this polynomial. All equivalent polynomials will be mapped to the same point. One can check for example that this point is  $(9, -2592, -34992)$  for both  $f_1$  and  $f_2$  defined above.

**1.2. Application in Neuroimaging.** DMRI (Diffusion Magnetic Resonance Imaging), developed since 1984, is a field where the problem of characterizing spherical shapes arises. DMRI measures non-invasively the amount of water diffusion in biological tissues. These diffusion properties are intimately related to the




FIGURE 1.2. Slice of a brain diffusion image (a). The yellow boxes indicate the regions that are zoomed in (b) showing the individual voxel shapes. From [PGD14].

structure of these tissues notably in fibrous structures such as muscles – e.g. the heart – or brain white matter. The measurements are made in a rectangular portion of the 3D space sampled by voxels (volumic picture element). At each voxel, the diffusion information is captured as a function  $f: S^2 \rightarrow \mathbb{R}_{\geq 0}$  of orientation  $v$ , sampled at several points  $v_i$  on the sphere  $S^2$ . This (discretized) spherical function  $f$  is called an ODF (diffusion Orientation Distribution Function) [TRWW03]. Here  $f(v)$ , for  $v \in S^2$ , is a positive value that characterizes the amount of diffusion in the direction  $v$  in a given amount of time  $T$  (which is a constant defined in the diffusion acquisition protocol).  $f$  satisfies  $f(v) = f(-v)$ , meaning that  $f$  is symmetric with respect to the origin. Figure 1.2 shows an example of such a diffusion image for brain white matter.

Integrating the discrete diffusion information into a global model is interesting for e.g. noise reduction or for computing mathematical properties of  $f$ . This global model was initially a simple a ternary quadratic function (or *tensor*), leading to what is called Diffusion Tensor Imaging (DTI). For DTI, all rotational invariants can be easily computed as functions of  $e_1, e_2, e_3$  given in (1.4). Biologically interesting quantities have been computed as combination of these invariants, with clinical validation showing their interest to detect various important diseases such as Alzheimer’s or Parkinson’s diseases, writer’s cramp disease, etc [DVW<sup>+</sup>09, LMM<sup>+</sup>10].

With diffusion MRI improvements, acquiring more and more samples on the sphere  $S^2$  is both possible and interesting (High Angular Resolution Diffusion-weighted Imaging - HARDI - technique [Tuc02]). This allows the study of complex fibrous structures, e.g. revealing places in the brain where white matter fiber bundles are crossing, with the potential of characterizing more completely such complex structures and their abnormalities which may occur in patients. Fully exploiting these images can only be realized using shapes for the diffusion information more complex than those described by quadratic forms.

One popular solution is to represent  $f$  using a spherical harmonic basis [Fra02]. Because of the symmetry property mentioned above, only spherical harmonics of even order are meaningful. Using spherical harmonic functions up to order  $2d$  is mathematically equivalent to representing  $f$  as a homogeneous polynomials of even degree  $2d$ :

$$f(x, y, z) = \sum_{i+j+k=2d} a_{ijk} x^i y^j z^k.$$

Of particular interest is the case of degree  $2d = 4$  introduced in [BVF08] as it is the simplest model generalizing the DTI case ( $2d = 2$ ) and as the positivity of  $f$  can be imposed using Hilbert’s ternary quartic theorem. Since then, numerous efforts have been made to obtain a complete invariant description for these homogeneous polynomials of degree  $2d = 4$  [BP07, GPD12a, GPD12b], with the prospect of finding more powerful generalizations of the biomarkers found in the DTI case. A maximal set of functionally independent local invariants were proposed in [PGD14]; they are algebraic functions of  $a_{ijk} \in \mathbb{R}$ . Functionally independent sets of polynomial invariants were also determined in [CV15] for the cases  $2d = 4$  and  $2d = 6$ .

The present article provides a complete (functionally independent and generating) set of rational invariants, for all even degrees  $2d$ .

## 2. PRELIMINARIES

In order to set the notation, we review the definitions of the action of  $O_3$  on the vector space of forms  $V_d := \mathbb{R}[x, y, z]_d$  and of its rational invariants. We then elaborate on the Harmonic Decomposition of  $V_{2d}$ .

**2.1. The action of the orthogonal group on homogeneous polynomials.** We start out with a few notational conventions. We denote by  $\mathbb{R}[x, y, z]_d$  the set of homogeneous polynomials

$$f = \sum_{i+j+k=d} a_{ijk} x^i y^j z^k$$

of degree  $d$  in the three variables  $x, y, z$  with real coefficients  $a_{ijk} \in \mathbb{R}$ . By  $O_3 \subset \mathbb{R}^{3 \times 3}$  we denote the group of orthogonal matrices and we consider the action of  $O_3$  on  $\mathbb{R}[x, y, z]$  given by

$$\begin{aligned} O_3 \times \mathbb{R}[x, y, z]_d &\rightarrow \mathbb{R}[x, y, z]_d \\ (g, f) &\mapsto f \circ g^{-1}. \end{aligned}$$

By  $f \circ g^{-1}$ , we mean the composition of the orthogonal transformation  $g^{-1}: \mathbb{R}^3 \rightarrow \mathbb{R}^3$  with the polynomial function  $f: \mathbb{R}^3 \rightarrow \mathbb{R}$ , resulting in a different polynomial function that we denote by  $gf$ . Note that  $gf$  is again homogeneous of degree  $d$ , so the above action is well-defined. As  $g^{-1} = g^T$ , the polynomial  $gf$  is obtained from  $f$  by applying the substitutions

$$x \mapsto g_{11}x + g_{21}y + g_{31}z, \quad y \mapsto g_{12}x + g_{22}y + g_{32}z, \quad z \mapsto g_{13}x + g_{23}y + g_{33}z,$$

when  $g = (g_{ij}) \in O_3$  is an orthogonal  $3 \times 3$ -matrix with entries  $g_{ij}$ . The use of the inverse  $g^{-1} (= g^T)$  instead of  $g$  in the above composition is only of notational importance, guaranteeing  $g_1(g_2f) = (g_1g_2)f$  for all  $g_1, g_2 \in O_3$ .

A counting argument shows that there are  $N := \binom{d+2}{2} = \frac{1}{2}(d+2)(d+1)$  monomials  $x^i y^j z^k$  of degree  $d$ , so formally,  $\mathbb{R}[x, y, z]_d$  is an  $N$ -dimensional vector space over  $\mathbb{R}$ . In order to stress this point of view, we shall denote

$$V_d := \mathbb{R}[x, y, z]_d$$

and, accordingly, we shall from now on typically denote elements of  $V_d$  by letters like  $v$  or  $w$ . The monomials  $\{x^i y^j z^k \mid i + j + k = d\}$  form a basis for this vector space. But alternative bases will prove useful in this paper.

As  $g(v + w) = gv + gw$  and  $g(\lambda v) = \lambda(gv)$  holds for all  $g \in O_3$ ,  $v, w \in V_d$  and  $\lambda \in \mathbb{R}$ , the action of  $O_3$  on  $V_d$  is a *linear* group action. If we fix a basis for  $V_d$ , there is polynomial map from  $O_3$  to the group of invertible  $N \times N$  matrices that describes the action of  $O_3$  on  $V_d$  in this basis.

**Example 2.1** (The action on quadratic forms). We show the matrix that provides the transformation by  $g = (g_{ij})_{1 \leq i, j \leq 3} \in O_3$  on  $V_2$  in two different bases, starting with the monomial basis.

If

$$f := a_{200}x^2 + a_{110}xy + a_{020}y^2 + a_{101}xz + a_{011}yz + a_{002}z^2$$

and

$$gf = b_{200}x^2 + b_{110}xy + b_{020}y^2 + b_{101}xz + b_{011}yz + b_{002}z^2,$$

then the relation between the coefficients of  $f$  and  $gf$  is given by the matricial equality:

$$\begin{pmatrix} b_{200} \\ b_{110} \\ b_{020} \\ b_{101} \\ b_{011} \\ b_{002} \end{pmatrix} = \begin{pmatrix} g_{11}^2 & g_{11}g_{12} & g_{12}^2 & g_{11}g_{13} & g_{12}g_{13} & g_{13}^2 \\ 2g_{21}g_{11} & g_{21}g_{12} + g_{11}g_{22} & 2g_{22}g_{12} & g_{21}g_{13} + g_{11}g_{23} & g_{22}g_{13} + g_{12}g_{23} & 2g_{23}g_{13} \\ g_{21}^2 & g_{21}g_{22} & g_{22}^2 & g_{21}g_{23} & g_{22}g_{23} & g_{23}^2 \\ 2g_{31}g_{11} & g_{31}g_{12} + g_{11}g_{32} & 2g_{32}g_{12} & g_{31}g_{13} + g_{11}g_{33} & g_{32}g_{13} + g_{12}g_{33} & 2g_{33}g_{13} \\ 2g_{31}g_{21} & g_{31}g_{22} + g_{21}g_{32} & 2g_{32}g_{22} & g_{31}g_{23} + g_{21}g_{33} & g_{23}g_{32} + g_{22}g_{33} & 2g_{33}g_{23} \\ g_{31}^2 & g_{31}g_{32} & g_{32}^2 & g_{31}g_{33} & g_{32}g_{33} & g_{33}^2 \end{pmatrix} \begin{pmatrix} a_{200} \\ a_{110} \\ a_{020} \\ a_{101} \\ a_{011} \\ a_{002} \end{pmatrix}$$

Let us denote by  $R(g)$  the matrix in the above equality.

Alternatively the set of polynomials  $\{x^2 + y^2 + z^2, yz, zx, xy, y^2 - z^2, z^2 - x^2\}$  forms a basis for  $V_2$ . Let  $P$  be the matrix of change of basis, i.e.

$$(x^2 + y^2 + z^2 \quad yz \quad zx \quad xy \quad y^2 - z^2 \quad z^2 - x^2) = (x^2 \quad xy \quad y^2 \quad zx \quad yz \quad z^2) P$$

If

$$f = a_1(x^2 + y^2 + z^2) + a_2 yz + a_3 zx + a_4 xy + a_5(y^2 - z^2) + a_6(z^2 - x^2)$$

and

$$gf = b_1(x^2 + y^2 + z^2) + b_2 yz + b_3 zx + b_4 xy + b_5(y^2 - z^2) + b_6(z^2 - x^2),$$

then the vector of coefficients  $b = (b_1 \dots b_6)^T$  is related to the vector of coefficients  $a = (a_1 \dots a_6)^T$  by the matrixial equality:

$$b = \tilde{R}(g) a, \quad \text{where } \tilde{R}(g) = P^{-1} R(g) P$$

Using that  $g \in O_3$ , one can compute that

$$\tilde{R}(g) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & g_{23}g_{32} + g_{22}g_{33} & g_{31}g_{23} + g_{21}g_{33} & g_{31}g_{22} + g_{21}g_{32} & 2g_{32}g_{22} - 2g_{33}g_{23} & 2g_{33}g_{23} - 2g_{31}g_{21} \\ 0 & g_{32}g_{13} + g_{12}g_{33} & g_{31}g_{13} + g_{11}g_{33} & g_{31}g_{12} + g_{11}g_{32} & 2g_{32}g_{12} - 2g_{33}g_{13} & 2g_{33}g_{13} - 2g_{31}g_{11} \\ 0 & g_{22}g_{13} + g_{12}g_{23} & g_{21}g_{13} + g_{11}g_{23} & g_{21}g_{12} + g_{11}g_{22} & 2g_{22}g_{12} - 2g_{23}g_{13} & 2g_{23}g_{13} - 2g_{21}g_{11} \\ 0 & g_{22}g_{23} & g_{21}g_{23} & g_{21}g_{22} & g_{22}^2 - g_{23}^2 & g_{23}^2 - g_{21}^2 \\ 0 & g_{32}g_{33} + g_{22}g_{23} & g_{31}g_{33} + g_{21}g_{23} & g_{31}g_{32} + g_{21}g_{22} & g_{22}^2 + g_{32}^2 - g_{23}^2 - g_{33}^2 & g_{23}^2 + g_{33}^2 - g_{21}^2 - g_{31}^2 \end{pmatrix}.$$

Hence the linear spaces generated by  $x^2 + y^2 + z^2$ , on one hand, and by  $\{yz, zx, xy, y^2 - z^2, z^2 - x^2\}$ , on the other hand, are both invariant under the linear action of  $O_3$  on  $V_2$ .

As we shall see in Section 2.3, the quadratic polynomial  $x^2 + y^2 + z^2 \in V_2$  also plays a special role in the action of  $O_3$  on  $V_d$  for  $d > 2$  due to the property that it is fixed by the action of  $O_3$ .

**2.2. Rational invariants and algorithmic problems.** We denote by  $\mathbb{R}(V_d)$  the set of rational functions  $p: V_d \dashrightarrow \mathbb{R}$  on the vector space  $V_d$ . Explicitly, if we denote elements of  $V_d$  as

$$\sum_{i+j+k=d} a_{ijk} x^i y^j z^k,$$

then

$$(2.1) \quad \mathbb{R}(V_d) = \mathbb{R}(a_{ijk} \mid i, j, k \geq 0, i + j + k = d)$$

is the field of rational expressions (i.e. quotients of polynomial expressions) in the variables  $a_{ijk}$ .

The explicit description of elements in  $\mathbb{R}(V_d)$  as expressions in the variables  $a_{ijk}$  however reflects the choice of the monomial basis  $\{x^i y^j z^k \mid i + j + k = d\}$  for  $V_d$ . Later on, we will work with a different basis for  $V_d$  giving rise to an alternative presentations of elements in  $\mathbb{R}(V_d)$ . We will therefore mostly avoid the description of  $\mathbb{R}(V_d)$  as in (2.1) in the following discussions.

When working with rational functions  $p = \frac{p_1}{p_0} \in \mathbb{R}(V_d)$  (where  $p_1, p_0: V_d \rightarrow \mathbb{R}$  are polynomial functions and  $p_2 \neq 0$ ), there is always an issue of division by zero: The function  $p = \frac{p_1}{p_0}$  is only defined *defined on a general point*, namely, on the set  $\{v \in V_d \mid p_0(v) \neq 0\}$ . To keep this in mind, the rational function is denoted by a dashed arrow  $p: V_d \dashrightarrow \mathbb{R}$ . Similarly, the equalities we shall write have to be understood wherever they are defined, i.e. where all denominators involved are non-zero.

More generally, it is said that a statement  $\mathcal{P}$  about points  $v \in V$  in a given  $\mathbb{R}$ -vector space  $V$  (e.g.  $V = V_d$ ) holds for a **general point** if there exists a non-zero polynomial function  $p_0: V \rightarrow \mathbb{R}$  such that  $\mathcal{P}$  holds for all points  $v \in V$  where  $p_0(v) \neq 0$ .

The set of **rational invariants** for the action of  $O_3$  on  $V_d$  is defined as

$$\mathbb{R}(V_d)^{O_3} := \{p \in \mathbb{R}(V_d) \mid p(v) = p(gv) \forall v \in V_d, g \in O_3\}.$$

For any  $d$  there exists a *finite set*  $\{p_1, \dots, p_m\} \subset \mathbb{R}(V_d)^{O_3}$  of rational invariants which generate  $\mathbb{R}(V_d)^{O_3}$  as a field extension of  $\mathbb{R}$ . This means that any other rational invariant  $q \in \mathbb{R}(V_d)^{O_3}$  can be written as a rational



expression in terms of  $p_1, \dots, p_m$ . We call such a finite set  $\{p_1, \dots, p_m\}$  a **generating set of rational invariants**.

In contrast to the ring of polynomial invariants, the fact that the field of rational invariants is finitely generated is just an instance of the elementary algebraic fact: Any subfield of a finitely generated field is again finitely generated (see for example [Isa09, Theorem 24.9]). A lower bound for the cardinality of a generating set of rational invariants is given by the following result, implied by [PV94, Corollary of Theorem 2.3]:

**Theorem 2.2.** *For  $d \geq 2$ , any generating set of rational invariants for the action of  $O_3$  on  $V_d$  consists of at least  $\dim V_d - \dim O_3 = \binom{d+2}{2} - 3$  elements.*

An important characterization of rational invariants is given by the following theorem, for which we refer to [PV94, Lemma 2.1, Theorem 2.3] or [Ros56].

**Theorem 2.3.** *Rational invariants  $p_1, \dots, p_m \in \mathbb{R}(V_d)^{O_3}$  form a generating set if and only if for general points  $v, w \in V_d$  the following holds:*

$$w = gv \text{ for some } g \in O_3 \Leftrightarrow p_i(v) = p_i(w) \quad \forall i \in \{1, \dots, m\}.$$

In this article, we explicitly construct a generating set of rational invariants for the action of  $O_3$  on  $V_{2d}$ , i.e. even degree forms, whose cardinality attains the lower bound indicated in Theorem 2.2. The results appear in Corollary 4.4 for  $V_4$  and in Corollary 5.13 for  $V_{2d}$ ,  $d \geq 2$ .

Theorem 2.3 gives a crucial justification for approaching Question 1.1 with rational invariants. It further addresses Question 1.2 of how to encode even degree polynomials up to orthogonal transformations: We may encode a general point  $v \in V_{2d}$  as the  $m$ -tuple  $(p_1(v), \dots, p_m(v)) \in \mathbb{R}^m$ . Then the  $m$ -tuples of  $v \in V_{2d}$  and  $w \in V_{2d}$  are equal if and only if  $v$  and  $w$  are equivalent under the action of  $O_3$ .

Associated with this approach are the following main Algorithmic Problems:

1. **Characterization:** Determine a set of generating rational invariants  $p_1, \dots, p_m \in \mathbb{R}(V_{2d})^{O_3}$ .
2. **Evaluation:** Evaluate  $p_1(v), \dots, p_m(v)$  for a given (general) point  $v \in V_{2d}$  in an efficient and robust way.
3. **Rewriting:** Given a rational invariant  $q \in \mathbb{R}(V_{2d})^{O_3}$ , express  $q$  as a rational expression in terms of  $p_1, \dots, p_m$ .
4. **Reconstruction:** Which  $m$ -tuples  $\mu = (\mu_1, \dots, \mu_m) \in \mathbb{R}^m$  lie in the image of the map

$$\pi: V_{2d} \dashrightarrow \mathbb{R}^m, \quad v \mapsto (p_1(v), \dots, p_m(v))?$$

If  $\mu$  lies in the image of  $\pi$ , find a representative  $v \in V_{2d}$  such that  $\pi(v) = \mu$ .

General algorithms for computing a generating set of rational invariants based on Gröbner basis algorithms exist [HK07a], [DK15, Section 4.10], but the complexity increases drastically with the dimension of  $V_{2d}$ , which in turn grows quadratically in  $d$ . Already for  $2d = 4$ , these general methods are far from a feasible computation. Furthermore, these algorithms typically do not produce a minimal generating set. In this article we shall demonstrate the efficiency of a more structural approach for describing a generating set of rational invariants with minimal cardinality. How to address the algorithmic challenges 2–4 will become more apparent from our construction of the generating rational invariants, and we will examine them in detail in Section 6.

The field of rational invariants  $\mathbb{R}(V_{2d})^{O_3}$  is the quotient field of the ring of polynomial invariants  $\mathbb{R}[V_{2d}]^{O_3}$  [PV94, Theorem 3.3]; Any rational invariant can be written as the quotient of two polynomial invariants. Yet determining a generating set of polynomial invariants is a somewhat more arduous task. In [OKA17] the author determined a minimal set of generating polynomial invariants for the action of  $O_3$  on  $\mathcal{H}_4 \oplus 2\mathcal{H}_2 \oplus 2\mathcal{H}_0$ , the space of the elasticity tensor. We can extract from this basis a set of 64 polynomial invariants that generate  $\mathbb{R}[V_4]^{O_3}$ . These invariants are computed thanks to Gordan’s algorithm [GY10, Oli16] after the problem is reduced to the action of  $SL(2, \mathbb{C})$  on binary forms through Cartan’s map. One has to observe though that polynomial invariants separate the real orbits of  $O_3$  [Sch01, Proposition 2.3], while rational invariants will only separate *general* orbits (Theorem 2.3).

**2.3. Harmonic Decomposition.** In the study of the action of  $O_3$  on  $V_d$ , the *Harmonic Decomposition* plays a central role. We start out by collecting some basic facts about the apolar inner product and harmonic polynomials.

**Definition 2.4.** The *apolar inner product*  $\langle \cdot, \cdot \rangle_d: V_d \times V_d \rightarrow \mathbb{R}$  is defined as follows: If  $v = \sum_{ijk} a_{ijk} x^i y^j z^k \in V_d$  and  $w = \sum_{ijk} b_{ijk} x^i y^j z^k \in V_d$ , we define

$$\langle v, w \rangle_d := \sum_{i,j,k} i!j!k! a_{ijk} b_{ijk}.$$

The apolar inner product arises naturally as follows: Identifying  $V_d = \mathbb{R}[x, y, z]_d$  with the space of symmetric tensors  $\text{Sym}^d(\mathbb{R}^3)^* = \text{Sym}^d(V_1)$ , it is (up to a rescaling constant) the inner product inherited from  $V_1^{\otimes d}$  via the embedding  $\text{Sym}^d(V_1) \hookrightarrow V_1^{\otimes d}$ . Here, the inner product on  $V_1^{\otimes d}$  is induced by the standard inner product on  $V_1 = (\mathbb{R}^3)^*$ .

Since the group  $O_3$  preserves the standard inner product on  $V_1$ , this viewpoint leads to the following fact.

**Proposition 2.5.** The apolar inner product is preserved by the group action of  $O_3$ , i.e. if  $v, w \in V_d$  and  $g \in O_3$ , then  $\langle gv, gw \rangle_d = \langle v, w \rangle_d$ .

Another intrinsic formulation of the apolar product [Veg00] is given as follows: For a polynomial function  $f(x, y, z) \in \mathbb{R}[x, y, z]_d$ , let  $f(\partial)$  be the differential operator obtained from  $f$  by replacing  $x, y, z$  respectively by  $\frac{\partial}{\partial x}, \frac{\partial}{\partial y}, \frac{\partial}{\partial z}$ . One then checks that for  $f_1, f_2 \in \mathbb{R}[x, y, z]_d$  the apolar inner product is given by

$$\langle f_1, f_2 \rangle_d = f_1(\partial)(f_2) = f_2(\partial)(f_1).$$

With this viewpoint, Proposition 2.5 follows by observing that  $\langle gf_1, f_2 \rangle_d = \langle f_1, g^T f_2 \rangle_d$  holds for all group elements  $g$ .

From now on, we denote

$$q := x^2 + y^2 + z^2 \in V_2.$$

This  $q \in V_2$  plays a special role as it is fixed by the action of  $O_3$ :  $gq = q$  for all  $g \in O_3$ .

**Definition 2.6.** For any  $d \geq 2$ , we consider the inclusion of vector spaces

$$V_{d-2} \hookrightarrow V_d, \quad v \mapsto q \cdot v.$$

and its image  $qV_{d-2} \subset V_d$ , which is given by those polynomials in  $V_d = \mathbb{R}[x, y, z]_d$  that are divisible by  $q$ .

We define the subspace  $\mathcal{H}_d \subset V_d$  of **harmonic polynomials** of degree  $d$  to be the orthogonal complement of  $qV_{d-2} \subset V_d$  with respect to the apolar inner product on  $V_d$ .

An immediate consequence of the invariance of  $q$  and Proposition 2.5 is the following observation.

**Proposition 2.7.** Let  $g \in O_3$  and  $v \in V_d$ . Then the following holds:

- (i) If  $v \in qV_{d-2} \subset V_d$ , then also  $gv \in qV_{d-2}$ .
- (ii) If  $v \in \mathcal{H}_d$ , then also  $gv \in \mathcal{H}_d$ .

Harmonic functions are typically introduced as the functions  $f$  such that  $\Delta(f) = 0$ , where  $\Delta$  is the Laplacian operator, i.e.  $\Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$ , [ABR01]. We can see that this is equivalent to Definition 2.6 as follows: Understanding the apolar inner product via differential operators as described above, we have

$$\langle f_1 f_3, f_2 \rangle_d = \langle f_1, f_3(\partial)(f_2) \rangle_k \quad \text{for all } f_1 \in \mathbb{R}[x, y, z]_k, f_2 \in \mathbb{R}[x, y, z]_d, f_3 \in \mathbb{R}[x, y, z]_{d-k}.$$

In particular,  $\langle f_1, qf_2 \rangle_d = \langle \Delta(f_1), f_2 \rangle_{d-2}$  holds for all  $f_1 \in \mathbb{R}[x, y, z]_d, f_2 \in \mathbb{R}[x, y, z]_{d-2}$ , so that  $f_1 \in \mathcal{H}_d$  if and only if  $\Delta(f_1) = 0$ .

So far, all formulations have been made for arbitrary degree  $d$ . However, we are ultimately interested in the case of even degree only. We therefore will from now on and for the remainder of the article only consider the case of degree  $2d$ .

By Definition 2.6, there is an orthogonal decomposition  $V_{2d} = \mathcal{H}_{2d} \oplus \mathfrak{q}V_{2d-2}$ . For  $d \geq 2$  we can also decompose  $V_{2d-2}$  in this manner, so we may iterate this decomposition which leads to the following observation, [ABR01, Theorem 5.7].

**Theorem 2.8** (Harmonic Decomposition). *For  $d \geq 2$  there is a decomposition*

$$V_{2d} = \mathcal{H}_{2d} \oplus \mathfrak{q}\mathcal{H}_{2d-2} \oplus \mathfrak{q}^2\mathcal{H}_{2d-4} \cdots \oplus \mathfrak{q}^{d-2}\mathcal{H}_4 \oplus \mathfrak{q}^{d-1}V_2,$$

*i.e. each  $v \in V_{2d}$  can uniquely be written as a sum*

$$v = h_{2d} + \mathfrak{q}h_{2d-2} + \mathfrak{q}^2h_{2d-4} + \dots + \mathfrak{q}^{d-2}h_4 + \mathfrak{q}^{d-1}v'$$

*where  $h_{2k} \in \mathcal{H}_{2k}$  and  $v' \in V_2$ .*

We mention at this point that it would be possible to refine Theorem 2.8 by further decomposing  $V_2 = \mathcal{H}_2 \oplus \mathbb{R}\mathfrak{q}$ , but this is not beneficial for our purpose.

Even though  $V_{2d} = \mathcal{H}_{2d} \oplus \mathfrak{q}V_{2d-2}$  is a decomposition into orthogonal subspaces with respect to the apolar inner product, we wish to highlight that the decomposition in Theorem 2.8 is *not* an orthogonal decomposition if  $2d > 4$ . This is due to the fact that in general  $\langle v, w \rangle_{2d-2} \neq \langle \mathfrak{q}v, \mathfrak{q}w \rangle_{2d}$  for  $v, w \in V_{2d-2}$ . On the other hand, it remains true that each of the subspaces in the Harmonic Decomposition is preserved under every element  $g$  of  $O_3$  as in Proposition 2.7.

Different bases for the vector spaces  $\mathcal{H}_{2d} \subset V_{2d}$  of harmonic polynomials are used in applications. A frequent choice in practice is the basis of *spherical harmonics* [AH12] which are usually given as functions in spherical coordinates. In Section 5, we shall construct another basis for  $\mathcal{H}_{2d}$  that exhibits certain symmetries with respect to the group of signed permutations.

### 3. THE SLICE METHOD

Our aim is to determine a generating set of rational invariants for the linear action of the orthogonal group  $O_3$  on the vector spaces  $V_{2d}$  of even degree ternary forms. The group  $O_3$  is infinite and of dimension 3 as an algebraic group. We reduce the problem to the simpler question of determining rational invariants for the linear action of a *finite* group  $B_3$  (contained in  $O_3$  as a subgroup) on a subspace  $\Lambda_{2d}$  of  $V_{2d}$ .

**3.1. The Slice Lemma.** We now introduce the general technique for the reduction mentioned above, called the *slice method*. For the formulation, we abstract from our specific setting: We consider a linear action of an algebraic group  $G$  on a finite-dimensional  $\mathbb{R}$ -vector space  $V$ , denoted

$$G \times V \rightarrow V, \quad (g, v) \mapsto gv.$$

In our particular case, we have  $G = O_3$  and  $V = V_{2d}$ , and the action is defined in Section 2.1. We denote by  $\mathbb{R}(V)$  the field of rational functions and by  $\mathbb{R}(V)^G$  the finitely generated subfield of rational invariants, as introduced already for  $V = V_{2d}$  and  $G = O_3$ .

The main technique for the announced reduction is known as the *Slice Method* [CTS07, Section 3.1], [Pop94]. It is based on the following definition.

**Definition 3.1.** *Consider a linear group action of an algebraic group  $G$  on a finite-dimensional  $\mathbb{R}$ -vector space  $V$ . A subspace  $\Lambda \subset V$  is called a **slice** for the group action, and the subgroup*

$$B := \{g \in G \mid gs \in \Lambda \ \forall s \in \Lambda\} \subset G$$

*is called its **stabilizer**, if the following two properties hold:*

- (i) *For a general point  $v \in V$  there exists  $g \in G$  such that  $gv \in \Lambda$ .*
- (ii) *For a general point  $s \in \Lambda$  the following holds: If  $g \in G$  is such that  $gs \in \Lambda$ , then  $g \in B$ .*

The Slice Lemma then states that rational invariants of the action of  $G$  on  $V$  are in one-to-one correspondence with rational invariants of the smaller group  $B \subset G$  on the slice  $\Lambda \subset V$ :

**Theorem 3.2** (Slice Lemma). *Let  $\Lambda$  be a slice of a linear action of an algebraic group  $G$  on a finite-dimensional  $\mathbb{R}$ -vector space  $V$ , and let  $B$  be its stabilizer. Then there is a field isomorphism over  $\mathbb{R}$  between rational invariants*

$$\varrho: \mathbb{R}(V)^G \xrightarrow{\cong} \mathbb{R}(\Lambda)^B, \quad p \mapsto p|_{\Lambda},$$

which restricts a rational invariant  $p: V \dashrightarrow \mathbb{R}$  to  $p|_{\Lambda}: \Lambda \dashrightarrow \mathbb{R}$ .

This observation goes back to [Ses62], and we refer to [CTS07, Theorem 3.1] for a proof. Explicitly, the inverse to  $\varrho: \mathbb{R}(V)^G \xrightarrow{\cong} \mathbb{R}(\Lambda)^B$  is given by

$$\begin{aligned} \varrho^{-1}: \mathbb{R}(\Lambda)^B &\rightarrow \mathbb{R}(V)^G, \\ q &\mapsto (V \dashrightarrow \mathbb{R}, \quad v \mapsto q(gv), \text{ where } g \in G \text{ such that } gv \in \Lambda). \end{aligned}$$

We will apply Theorem 3.2 for  $G = O_3$ ,  $V = V_{2d}$  and a suitable choice for the slice  $\Lambda$ . The consequence of Theorem 3.2 for the construction of a generating set of rational invariants is the following.

**Corollary 3.3.** *Let  $\Lambda$  be a slice of a linear action of an algebraic group  $G$  on a finite-dimensional  $\mathbb{R}$ -vector space  $V$ , and let  $B$  be its stabilizer. If  $\mathcal{I} = \{p_1, \dots, p_m\}$  is a generating set of rational invariants for the action of  $B$  on  $\Lambda$ , then  $\mathcal{J} := \{\varrho^{-1}(p_1), \dots, \varrho^{-1}(p_m)\}$  is a generating set of rational invariants for the action of  $G$  on  $V$  (where  $\varrho$  is given as above).*

*Proof.* Let  $p_0 \in \mathbb{R}(V)^G$ . By assumption,  $\varrho(p_0) \in \mathbb{R}(\Lambda)^B$  can be written as a rational expression in the generators  $p_1, \dots, p_m$ . Since  $\varrho$  is a field isomorphism,  $p_0 = \varrho^{-1}(\varrho(p_0))$  is the same rational expression in  $\varrho^{-1}(p_1), \dots, \varrho^{-1}(p_m)$ .  $\square$

A corresponding statement for polynomial invariants requires much stronger hypotheses on the slice. In particular, even if the generating set for  $\mathbb{R}(\Lambda)^B$  consists of *polynomial expressions*  $p_1, \dots, p_m \in \mathbb{R}[\Lambda]^B$ , the construction described above typically introduces denominators, so that  $\varrho^{-1}(p_1), \dots, \varrho^{-1}(p_m) \in \mathbb{R}(V)^G$  become *rational expressions*.

**3.2. A slice for  $V_{2d}$ .** We now describe a slice  $\Lambda_{2d} \subset V_{2d}$  for the action of  $O_3$  on  $V_{2d}$  for any  $d \geq 1$ .

We recall from Section 1.1 the description of the action of  $O_3$  on  $V_2$ : Elements of  $V_2$  are ternary quadratic forms and they can be identified with symmetric  $3 \times 3$ -matrices as in (1.3). If the associated symmetric matrix of  $v \in V_2$  is  $A$ , then for any  $g \in O_3 \subset \mathbb{R}^{3 \times 3}$ , the associated symmetric matrix of  $gv \in V_2$  is the matrix product  $gAg^T$ .

**Definition 3.4.** *Let  $\Lambda_2 \subset V_2$  denote the subspace of quadratic forms whose associated symmetric matrix is diagonal. Explicitly,*

$$\Lambda_2 = \{\lambda_1 x^2 + \lambda_2 y^2 + \lambda_3 z^2 \in V_2 \mid \lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}\}.$$

*For  $d \geq 2$  we consider the Harmonic Decomposition of  $V_{2d}$  from Theorem 2.8 and define  $\Lambda_{2d} \subset V_{2d}$  to be the subspace*

$$\Lambda_{2d} := \mathcal{H}_{2d} \oplus \mathfrak{q}\mathcal{H}_{2d-2} \oplus \dots \oplus \mathfrak{q}^{d-2}\mathcal{H}_4 \oplus \mathfrak{q}^{d-1}\Lambda_2.$$

In other words, elements of the subspace  $\Lambda_{2d}$  are those  $v \in V_{2d}$  that can be written as

$$v = h_{2d} + \mathfrak{q}h_{2d-2} + \mathfrak{q}^2h_{2d-4} + \dots + \mathfrak{q}^{d-2}h_4 + \mathfrak{q}^{d-1}v'$$

with  $h_{2k} \in \mathcal{H}_{2k}$  and  $v' \in \Lambda_2$  a quadratic form whose associated symmetric matrix is diagonal. The main observation is now the following:

**Proposition 3.5.** *Let  $d \geq 1$ . The subspace  $\Lambda_{2d} \subset V_{2d}$  is a slice for the action of  $O_3$  on  $V_{2d}$  and its stabilizer is the group  $B_3 \subset O_3$  of signed permutation matrices. In particular, there is a one-to-one correspondence between rational invariants*

$$\varrho: \mathbb{R}(V_{2d})^{O_3} \xrightarrow{\cong} \mathbb{R}(\Lambda_{2d})^{B_3}$$

given by the restriction of rational functions.

We recall that a *signed permutation matrix* is a matrix for which each row and each column contain only one non-zero entry and this entry is either 1 or -1. Below we will remark on further structural descriptions of the group  $B_3$ .

*Proof of Proposition 3.5.* The second statement is a consequence of the first statement by Theorem 3.2.

Let

$$v = h_{2d} + qh_{2d-2} + q^2h_{2d-4} + \dots + q^{d-2}h_4 + q^{d-1}v'$$

be the Harmonic Decomposition of an element  $v \in V_{2d}$  and let  $A \in \mathbb{R}^{3 \times 3}$  be the symmetric matrix associated to the quadratic form  $v' \in V_2$ . By the Spectral Theorem for symmetric matrices, there exists an orthogonal matrix  $g \in O_3 \subset \mathbb{R}^{3 \times 3}$  such that  $gAg^T$  is a diagonal matrix (whose diagonal entries are the eigenvalues of  $A$ ). Since  $gAg^T$  is the associated symmetric matrix of  $gv' \in V_2$ , this means that

$$gv = (gh_{2d}) + q(gh_{2d-2}) + q^2(gh_{2d-4}) + \dots + q^{d-2}(gh_4) + q^{d-1}(gv')$$

is contained in  $\Lambda_{2d}$ . This verifies property (i) of Definition 3.1 for  $\Lambda_{2d} \subset V_{2d}$ .

Note that an orthogonal matrix  $g \in O_3$  is an element of the stabilizer  $B := \{g \in O_3 \mid gv \in \Lambda_{2d} \forall v \in \Lambda_{2d}\}$  if and only if the matrix product  $g \operatorname{diag}(\lambda_1, \lambda_2, \lambda_3)g^T$  is again a diagonal matrix for all values of  $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$ . This means that the rows of the matrix  $g$  are orthonormal eigenvectors of  $\operatorname{diag}(\lambda_1, \lambda_2, \lambda_3)$  for all values of  $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$ . In particular, this holds when  $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$  are *distinct*, in which case the only unit eigenvectors are  $\pm e_i$  (where  $e_1, e_2, e_3$  is the standard basis of  $\mathbb{R}^3$ ). Hence,  $g$  is a signed permutation matrix. This establishes the description of the stabilizer  $B = B_3$ .

Similarly, to verify property (ii) of Definition 3.1, we only need to show: If  $g \in O_3$  is such that for a *general point*  $(\lambda_1, \lambda_2, \lambda_3) \in \mathbb{R}^3$ , the matrix product  $g \operatorname{diag}(\lambda_1, \lambda_2, \lambda_3)g^T$  is a diagonal matrix, then  $g \in B_3$ . Indeed, we have just seen that this property holds whenever  $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}$  are distinct values, i.e. wherever the polynomial function

$$\mathbb{R}^3 \rightarrow \mathbb{R}, \quad (\lambda_1, \lambda_2, \lambda_3) \mapsto (\lambda_1 - \lambda_2)(\lambda_2 - \lambda_3)(\lambda_3 - \lambda_1)$$

does not vanish. □

We note that in the proof we verified property (i) of Definition 3.1 for *all*  $v \in V_{2d}$ , while for property (ii) we had to make use of the notion of a *general point*.

Above, we introduced the group  $B_3 \subset O_3$  of signed permutation matrices. This is a *finite* group with  $2^3 \cdot 3! = 48$  elements and is known as the *hyperoctahedral group* or as the wreath product  $\mathfrak{S}_2 \wr \mathfrak{S}_3$ , i.e. as the semidirect product of the symmetric group  $\mathfrak{S}_3$  with the abelian group  $\{1, -1\}^3$ . For an explicit description, we introduce the following notation.

**Notation 3.6.** For  $\sigma \in \mathfrak{S}_3$  we denote by  $g_\sigma \in O_3$  the permutation matrix such that

$$g_\sigma \begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \alpha_3 \end{pmatrix} = \begin{pmatrix} \alpha_{\sigma^{-1}(1)} \\ \alpha_{\sigma^{-1}(2)} \\ \alpha_{\sigma^{-1}(3)} \end{pmatrix}$$

For  $\tau = (\tau_1, \tau_2, \tau_3) \in \{1, -1\}^3$  we write  $g_\tau := \operatorname{diag}(\tau_1, \tau_2, \tau_3) \in O_3$  and we will call these matrices **sign-change matrices**.

Then  $B_3$  is the smallest subgroup of  $O_3$  containing all permutation matrices and all sign-change matrices: Each signed permutation matrix  $g \in B_3$  can uniquely be written as  $g = g_\tau g_\sigma$  with  $\tau \in \{1, -1\}^3$ ,  $\sigma \in \mathfrak{S}_3$ . Indeed,  $\tau_i$  must be 1 or  $-1$  corresponding to the sign of the unique non-zero entry in the  $i$ -th row of the matrix  $g$ . For this  $\tau$ , the matrix  $g_\sigma := g_\tau^{-1}g$  is a permutation matrix. Analogously, we can also write each  $g \in B_3$  uniquely as  $g = g_\sigma g_{\tau'}$  with  $\tau' \in \{1, -1\}^3$ ,  $\sigma \in \mathfrak{S}_3$ . However,  $\tau' \neq \tau$  in general.

**3.3. Illustration on quadratic forms.** We illustrate the Slice Method on the well-known case of quadratic forms: The slice  $\Lambda_2 \subset V_2$  is given as

$$\Lambda_2 = \{\lambda_1 x^2 + \lambda_2 y^2 + \lambda_3 z^3 \mid \lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}\}.$$

By Theorem 3.2, the field of rational  $O_3$ -invariants on  $V_2$  is isomorphic to the field of rational  $B_3$ -invariants on  $\Lambda_2$ .

We observe that sign-change matrices in  $B_3$  act trivially on  $\Lambda_2$ , so

$$\mathbb{R}(\Lambda_2)^{B_3} = \mathbb{R}(\Lambda_2)^{\mathfrak{S}_3} = \mathbb{R}(\lambda_1, \lambda_2, \lambda_3)^{\mathfrak{S}_3},$$

where the symmetric group  $\mathfrak{S}_3 \subset B_3$  acts by permuting  $\lambda_1, \lambda_2, \lambda_3$  (which we view as coordinates on  $\Lambda_2$ ). By the Fundamental Theorem of symmetric polynomials, we can thus choose the set of elementary symmetric polynomials

$$\mathcal{I} = \{\lambda_1 + \lambda_2 + \lambda_3, \lambda_1 \lambda_2 + \lambda_2 \lambda_3 + \lambda_3 \lambda_1, \lambda_1 \lambda_2 \lambda_3\}$$

as a generating set of rational invariants for  $\mathbb{R}(\Lambda_2)^{B_3}$ . By Corollary 3.3, there exist unique rational  $O_3$ -invariants  $\mathcal{J} \subset \mathbb{R}(V_2)^{O_3}$  on  $V_2$  restricting to these invariants on the subspace  $\Lambda_2$ , and they form a generating set for  $\mathbb{R}(V_2)^{O_3}$ . By uniqueness, the generating set  $\mathcal{J}$  consists up to scalars precisely of the polynomial invariants  $e_1, e_2, e_3 \in \mathbb{R}[V_2]^{O_3}$  described in (1.4), arising as the coefficients of the characteristic polynomial of the symmetric matrix corresponding to a quadratic form.

In this case, one can in fact show that  $\mathcal{J}$  even generates the ring of *polynomial* invariants  $\mathbb{R}[V_2]^{O_3}$ . This will however no longer be true for our construction of generating rational invariants in higher degree. Furthermore, the construction of  $B_3$ -invariants on the slice will be more involved, especially because the sign-change matrices in  $B_3$  no longer act trivially on  $\Lambda_{2d}$  for  $d \geq 2$ .

#### 4. INVARIANTS OF TERNARY QUARTICS

In this section we implicitly describe a generating set of rational invariants of minimal cardinality for ternary quartics under the action of  $O_3$ , i.e. for the case  $2d = 4$ . Following the approach of Section 3.1 this set of rational invariants is uniquely determined by a set of rational invariants for the action of  $B_3$  on the slice  $\Lambda_4$ . The first step is to construct an appropriate basis of  $\Lambda_4$  that is equivariant. In this basis, a minimal generating set of  $B_3$ -invariants takes a particularly compact form, and can be chosen to consist of polynomial invariants.

We provide an additional (near minimal) generating set of  $\mathbb{R}(\Lambda_4)^{B_3}$  that extends the invariants for quadratic forms. For both choices of generating invariants, we make explicit how to write any other invariant in terms of these, following [HK07a].

The construction of this section serves as a model for invariants of ternary forms in higher degree  $2d > 4$  which are treated in Section 5.

**4.1. A  $B_3$ -equivariant basis for harmonic quartics.** In order to construct  $B_3$ -invariants on the vector space  $\Lambda_4 = \mathcal{H}_4 \oplus \mathfrak{q}\Lambda_2$ , we introduce a basis of  $\mathcal{H}_4$  that exhibits certain symmetries with respect to  $B_3$ . In Section 5 we show how to construct bases of  $\mathcal{H}_{2d}$  with analogous symmetry properties for arbitrary  $d \geq 2$ .

**Proposition 4.1.** *The following nine ternary quartics form a basis for the  $\mathbb{R}$ -vector space  $\mathcal{H}_4$ :*

$$\begin{aligned} r_1 &:= y^4 - 6y^2 z^2 + z^4, & s_1 &:= y^3 z - y z^3, & t_1 &:= 6x^2 y z - y^3 z - y z^3, \\ r_2 &:= z^4 - 6z^2 x^2 + x^4, & s_2 &:= z^3 x - z x^3, & t_2 &:= 6y^2 z x - z^3 x - z x^3, \\ r_3 &:= x^4 - 6x^2 y^2 + y^4, & s_3 &:= x^3 y - x y^3, & t_3 &:= 6z^2 x y - x^3 y - x y^3. \end{aligned}$$




FIGURE 4.1. Harmonic quartics

The group  $B_3 \subset O_3$  acts on  $\mathcal{H}_4$  with respect to this basis as follows: For  $\sigma \in \mathfrak{S}_3$  and  $\tau \in \{1, -1\}^3$ , the corresponding permutation and sign-change matrices  $g_\sigma$  and  $g_\tau$  act by

$$\begin{aligned} g_\sigma r_i &= r_{\sigma(i)}, & g_\sigma s_i &= \text{sgn}(\sigma) s_{\sigma(i)}, & g_\sigma t_i &= t_{\sigma(i)}, \\ g_\tau r_i &= r_i, & g_\tau s_i &= \frac{\tau_1 \tau_2 \tau_3}{\tau_i} s_i, & g_\tau t_i &= \frac{\tau_1 \tau_2 \tau_3}{\tau_i} t_i. \end{aligned}$$

Figure 4.1 illustrates this basis of  $\mathcal{H}_4$  as explained in Section 1.1, similar to Figure 1.1. Here, different colors within one picture correspond to different signs of the harmonic polynomial at corresponding points.

*Proof.* One can first check that the polynomials  $r_i, s_i, t_i$  ( $1 \leq i \leq 3$ ) are in the kernel of the Laplacian operator  $\Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$ . As  $\mathcal{H}_4$  is a 9-dimensional  $\mathbb{R}$ -vector space, it suffices to check that all  $r_i, s_i, t_i$  ( $1 \leq i \leq 3$ ) are linearly independent. This easily follows from examining the monomials occurring in their expressions.

Note that a permutation matrix in  $B_3$  acts by applying the corresponding permutation to the variables  $x, y, z$ , and a sign-change matrix acts by replacing some of the variables  $x, y, z$  by their negatives. The claim about the action of  $B_3$  is then read off the formulas for the basis elements.  $\square$

A polynomial  $v$  in  $\Lambda_4 = \mathfrak{q}\Lambda_2 \oplus \mathcal{H}_4$  can thus be written as

$$(4.1) \quad v = (x^2 + y^2 + z^2) \sum_{i=1}^3 \lambda_i x_i^2 + \sum_{i=1}^3 (\alpha_i r_i + \beta_i s_i + \gamma_i t_i).$$

We introduce the vector of coefficients  $\lambda = (\lambda_1 \ \lambda_2 \ \lambda_3)^T$  and similarly  $\alpha, \beta$  and  $\gamma$ . Then Proposition 4.1 shows: For  $\sigma \in \mathfrak{S}_3$ , if  $g_\sigma v$  has coefficients  $(\tilde{\lambda}, \tilde{\alpha}, \tilde{\beta}, \tilde{\gamma})$ , then

$$\tilde{\lambda} = g_\sigma \lambda, \quad \tilde{\alpha} = g_\sigma \alpha, \quad \tilde{\beta} = \text{sgn}(\sigma) g_\sigma \beta, \quad \tilde{\gamma} = g_\sigma \gamma.$$

For  $\tau \in \{1, -1\}^3$ , if  $g_\tau v$  is determined by  $(\bar{\lambda}, \bar{\alpha}, \bar{\beta}, \bar{\gamma})$ , then

$$\bar{\lambda} = \lambda, \quad \bar{\alpha} = \alpha, \quad \bar{\beta} = \det(g_\tau) g_\tau \beta, \quad \bar{\gamma} = \det(g_\tau) g_\tau \gamma.$$

**4.2.  $B_3$ -invariants on the slice.** The rational invariants of  $B_3$  on  $\Lambda_4$  can be obtained computationally by applying the general algorithm for generating sets of rational invariants presented in [HK07a]. The results obtained with this approach allow to suspect nice structures, when the basis of Proposition 4.1 is used. We accordingly present generating sets of invariants as the results of the composition of equivariant maps. We first present a minimal generating set, which consists of 12 algebraically independent polynomials, and then a generating set that consist of 13 rational invariants, including the invariants for  $\Lambda_2$ .

**Lemma 4.2.** *The maps  $W_1, W_2: \Lambda_2 \rightarrow \mathbb{R}^{3 \times 3}$  and  $W_3, W_4: \Lambda_4 \rightarrow \mathbb{R}^3$  whose values at  $v$  as in (4.1) are respectively given by*

$$W_1(v) = \begin{pmatrix} 1 & \lambda_1 & \lambda_1^2 \\ 1 & \lambda_2 & \lambda_2^2 \\ 1 & \lambda_3 & \lambda_3^2 \end{pmatrix}, \quad W_2(v) = \begin{pmatrix} 1 & \gamma_1^2 & \gamma_1^4 \\ 1 & \gamma_2^2 & \gamma_2^4 \\ 1 & \gamma_3^2 & \gamma_3^4 \end{pmatrix}, \quad W_3(v) = \begin{pmatrix} \beta_1 \gamma_1 [\lambda] \\ \beta_2 \gamma_2 [\lambda] \\ \beta_3 \gamma_3 [\lambda] \end{pmatrix} \quad \text{and} \quad W_4(v) = \begin{pmatrix} \beta_1 \gamma_1 [\gamma^2] \\ \beta_2 \gamma_2 [\gamma^2] \\ \beta_3 \gamma_3 [\gamma^2] \end{pmatrix},$$

where

$$[\lambda] := (\lambda_1 - \lambda_2)(\lambda_2 - \lambda_3)(\lambda_3 - \lambda_1) = \det W_1(v) \quad \text{and} \quad [\gamma^2] := (\gamma_1^2 - \gamma_2^2)(\gamma_2^2 - \gamma_3^2)(\gamma_3^2 - \gamma_1^2) = \det W_2(v),$$

are equivariant in the sense that  $W_i(g_\sigma v) = g_\sigma W_i(v)$  and  $W_i(g_\tau v) = W_i(v)$ .

*Proof.* This is a direct consequence of the discussion in Section 4.1 of the action of  $B_3$  on the basis of Proposition 4.1.  $\square$

It follows that the entries of  $W_i^T W_j$ , for  $i, j \in \{1, 2, 3, 4\}$ , and of  $W_i^{-1} W_j$ , for  $i \in \{1, 2\}$  and  $j \in \{1, 2, 3, 4\}$ , are rational invariants on  $\Lambda_4$ . With this mechanism, we provide a generating set of invariants of minimal cardinality.

**Theorem 4.3.** *A generating set of rational invariants for  $\mathbb{R}(\Lambda_4)^{B_3}$  is given by the 12 polynomial functions  $\mathcal{I}_4 = \{\ell_i, a_i, b_i, c_i \mid 1 \leq i \leq 3\}$ , whose values at  $v$  as in (4.1) are given by*

$$\begin{aligned} c_1(v) &:= \gamma_1^2 + \gamma_2^2 + \gamma_3^2, \\ c_2(v) &:= \gamma_1 \gamma_2 \gamma_3, \\ c_3(v) &:= \gamma_1^4 + \gamma_2^4 + \gamma_3^4, \end{aligned}$$

and the entries of the  $3 \times 3$  matrix

$$\begin{pmatrix} \ell_1(v) & a_1(v) & b_1(v) \\ \ell_2(v) & a_2(v) & b_2(v) \\ \ell_3(v) & a_3(v) & b_3(v) \end{pmatrix} := \begin{pmatrix} 1 & 1 & 1 \\ \gamma_1^2 & \gamma_2^2 & \gamma_3^2 \\ \gamma_1^4 & \gamma_2^4 & \gamma_3^4 \end{pmatrix} \begin{pmatrix} \lambda_1 & \alpha_1 & \beta_1 \gamma_1 [\gamma^2] \\ \lambda_2 & \alpha_2 & \beta_2 \gamma_2 [\gamma^2] \\ \lambda_3 & \alpha_3 & \beta_3 \gamma_3 [\gamma^2] \end{pmatrix}$$

where  $[\gamma^2] := (\gamma_1^2 - \gamma_2^2)(\gamma_2^2 - \gamma_3^2)(\gamma_3^2 - \gamma_1^2)$ .

*Proof.* We first observe that the polynomials  $\ell_i, a_i, b_i \in \mathbb{R}[\Lambda_4]$  are the entries of the matrix  $W_2^T(v)W(v)$ , where  $W_2$  was introduced in Lemma 4.2 and  $W: \Lambda_2 \rightarrow \mathbb{R}^{3 \times 3}$  is given by

$$W(v) := \begin{pmatrix} \lambda_1 & \alpha_1 & \beta_1 \gamma_1 [\gamma^2] \\ \lambda_2 & \alpha_2 & \beta_2 \gamma_2 [\gamma^2] \\ \lambda_3 & \alpha_3 & \beta_3 \gamma_3 [\gamma^2] \end{pmatrix}.$$

With Lemma 4.2 one easily checks that  $W$  is equivariant in the sense that  $W(\sigma v) = g_\sigma W(v)$  and  $W(\tau v) = W(v)$ . Hence the entries of  $W_2^T(v)W(v)$  are invariants. It is also straightforward to check that  $c_1, c_2, c_3 \in \mathbb{R}[\Lambda_4]$  are polynomial invariants.



We now show in three steps that any rational invariant can be written as a rational expression in terms of  $c_i, \ell_i, a_i, b_i$ , for  $i \in \{1, 2, 3\}$ .

*Step 1: Reducing the problem to rewriting invariant polynomials.*

For a finite group, the field of rational invariants is the quotient field of the ring of invariants [PV94, Theorem 3.3]. That means that for any rational invariant  $p \in \mathbb{R}(\Lambda_4)^{\mathbb{B}_3}$  there exist  $p_1, p_0 \in \mathbb{R}[\Lambda_4]^{\mathbb{B}_3}$  such that  $p = \frac{p_1}{p_0}$ . We are thus left to show that any polynomial invariant can be written as a rational expression of  $\ell_i, a_i, b_i, c_i$ , for  $i \in \{1, 2, 3\}$ .

*Step 2: If  $p \in \mathbb{R}[\Lambda_4]^{\mathbb{B}_3}$  is an invariant whose value at  $v$  as in (4.1) is given by a polynomial expression in  $\gamma_1, \gamma_2, \gamma_3$  only, then  $p$  can be written polynomially in terms of  $c_1, c_2, c_3$ .*

First, we consider the monomials  $\gamma_1^i \gamma_2^j \gamma_3^k$  of the polynomial expression  $p$ . By Proposition 4.1, a sign-change matrix  $g_\tau = \text{diag}(\tau_1, \tau_2, \tau_3) \in \mathbb{B}_3$  acts on  $q$  by replacing  $\gamma_1^i \gamma_2^j \gamma_3^k$  by

$$\tau_1^{j+k} \tau_2^{i+k} \tau_3^{i+j} \gamma_1^i \gamma_2^j \gamma_3^k,$$

so  $p$  can only be invariant with respect to all sign-change matrices if for all its monomials  $\gamma_1^i \gamma_2^j \gamma_3^k$ , the numbers  $i + j$ ,  $i + k$  and  $j + k$  are even numbers, i.e.  $i \equiv j \equiv k \pmod{2}$ . In particular, we can write  $p$  as a polynomial in

$$c_2 = \gamma_1 \gamma_2 \gamma_3, \quad \delta_1 := \gamma_1^2, \quad \delta_2 := \gamma_2^2, \quad \delta_3 := \gamma_3^2.$$

If  $g_\sigma \in \mathbb{B}_3$  is a permutation matrix, then  $g_\sigma$  acts according to Proposition 4.1 on  $p$  by replacing  $\delta_i$  by  $\delta_{\sigma_i}$ . Therefore,  $p$  is a symmetric polynomial expression in the three variables  $\delta_1, \delta_2, \delta_3$ . By the Fundamental Theorem of symmetric functions,  $p$  can therefore be written as a polynomial expression in the three symmetric power sum polynomials

$$\begin{aligned} \delta_1 + \delta_2 + \delta_3 &= c_1, \\ \delta_1^2 + \delta_2^2 + \delta_3^2 &= c_3 \text{ and} \\ \delta_1^3 + \delta_2^3 + \delta_3^3 &= \frac{3}{2} c_1 c_3 - \frac{1}{2} c_1^3 + 3c_2^2. \end{aligned}$$

With this, we have expressed  $p$  as a polynomial expression in terms of  $c_1, c_2, c_3$ .

*Step 3: If  $p \in \mathbb{R}[\Lambda_4]^{\mathbb{B}_3}$  is a polynomial expression in  $\lambda, \alpha, \beta, \gamma$ , then  $p$  can be written as a rational expression in the invariants  $\ell, a, b, c$ .*

We have

$$\begin{pmatrix} \lambda_1 & \alpha_1 & \beta_1 \gamma_1 [\gamma^2] \\ \lambda_2 & \alpha_2 & \beta_2 \gamma_2 [\gamma^2] \\ \lambda_3 & \alpha_3 & \beta_3 \gamma_3 [\gamma^2] \end{pmatrix} = W_2^{-T}(v) \begin{pmatrix} \ell_1(v) & a_1(v) & b_1(v) \\ \ell_2(v) & a_2(v) & b_2(v) \\ \ell_3(v) & a_3(v) & b_3(v) \end{pmatrix},$$

where  $W_2(v)$  is a matrix that only involves the variables  $\gamma_1, \gamma_2, \gamma_3$ . With this, we can replace each occurrence of  $\lambda_i, \alpha_i, \beta_i$  for  $i \in \{1, 2, 3\}$ , by a linear combination of  $\ell_i, a_i, b_i$  with coefficients that are rational expressions in  $\gamma_1, \gamma_2, \gamma_3$ . Hence,  $p$  is written as a polynomial in  $\ell_i, a_i, b_i$  with coefficients that are rational expressions in  $\gamma_1, \gamma_2, \gamma_3$ . Since the  $\ell_i, a_i, b_i$  are algebraically independent, these rational expressions of  $\gamma_1, \gamma_2, \gamma_3$  must be invariant. By Step 1 and 2 they can be written as rational expressions of  $c_1, c_2, c_3$ .  $\square$

Except maybe for *Step 1*, the proof above shows how to rewrite any rational invariants in terms of the minimal generating set. One could alternatively rely on a computational proof following [HK07a]; that way shows that rewriting a rational invariant in terms of this minimal generating set of invariants can be done by applying the following rewrite rules to both the numerator and denominator. They indeed reflect a (non-reduced) Gröbner basis of the ideal of the generic orbit of the action.

$$\begin{pmatrix} \lambda_1 \\ \lambda_2 \\ \lambda_3 \end{pmatrix} \longrightarrow \frac{[\gamma^2]}{c_0} \begin{pmatrix} (\gamma_3^2 - \gamma_2^2) \gamma_2^2 \gamma_3^2 & \gamma_2^4 - \gamma_3^4 & \gamma_3^2 - \gamma_2^2 \\ (\gamma_1^2 - \gamma_3^2) \gamma_3^2 \gamma_1^2 & \gamma_3^4 - \gamma_1^4 & \gamma_1^2 - \gamma_3^2 \\ (\gamma_2^2 - \gamma_1^2) \gamma_1^2 \gamma_2^2 & \gamma_1^4 - \gamma_2^4 & \gamma_2^2 - \gamma_1^2 \end{pmatrix} \begin{pmatrix} \ell_1 \\ \ell_2 \\ \ell_3 \end{pmatrix},$$

$$\begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \alpha_3 \end{pmatrix} \longrightarrow \frac{[\gamma^2]}{c_0} \begin{pmatrix} (\gamma_3^2 - \gamma_2^2)\gamma_2^2\gamma_3^2 & \gamma_2^4 - \gamma_3^4 & \gamma_3^2 - \gamma_2^2 \\ (\gamma_1^2 - \gamma_3^2)\gamma_3^2\gamma_1^2 & \gamma_3^4 - \gamma_1^4 & \gamma_1^2 - \gamma_3^2 \\ (\gamma_2^2 - \gamma_1^2)\gamma_1^2\gamma_2^2 & \gamma_1^4 - \gamma_2^4 & \gamma_2^2 - \gamma_1^2 \end{pmatrix} \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix},$$

$$\begin{pmatrix} \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix} \longrightarrow \frac{1}{c_0 c_2} \begin{pmatrix} \gamma_2\gamma_3 & 0 & 0 \\ 0 & \gamma_3\gamma_1 & 0 \\ 0 & 0 & \gamma_1\gamma_2 \end{pmatrix} \begin{pmatrix} (\gamma_3^2 - \gamma_2^2)\gamma_2^2\gamma_3^2 & \gamma_2^4 - \gamma_3^4 & \gamma_3^2 - \gamma_2^2 \\ (\gamma_1^2 - \gamma_3^2)\gamma_3^2\gamma_1^2 & \gamma_3^4 - \gamma_1^4 & \gamma_1^2 - \gamma_3^2 \\ (\gamma_2^2 - \gamma_1^2)\gamma_1^2\gamma_2^2 & \gamma_1^4 - \gamma_2^4 & \gamma_2^2 - \gamma_1^2 \end{pmatrix} \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix},$$

$$\gamma_1 \longrightarrow \frac{1}{c_2}(c_1\gamma_2\gamma_3 - \gamma_2^3\gamma_3 - \gamma_2\gamma_3^3), \quad \gamma_2^4 \longrightarrow c_1\gamma_3^2 - \gamma_2^2\gamma_3^2 + c_1\gamma_3^2 - \gamma_3^4 + \frac{1}{2}(c_3 - c_1^2), \quad \gamma_3^6 \longrightarrow \frac{1}{2}(c_3 - c_1^2)\gamma_3^2 + c_1\gamma_3^4 + c_2^2,$$

where

$$c_0 := [\gamma^2]^2 = (\gamma_1^2 - \gamma_2^2)^2(\gamma_2^2 - \gamma_3^2)(\gamma_3^2 - \gamma_1^2)^2 = \frac{1}{2}c_3^3 - \frac{1}{4}c_1^6 - 27c_2^4 + c_1^4c_3 - \frac{5}{4}c_1^2c_2^2 - 9c_2^2c_1c_3 + 5c_2^2c_1^3.$$

The field of rational functions  $\mathbb{R}(\Lambda_4)$  has transcendence degree  $\dim \Lambda_4 = 12$  over  $\mathbb{R}$ , and since  $B_3$  is a finite group, the same is true for the field of invariants  $\mathbb{R}(\Lambda_4)^{B_3}$ . Therefore, the generating set  $\mathcal{I}_4 \subset \mathbb{R}(\Lambda_4)^{B_3}$  of rational invariants specified in Theorem 4.3 is of minimal cardinality. By Corollary 3.3 and Proposition 3.5, there are 12 unique rational invariants in  $\mathbb{R}(V_4)^{O_3}$  restricting to the  $B_3$ -invariants  $\mathcal{I}_4$  on the subspace  $\Lambda_4$ , and they form a generating set for  $\mathbb{R}(V_4)^{O_3}$ . In particular, the cardinality of this generating set attains the lower bound given in Theorem 2.2:

**Corollary 4.4.** *There exists a generating set of  $\dim V_4 - \dim O_3 = 12$  rational invariants  $\mathcal{J}_4 \subset \mathbb{R}(V_4)^{O_3}$  restricting to the invariants  $\mathcal{I}_4 \subset \mathbb{R}[\Lambda_4]^{B_3}$  on the subspace  $\Lambda_4 \subset V_4$  specified in Theorem 4.3.*

While the above generating set  $\mathcal{J}_4 \subset \mathbb{R}(V_4)^{O_3}$  (respectively,  $\mathcal{I}_4 \subset \mathbb{R}(\Lambda_4)^{B_3}$ ) is minimal, we observe that it does not directly contain the three invariants for quadratic forms from Section 3.3, which can be considered as invariants on  $V_4$  via the decomposition  $V_4 = \mathcal{H}_4 \oplus qV_2$  (respectively on  $\Lambda_4$  via  $\Lambda_4 = \mathcal{H}_4 \oplus q\Lambda_2$ ). We therefore now introduce an alternative, non-minimal, generating set of rational  $B_3$ -invariants on  $\Lambda_4$ , such that a generating set of invariants of  $B_3$  on  $\Lambda_2$  is obtained by restriction. We also make explicit how to rewrite any other invariants in terms of these.

**Theorem 4.5.** *A generating set of rational invariants for  $\mathbb{R}(\Lambda_4)^{B_3}$  is given by the 13 rational functions  $\tilde{\ell}_i, \tilde{a}_i, \tilde{b}_i, \tilde{c}_i$ , for  $i \in \{1, 2, 3\}$ , and  $\tilde{c}_0$  whose values at  $p$  as in (4.1) are given by*

$$\begin{aligned} \tilde{c}_0(v) &= \gamma_1\gamma_2\gamma_3, \\ \tilde{\ell}_1(v) &= \lambda_1 + \lambda_2 + \lambda_3, \quad \tilde{\ell}_2(v) = \lambda_1^2 + \lambda_2^2 + \lambda_3^2, \quad \tilde{\ell}_3(v) = \lambda_1^3 + \lambda_2^3 + \lambda_3^3, \end{aligned}$$

and the entries of the  $3 \times 3$  matrix

$$\begin{pmatrix} \tilde{a}_1(v) & \tilde{b}_1(v) & \tilde{c}_1(v) \\ \tilde{a}_2(v) & \tilde{b}_2(v) & \tilde{c}_2(v) \\ \tilde{a}_3(v) & \tilde{b}_3(v) & \tilde{c}_3(v) \end{pmatrix} := \begin{pmatrix} 1 & \lambda_1 & \lambda_1^2 \\ 1 & \lambda_2 & \lambda_2^2 \\ 1 & \lambda_3 & \lambda_3^2 \end{pmatrix}^{-1} \begin{pmatrix} \alpha_1 & \beta_1\gamma_1[\lambda] & \gamma_1^2 \\ \alpha_2 & \beta_2\gamma_2[\lambda] & \gamma_2^2 \\ \alpha_3 & \beta_3\gamma_3[\lambda] & \gamma_3^2 \end{pmatrix}.$$

where  $[\lambda] := (\lambda_1 - \lambda_2)(\lambda_2 - \lambda_3)(\lambda_3 - \lambda_1)$ .

Let

$$\tilde{\ell}_0 := (\lambda_1 - \lambda_2)^2(\lambda_2 - \lambda_3)^2(\lambda_3 - \lambda_1)^2 = \frac{3}{2}\tilde{\ell}_2\tilde{\ell}_1^4 - \frac{1}{6}\tilde{\ell}_1^6 + 6\tilde{\ell}_3\tilde{\ell}_2\tilde{\ell}_1 - \frac{4}{3}\tilde{\ell}_3\tilde{\ell}_1^3 - \frac{7}{2}\tilde{\ell}_2^2\tilde{\ell}_1^2 - 3\tilde{\ell}_3^2 + \frac{1}{2}\tilde{\ell}_2^3.$$

Then any rational invariant  $p = p_0/p_1 \in \mathbb{R}(\Lambda_4)^{B_3}$  can be written in terms of the above generating set by applying the following rewrite rules to both the numerator  $p_0$  and the denominator  $p_1$ :

$$\begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \alpha_3 \end{pmatrix} \longrightarrow \begin{pmatrix} 1 & \lambda_1 & \lambda_1^2 \\ 1 & \lambda_2 & \lambda_2^2 \\ 1 & \lambda_3 & \lambda_3^2 \end{pmatrix} \begin{pmatrix} \tilde{a}_1 \\ \tilde{a}_2 \\ \tilde{a}_3 \end{pmatrix},$$

$$\begin{pmatrix} \beta_1 \\ \beta_2 \\ \beta_3 \end{pmatrix} \longrightarrow \frac{[\lambda]}{\tilde{c}_0\tilde{\ell}_0} \begin{pmatrix} \gamma_2\gamma_3 & 0 & 0 \\ 0 & \gamma_3\gamma_1 & 0 \\ 0 & 0 & \gamma_1\gamma_2 \end{pmatrix} \begin{pmatrix} 1 & \lambda_1 & \lambda_1^2 \\ 1 & \lambda_2 & \lambda_2^2 \\ 1 & \lambda_3 & \lambda_3^2 \end{pmatrix} \begin{pmatrix} \tilde{b}_1 \\ \tilde{b}_2 \\ \tilde{b}_3 \end{pmatrix},$$

$$\gamma_1\gamma_2\gamma_3 \longrightarrow \tilde{c}_0,$$

$$\begin{pmatrix} \gamma_1^2 \\ \gamma_2^2 \\ \gamma_3^2 \end{pmatrix} \longrightarrow \begin{pmatrix} 1 & \lambda_1 & \lambda_1^2 \\ 1 & \lambda_2 & \lambda_2^2 \\ 1 & \lambda_3 & \lambda_3^2 \end{pmatrix} \begin{pmatrix} \tilde{c}_1 \\ \tilde{c}_2 \\ \tilde{c}_3 \end{pmatrix}, \quad \begin{pmatrix} \gamma_2 \gamma_3 \\ \gamma_1 \gamma_3 \\ \gamma_1 \gamma_2 \end{pmatrix} \longrightarrow \frac{1}{\tilde{c}_0} \begin{pmatrix} \gamma_1(\tilde{c}_1 + \lambda_2 \tilde{c}_2 + \lambda_2^2 \tilde{c}_3)(\tilde{c}_1 + \lambda_3 \tilde{c}_2 + \lambda_3^2 \tilde{c}_3) \\ \gamma_2(\tilde{c}_1 + \lambda_1 \tilde{c}_2 + \lambda_1^2 \tilde{c}_3)(\tilde{c}_1 + \lambda_3 \tilde{c}_2 + \lambda_3^2 \tilde{c}_3) \\ \gamma_3(\tilde{c}_1 + \lambda_1 \tilde{c}_2 + \lambda_1^2 \tilde{c}_3)(\tilde{c}_1 + \lambda_2 \tilde{c}_2 + \lambda_2^2 \tilde{c}_3) \end{pmatrix},$$

and finally

$$\lambda_1 \longrightarrow \ell_1 - \lambda_3 - \lambda_2, \lambda_2^2 \longrightarrow \ell_1 \lambda_2 + \ell_1 \lambda_3 - \lambda_3 \lambda_2 - \lambda_3^2 + \frac{1}{2} \ell_2 - \frac{1}{2} \ell_1^2, \lambda_3^3 \longrightarrow \frac{1}{3} \ell_3 + \frac{1}{6} \ell_1^3 - \frac{1}{2} \ell_2 \ell_1 + \frac{1}{2} (\ell_2 - \ell_1^2) \gamma_3 + \ell_1 \gamma_3^2.$$

One observes that with this generating set the rewriting of a polynomial invariant only introduces powers of  $\tilde{c}_0$  and  $\tilde{\ell}_0$  as denominator. This result was first obtained by applying the construction in [HK07a, Theorem 2.16]. This latter theorem shows that the coefficients of the reduced Gröbner basis of the generic orbit ideal form a generating set and how to rewrite any other rational invariants in terms of these. The rewrite rules given above can be recognized as a Gröbner basis of the generic orbit ideal. We do not present the reduced Gröbner basis as it is rather cumbersome.

## 5. HARMONIC BASES WITH $B_3$ -SYMMETRIES AND INVARIANTS FOR HIGHER DEGREE

In this section, we extend the description of generating rational invariants for ternary quartics given in Section 4 to ternary forms of arbitrary even degree. According to Section 3, the  $O_3$ -invariants on  $V_{2d}$  are uniquely determined by the  $B_3$ -invariants on  $\Lambda_{2d}$ .

A crucial step in order to describe invariants of the  $B_3$ -action on  $\Lambda_{2d}$  consists in constructing a basis of the vector space  $\mathcal{H}_{2d}$  exhibiting certain symmetries with respect to the group  $B_3$  of signed permutations. We make precise what we mean by this in Section 5.1, and subsequently we give an explicit construction of such a  $B_3$ -equivariant basis.

Because of the wide-range of applications that involve harmonic functions, this  $B_3$ -equivariant basis is of independent interest. We therefore devote Section 5.3 to an illustration of the constructed harmonic basis and briefly discuss connections to other bases for  $\mathcal{H}_{2d}$ . We mention at this point that our construction of  $B_3$ -equivariant bases for harmonic polynomials would work analogously for the case of odd degree  $\mathcal{H}_{2d+1}$ , but with a view toward rational  $B_3$ -invariants on  $\Lambda_{2d}$ , we restrict our treatment to the case of even degree  $\mathcal{H}_{2d}$ .

Finally, in Section 5.5, we deduce from the  $B_3$ -equivariant basis for  $\mathcal{H}_{2d}$  a generating set of rational  $B_3$ -invariants on  $\Lambda_{2d}$ , as a natural extension of Theorem 4.3.

**5.1.  $B_3$ -equivariant bases.** We construct a basis  $\mathcal{H}_{2d}$  ( $d \geq 2$ ) that essentially splits into subsets of three polynomials that are obtained from one another by permutation of the variables. Each of these subsets spans a  $B_3$ -invariant subspace and the action of  $B_3$  on this subspace is given by signed permutations on the subset. We introduce the following notation to be more precise.

**Definition 5.1.** *Let  $k \in \mathbb{N}$  and consider two maps  $\zeta, \xi: \{0, \dots, k-1\} \rightarrow \{0, 1\}$ . An indexed set*

$$\{u_{i,j} \mid 1 \leq i \leq 3, 0 \leq j < k\} \subset V_{2d}$$

*of  $3k$  elements of  $V_{2d}$  is called a  $B_3$ -equivariant subset of  $V_{2d}$  with respect to  $\zeta, \xi$  (or, a  $(\zeta, \xi)$ -equivariant subset) if the action of  $B_3$  on this set is given as follows: For  $\sigma \in \mathfrak{S}_3$  and  $\tau \in \{1, -1\}^3$ , the permutation and sign-change matrices  $g_\sigma$  and  $g_\tau$  act by*

$$\forall i, j: \quad g_\sigma u_{i,j} = \text{sgn}(\sigma)^{\zeta(j)} u_{\sigma(i),j} \quad \text{and} \quad g_\tau u_{i,j} = \left( \frac{\tau_1 \tau_2 \tau_3}{\tau_i} \right)^{\xi(j)} u_{i,j}.$$

Our aim is to find a  $B_3$ -equivariant basis for the vector space  $\mathcal{H}_{2d}$  for arbitrary  $d \geq 2$ , similar to Proposition 4.1 in the case of  $\mathcal{H}_4$ . For  $d \not\equiv 2 \pmod{3}$ , however, we need to slightly relax this aim, and instead of a basis, we shall determine a  $B_3$ -equivariant spanning set of the vector space  $\mathcal{H}_{2d}$  together with the linear relationships satisfied.

**Theorem 5.2.** *Let  $d \geq 2$  and let  $k := \lceil \frac{4d+1}{3} \rceil$ . Then there exist  $3k$  harmonic polynomials  $u_{i,j} \in \mathcal{H}_{2d}$  for  $i \in \{1, 2, 3\}, j \in \{0, \dots, k-1\}$  forming a  $(\zeta, \xi)$ -equivariant set for some maps  $\zeta, \xi: \{0, \dots, k-1\} \rightarrow \{0, 1\}$ , such that:*

- (i) *The harmonic polynomials  $u_{i,j} \in \mathcal{H}_{2d}$  for  $i \in \{1, 2, 3\}, j \in \{0, \dots, k-1\}$  span the vector space  $\mathcal{H}_{2d}$ .*
- (ii) (a) *If  $d \equiv 2 \pmod{3}$ , then the  $u_{i,j}$  form a basis of  $\mathcal{H}_{2d}$ .*  
 (b) *If  $d \equiv 1 \pmod{3}$ , then  $\dim \mathcal{H}_{2d} = 3k - 1$  and the  $u_{i,j}$  satisfy the linear relation*

$$u_{1,0} + u_{2,0} + u_{3,0} = 0.$$

- (c) *If  $d \equiv 0 \pmod{3}$ , then  $\dim \mathcal{H}_{2d} = 3k - 2$  and the  $u_{i,j}$  satisfy the relations*

$$u_{1,0} = u_{2,0} = u_{3,0}.$$

We observe that Theorem 5.2 generalizes Proposition 4.1: For  $2d = 4$ , we have  $k = 3$  and the harmonic polynomials  $u_{i,0} := r_i, u_{i,1} := s_i$  and  $u_{i,2} := t_i$  ( $i \in \{1, 2, 3\}$ ) form a basis of  $\mathcal{H}_{2d}$  satisfying the assertions of Theorem 5.2 for

$$\zeta(j) = \begin{cases} 1 & \text{if } j = 1, \\ 0 & \text{otherwise} \end{cases} \quad \text{and} \quad \xi(j) = \begin{cases} 1 & \text{if } j \in \{1, 2\}, \\ 0 & \text{otherwise.} \end{cases}$$

We formulated Theorem 5.2 as an existence result, but the crucial part for applications is the constructive proof given below. We provide explicit closed formulas for the elements  $u_{i,j}$ , in addition to algorithmic constructions that appear as proofs.

**5.2. Construction of  $B_3$ -equivariant harmonic bases.** We prove Theorem 5.2 by constructing a spanning set with the asserted properties.

For nonnegative integers  $i, j, k$  and  $r = i + j + k$ , we denote the *multinomial coefficient*

$$\binom{r}{i, j, k} := \frac{r!}{i!j!k!}.$$

The following result is a simple characterization of harmonic polynomials.

**Lemma 5.3.** *Let  $d \geq 2$ . A ternary form*

$$v = \sum_{i+j+k=2d} \binom{2d}{i, j, k} \beta_{i,j,k} x^i y^j z^k \in V_{2d}$$

*lies in  $\mathcal{H}_{2d}$  if and only if  $\beta_{i+2,j,k} + \beta_{i,j+2,k} + \beta_{i,j,k+2} = 0$  holds for all  $i, j, k \in \mathbb{N}$  such that  $i + j + k = 2d - 2$ .*

*Proof.* Recall that  $\mathcal{H}_{2d}$  is the orthogonal complement of  $qV_{2d-2}$  in  $V_{2d}$  with respect to the apolar product, where  $q = x^2 + y^2 + z^2$ . Hence,  $v \in \mathcal{H}_{2d}$  if and only if  $\langle v, qx^i y^j z^k \rangle = 0$  for all  $i, j, k$  such that  $i + j + k = 2d - 2$ . From

$$\langle x^{i'} y^{j'} z^{k'}, x^i y^j z^k \rangle = \begin{cases} i!j!k! & \text{if } i = i', j = j', k = k' \\ 0 & \text{otherwise.} \end{cases}$$

we see that

$$\langle v, qx^i y^j z^k \rangle = (2d)! \cdot (\beta_{i+2,j,k} + \beta_{i,j+2,k} + \beta_{i,j,k+2}),$$

so the claim follows.  $\square$

**Lemma 5.4.** *Let  $d \geq 2$  and  $0 \leq \ell \leq d$ . There exists a harmonic polynomial  $\theta_\ell \in \mathcal{H}_{2d}$ , unique up to scaling, with the following properties:*

- (1) *Each monomial of  $\theta_\ell$  is of even degree in each of the variables  $x, y$  and  $z$ .*
- (2) *The highest degree in which the variable  $x$  occurs in  $\theta_\ell$  is  $2\ell$ .*
- (3) *The monomial  $y^{2j} z^{2k}$  (for  $j + k = d$ ) does not occur in the expression  $\theta_\ell$  if  $|j - k| < \ell$ .*
- (4) *Permuting the variables  $y$  and  $z$  in the expression  $\theta_\ell$  gives  $(-1)^{d-\ell} \theta_\ell$ .*

*Then all monomials in  $V_{2d}$  of degree  $2\ell$  in  $x$  and even degree in each  $y$  and  $z$  occur in  $\theta_\ell$ .*

*Proof.* By Lemma 5.3, determining the expression

$$\theta_\ell = \sum_{i+j+k=d} \binom{2d}{2i, 2j, 2k} \beta_{i,j,k} x^{2i} y^{2j} z^{2k}$$

amounts to finding  $\beta_{i,j,k} \in \mathbb{R}$  for  $i, j, k \geq 0$ ,  $i + j + k = d$  such that

- (i)  $\beta_{i+1,j,k} + \beta_{i,j+1,k} + \beta_{i,j,k+1} = 0$  for all  $i, j, k \geq 0$ ,  $i + j + k = d - 1$ ,
- (ii)  $\beta_{i,j,k} = 0$  if  $i > \ell$ ,
- (iii)  $\beta_{\ell,j,k} \neq 0$  for some  $j, k$ ,
- (iv)  $\beta_{i,j,k} = (-1)^{d-\ell} \beta_{i,k,j}$  for all  $i + j + k = d$ ,
- (v)  $\beta_{0,j,k} = 0$  if  $|j - k| < \ell$ .

From this, the coefficients  $\beta_{i,j,k}$  can be determined iteratively. Combining (v) with (i), one shows by induction on  $i$  that

$$(5.1) \quad \beta_{i,j,k} = 0 \quad \text{if } i + |j - k| < \ell.$$

Moreover, (ii) imposes  $\beta_{i,j,k} = 0$  if  $i > \ell$ . Then (i) implies that  $\beta_{\ell,j+1,k} = -\beta_{\ell,j,k+1}$  for all  $j, k \geq 0$  with  $\ell + j + k = d - 1$ . Hence,  $\beta_{\ell,j,k} = (-1)^j \beta_{\ell,0,d-\ell}$ . Because of (iii), we may assume (after rescaling all  $\beta_{i,j,k}$ ) that  $\beta_{\ell,j,k} = (-1)^j$  for all  $j + k = d - \ell$ .

If  $d - \ell$  is even and  $s := \frac{d-\ell}{2}$ , we additionally have  $\beta_{\ell-1,s,s+1} + \beta_{\ell-1,s+1,s} + \beta_{\ell,s,s} = 0$  by (i). From  $\beta_{\ell,s,s} = (-1)^s$ , we conclude with (iv) that

$$(5.2) \quad \beta_{\ell-1,s,s+1} = \beta_{\ell-1,s+1,s} = \frac{(-1)^{s+1}}{2} \quad (\neq 0).$$

At this stage we have determined all  $\beta_{i,j,k}$  for  $i + j + k = d$  such that  $i \geq \ell$  or  $|j - k| < \max\{\ell - i, 2\}$  and they are compatible with (i)–(v) in the sense that all statements from (i)–(v) only involving those  $\beta_{i,j,k}$  are satisfied. We now proceed to determine  $\beta_{\ell-K,j,k}$  for  $K \in \mathbb{N}$  increasing from 1 to  $\ell$ , assuming that the values  $\beta_{i,j,k}$  for  $i > \ell - K$  are already known and compatible with (i)–(v).

Recall that from (5.1) and (5.2) we already know the values  $\beta_{\ell-K,j,k}$  whenever  $|j - k| < \max\{K, 2\}$ . For increasing values of  $|j - k|$ , we then iteratively obtain the values  $\beta_{\ell-K,j,k}$  for  $j > k$  from the recursion

$$\beta_{\ell-K,j+1,k} = -\beta_{\ell-(K-1),j,k} - \beta_{\ell-K,j,k+1}$$

and for  $k > j$  from the recursion

$$\beta_{\ell-K,j,k+1} = -\beta_{\ell-(K-1),j,k} - \beta_{\ell-K,j+1,k}$$

(both resulting from (i)). This determines all  $\beta_{\ell-K,j,k}$ . Note that  $\beta_{\ell-K,j,k} = (-1)^{d-\ell} \beta_{\ell-K,k,j}$  follows iteratively from the assumption that property (iv) holds for all previously constructed  $\beta_{i,j,k}$ . Hence, we determined the unique values for  $\beta_{\ell-K,j,k}$  such that the properties (i)–(v) are still preserved. This concludes the construction.  $\square$

**Remark 5.5.** We just saw an iterative construction of the coefficients  $\beta_{i,j,k}$  defining  $\theta_\ell$  in the proof above. We now give an explicit formula: If  $d - \ell$  is odd and  $s := \frac{d-\ell+1}{2}$ , then the values

$$\beta_{i,j,k} = (-1)^j \binom{j-s}{\ell-i}$$

are easily checked to satisfy (i)–(v). Here, we employ for  $a, b \in \mathbb{Z}$  the convention

$$\binom{a}{b} := \begin{cases} \frac{a(a-1)\dots(a-b+1)}{b!} & \text{if } b \geq 0, \\ 0 & \text{otherwise} \end{cases}$$

and use the identities  $\binom{a+1}{b+1} = \binom{a}{b} + \binom{a}{b+1}$  and  $\binom{a}{b} = (-1)^b \binom{b-a-1}{b}$ . On the other hand, if  $d - \ell$  is even and  $s := \frac{d-\ell}{2}$ , then

$$\beta_{i,j,k} = (-1)^j \binom{j-s}{\ell-i} + (-1)^j \binom{j-s-1}{\ell-i}$$

satisfy the desired properties. This gives an explicit closed formula for the harmonic polynomials  $\theta_\ell$ .

Similar to Lemma 5.4, we also obtain the following result:

**Lemma 5.6.** *Let  $d \geq 2$  and  $0 \leq \ell \leq d-1$ . Then there exists a harmonic polynomial  $\eta_\ell \in \mathcal{H}_{2d}$ , unique up to scaling, with the following properties:*

- (1) Each monomial of  $\eta_\ell$  is of even degree in  $x$  and of odd degree in each of the variables  $y$  and  $z$ .
- (2) The highest degree in which the variable  $x$  occurs in  $\eta_\ell$  is  $2\ell$ .
- (3) The monomial  $y^{2j+1}z^{2k+1}$  (for  $j+k=d-1$ ) does not occur in the expression  $\eta_\ell$  if  $|j-k| < \ell$ .
- (4) Permuting the variables  $y$  and  $z$  in the expression  $\eta_\ell$  gives  $(-1)^{d-\ell+1}\eta_\ell$ .

Then all monomials in  $V_{2d}$  of degree  $2\ell$  in  $x$  and odd degree in each  $y$  and  $z$  occur in  $\eta_\ell$ .

*Proof.* Writing

$$\eta_\ell = \sum_{i+j+k=d-1} \binom{2d}{2i, 2j+1, 2k+1} \beta_{i,j,k} x^{2i} y^{2j+1} z^{2k+1},$$

the coefficients  $\beta_{i,j,k}$  for  $i+j+k=d-1$  have to meet the same conditions as before (replacing  $d$  by  $d-1$ ), so the construction is the same as before (and the corresponding explicit formulas from Remark 5.5 apply).  $\square$

While the harmonic polynomials  $\theta_\ell, \eta_\ell$  are (anti-)symmetric with respect to the variables  $y$  and  $z$ , the variable  $x$  plays a special role. We therefore now consider the expressions that arise from  $\theta_\ell, \eta_\ell$  by cyclically permuting the variables  $x, y$  and  $z$ . For this, we consider the cycle  $c = (123) \in \mathfrak{S}_3$  and its associated permutation matrix

$$g_c = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix} \in B_3.$$

**Lemma 5.7.** *Let  $d \geq 2$  and consider  $\eta_\ell, \theta_\ell \in \mathcal{H}_{2d}$  as in Lemmas 5.6 and 5.4. Then the harmonic polynomials*

$$\begin{aligned} &\eta_0, g_c \eta_0, g_c^2 \eta_0, \quad \eta_1, g_c \eta_1, g_c^2 \eta_1, \quad \dots, \quad \eta_{d-1}, g_c \eta_{d-1}, g_c^2 \eta_{d-1}, \\ &\theta_0, g_c \theta_0, g_c^2 \theta_0, \quad \theta_1, g_c \theta_1, g_c^2 \theta_1, \quad \dots, \quad \theta_r, g_c \theta_r, g_c^2 \theta_r \end{aligned}$$

are linearly independent in  $\mathcal{H}_{2d}$  for  $r := \lfloor \frac{d-2}{3} \rfloor$ .

*Proof.* The monomials in  $\theta_\ell, g_c \theta_\ell, g_c^2 \theta_\ell$  are of the form  $x^{2i} y^{2j} z^{2k}$ , in  $\eta_\ell$  of the form  $x^{2i} y^{2j+1} z^{2k+1}$ , in  $g_c \eta_\ell$  of the form  $x^{2i+1} y^{2j} z^{2k+1}$ , and in  $g_c^2 \eta_\ell$  of the form  $x^{2i+1} y^{2j+1} z^{2k}$ . Therefore, we can show the linear independence of those families of harmonic polynomials separately.

The linear independence of  $\eta_0, \eta_1, \dots, \eta_{d-1} \in \mathcal{H}_{2d}$  is a consequence of property (2) in Lemma 5.6. The same for  $g_c \eta_0, \dots, g_c \eta_{d-1}$  and for  $g_c^2 \eta_0, \dots, g_c^2 \eta_{d-1}$  follows directly.

To see that  $\{g_c^{i-1} \theta_\ell \mid 1 \leq i \leq 3, 0 \leq \ell \leq r\}$  is a linearly independent set, suppose that  $\sum_{i=1}^3 \sum_{\ell=0}^r \alpha_{i,\ell} g_c^{i-1} \theta_\ell = 0$  for some  $\alpha_{i,\ell} \in \mathbb{R}$ , not all equal to zero. Let  $s \in \{0, \dots, r\}$  be maximal such that  $\alpha_{i,s} \neq 0$  for some  $i \in \{1, 2, 3\}$ . We may assume that  $\alpha_{1,s} \neq 0$ . Note that  $s \leq r = \lfloor \frac{d-2}{3} \rfloor$  implies that there exist integers  $j, k > s$  such that  $j+k+s=d$ . It follows from property (2) in Lemma 5.4 that the coefficient of  $x^{2s} y^{2j} z^{2k}$  in the expression  $\sum_{i=1}^3 \sum_{\ell=0}^s \alpha_{i,\ell} g_c^{i-1} \theta_\ell$  is  $\alpha_{1,\ell} \lambda_1$ , where  $\lambda_1 \neq 0$  is the coefficient of  $x^{2s} y^{2j} z^{2k}$  in  $\theta_s$ . This implies  $\sum_{i=1}^3 \sum_{\ell=0}^s \alpha_{i,\ell} g_c^{i-1} \theta_\ell \neq 0$ , contrary to the assumption.  $\square$

**Lemma 5.8.** *Let  $d = 3k_0 + 1$  for some  $k_0 \geq 1$  and let  $g_c \in B_3$  and  $\theta_\ell$  as before. Then*

$$\theta_{k_0} + g_c \theta_{k_0} + g_c^2 \theta_{k_0} = 0.$$

*Proof.* From Lemma 5.7 we know that the set  $\{g_c^{i-1} \theta_\ell \mid 1 \leq i \leq 3, 0 \leq \ell \leq k_0-1\}$  forms a basis for a  $3k_0$ -dimensional subspace  $W \subset \mathcal{H}_{2d}$ . By properties (2) and (1) in Lemma 5.4, all monomials occurring in any  $g_c^{i-1} \theta_\ell$  for  $1 \leq i \leq 3, 0 \leq \ell \leq k_0-1$  must be of the form  $x^{2i} y^{2j} z^{2k}$  with  $\min\{i, j, k\} \leq k_0-1$ , so

$W$  is contained in the subspace  $U$  of  $V_{2d}$  spanned by the monomials of this form. Lemma 5.3 implies that  $\dim U \cap \mathcal{H}_{2d} = d - 1 = 3k_0$ , so in fact  $W = U \cap \mathcal{H}_{2d}$ .

By properties (2) and (4) in Lemma 5.4 we have  $\theta_{k_0} = \lambda x^{2k_0} y^{2k_0+2} z^{2k_0} - \lambda x^{2k_0} y^{2k_0+2} z^{2k_0} + u$  for some  $u \in U$  and  $\lambda \in \mathbb{R}$ . Therefore,

$$\theta_{k_0} + g_c \theta_{k_0} + g_c^2 \theta_{k_0} = u + g_c u + g_c^2 u =: u' \in U \cap \mathcal{H}_{2d}$$

and we can write  $u' = \sum_{i=1}^3 \sum_{\ell=0}^{k_0-1} \alpha_{i,\ell} g_c^{i-1} \theta_\ell$ . Suppose for contradiction that not all  $\alpha_{i,\ell}$  are zero and let  $s \in \{0, \dots, k_0 - 1\}$  be minimal such that  $\alpha_{i,s} \neq 0$  for some  $i \in \{1, 2, 3\}$ . By symmetry, we may assume  $\alpha_{1,s} \neq 0$ .

If  $d + s$  is even, then the monomial  $y^{d+s} z^{d-s}$  occurs in  $\theta_s$  with a non-zero coefficient  $\mu \in \mathbb{R}$ , but not in  $\theta_\ell$  for  $\ell > s$  by property (3) in Lemma 5.4. Observe also that the  $y$ -degree and the  $z$ -degree of this monomial are both larger than  $2k_0$ , so  $y^{d+s} z^{d-s}$  is not contained as a monomial in any of the expressions  $g_c \theta_\ell$  or  $g_c^2 \theta_\ell$  for  $\ell \leq k_0$ . Therefore, the monomial  $y^{d+s} z^{d-s}$  occurs in  $u'$  with the non-zero coefficient  $\alpha_{1,s} \mu$ , but not in  $\theta_{k_0}$  or  $g_c \theta_{k_0}$  or  $g_c^2 \theta_{k_0}$ , a contradiction.

If  $d + s$  is odd, then  $s \leq k_0 - 2$  and we consider the monomial  $y^{d+s+1} z^{d-s-1}$ . As before, this monomial does not occur in  $\theta_\ell$  for  $\ell > s + 1$  or in any of the expressions  $g_c \theta_\ell$  or  $g_c^2 \theta_\ell$  for  $0 \leq \ell \leq k_0$ . Hence, its coefficient in  $u'$  is  $\alpha_{1,s} \mu + \alpha_{1,s+1} \mu' = 0$ , where  $\mu, \mu' \neq 0$  are the coefficients of  $y^{d+s+1} z^{d-s-1}$  in  $\theta_s$  and  $\theta_{s+1}$ , respectively. In the same way we can consider the monomial  $y^{d-s-1} z^{d+s+1}$ , which then gives  $-\alpha_{1,s} \mu + \alpha_{1,s+1} \mu' = 0$  by the (anti-)symmetry of  $\theta_s$  and  $\theta_{s+1}$  due to property (4) in Lemma 5.4. From both equations we deduce  $\alpha_{1,s} = 0$ , a contradiction.  $\square$

*Proof of Theorem 5.2.* We use the notations of the previous Lemmas and denote

$$k_0 := \left\lceil \frac{d-2}{3} \right\rceil \quad \text{and} \quad k := \left\lceil \frac{4d+1}{3} \right\rceil = k_0 + d + 1.$$

For  $i \in \{1, 2, 3\}$  and  $j \in \{1, \dots, k-1\}$ , we define

$$u_{i,j} := \begin{cases} g_c^{i-1} \theta_{k_0-j}, & \text{if } 1 \leq j \leq k_0, \\ g_c^{i-1} \eta_{j-k_0-1}, & \text{if } k_0 + 1 \leq j < k. \end{cases}$$

Note that we have only defined  $u_{i,j}$  for  $j \geq 1$ . Additionally, let

$$u_{i,0} := \begin{cases} g_c^{i-1} \theta_{k_0}, & \text{if } d \equiv 1 \text{ or } 2 \pmod{3}, \\ \theta_{k_0} + g_c \theta_{k_0} + g_c^2 \theta_{k_0}, & \text{if } d \equiv 0 \pmod{3}. \end{cases}$$

In the following, we show that these harmonic polynomials  $u_{i,j} \in \mathcal{H}_{2d}$  for  $i \in \{1, 2, 3\}$ ,  $j \in \{0, \dots, k\}$  satisfy the properties stated in Theorem 5.2 if we define  $\zeta, \xi: \{0, \dots, k-1\} \rightarrow \{0, 1\}$  as

$$\zeta(j) := \begin{cases} 0 & \text{if } d + k_0 + j \text{ even,} \\ 1 & \text{otherwise,} \end{cases} \quad \text{and} \quad \xi(j) = \begin{cases} 0 & \text{if } 0 \leq j \leq k_0, \\ 1 & \text{if } k_0 + 1 \leq j < k. \end{cases}$$

First, we check the  $(\zeta, \xi)$ -equivariance of the harmonic polynomials  $u_{i,j}$ . The symmetric group  $\mathfrak{S}_3$  is generated by the cycle  $c = (123)$  together with the transposition  $t = (23)$ , so in order to show  $g_\sigma u_{i,j} = \text{sgn}(\sigma)^{\zeta(j)} u_{\sigma(i),j}$  for all  $\sigma \in \mathfrak{S}_3$ , it suffices to check this for  $\sigma = c$  and  $\sigma = t$ . By definition of  $u_{i,j}$  we have

$$g_c u_{1,j} = u_{2,j}, \quad g_c u_{2,j} = u_{3,j}, \quad g_c u_{3,j} = u_{1,j},$$

i.e.  $g_c u_{i,j} = u_{c(i),j}$ . Note that  $\text{sgn}(c) = 1$ . Since  $g_t$  acts by interchanging the variables  $y$  and  $z$ , it follows from property (4) in Lemmas 5.4 and 5.6 that  $g_t u_{1,j} = (-1)^{\zeta(j)} u_{1,j}$ . Indeed, for  $j \geq 1$  or  $d \not\equiv 0 \pmod{3}$ , this is immediate from the definition of  $u_{1,j}$ , while for  $j = 0$  and  $d \equiv 0 \pmod{3}$ , it follows after using the identities  $g_t g_c = g_c^2 g_t$  and  $g_t g_c^2 = g_c g_t$  and observing that  $\zeta(0) = 0$  in this case. After that, we deduce

$$g_t u_{2,j} = g_t (g_c u_{1,j}) = g_c^2 (g_t u_{1,j}) = (-1)^{\zeta(j)} g_c^2 (u_{1,j}) = (-1)^{\zeta(j)} u_{3,j},$$

(where we used again  $g_t g_c = g_c^2 g_t$ ) and therefore also  $g_t u_{3,j} = (-1)^{\zeta(j)} u_{2,j}$ . All together, we have  $g_t u_{i,j} = (-1)^{\zeta(j)} u_{t(i),j}$ . Note that  $\text{sgn} t = -1$ . With this, we have verified the equivariance for permutation matrices.

Now let  $\tau \in \{1, -1\}^3$ . Note that property (1) of Lemmata 5.4 and 5.6 ensures that  $g_\tau u_{1,j} = (\tau_2 \tau_3)^{\xi(j)} u_{1,j}$ . Using

$$g_\tau g_c = \text{diag}(\tau_2, \tau_3, \tau_1) \quad \text{and} \quad g_\tau g_c^2 = \text{diag}(\tau_3, \tau_1, \tau_2),$$

we also see that  $g_\tau u_{2,j} = (\tau_3 \tau_1)^{\xi(j)} u_{1,j}$  and  $g_\tau u_{3,j} = (\tau_1 \tau_2)^{\xi(j)} u_{1,j}$ . This concludes the proof of the  $(\zeta, \xi)$ -equivariance.

If  $d \equiv 2 \pmod{3}$ , then  $k = \frac{4d+1}{3}$ , so  $\dim \mathcal{H}_{2d} = 3k$ . The  $3k$  harmonic polynomials  $u_{i,j}$  are linearly independent by Lemma 5.7, hence they form a basis of  $\mathcal{H}_{2d}$ . This verifies properties (i) and (ii) for  $d \equiv 2 \pmod{3}$ .

If  $d \equiv 0 \pmod{3}$ , then  $\dim \mathcal{H}_{2d} = 4d + 1 = 3k - 2$  and the identities  $u_{1,0} = u_{2,0} = u_{3,0}$  hold by definition of  $u_{i,0}$ . This shows property (ii). The  $3k - 3$  harmonic polynomials  $u_{i,j}$  for  $i \in \{1, 2, 3\}$ ,  $j \in \{1, \dots, k-1\}$  are linearly independent by Lemma 5.7, and the subspace of  $\mathcal{H}_{2d}$  spanned by them does not contain the harmonic polynomial  $u_{1,0} = u_{2,0} = u_{3,0}$ , because the latter contains the monomial  $x^{\frac{2d}{3}} y^{\frac{2d}{3}} z^{\frac{2d}{3}}$ , which does not occur in  $u_{i,j}$  for  $j \geq 1$ . This also establishes (i) for  $d \equiv 0 \pmod{3}$ .

We are left with the case  $d \equiv 1 \pmod{3}$ . Note that then  $\dim \mathcal{H}_{2d} = 4d + 1 = 3k - 1$ . By Lemma 5.7, the  $3(k-1)$  polynomials  $u_{i,j}$  for  $i \in \{1, 2, 3\}$ ,  $j \in \{1, \dots, k-1\}$  span a  $3(k-1)$ -dimensional subspace of  $\mathcal{H}_{2d}$ . From considering the (non-)occurrence of the monomials

$$x^{\frac{2d+2}{3}} y^{\frac{2d-1}{3}} z^{\frac{2d-1}{3}}, \quad x^{\frac{2d-1}{3}} y^{\frac{2d+2}{3}} z^{\frac{2d-1}{3}} \quad \text{and} \quad x^{\frac{2d-1}{3}} y^{\frac{2d-1}{3}} z^{\frac{2d+2}{3}}$$

in the expressions  $u_{i,j}$ , it follows that the subspace of  $\mathcal{H}_{2d}$  spanned by all the  $3k$  harmonic polynomials  $u_{i,j}$  must be at least of dimension  $3k - 1$  and must hence coincide with  $\mathcal{H}_{2d}$ . Together with Lemma 5.8, this verifies (i) and (ii) in this last case, concluding the proof.  $\square$

**5.3. Illustrations and examples.** For degree 4, the construction above reproduces the basis for  $\mathcal{H}_4$  given in Proposition 4.1 (up to scalars). For arbitrary  $d$ , the proof of Theorem 5.2 together with Remark 5.5 give explicit closed formulas for elements  $u_{i,j}$  of a  $B_3$ -equivariant spanning set of  $\mathcal{H}_{2d}$ , by means of several case distinctions.

For example, tracing back the definitions gives: If  $1 \leq j \leq k_0 = \lceil \frac{d-2}{3} \rceil$  and  $d + j - k_0$  is odd, then

$$u_{1,j} = \sum_{i_1+i_2+i_3=d} (-1)^{i_2} \binom{i_2 - s}{k_0 - j - i_1} x^{2i_1} y^{2i_2} z^{2i_3},$$

where  $s := \frac{1}{2}(d - k_0 + j + 1)$ ; and  $u_{2,j}$ ,  $u_{3,j}$  are obtained from this by cyclically permuting  $x, y, z$ . For the other elements  $u_{i,j}$ , similar formulas can be written out.

We give examples for  $B_3$ -equivariant spanning sets of  $\mathcal{H}_{2d}$  for  $d = 3$  and 4. For this, we will denote the constructed harmonic polynomials  $u_{i,j} \in \mathcal{H}_{2d}$  from Theorem 5.2 by  $u_{i,j}^{(2d)}$  to avoid confusion between different values of  $d$ .

In degree 6, the following 13 harmonic polynomials form the  $B_3$ -equivariant basis for  $\mathcal{H}_6$  constructed in Theorem 5.2:

$$\begin{aligned} u_{1,0}^{(6)} = u_{2,0}^{(6)} = u_{3,0}^{(6)} &= -2x^6 - 2z^6 - 2y^6 + 15x^4y^2 + 15y^4z^2 + 15z^4x^2 + 15x^4z^2 + 15y^4x^2 + 15z^4y^2 - 180x^2y^2z^2, \\ u_{1,1}^{(6)} &= -y^6 + 15y^4z^2 - 15y^2z^4 + z^6, & u_{1,2}^{(6)} &= 3y^5z - 10y^3z^3 + 3yz^5, \\ u_{2,1}^{(6)} &= -z^6 + 15z^4x^2 - 15z^2x^4 + x^6, & u_{2,2}^{(6)} &= 3z^5x - 10z^3x^3 + 3zx^5, \\ u_{3,1}^{(6)} &= -x^6 + 15x^4y^2 - 15x^2y^4 + y^6, & u_{3,2}^{(6)} &= 3x^5y - 10x^3y^3 + 3xy^5, \\ u_{1,3}^{(6)} &= -10x^2y^3z + 10x^2yz^3 + y^5z - yz^5, & u_{1,4}^{(6)} &= 10x^4yz - 10x^2y^3z - 10x^2yz^3 + y^5z + yz^5, \\ u_{2,3}^{(6)} &= -10y^2z^3x + 10y^2zx^3 + z^5x - zx^5, & u_{2,4}^{(6)} &= 10y^4zx - 10y^2z^3x - 10y^2zx^3 + z^5x + zx^5, \\ u_{3,3}^{(6)} &= -10z^2x^3y + 10z^2xy^3 + x^5y - xy^5, & u_{3,4}^{(6)} &= 10z^4xy - 10z^2x^3y - 10z^2xy^3 + x^5y + xy^5. \end{aligned}$$

The five harmonic polynomials  $u_{1,j}^{(6)}$  for  $j \in \{0, \dots, 4\}$  are illustrated in Figure 5.3, presented in the same way as previously for quartics in Figure 4.1. The remaining basis elements  $u_{2,j}^{(6)}$  and  $u_{3,j}^{(6)}$  not depicted arise from these by permuting the coordinates.






FIGURE 5.1. Illustration of the elements  $u_{1,j}^{(6)}$  in the  $B_3$ -symmetric basis of  $\mathcal{H}_6$




FIGURE 5.2. Illustration of the elements  $u_{1,j}^{(8)}$  in the  $B_3$ -symmetric spanning set of  $\mathcal{H}_8$

For degree 8, the following polynomials form the  $B_3$ -equivariant spanning set for  $\mathcal{H}_8$ :

$$\begin{aligned} u_{1,0}^{(8)} &= -14x^2y^6 + 210z^2x^2y^4 - 210y^2z^4x^2 + 14z^6x^2 + y^8 - 14y^6z^2 + 14y^2z^6 - z^8, \\ u_{2,0}^{(8)} &= -14y^2z^6 + 210y^2z^4x^2 - 210z^2x^4y^2 + 14x^6y^2 + z^8 - 14z^6x^2 + 14z^2x^6 - x^8, \\ u_{3,0}^{(8)} &= -14z^2x^6 + 210z^2x^4y^2 - 210z^2x^2y^4 + 14y^6z^2 + x^8 - 14x^6y^2 + 14x^2y^6 - y^8, \end{aligned}$$

$$\begin{aligned} u_{1,1}^{(8)} &= y^8 - 28y^6z^2 + 70y^4z^4 - 28y^2z^6 + z^8, & u_{1,2}^{(8)} &= -y^7z + 7y^5z^3 - 7y^3z^5 + yz^7, \\ u_{2,1}^{(8)} &= z^8 - 28z^6x^2 + 70z^4x^4 - 28z^2x^6 + x^8, & u_{2,2}^{(8)} &= -z^7x + 7z^5x^3 - 7z^3x^5 + zx^7, \\ u_{3,1}^{(8)} &= x^8 - 28x^6y^2 + 70x^4y^4 - 28x^2y^6 + y^8, & u_{3,2}^{(8)} &= -x^7y + 7x^5y^3 - 7x^3y^5 + xy^7, \end{aligned}$$

$$\begin{aligned} u_{1,3}^{(8)} &= 42x^2y^5z - 140x^2y^3z^3 + 42x^2yz^5 - 3y^7z + 7y^5z^3 + 7y^3z^5 - 3yz^7, \\ u_{2,3}^{(8)} &= 42y^2z^5x - 140y^2z^3x^3 + 42y^2zx^5 - 3z^7x + 7z^5x^3 + 7z^3x^5 - 3zx^7, \\ u_{3,3}^{(8)} &= 42z^2x^5y - 140z^2x^3y^3 + 42z^2xy^5 - 3x^7y + 7x^5y^3 + 7x^3y^5 - 3xy^7, \end{aligned}$$

$$\begin{aligned} u_{1,4}^{(8)} &= -35x^4y^3z + 35x^4yz^3 + 21x^2y^5z - 21x^2yz^5 - y^7z + yz^7, \\ u_{2,4}^{(8)} &= -35y^4z^3x + 35y^4zx^3 + 21y^2z^5x - 21y^2zx^5 - z^7x + zx^7, \\ u_{3,4}^{(8)} &= -35z^4x^3y + 35z^4xy^3 + 21z^2x^5y - 21z^2xy^5 - x^7y + xy^7, \end{aligned}$$

$$\begin{aligned} u_{1,5}^{(8)} &= 14x^6yz - 35x^4y^3z - 35x^4yz^3 + 21x^2y^5z + 21x^2yz^5 - y^7z - yz^7, \\ u_{2,5}^{(8)} &= 14y^6zx - 35y^4z^3x - 35y^4zx^3 + 21y^2z^5x + 21y^2zx^5 - z^7x - zx^7, \\ u_{3,5}^{(8)} &= 14z^6xy - 35z^4x^3y - 35z^4xy^3 + 21z^2x^5y + 21z^2xy^5 - x^7y - xy^7. \end{aligned}$$

They satisfy the linear relation

$$u_{1,0}^{(8)} + u_{2,0}^{(8)} + u_{3,0}^{(8)} = 0$$

and are otherwise linearly independent. This spanning set is illustrated in Figure 5.3.

Generating these expression is straightforward, as the definition of  $u_{i,j}^{(2d)}$  in the proof of Theorem 5.2 and the formulas for  $\theta_\ell$  and  $\eta_\ell$  in Remark 5.5 give explicit formulas for the coefficients in  $u_{i,j}^{(2d)}$ . A straightforward implementation in `Maple`<sup>1</sup> produces the  $B_3$ -equivariant basis of  $\mathcal{H}_{100}$  in a few seconds.

<sup>1</sup>Maple 2016. Maplesoft, a division of Waterloo Maple Inc., Waterloo, Ontario.

*Representation-theoretic viewpoint and cubic harmonics.* While *spherical harmonics* form the most commonly used basis of harmonic polynomials, another basis is given by the *cubic harmonics* [Mug72, FK77], which correspond to the decomposition of the space of harmonic polynomials into irreducible  $B_3$ -representation spaces. With the following representation theoretic viewpoint on Theorem 5.2, one can see how our explicitly constructed  $B_3$ -equivariant basis are related to these cubic harmonics. Indeed, the construction of the spanning set  $u_{i,j}^{(2d)}$  of  $\mathcal{H}_{2d}$  can be seen as decomposing  $\mathcal{H}_{2d}$  into  $B_3$ -invariant subspaces endowed with one out of six possible representations of dimension 3, 2 or 1.

For  $\zeta, \xi \in \{0, 1\}$ , there is a three-dimensional representation space  $W_{\zeta, \xi}$  of  $B_3$  given by

$$B_3 \rightarrow \mathrm{GL}(\mathbb{R}^3) \subset \mathbb{R}^{3 \times 3}, \quad g = g_\tau g_\sigma \mapsto (\mathrm{sgn} \sigma)^\zeta (\det g_\tau)^\xi g_\tau^\xi g_\sigma,$$

where we (uniquely) write  $g \in B_3$  as a product of a sign-change matrix  $g_\tau$  and a permutation matrix  $g_\sigma$ .  $W_{0,1}$  and  $W_{1,1}$  are irreducible representations of  $B_3$ , but

$$W_{0,0} \cong W_{\mathrm{triv}} \oplus W_{\mathrm{std}}, \quad W_{1,0} = W_{\mathrm{sgn}} \oplus W_{\mathrm{std}},$$

where  $W_{\mathrm{triv}}$  is the one-dimensional trivial representation,  $W_{\mathrm{std}}$  the two-dimensional irreducible representation of the symmetric group, and  $W_{\mathrm{sgn}}$  the one-dimensional alternating representation of the symmetric group. The five irreducible  $B_3$ -representations  $W_{\mathrm{triv}}$ ,  $W_{\mathrm{sgn}}$ ,  $W_{\mathrm{std}}$ ,  $W_{0,1}$ ,  $W_{1,1}$  are all the irreducible  $B_3$ -representations such that  $-\mathrm{id} \in B_3$  lies in the kernel of the corresponding group homomorphism  $B_3 \rightarrow \mathrm{GL}(\mathbb{R}^k)$  [GW09].

For  $d \equiv 2 \pmod{3}$ , Theorem 5.2 then corresponds to a decomposition of  $\mathcal{H}_{2d}$  into the four types of representations  $W_{0,0}$ ,  $W_{1,0}$ ,  $W_{0,1}$  and  $W_{1,1}$  as

$$\mathcal{H}_{2d} \cong \bigoplus_{j=0}^{k-1} W_{\zeta(j), \xi(j)},$$

and we have given an explicit construction of a basis  $u_{i,j}$  of  $\mathcal{H}_{2d}$  corresponding to this decomposition. Indeed, for each  $j \in \{0, \dots, k-1\}$ , the three-dimensional subspace of  $\mathcal{H}_{2d}$  spanned by  $u_{1,j}, u_{2,j}, u_{3,j}$  corresponds to the representation  $W_{\zeta(j), \xi(j)}$ .

If  $d \not\equiv 2 \pmod{3}$ , then only for  $j \geq 1$  is it true that the subspace spanned by  $u_{1,j}, u_{2,j}, u_{3,j}$  is the three-dimensional representation  $W_{\zeta(j), \xi(j)}$ . For  $j = 0$ , we get the one-dimensional trivial representation  $W_{\mathrm{triv}}$  if  $d \equiv 0 \pmod{3}$ , or the two-dimensional irreducible representation  $W_{\mathrm{std}}$  of the symmetric group if  $d \equiv 1 \pmod{3}$ . Precise counting gives the following decompositions of  $\mathcal{H}_{2d}$  into the mentioned  $B_3$ -representations:

$$\mathcal{H}_{2d} \cong \begin{cases} \left[ \frac{d+1}{6} \right] W_{0,0} \oplus \left[ \frac{d+1}{6} \right] W_{1,0} \oplus \left[ \frac{d}{2} \right] W_{0,1} \oplus \left[ \frac{d}{2} \right] W_{1,1} & \text{if } d \equiv 2 \pmod{3}, \\ W_{\mathrm{triv}} \oplus \left[ \frac{d}{6} \right] W_{0,0} \oplus \left[ \frac{d}{6} \right] W_{1,0} \oplus \left[ \frac{d}{2} \right] W_{0,1} \oplus \left[ \frac{d}{2} \right] W_{1,1} & \text{if } d \equiv 1 \pmod{3}, \\ W_{\mathrm{std}} \oplus \left[ \frac{d-1}{6} \right] W_{0,0} \oplus \left[ \frac{d-1}{6} \right] W_{1,0} \oplus \left[ \frac{d}{2} \right] W_{0,1} \oplus \left[ \frac{d}{2} \right] W_{1,1} & \text{if } d \equiv 0 \pmod{3}. \end{cases}$$

**5.4.  $B_3$ -equivariant bases for  $\Lambda_{2d}$ .** In order to describe rational  $B_3$ -invariants on  $\Lambda_{2d}$ , we now describe a basis for  $\Lambda_{2d} = \mathcal{H}_{2d} \oplus \mathfrak{q}\mathcal{H}_{2d-2} \oplus \dots \oplus \mathfrak{q}^{d-1}\Lambda_2$  with  $B_3$ -symmetries. The main observation is that we can combine the linear spanning sets of the spaces  $\mathcal{H}_{2d+2}$  and  $\mathcal{H}_{2d}$ , when  $d \equiv 0 \pmod{3}$ , from Theorem 5.2, so as to remove the linear dependencies. Specifically, we observe:

**Lemma 5.9.** *Let  $m \geq 1$  and  $k := 8m + 2$ . The  $3k$ -dimensional subspace  $\mathcal{H}_{6m+2} \oplus \mathfrak{q}\mathcal{H}_{6m} \subset V_{6m+2}$  has a  $B_3$ -equivariant basis  $\tilde{u}_{i,j}$  ( $1 \leq i \leq 3, 0 \leq j < k$ ) with respect to some maps  $\zeta, \xi: \{0, \dots, k-1\} \rightarrow \{0, 1\}$ .*

*Proof.* A  $B_3$ -equivariant basis is given by

$$\{\mathfrak{q}u_{i,0}^{(6m)} + u_{i,0}^{(6m+2)} \mid 1 \leq i \leq 3\} \cup \{\mathfrak{q}u_{i,j}^{(6m)} \mid i, j \geq 1\} \cup \{u_{i,j}^{(6m+2)} \mid i, j \geq 1\}$$

where  $\{u_{i,j}^{(6m)}\}$  and  $\{u_{i,j}^{(6m+2)}\}$  are respectively the spanning sets of  $\mathcal{H}_{6m}$  and  $\mathcal{H}_{6m+2}$  constructed in Theorem 5.2 and the maps  $\zeta$  and  $\xi$  are defined accordingly.  $\square$

**Theorem 5.10.** *Let  $d \geq 1$  and let  $k := \left\lfloor \frac{(d+1)(2d+1)}{3} \right\rfloor - 1$ . Then the vector space  $\Lambda_{2d}$  contains  $3k$  linearly independent elements  $w_{i,j} \in \Lambda_{2d}$  for  $1 \leq i \leq 3, 0 \leq j < k$  forming a  $B_3$ -equivariant set with respect to some maps  $\zeta, \xi: \{0, \dots, k-1\} \rightarrow \{0, 1\}$ .*

*If  $d$  is not divisible by 3, then these  $w_{i,j}$  form a basis of  $\Lambda_{2d}$ . If  $d \equiv 0 \pmod{3}$ , the set  $\{w_{i,j}\}$  can be extended to a basis of  $\mathcal{H}_{2d}$  by adding an element  $w_\infty \in \Lambda_{2d}$  which is fixed under the action of  $B_3$ .*

*Proof.* We recall that  $\Lambda_{2d} = \mathcal{H}_{2d} \oplus q\mathcal{H}_{2d-2} \oplus \dots \oplus q^{d-2}\mathcal{H}_4 \oplus q^{d-1}\Lambda_2$ , where  $q^{d-1}\Lambda_2 \subset V_{2d}$  is a three-dimensional subspace with basis  $w_{1,0} := q^{d-1}x^2, w_{2,0} := q^{d-1}y^2, w_{3,0} := q^{d-1}z^2$ . For  $i \in \{1, 2, 3\}$  we observe that  $gw_{i,0} = w_{\sigma(i),0}$  if  $g \in B_3$  is a  $3 \times 3$  permutation matrix with corresponding permutation  $\sigma$  and  $gw_{i,0} = w_{i,0}$  if  $g \in B_3$  is a  $3 \times 3$  sign-change matrix.

From Theorem 5.2 and Lemma 5.9 we obtain bases  $\{u_{i,j}^{(6m-2)}\} \subset \mathcal{H}_{6m-2}$  and  $\{\tilde{u}_{i,j}^{(6m+2)}\} \subset \mathcal{H}_{6m+2} \oplus q\mathcal{H}_{6m}$  with the desired  $B_3$ -equivariance property. By multiplying with appropriate powers of  $q$ , we obtain the desired elements  $w_{i,j} \in \Lambda_{2d}$ . Explicitly,

$$\{w_{i,j}\} = \left\{ q^{d-3m-1} \tilde{u}_{i,j}^{(6m+2)} \mid 1 \leq m \leq \frac{d-2}{6} \right\} \cup \left\{ q^{d-3m+1} u_{i,j}^{(6m-2)} \mid 1 \leq m \leq \frac{d+1}{3} \right\} \cup \left\{ u_{i,j}^{(6m)} \mid m = \frac{d}{3}, j \geq 1 \right\},$$

additionally to the elements  $w_{i,0}$  from above. Note that the last set in this union is empty if  $d \not\equiv 0 \pmod{3}$ , and otherwise consists of those elements  $u_{i,j}^{(2d)} \in \mathcal{H}_{2d}$  that are linearly independent.

From Theorem 5.2 and Lemma 5.9 it is clear that this set  $\{w_{i,j}\}$  is  $B_3$ -equivariant (adequately inheriting the definition of the maps  $\zeta, \xi$  from Theorem 5.2 and Lemma 5.9). In the case  $d \equiv 0 \pmod{3}$ , we do not yet have a basis of  $\Lambda_{2d}$ , because we left out the element

$$w_\infty := u_{1,0}^{(2d)} = u_{2,0}^{(2d)} = u_{3,0}^{(2d)}$$

from the linear spanning set  $u_{i,j}^{(2d)}$  of the subspace  $\mathcal{H}_{2d} \subset \Lambda_{2d}$  described in Theorem 5.2. Note that the relation  $u_{1,0}^{(2d)} = u_{2,0}^{(2d)} = u_{3,0}^{(2d)}$  and the description of the  $B_3$ -action on  $u_{1,0}^{(2d)}, u_{2,0}^{(2d)}, u_{3,0}^{(2d)}$  in Theorem 5.2 imply that  $gw_\infty = w_\infty$  for all  $g \in B_3$ . This concludes the proof.  $\square$

Again it is important for applications to not consider Theorem 5.10 a pure existence result, but to observe how to immediately obtain the basis  $\{w_{i,j}\}$  (resp.  $\{w_{i,j}, w_\infty\}$ ) of  $\Lambda_{2d}$  from harmonic polynomials  $u_{i,j}^{(2k)} \in \mathcal{H}_{2k}$  as in Theorem 5.2. In particular, one can easily write out such a basis. Slightly deviating from Definition 5.1, we will call the constructed basis  $B_3$ -equivariant, even in the case  $d \equiv 0 \pmod{3}$ , when the basis includes an element  $w_\infty$  fixed under  $B_3$ .

**Remark 5.11.** We observe that for  $d \geq 2$  there exists an index  $j_0 \in \{0, \dots, k-1\}$  such that  $\zeta(j_0) = 0$  and  $\xi(j_0) = 1$  (and may assume after permuting the basis elements that this holds for  $j = 0$ ). Indeed, the basis contains – up to multiplication with a power of  $q$  – the elements  $u_{i,j}^{(4)}$  forming a basis of  $\mathcal{H}_4$ . This basis of  $\mathcal{H}_4$ , given in Proposition 4.1, satisfies the desired property, as remarked after Theorem 5.2.

**5.5. Rational invariants for ternary forms of arbitrary even degree.** With the construction of the  $B_3$ -equivariant basis for the slice  $\Lambda_{2d} \subset V_{2d}$  established for arbitrary  $d$ , we can now turn to the construction of a generating set of rational invariants for the action of  $B_3$  on  $\Lambda_{2d}$ , generalizing the case  $2d = 4$  described in Section 4.

Let  $d \geq 2$  and consider a basis  $\{w_{i,j} \mid 1 \leq i \leq 3, 0 \leq j < k\} \cup \{w_\infty\}$  of  $\Lambda_{2d}$  and corresponding maps  $\zeta, \xi: \{0, \dots, k-1\} \rightarrow \{0, 1\}$  as described in Theorem 5.10. Here, and from now on, we distinguish between the case  $d \not\equiv 0 \pmod{3}$  and the case  $d \equiv 0 \pmod{3}$ , where there is an additional basis element  $w_\infty \in \Lambda_{2d}$ , with square brackets, whenever we are confident that no confusion arises from this.

As observed in Remark 5.11, we can assume  $\zeta(0) = 0, \xi(0) = 1$ . With this chosen basis, an element  $v \in \Lambda_{2d}$  can be uniquely expressed as

$$(5.3) \quad v = \sum_{i=1}^3 \gamma_i w_{i,0} + \sum_{i=1}^3 \sum_{j=1}^{k-1} \alpha_{i,j} w_{i,j} [+ \alpha_\infty w_\infty]$$

for  $\gamma_i, \alpha_{i,j} [, \alpha_\infty] \in \mathbb{R}$ . In this way, we identify  $\mathbb{R}(\Lambda_{2d})$  with the field of rational expressions in variables  $\gamma_i, \alpha_{i,j} [, \alpha_\infty]$ , i.e.

$$\mathbb{R}(\Lambda_{2d}) = \mathbb{R}(\gamma_i, \alpha_{i,j} [, \alpha_\infty] \mid 1 \leq i \leq 3, 0 \leq j < k).$$

**Theorem 5.12.** *With the notations as above, a minimal set of generating rational invariants for  $\mathbb{R}(\Lambda_{2d})^{\mathbb{B}_3}$  is given by the  $2d^2 + 3d - 2 = \dim \Lambda_{2d}$  polynomial functions  $\mathcal{I}_{2d} = \{p_{i,j}\} \cup \{p_\infty\}$ , whose values at  $v$  as in (5.3) are given as follows:*

- Three invariants are given by

$$p_{1,0}(v) := \gamma_1^2 + \gamma_2^2 + \gamma_3^2,$$

$$p_{2,0}(v) := \gamma_1 \gamma_2 \gamma_3,$$

$$p_{3,0}(v) := \gamma_1^4 + \gamma_2^4 + \gamma_3^4.$$

- In the case  $d \equiv 0 \pmod{3}$ , one invariant is given by

$$p_\infty(v) := \alpha_\infty.$$

- The remaining invariants in the generating set are given as the entries of the  $3 \times (k-1)$ -matrix

$$\begin{pmatrix} p_{1,1}(v) & \cdots & p_{1,k-1}(v) \\ p_{2,1}(v) & \cdots & p_{2,k-1}(v) \\ p_{3,1}(v) & \cdots & p_{3,k-1}(v) \end{pmatrix} := \begin{pmatrix} 1 & 1 & 1 \\ \gamma_1^2 & \gamma_2^2 & \gamma_3^2 \\ \gamma_1^4 & \gamma_2^4 & \gamma_3^4 \end{pmatrix} M(\alpha),$$

where  $k = \lfloor \frac{2d^2+3d-5}{3} \rfloor$  and  $M(\alpha)$  is the  $3 \times (k-1)$ -matrix whose  $(i, j)$ -th entry is

$$M(\alpha)_{i,j} = \gamma_i^{\xi(j)} ((\gamma_1^2 - \gamma_2^2)(\gamma_2^2 - \gamma_3^2)(\gamma_3^2 - \gamma_1^2))^{\zeta(j)} \alpha_{i,j}.$$

*Proof.* With Theorem 5.10 established, the proof is analogous to the one for Theorem 4.3: One checks that  $\mathcal{I}_{2d}$  consists of  $\mathbb{B}_3$ -invariants (for example, by the use of equivariant maps as before, or directly from the definitions). In order to express a given rational invariant  $\hat{p} \in \mathbb{R}(\Lambda_{2d})^{\mathbb{B}_3}$  as a rational combination in terms of  $\mathcal{I}_{2d}$ , we may first replace each occurrence of the variables  $\alpha_{i,j}$  in  $\hat{p}$  by a rational expression in terms of invariants in  $\mathcal{I}_{2d}$  and  $\gamma_1, \gamma_2, \gamma_3$  by using that  $M(\alpha)_{i,j}$  is the  $(i, j)$ -th entry of the matrix product

$$\begin{pmatrix} 1 & 1 & 1 \\ \gamma_1^2 & \gamma_2^2 & \gamma_3^2 \\ \gamma_1^4 & \gamma_2^4 & \gamma_3^4 \end{pmatrix}^{-1} \begin{pmatrix} p_{1,1}(v) & \cdots & p_{1,k-1}(v) \\ p_{2,1}(v) & \cdots & p_{2,k-1}(v) \\ p_{3,1}(v) & \cdots & p_{3,k-1}(v) \end{pmatrix}.$$

If  $d \equiv 0 \pmod{3}$ , we also replace each occurrence of  $\alpha_\infty$  by  $p_\infty$ . As in the proof of Theorem 4.3, we express the remaining  $\mathbb{B}_3$ -invariant rational expression in  $\gamma_1, \gamma_2, \gamma_3$  in terms of  $p_{1,0}, p_{2,0}, p_{3,0}$  – note that  $\gamma_1, \gamma_2, \gamma_3$  play the exact same role here as in Theorem 4.3.  $\square$

Explicitly, as in Section 4.2, we again have a routine for expressing rational  $\mathbb{B}_3$ -invariants on  $\Lambda_{2d}$  in terms of  $\mathcal{I}_{2d}$ , given by the following rewrite rules:

$$\begin{pmatrix} \alpha_{1,j} \\ \alpha_{2,j} \\ \alpha_{3,j} \end{pmatrix} \longrightarrow \frac{\delta^{1-\zeta(j)}}{c_0 p_{2,0}^{\xi(j)}} \begin{pmatrix} \gamma_2 \gamma_3 & 0 & 0 \\ 0 & \gamma_3 \gamma_1 & 0 \\ 0 & 0 & \gamma_1 \gamma_2 \end{pmatrix}^{\xi(j)} \begin{pmatrix} (\gamma_3^2 - \gamma_2^2) \gamma_2^2 \gamma_3^2 & \gamma_2^4 - \gamma_3^4 & \gamma_3^2 - \gamma_2^2 \\ (\gamma_1^2 - \gamma_3^2) \gamma_3^2 \gamma_1^2 & \gamma_3^4 - \gamma_1^4 & \gamma_1^2 - \gamma_3^2 \\ (\gamma_2^2 - \gamma_1^2) \gamma_1^2 \gamma_2^2 & \gamma_1^4 - \gamma_2^4 & \gamma_2^2 - \gamma_1^2 \end{pmatrix} \begin{pmatrix} p_{1,j} \\ p_{2,j} \\ p_{3,j} \end{pmatrix} \quad \forall j \geq 1,$$

$$\alpha_\infty \longrightarrow p_\infty, \quad \gamma_1 \longrightarrow \frac{1}{p_{3,0}} (p_{1,0} \gamma_2 \gamma_3 - \gamma_2^3 \gamma_3 - \gamma_2 \gamma_3^3),$$

$$\gamma_2^4 \longrightarrow p_{1,0} \gamma_2^2 - \gamma_2^2 \gamma_3^2 + p_{1,0} \gamma_3^2 - \gamma_3^4 + \frac{1}{2} (p_{2,0} - p_{1,0}^2), \quad \gamma_3^6 \longrightarrow \frac{1}{2} (p_{2,0} - p_{1,0}^2) \gamma_3^2 + p_{1,0} \gamma_3^4 + p_{3,0}^2,$$

where

$$\delta := (\gamma_1^2 - \gamma_2^2)(\gamma_2^2 - \gamma_3^2)(\gamma_3^2 - \gamma_1^2) \quad \text{and} \\ c_0 := \delta^2 = \frac{1}{2} p_{2,0}^3 - \frac{1}{4} p_{1,0}^6 - 27 p_{3,0}^4 + p_{1,0}^4 p_{2,0} - \frac{5}{4} p_{1,0}^2 p_{2,0}^2 - 9 p_{3,0}^2 p_{1,0} p_{2,0} + 5 p_{3,0}^2 p_{1,0}^3.$$

As in Section 4.2, we notice that the generating set  $\mathcal{I}_{2d} \subset \mathbb{R}(\Lambda_{2d})^{\mathbb{B}_3}$  is a transcendence basis for  $\mathbb{R}(\Lambda_{2d})^{\mathbb{B}_3}$  as a field extension over  $\mathbb{R}$ , since  $|\mathcal{I}_{2d}| = \dim \Lambda_{2d}$ . In particular, this generating set is of minimal cardinality.

By Corollary 3.3, there uniquely exists a corresponding generating set  $\mathcal{J}_{2d}$  of  $O_3$ -invariants on  $V_{2d}$  with  $|\mathcal{J}_{2d}| = |\mathcal{I}_{2d}| = \dim \Lambda_{2d}$ , attaining the lower bound from Theorem 2.2 because of  $\dim V_{2d} = \Lambda_{2d} + \dim O_3$ .

**Corollary 5.13.** *There exists a generating set of  $\dim V_{2d} - \dim O_3 = 2d^2 + 3d - 2$  rational invariants  $\mathcal{J}_{2d} \subset \mathbb{R}(V_{2d})^{O_3}$  uniquely determined by their restrictions to  $\Lambda_{2d} \subset V_{2d}$  that are given by the invariants  $\mathcal{I}_{2d} \subset \mathbb{R}[\Lambda_{2d}]^{B_3}$  specified in Theorem 5.12.*

## 6. SOLVING THE MAIN ALGORITHMIC CHALLENGES

In Sections 4 and 5, in Corollaries 4.4 and 5.13, we identified, for any  $d \geq 2$ , a finite set of rational invariants  $\mathcal{J}_{2d} \subset \mathbb{R}(V_{2d})^{O_3}$  generating  $\mathbb{R}(V_{2d})^{O_3}$ . We denote the number of these rational invariants as  $m := |\mathcal{J}_{2d}|$ . Recall that  $m$  is minimal, that is

$$m = \dim V_{2d} - \dim O_3 = \binom{2d+2}{2} - 3.$$

Instead of being given by a closed form formula, each invariant  $\tilde{p} \in \mathcal{J}_{2d}$  is uniquely determined, in viture of Theorem 3.2, by the restriction  $p = \tilde{p}|_{\Lambda_{2d}} \in \mathbb{R}(\Lambda_{2d})^{B_3}$  of the rational map  $\tilde{p}: V_{2d} \dashrightarrow \mathbb{R}$  to the subspace  $\Lambda_{2d} \subset V_{2d}$ . We shall in this section examine the practical implications of Theorem 3.2 in addressing the algorithmic challenges formulated in Section 2.2. For these, we provide algorithms that rely only on the explicit knowledge of the restricted invariants  $\mathcal{I}_{2d} = \{\tilde{p}|_{\Lambda_{2d}} \mid \tilde{p} \in \mathcal{J}_{2d}\}$ .

**6.1. The Evaluation Problem.** Having identified a finite generating set of rational invariants  $\mathcal{J}_{2d} \subset \mathbb{R}(V_{2d})^{O_3}$ , the most basic algorithmic question is: *How can we evaluate each of the generating invariants  $\tilde{p} \in \mathcal{J}_{2d}$  at a given point  $v \in V_{2d}$ ?*

Let  $\tilde{p} \in \mathcal{J}_{2d}$  and consider the restriction  $p := \tilde{p}|_{\Lambda_{2d}} \in \mathcal{I}_{2d}$  of  $\tilde{p}: V_{2d} \dashrightarrow \mathbb{R}$  to the subspace  $\Lambda_{2d} \subset V_{2d}$ . By construction of the invariants, we know explicit expressions for  $p \in \mathbb{R}(\Lambda_{2d})^{B_3}$ , which allow us to evaluate  $\tilde{p} \in \mathcal{J}_{2d}$  at any point of the slice  $u \in \Lambda_{2d}$  by computing  $\tilde{p}(u) = p(u) \in \mathbb{R}$ .

When we want to evaluate  $\tilde{p}(v)$  for arbitrary  $v \in V_{2d}$ , we observe that  $\tilde{p}(gv) = \tilde{p}(v)$  for all  $g \in O_3$ , since  $\tilde{p}$  is an invariants for the action of  $O_3$ . By Proposition 3.5, we know that for general  $v \in V_{2d}$  there exists an orthogonal transformation  $g \in O_3$  such that  $gv \in \Lambda_{2d}$ . For such  $g \in O_3$ , we may then compute

$$\tilde{p}(v) = \tilde{p}(gv) = p(gv).$$

Recalling the definition of the slice  $\Lambda_{2d}$ , this idea leads to Algorithm 1.

**Input** :  $v \in V_{2d} = \mathbb{R}[x, y, z]_{2d}$ ,  $d \geq 2$   
**Output**:  $\tilde{p}(v) \in \mathbb{R}$  for each  $\tilde{p} \in \mathcal{J}_{2d}$

- 1 Compute  $v' \in V_2$  in the Harmonic Decomposition
 
$$v = h_{2d} + qh_{2d-2} + q^2h_{2d-4} + \dots + q^{d-2}h_4 + q^{d-1}v', \quad \text{where } h_{2k} \in \mathcal{H}_{2k}.$$
- 2 Let  $A \in \mathbb{R}^{3 \times 3}$  be the symmetric matrix corresponding to the quadratic form  $v' \in V_2$  as in (1.3).
- 3 Determine  $g \in O_3$  such that the matrix product  $gAg^T$  is diagonal:
 
$$gAg^T = \text{diag}(\lambda_1, \lambda_2, \lambda_3) \quad \text{for some } \lambda_1, \lambda_2, \lambda_3 \in \mathbb{R}.$$
- 4 If  $\lambda_i = \lambda_j$  for some  $i \neq j$ , output “*undefined at v*” and stop.
- 5 Compute  $u := gv \in \Lambda_{2d}$ .
- 6 Compute and output the values  $p(u)$  for each  $p \in \mathcal{I}_{2d}$ .

**Algorithm 1:** Evaluation Algorithm

In the following, we comment on the validity and on the computational realization of the various steps of Algorithm 1.

*Validity of Algorithm 1.* First, we describe how the formulation in Algorithm 1 corresponds to the idea described above of evaluating  $\tilde{p}(v)$  as  $p(gv)$  where  $g \in \mathcal{O}_3$  is some orthogonal transformation such that  $gv \in \Lambda_{2d}$ . We recall that if

$$v = h_{2d} + qh_{2d-2} + q^2h_{2d-4} + \dots + q^{d-2}h_4 + q^{d-1}v'$$

is the Harmonic Decomposition of  $v$  computed in Step 1, then the Harmonic Decomposition of  $gv$  is given as

$$gv = gh_{2d} + q(gh_{2d-2}) + q^2(gh_{2d-4}) + \dots + q^{d-2}(gh_4) + q^{d-1}(gv')$$

by Proposition 2.7. Then the definition of  $\Lambda_{2d}$  gives:  $gv$  is contained in  $\Lambda_{2d}$  if and only if  $gv' \in V_2$  lies in  $\Lambda_2$ , i.e. the symmetric matrix associated to the quadratic form  $gv'$  is diagonal. This matrix is  $gAg^T$  (where  $A$  is the matrix of the quadratic form  $v' \in V_2$ ).

To validate Step 4 in Algorithm 1, we recall that even when the restricted invariant  $p \in \mathbb{R}(\Lambda_{2d})^{\mathcal{B}_3}$  is known to be a polynomial invariant, the corresponding invariant  $\tilde{p} \in \mathbb{R}(V_{2d})^{\mathcal{O}_3}$  is typically only a rational function, defined at a general point only. In fact, going back to the proof of Proposition 3.5, we see that the rational function  $\tilde{p}: V_{2d} \dashrightarrow \mathbb{R}$  which we obtain from  $p \in \mathbb{R}(\Lambda_{2d})^{\mathcal{B}_3}$  is only defined at those points  $v \in V_{2d}$  whose quadratic part  $v' \in V_2$  (in the Harmonic Decomposition) does not have repeated eigenvalues in the matrix representation (1.3). This precisely corresponds to Step 4. We observe that leaving out Step 4, Algorithm 1 would still output a (meaningless) value for the non-defined cases, but that value would depend on the choice of  $g$  in Step 3.

*Computational realization of Algorithm 1.* First, we discuss the implementation of Step 1: In order to compute  $v' \in V_2$  in the Harmonic Decomposition of  $v \in V_{2d}$ , we can use the explicit projection operators on  $\mathcal{H}_{2d}, \dots, \mathcal{H}_4$  given in [AR95]. Alternatively we express the given element  $v = \sum_{ijk} a_{ijk}x^i y^j z^k$  in terms of a basis of  $V_{2d}$  that reflects the Harmonic Decomposition

$$V_{2d} = \mathcal{H}_{2d} \oplus \dots \oplus q^{d-2}\mathcal{H}_4 \oplus q^{d-1}V_2.$$

Say,  $\mathcal{B}_{2k} \subset \mathcal{H}_{2k}$  is a basis of the space  $\mathcal{H}_{2k}$  of harmonic polynomials of degree  $2k$  (for  $2 \leq k \leq d$ ). Then

$$\mathbb{B}_{2d} := \bigcup_{k=2}^d q^{d-k}\mathcal{B}_{2k} \cup \{q^{d-1}x^2, q^{d-1}y^2, q^{d-1}z^2, q^{d-1}xy, q^{d-1}yz, q^{d-1}zx\}$$

is a basis of  $V_{2d}$  and we can uniquely write

$$(6.1) \quad v = \sum_{k=2}^d \sum_{b \in \mathcal{B}_{2k}} \alpha_b q^{d-k} b + \beta_{200} q^{d-1} x^2 + \beta_{020} q^{d-1} y^2 + \beta_{002} q^{d-1} z^2 + \beta_{110} q^{d-1} xy + \beta_{011} q^{d-1} yz + \beta_{101} q^{d-1} zx$$

with  $\alpha_b, \beta_{ijk} \in \mathbb{R}$ . Then the Harmonic Decomposition of  $v$  as in Step 1 is given by

$$h_{2k} = \sum_{b \in \mathcal{B}_{2k}} \alpha_b b \in \mathcal{H}_{2k}, \quad v' = \beta_{200} x^2 + \beta_{020} y^2 + \beta_{002} z^2 + \beta_{110} xy + \beta_{011} yz + \beta_{101} zx \in V_2.$$

For  $\mathcal{B}_{2k} \subset \mathcal{H}_{2k}$ , we might as well use the elements  $u_{i,j}^{(2k)}$  explicitly constructed in Theorem 5.2 (by selecting a linearly independent subset of these in the cases  $k \not\equiv 2 \pmod{3}$ ), though it is not essential at this stage. For chosen bases  $\mathcal{B}_{2k} \subset \mathcal{H}_{2k}$ , computing the expression (6.1) for a given  $v = \sum_{ijk} a_{ijk} x^i y^j z^k$  corresponds to converting from the monomial basis  $\{x^i y^j z^k \mid i + j + k = 2d\} \subset V_{2d}$  to the basis  $\mathbb{B}_{2d} \subset V_{2d}$ . The corresponding base-change matrix is the inverse of the matrix whose entries are given by the coefficients of the expressions  $b \in \mathbb{B}_{2d}$  in the variables  $x, y, z$ . This base-change matrix may be precomputed for fixed  $d$ . If we use bases  $\mathcal{B}_{2k} \subset \mathcal{H}_{2k}$  built from elements  $u_{i,j}^{(2k)} \in \mathcal{H}_{2k}$  as constructed in Theorem 5.2, the base change transformation could in fact be described explicitly by a careful analysis of the proofs in Section 5.2.

Step 3 is the problem of computing the *eigendecomposition* of the symmetric matrix  $A$  and is equivalent to finding the eigenvalues and eigenvectors of  $A$ . *Exact (symbolic)* algorithmic solutions to this problem would involve introducing (nested) square roots and cubic roots, but such expressions are typically very undesirable from a practical standpoint. However, there are well-established numerical methods for computing the

eigendecomposition of a symmetric matrix – with the additional benefit that the numerical stability of these methods is well-studied [Kre05], [GVL13, Chapter 8].

Realizing Step 5 is straightforward by the definition of the action of  $g = (g_{ij}) \in O_3$  on  $V_{2d}$ : Applying the substitutions

$$x \mapsto g_{11}x + g_{21}y + g_{31}z, \quad y \mapsto g_{12}x + g_{22}y + g_{32}z, \quad z \mapsto g_{13}x + g_{23}y + g_{33}z,$$

to  $v \in V_{2d} = \mathbb{R}[x, y, z]_{2d}$  and expanding the resulting expression gives  $u = gv \in \Lambda_{2d}$ . This expansion may also be precomputed symbolically for fixed  $d$  such that it is only necessary to evaluate with the entries  $g_{ij}$ .

Finally, for Step 6, we want to evaluate at  $u \in \Lambda_{2d}$  the expressions for the rational invariants  $\mathcal{I}_{2d} \subset \Lambda_{2d}$  described in Theorem 5.12. For that, we express  $u = \sum_{i=1}^3 \gamma_i w_{i,0} + \sum_{i=1}^3 \sum_{j=1}^k \alpha_{i,j} w_{i,j} [+ \alpha_\infty w_\infty]$ , where  $\{w_{i,j} \mid 1 \leq i \leq 3, 0 \leq j \leq k\} \cup \{w_\infty\}$  is the basis of  $\Lambda_{2d}$  from Theorem 5.10. This can again be realized by an appropriate linear base change transformation. We then evaluate  $p(u)$  for each  $p \in \mathcal{I}_{2d}$  with the formulas for the invariants given in Theorem 5.12.

**6.2. The Rewriting Problem.** In this section, we discuss how to address the Rewriting Problem specified in Section 2.2.

For notational simplification, we now assume that the rational invariants in  $\mathcal{J}_{2d} \in \mathbb{R}(V_{2d})^{O_3}$  are indexed as  $\mathcal{J}_{2d} = \{\tilde{p}_1, \dots, \tilde{p}_m\}$  and their restrictions to  $\Lambda_{2d}$  are  $\mathcal{I}_{2d} = \{p_1, \dots, p_m\}$ . Since  $\tilde{p}_1, \dots, \tilde{p}_m \in \mathbb{R}(V_{2d})^{O_3}$  form a set of generating rational invariants, it is possible to express any other rational invariant  $\tilde{p}_0 \in \mathbb{R}(V_{2d})^{O_3}$  as a rational combination of  $\tilde{p}_1, \dots, \tilde{p}_m$ , i.e. there exists a rational expression in  $m$  variables  $r(T_1, \dots, T_m) \in \mathbb{R}(T_1, \dots, T_m)$  such that

$$(6.2) \quad \tilde{p}_0 = r(\tilde{p}_1, \dots, \tilde{p}_m).$$

Determining such  $r(T_1, \dots, T_m) \in \mathbb{R}(T_1, \dots, T_m)$  for any given  $\tilde{p}_0 \in \mathbb{R}(V_{2d})^{O_3}$  is a problem that can be reduced to the corresponding problem for  $B_3$ -invariants on the subspace  $\Lambda_{2d} \subset V_{2d}$ .

Note that restricting the equality (6.2) to the subspace  $\Lambda_{2d} \subset V_{2d}$  gives:

$$\tilde{p}_0|_{\Lambda_{2d}} = r(p_1, \dots, p_m).$$

Hence, to determine the rational expression  $r$ , it is sufficient to rewrite  $p_0 := \tilde{p}_0|_{\Lambda_{2d}} \in \mathbb{R}(\Lambda_{2d})^{B_3}$  in terms of the restricted generating rational invariants  $p_1, \dots, p_m \in \mathbb{R}(\Lambda_{2d})^{B_3}$ . This leads to Algorithm 2.

**Input** :  $\tilde{p}_0 \in \mathbb{R}(V_{2d})^{O_3}$  for some  $d \geq 2$   
**Output**: A rational expression  $r \in \mathbb{R}(T_1, \dots, T_m)$  such that  $\tilde{p}_0 = r(\tilde{p}_1, \dots, \tilde{p}_m)$

- 1 Let  $p_0 := \tilde{p}_0|_{\Lambda_{2d}} \in \mathbb{R}(\Lambda_{2d})^{B_3}$ .
- 2 Determine  $r \in \mathbb{R}(T_1, \dots, T_m)$  such that  $p_0 = r(p_1, \dots, p_m)$ .
- 3 Output  $r$ .

**Algorithm 2:** Rewriting Algorithm

Step 2 is the problem of rewriting a rational  $B_3$ -invariant on  $\Lambda_{2d}$  in terms of the generating rational invariants  $p_1, \dots, p_m \in \mathbb{R}(\Lambda_{2d})^{B_3}$ . The realization of this is addressed in the proofs of Theorems 4.3 and 5.12 and made explicit by the “rewrite rules” given in Sections 4.2 and 5.5.

**6.3. The Reconstruction Problem.** By construction,

$$\mathcal{J}_{2d} = \{\tilde{p}_{i,j} \mid 1 \leq i \leq 3, 0 \leq j < k\} \cup \{\tilde{p}_\infty\}$$

is the set of rational  $O_3$ -invariants on  $V_{2d}$  such that

$$\tilde{p}_{i,j}|_{\Lambda_{2d}} = p_{i,j} \quad \forall i, j \quad [\text{and} \quad \tilde{p}_\infty|_{\Lambda_{2d}} = p_\infty]$$

are the  $B_3$ -invariants on  $\Lambda_{2d}$  from Theorem 5.12.

In Section 6.1, we saw how to numerically evaluate  $\tilde{p}_{i,j}(v) \in \mathbb{R}$  [and  $\tilde{p}_\infty(v) \in \mathbb{R}$ ] at a general point  $v \in V_{2d}$ . Now we consider the inverse algorithmic problem: Given real values  $\mu_{i,j} \in \mathbb{R}$  for  $1 \leq i \leq 3, 0 \leq j < k$  [and  $\mu_\infty \in \mathbb{R}$ ], we want to compute  $v \in V_{2d}$  such that  $\tilde{p}_{i,j}(v) = \mu_{i,j}$  for all  $i, j$  [and  $\tilde{p}_\infty = \mu_\infty$ ]. This may not be possible for all  $m$ -tuples  $\mu = (\mu_{i,j} [ , \mu_\infty] \mid 1 \leq i \leq 3, 0 \leq j < k) \in \mathbb{R}^m$ , so we are also interested in the following question:

*For which  $m$ -tuples  $\mu = (\mu_{i,j} [ , \mu_\infty] \mid 1 \leq i \leq 3, 0 \leq j < k) \in \mathbb{R}^m$  does there exist  $v \in V_{2d}$  such that  $\tilde{p}_{i,j}(v) = \mu_{i,j}$  for all  $i, j$  [and  $\tilde{p}_\infty = \mu_\infty$ ]?*

Note that the reconstructed  $v \in V_{2d}$  is not uniquely determined, since for any orthogonally equivalent  $w \in V_{2d}$  (i.e.  $w = gv$  for some  $g \in O_3$ ), the invariants  $\tilde{p} \in \mathcal{J}_{2d}$  take the same values for  $v$  and  $w$ . Theorem 2.3 implies that generically, the reconstructed  $v$  is unique up to orthogonal transformations, i.e. any different reconstructed  $w \in V_{2d}$  is orthogonally equivalent to  $v$ .

By Proposition 3.5, for any  $v \in V_{2d}$  there exists  $g \in O_3$  such that  $u := gv \in \Lambda_{2d}$ . Then

$$\tilde{p}(v) = \tilde{p}(gv) = p(u) \quad \forall \tilde{p} \in \mathcal{J}_{2d}, p := \tilde{p}|_{\Lambda_{2d}}.$$

In particular, we can always choose to reconstruct an element  $v$  that lies in the subspace  $\Lambda_{2d}$ .

With respect to the basis  $\{w_{i,j} \mid 1 \leq i \leq 3, 0 \leq j < k\} \cup \{w_\infty\}$  of  $\Lambda_{2d}$  from Theorem 5.10, we therefore want to determine  $\alpha_{i,j}, \gamma_i, [\alpha_\infty] \in \mathbb{R}$  such that

$$v = \sum_{i=1}^3 \gamma_i w_{i,0} + \sum_{i=1}^3 \sum_{j=1}^{k-1} \alpha_{i,j} w_{i,j} [+ \alpha_\infty w_\infty] \in \Lambda_{2d}$$

satisfies  $p_{i,j}(v) = \mu_{i,j} \forall i, j$  [and  $p_\infty(v) = \mu_\infty$ ]. With the explicit formulas for the invariants given in Theorem 5.12, this leads to the problem of solving the following system of polynomial equations in unknowns  $\alpha_{i,j}, \gamma_i [ , \alpha_\infty] \in \mathbb{R}$ :

$$(6.3) \quad \begin{aligned} \gamma_1^2 + \gamma_2^2 + \gamma_3^2 &= \mu_{1,0}, & \gamma_1 \gamma_2 \gamma_3 &= \mu_{2,0}, & \gamma_1^4 + \gamma_2^4 + \gamma_3^4 &= \mu_{3,0}, & [\alpha_\infty = \mu_\infty], \\ \begin{pmatrix} 1 & 1 & 1 \\ \gamma_1^2 & \gamma_2^2 & \gamma_3^2 \\ \gamma_1^4 & \gamma_2^4 & \gamma_3^4 \end{pmatrix} \cdot M(\alpha) &= \begin{pmatrix} \mu_{1,1} & \cdots & \mu_{1,k-1} \\ \mu_{2,1} & \cdots & \mu_{2,k-1} \\ \mu_{3,1} & \cdots & \mu_{3,k-1} \end{pmatrix}, \end{aligned}$$

where the  $(i, j)$ -th entry of the matrix  $M(\alpha)$  is  $M(\alpha)_{i,j} = \alpha_{i,j} \gamma_i^{\xi(j)} ((\gamma_1^2 - \gamma_2^2)(\gamma_2^2 - \gamma_3^2)(\gamma_3^2 - \gamma_1^2))^{\zeta(j)}$ .

The crucial part for the resolution of the polynomial system (6.3) lies in solving the first three equation for  $\gamma_1, \gamma_2, \gamma_3$ . Once values for  $\gamma_1, \gamma_2, \gamma_3 \in \mathbb{R}$  are known, we are left with a system of *linear* equations in the remaining variables. Therefore, the following observations are essential:

**Lemma 6.1.** *Let  $\mu_{1,0}, \mu_{2,0}, \mu_{3,0} \in \mathbb{C}$ . If  $\gamma_1, \gamma_2, \gamma_3 \in \mathbb{C}$  are a solution of the system*

$$\gamma_1^2 + \gamma_2^2 + \gamma_3^2 = \mu_{1,0}, \quad \gamma_1 \gamma_2 \gamma_3 = \mu_{2,0}, \quad \gamma_1^4 + \gamma_2^4 + \gamma_3^4 = \mu_{3,0},$$

*then the squares  $\gamma_1^2, \gamma_2^2, \gamma_3^2 \in \mathbb{C}$  are the zeroes (with multiplicities) of the cubic polynomial*

$$T^3 - \mu_{1,0} T^2 + \frac{\mu_{1,0}^2 - \mu_{3,0}}{2} T - \mu_{2,0}^2 \in \mathbb{C}[T].$$

*Proof.* Let  $\gamma_1, \gamma_2, \gamma_3 \in \mathbb{C}$  be solution of the system and let  $f \in \mathbb{C}[T]$  be the polynomial whose zeroes (with multiplicities) are  $\gamma_1^2, \gamma_2^2, \gamma_3^2$ . Then

$$\begin{aligned} f(T) &= (T - \gamma_1^2)(T - \gamma_2^2)(T - \gamma_3^2) = T^3 - (\gamma_1^2 + \gamma_2^2 + \gamma_3^2)T^2 + (\gamma_1^2 \gamma_2^2 + \gamma_2^2 \gamma_3^2 + \gamma_3^2 \gamma_1^2)T - \gamma_1^2 \gamma_2^2 \gamma_3^2 \\ &= T^3 - \mu_{1,0} T^2 + \frac{\mu_{1,0}^2 - \mu_{3,0}}{2} T - \mu_{2,0}^2. \end{aligned}$$

□

The following classical fact then characterizes when the solution in Lemma 6.1 has real solutions:



**Lemma 6.2.** *A cubic polynomial  $f(T) = T^3 - aT^2 + bT - c \in \mathbb{R}[T]$  has three distinct positive real solutions if and only if*

$$a, b, c > 0 \quad \text{and} \quad a^2b^2 - 4b^3 - 4a^3c - 27c^2 + 18abc > 0.$$

*Proof.* The expression  $a^2b^2 - 4b^3 - 4a^3c - 27c^2 + 18abc$  is the discriminant of the cubic polynomial  $f$ , which is positive if and only if  $f$  has three distinct real solutions. Descartes' rule of signs implies that  $f$  has no negative solutions if and only if the signs of the coefficients of  $f$  alternate, i.e.  $a, b, c > 0$ .  $\square$

Combining these two results, we obtain Algorithm 3.

**Input** :  $\mu_{i,j} \in \mathbb{R}$  for  $1 \leq i \leq 3, 0 \leq j < k$  [plus  $\mu_\infty \in \mathbb{R}$ ]  
**Output**: A ternary form  $v \in V_{2d}$  such that  $\tilde{p}_{i,j}(v) = \mu_{i,j}$  for all  $i, j$  [and  $\tilde{p}_\infty(v) = \mu_\infty$ ], if possible

```

1 if  $d$  is a multiple of 3 then
2   |  $\alpha_\infty := \mu_\infty$ 
3 end
4  $a := \mu_{1,0}, b := \frac{\mu_{1,0}^2 - \mu_{3,0}}{2}, c := \mu_{2,0}^2$ 
5 if  $a, b, c > 0$  and  $a^2b^2 - 4b^3 - 4a^3c - 27c^2 + 18abc > 0$  then
6   | Compute the (distinct) roots  $r_1, r_2, r_3 > 0$  of the polynomial  $T^3 - aT^2 + bT - c \in \mathbb{R}[T]$ .
7   |  $\gamma_1 := \pm\sqrt{r_1}, \gamma_2 := \pm\sqrt{r_2}, \gamma_3 := \pm\sqrt{r_3}$ , where we choose signs such that  $\gamma_1\gamma_2\gamma_3 = \mu_{2,0}$ .
8   | Calculate the matrix product  $M := \begin{pmatrix} 1 & 1 & 1 \\ r_1 & r_2 & r_3 \\ r_1^2 & r_2^2 & r_3^2 \end{pmatrix}^{-1} \begin{pmatrix} \mu_{1,1} & \dots & \mu_{1,k-1} \\ \mu_{2,1} & \dots & \mu_{2,k-1} \\ \mu_{3,1} & \dots & \mu_{3,k-1} \end{pmatrix}$ .
9   | for  $1 \leq i \leq 3, 1 \leq j < k$  do
10  |   |  $\alpha_{i,j} := \frac{M_{i,j}}{\gamma_i^{\xi(j)}((r_1-r_2)(r_2-r_3)(r_3-r_1))^{\zeta(j)}}$ .
11  |   end
12  |   Output the ternary form  $v = \sum_{i=1}^3 \gamma_i w_{i,0} + \sum_{i=1}^3 \sum_{j=1}^k \alpha_{i,j} w_{i,j}$  [ $+ \alpha_\infty w_\infty$ ]  $\in \Lambda_{2d}$ .
13 else
14 |   Output “The values  $\mu_{i,j}$  allow no unambiguous reconstruction in  $V_{2d}$ ”.
15 end

```

**Algorithm 3:** Reconstruction Algorithm.

For Step 14 in Algorithm 3, it should be observed that an unambiguous reconstruction requires  $\gamma_1^2, \gamma_2^2, \gamma_3^2$  to be distinct and non-zero. Then the validity of the algorithm follows from the discussion above.

## REFERENCES

- [ABR01] S. Axler, P. Bourdon, and W. Ramey. *Harmonic function theory*, volume 137 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 2001.
- [AH12] K. Atkinson and W. Han. *Spherical harmonics and approximations on the unit sphere: an introduction*, volume 2044 of *Lecture Notes in Mathematics*. Springer, Heidelberg, 2012.
- [AR95] S. Axler and W. Ramey. Harmonic polynomials and Dirichlet-type problems. *Proc. Amer. Math. Soc.*, 123(12):3765–3773, 1995.
- [BP07] P. Basser and S. Pajevic. Spectral decomposition of a 4th-order covariance tensor : Applications to diffusion tensor mri. *Signal Processing*, 87(2):220–236, February 2007.
- [BVF08] A. Barmpoutis, B.C. Vemuri, and J.R. Forder. Fast displacement probability profile approximation from hardi using 4th-order tensors. In *Proceedings of ISBI: IEEE International Symposium on Biomedical Imaging*, pages 911–914, May 2008.
- [CTS07] J.-L. Colliot-Thélène and J.-J. Sansuc. The rationality problem for fields of invariants under linear algebraic groups (with special regards to the Brauer group). In *Algebraic groups and homogeneous spaces*, volume 19 of *Tata Inst. Fund. Res. Stud. Math.*, pages 113–186. Tata Inst. Fund. Res., Mumbai, 2007.
- [CV15] E. Caruyer and R. Verma. On facilitating the use of hardi in population studies by creating rotation-invariant markers. *Medical Image Analysis*, 20(1):87 – 96, 2015.

- [DK02] H. Derksen and G. Kemper. *Computational invariant theory*. Invariant Theory and Algebraic Transformation Groups, I. Springer-Verlag, Berlin, 2002. Encyclopaedia of Mathematical Sciences, 130.
- [DK15] H. Derksen and G. Kemper. *Computational invariant theory*. Springer-Verlag, 2 edition, 2015.
- [DVW<sup>+</sup>09] C. Delmaire, M. Vidailhet, Demian Wassermann, Maxime Descoteaux, Romain Valabregue, F. Bourdain, Christophe Lenglet, S. Sangla, A. Terrier, Rachid Deriche, and Stéphane Lehericy. Diffusion abnormalities in the primary sensorimotor pathways in writer’s cramp. *Archives of Neurology*, 66(4), April 2009.
- [FK77] K. Fox and B. Krohn. Computation of cubic harmonics. *J. Computational Phys.*, 25(4):386–408, 1977.
- [Fra02] L.R. Frank. Characterization of anisotropy in high angular resolution diffusion-weighted MRI. *Magnetic Resonance in Medicine*, 47(6):1083–1099, 2002.
- [GPD12a] A. Ghosh, T. Papadopoulos, and R. Deriche. Biomarkers for hardi: 2nd & 4th order tensor invariants. In *IEEE International Symposium on Biomedical Imaging: From Nano to Macro - 2012*, Barcelona, Spain, May 2012.
- [GPD12b] A. Ghosh, T. Papadopoulos, and R. Deriche. Generalized Invariants of a 4th order tensor: Building blocks for new biomarkers in dMRI. In *Computational Diffusion MRI Workshop (CDMRI), MICCAI*, pages 165–173, Nice, France, 2012.
- [GVL13] G. Golub and C. Van Loan. *Matrix computations*. Johns Hopkins Studies in the Mathematical Sciences. Johns Hopkins University Press, Baltimore, MD, fourth edition, 2013.
- [GW09] R. Goodman and N. R. Wallach. *Symmetry, representations, and invariants*, volume 255 of *Graduate Texts in Mathematics*. Springer, Dordrecht, 2009.
- [GY10] J. Grace and A. Young. *The algebra of invariants*. Cambridge Library Collection. Cambridge University Press, Cambridge, 2010. Reprint of the 1903 original.
- [HK07a] E. Hubert and I. Kogan. Rational invariants of a group action. Construction and rewriting. *J. Symbolic Comput.*, 42(1-2):203–217, 2007.
- [HK07b] E. Hubert and I. Kogan. Smooth and algebraic invariants of a group action. Local and global constructions. *Foundations of Computational Mathematics*, 7(4):355–393, 2007.
- [HL12] E. Hubert and G. Labahn. Rational invariants of scalings from Hermite normal forms. In *ISSAC 2012*, pages 219–226. ACM Press, 2012.
- [HL13] E. Hubert and G. Labahn. Scaling invariants and symmetry reduction of dynamical systems. *Foundations of Computational Mathematics*, 13(4):479–516, 2013.
- [HL16] E. Hubert and G. Labahn. Computation of the invariants of finite abelian groups. *Mathematics of Computations*, 85(302):3029–3050, 2016.
- [Hub12] E. Hubert. Algebraic and differential invariants. In F. Cucker, T. Krick, A. Pinkus, and A. Szanto, editors, *Foundations of computational mathematics, Budapest 2011*, number 403 in London Mathematical Society Lecture Note Series, pages 168–190. Cambridge University Press, 2012.
- [Isa09] I. Isaacs. *Algebra: a graduate course*, volume 100 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2009.
- [Kre05] D. Kressner. *Numerical methods for general and structured eigenvalue problems*, volume 46 of *Lecture Notes in Computational Science and Engineering*. Springer-Verlag, Berlin, 2005.
- [LMM<sup>+</sup>10] X. Li, A. Messé, G. Marrelec, M. Péligrini-Issac, and H. Benali. An enhanced voxel-based morphometry method to investigate structural changes: application to alzheimer’s disease. *Neuroradiology*, 52(3):203–213, 2010.
- [Mug72] J. Mugli. Cubic harmonics as linear combinations of spherical harmonics. *Z. Angew. Math. Phys.*, 23:311–317, 1972.
- [OKA17] M. Olive, B. Kolev, and N. Auffray. A minimal integrity basis for the elasticity tensor. *Archive for Rational Mechanics and Analysis*, 2017.
- [Oli16] M. Olive. About gordan’s algorithm for binary forms. *Foundations of Computational Mathematics*, pages 1–60, 2016.
- [PGD14] T. Papadopoulos, A. Ghosh, and R. Deriche. Complete Set of Invariants of a 4th Order Tensor: The 12 Tasks of HARDI from Ternary Quartics. In Polina Golland, Nobuhiko Hata, Christian Barillot, Joachim Hornegger, and Robert Howe, editors, *Medical Image Computing and Computer-Assisted Intervention – MICCAI 2014*, volume 8675 of *Lecture Notes in Computer Science*, pages 233 – 240, Boston, United States, September 2014.
- [Pop94] V. Popov. Sections in invariant theory. In *The Sophus Lie Memorial Conference (Oslo, 1992)*, pages 315–361. Scand. Univ. Press, Oslo, 1994.
- [PV94] V. Popov and È. Vinberg. *Invariant theory*, volume 55 of *Encyclopaedia of Mathematical Sciences*. Springer-Verlag, Berlin, 1994. A translation of it Algebraic geometry. 4 (Russian), Akad. Nauk SSSR Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1989, Translation edited by A. N. Parshin and I. R. Shafarevich.
- [Ros56] M. Rosenlicht. Some basic theorems on algebraic groups. *American Journal of Mathematics*, 78:401–443, 1956.
- [Sch01] G. Schwarz. Algebraic quotients of compact group actions. *J. Algebra*, 244(2):365–378, 2001.
- [Ses62] C. Seshadri. On a theorem of Weitzenböck in invariant theory. *J. Math. Kyoto Univ.*, 1:403–409, 1961/1962.
- [Stu08] B. Sturmfels. *Algorithms in invariant theory*. Texts and Monographs in Symbolic Computation. SpringerWien-NewYork, Vienna, second edition, 2008.
- [TRWW03] D.S. Tuch, T.G. Reese, M.R. Weigel, and V.J. Wedeen. Diffusion mri of complex neural architecture. *Neuron*, 40:885–895, December 2003.
- [Tuc02] David S. Tuch. *Diffusion MRI of Complex Tissue Structure*. PhD thesis, Harvard University and Massachusetts Institute of Technology, 2002.

[Veg00] G. Vegter. The apolar bilinear form in geometric modeling. *Math. Comp.*, 69(230):691–720, 2000.

INRIA MÉDITERRANÉE, MPI MIS LEIPZIG

*E-mail address:* `goerlach@mis.mpg.de`

INRIA MÉDITERRANÉE

*E-mail address:* `evelyne.hubert@inria.fr`

INRIA MÉDITERRANÉE

*E-mail address:* `theodore.papadopoulo@inria.fr`