

Genome Sequence of the Yeast Clavispora lusitaniae Type Strain CBS 6936

Pascal Durrens, Christophe C. Klopp, Nicolas Biteau, Valérie Fitton-Ouhabi, Karine Dementhon, Isabelle Accoceberry, David J. Sherman, Thierry Noël

▶ To cite this version:

Pascal Durrens, Christophe C. Klopp, Nicolas Biteau, Valérie Fitton-Ouhabi, Karine Dementhon, et al.. Genome Sequence of the Yeast Clavispora lusitaniae Type Strain CBS 6936. Genome Announcements, 2017, 5 (31), pp.e00724-17. 10.1128/genomeA.00724-17. hal-01583349

HAL Id: hal-01583349 https://inria.hal.science/hal-01583349

Submitted on 7 Sep 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

1	The genome of the yeast Clavispora lusitaniae type strain CBS 6936
2	
3	Pascal Durrens ^{#abc} , Christophe Klopp ^e , Nicolas Biteau ^{bd} , Valérie Fitton-Ouhabi ^{bd} ,
4	Karine Dementhon ^{bd} , Isabelle Accoceberry ^{bd} , David J. Sherman ^{cab} , Thierry Noël ^{bd}
5	
6	Laboratoire Bordelais de Recherche en Informatique (LaBRI), UMR-CNRS 5800,
7	Talence, France ^a ; Université de Bordeaux, Bordeaux, France ^b ; INRIA joint project
8	team PLEIADE ^c ; Laboratoire de Microbiologie Fondamentale et Pathogénicité UMR-
9	CNRS 5234, Bordeaux, France ^d ; Genotoul Bioinfo, MIAT, INRA Toulouse, Castanet-
10	Tolosan, France ^e
11	
12	
13	Running Head: Genome of C. lusitaniae type strain CBS 6936
14	
15	#Address correspondence to Pascal Durrens, pascal.durrens@labri.fr
16	

17 Abstract

18

19 Clavispora lusitaniae, an environmental saprophytic yeast belonging to the CTG clade 20 of Candida, can behave occasionally as an opportunistic pathogen in humans. We 21 report the genome sequence of the type strain CBS 6936. Comparison with sequences 22 of strain ATCC 42720 indicates conservation of chromosomal structure but significant 23 nucleotide divergence.

Clavispora lusitaniae, teleomorph of Candida lusitaniae, is an ubiquist environmental ascomycetous yeast, with no known specific ecological niche. It can be isolated from different substrates, such as soils, waters, plants, and gastrointestinal tracts of many animals including birds, mammals and humans. In immunocompromised hosts, C. *lusitaniae* can be pathogenic and is responsible for about 1% of invasive candidiasis, particularly in pediatric and onco-haematology patients (1).

31

32 So far, two strains have had their genomes sequenced : ATCC 42720, isolated from the blood of a patient with myeloid leukemia (2), and MTCC 1001, a self-fertile strain 33 34 isolated from citrus (3). We report here the genome sequencing and assembly of the C. *lusitaniae* type strain CBS 6936 (4), isolated from citrus peel juice. Genomic DNA was 35 36 isolated from a 50 ml YPD yeast culture, after sphaeroplast osmotic lysis and glass rod purification of ethanol-precipitated nucleic acids. The DNA library was prepared from 37 1 µg according to the NEBNext DNA Library Prep Master Mix Set for Illumina 38 39 (E6040) protocol with an insert size of 368±122nt generated with an ultrasonicator 40 Covaris.

41

The library was sequenced on a MiSeq Illumina v1.18.54 producing *circa* 1.5 million paired end reads, representing a 28X coverage. A *de novo* assembly of the reads was performed with the aim to compare the contigs to those of the ATCC 42720 strain. We used SPAdes v3.9.0 with default parameters (5) to perform the assembly. Only the 53 scaffolds longer than 1kb were retained as the GC % of many shorter ones deviated from the average.

49 The scaffolds represent an overall length of 12.0Mb which is close to the 12.11Mb of the ATCC 42720 genome. The GC% amounts to 44.5% which is identical to that of 50 51 ATCC 42720 strain. The alignment of CBS 6936 scaffolds against ATCC 42720 contigs by Nucmer (6) shows almost co-linearity between the two sets, apart from a 66kb long 52 inversion. The number of SNPs between CBS 6936 and ATCC 42720 amounts to 53 132141 SNPs with Phred quality higher than 30, according to BWA (7) alignment of 54 55 reads and Freebayes detection (8). This SNP density of 1 SNP per 90bp is twice the level observed between the strains SC5314 and WO-1 of Candida albicans, members 56 of different subgroups in the species and qualified as relatively recent divergent (9). 57

58

59 Genes for 4 rRNA, 197 tRNA and 5539 proteins were predicted by alignment, 60 TRNAscan-SE and a combination of Augustus (10), Snap (11) and Genmark-ES (12), 61 respectively. The average identity of proteins between the two strains amounts to 62 89.9% ± 23.2, based on BLASTp (13) best hit alignments.

63

Altogether, these results indicate that, in spite of a conserved genome structure, the
sequences are different, meaning that the strains underwent a significant divergence.

67 Accession numbers. This whole genome shotgun project has been deposited at 68 DDBJ/EMBL/GenBank under the accession numbers SRP075809 for reads and 69 LYUB00000000 for scaffolds. The second version of the assembly of CBS 6936 is 70 version LYUB02000000 and is described here.

72 Acknowledgements

73

The library and nucleotide sequencing were performed at the Genomic and Transcriptomic Facility of Bordeaux (grants from the Conseil Regional d'Aquitaine n°20030304002FA and 20040305003FA and from the European Union, FEDER n°2003227 and from Investissements d'avenir, Convention attributive d'aide N°ANR-10-EQPX-16-01). We would like to thank Christophe Hubert for its expertise and technical assistance.

The development of the bioinformatic pipeline was carried out using the PLAFRIM experimental testbed, being developed under the Inria PlaFRIM development action with support from Bordeaux INP, LABRI and IMB and other entities: Conseil Régional d'Aquitaine, Université de Bordeaux and CNRS (and ANR in accordance to the programme d'investissements d'Avenir (see https://plafrim.fr/).

88	1.	Favel A, Michel-Nguyen A, Peyron F, Martin C, Thomachot L, Datry A,
89		Bouchara JP, Challier S, Noël T, Chastin C, Regli P. 2003. Colony
90		morphology switching of Candida lusitaniae and acquisition of multidrug
91		resistance during treatment of a renal infection in a newborn: case report and
92		review of the literature. Diagn. Microbiol. Infect. Dis. 47:331-339.
93	2.	Pappagianis D, Collins MS, Hector R, Remington J. 1979. Development of
94		resistance to amphotericin B in Candida lusitaniae infecting a human.
95		Antimicrob Agents Chemother. 16:123-126.
96	3.	Lachance M, Daniel H, Meyer W, Prasad G, Gautam S, Boundymills K.
97		2003. The D1/D2 domain of the large-subunit rDNA of the yeast species is
98		unusually polymorphic. FEMS Yeast Res 4:253–258.
99	4.	Rodrigues de Miranda L. 1979. Clavispora, a new yeast genus of
100		Saccharomycetales. Antonie van Leeuwenhoek. 45:479-483.
101	5.	Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS,
102		Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin
103		AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: A New
104		Genome Assembly Algorithm and Its Applications to Single-Cell Sequencing. J
105		Comput Biol. 5:455-477.
106	6.	Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C,
107		Salzberg SL. 2004. Versatile and open software for comparing large genomes.
108		Genome Biology, 5:R12.
109	7.	Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-

- 110 Wheeler Transform. Bioinformatics, 25:1754-1760.
- 8. Garrison E, Marth G. 2012. Haplotype-based variant detection from shortread sequencing. arXiv preprint arXiv:1207.3907 [q-bio.GN].
- 113 9. Butler G, Rasmussen MD, Lin MF, , Santos MAS, Sakthikumar S, Munro
- 114 CA, Rheinbay E, Grabherr M, Forche A, Reedy JL, Agrafioti I, Arnaud
- 115 MB, Bates S, Brown AJP, Brunke S, Costanzo MC, Fitzpatrick DA, de
- 116 Groot PWJ, Harris D, Hoyer LL, Hube B, Klis FM, Kodira C, Lennard N,
- 117 Logue ME, Martin R, Neiman AM, Nikolaou E, Quail MA, Quinn J,
- 118 Santos MC, Schmitzberger FF, Sherlock G, Shah P, Silverstein K,
- 119 Skrzypek MS, Soll D, Staggs R, Stansfield I, Stumpf MPH, Sudbery PE,
- 120 Thyagarajan S, Zeng Q, Berman J, Berriman M, Heitman J, Gow NAR,
- Lorenz MC, Birren BW, Kellis M, Cuomo CA. 2009. Evolution of
 pathogenicity and sexual reproduction in eight *Candida* genomes. Nature
 459:657-662.
- 124 10. Stanke M, Schöffmann O, Morgenstern B, Waack S. 2006. Gene prediction
 in eukaryotes with a generalized hidden Markov model that uses hints from
 external sources. BMC Bioinformatics 7, 62.
- 127 11. Korf I. 2004. Gene finding in novel genomes. BMC Bioinformatics 5, 59.
- 128 12. Ter-Hovhannisyan V, Lomsadze A, Chernoff YO, Borodovsky M. 2008.
 129 Gene prediction in novel fungal genomes using an *ab initio* algorithm with
 130 unsupervised training. Genome Res. 18:1979-1990.
- 131 13. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K,
- Madden TL. 2008 BLAST+: architecture and applications. BMC
 Bioinformatics 10:421.