





A Three-Dimensional Computational Model of Action Potential Propagation Through a Network of Individual Cells.

Pierre-Elliott Bécue, Mark Potse, Yves Coudière

State of the art and current issues

- · Well-known models (bidomain, monodomain) are homogenized;
- Assumed periodicity;
- Some specific issues (disorganization in the tissue) at the microscopic scale are arrhythmia-prone (DeBakker 1993);
- Need for a "microscopic" computational model.

Geometry of the problem and equations



Figure: Basic geometry of the problem

$$-\sigma_i \Delta u_i = 0$$
 Ω_i

$$-\sigma_e \Delta u_e = 0$$
 Ω_e

$$-\sigma_i \nabla u_i \cdot n_i = \sigma_e \nabla u_e \cdot n_e \qquad \Gamma$$

$$\mathcal{L}_m \partial_t(\mathbf{v}) + I_{\mathrm{ion}}(\mathbf{v}) = -\sigma_i \nabla u_i \cdot n_i \quad \Gamma$$

 $\sigma_e \nabla u_e \cdot n_e = 0 \quad \Gamma_e$
(1)

- Nonlinearities on the border of the cell (ionic model)
- Only v⁰ provided as initial data
- Pure Neumann problem

This problem has a weak solution

Assembly 1/2 - Variational formulation

We use a P1-Lagrange finite element method with semi-implicit Euler time-stepping method

$$\begin{cases} \int_{\Omega_{i}} \sigma_{i} \nabla \boldsymbol{u}_{i}^{n} \cdot \nabla \varphi_{i} \mathrm{d}\boldsymbol{x} + \int_{\Sigma} \left[c_{m} \frac{\boldsymbol{u}_{i}^{n} - \boldsymbol{u}_{e}^{n}}{\delta t} + I_{\mathrm{ion}}(\boldsymbol{v}^{n-1}) \right] \varphi_{i} \mathrm{d}\boldsymbol{s} \\ = \int_{\Sigma} c_{m} \frac{\boldsymbol{v}^{n-1}}{\delta t} \varphi_{i} \mathrm{d}\boldsymbol{s} \\ \int_{\Omega_{e}} \sigma_{e} \nabla \boldsymbol{u}_{e}^{n} \cdot \nabla \varphi_{e} \mathrm{d}\boldsymbol{x} - \int_{\Sigma} \left[c_{m} \frac{\boldsymbol{u}_{i}^{n} - \boldsymbol{u}_{e}^{n}}{\delta t} + I_{\mathrm{ion}}(\boldsymbol{v}_{e}^{n-1}) \right] \varphi_{e} \mathrm{d}\boldsymbol{s} \\ = -\int_{\Sigma} c_{m} \frac{\boldsymbol{v}^{n-1}}{\delta t} \varphi_{e} \mathrm{d}\boldsymbol{s} \end{cases}$$

Assembly 2/2 - Linear system

$$\begin{pmatrix} A_{i,i}^n & A_{e,i}^n \\ A_{i,e}^n & A_{e,e}^n \end{pmatrix} \begin{pmatrix} U_i^n \\ U_e^n \end{pmatrix} = F_{\text{ion}}^{n-1} + F_{\text{time}}^{n-1}$$

- Combination of stiffness part on the volume, mass part on the membrane and coupled terms (on A_{i,e} and A_{e,i});
- Ill-conditionned problem;
- We split the mesh in two parts (intra- and extracellular) and duplicate nodes on the membrane;

Simulation protocol



- Arbitrary stimulation at t = 0.1ms for 0.02ms between an anode and a cathode;
- Use of Mitchell-Schaeffer model;
- 2D cases: cells are 100 \times 20µm², 3D case: three cells embedded in a domain of size 300 \times 100 \times 70µm³;
- Extra/intracellular conductivity ratio is 1.75.

2D simulation - Single Cell



(a) Potential field at t = 0.15ms

(b) Potential field at t = 0.40ms



(c) Potential field at t = 5.0ms

Figure: First test case with a single cell, with an initial stimulation of intensity I = 4.6.

2D simulation - Two connected cells



(b) Potential field at t = 5.0ms

Figure: Second test case with two connected cells, with an initial stimulation of intensity I = 11.0. The channel dimensions are $10 \times 2\mu m^2$.

nría

3D simulation



(a) Surface of the cells for the 3D case.



(c) Depolarization process on the 4 colored points.

(b) Potential field at t = 0.4ms

Solution

5,000e-01

0.25

- 1.2M elements;
- 214k nodes;

-5,000e-01

- 40 000 steps;
- 4 3.3 GHz CPUs;
- 53.2h (whole AP).

-0.25

Conclusion and further work

- A numerical solver of the microscopic bidomain model;
- It works in 2D and 3D;
- CPU time is reasonable.
- Build larger network of cells (in 2D) to test the propagation of the action potential;
- Implement a more accurate gap-junction model (Davidovic, CinC 2015);
- Test the scaling on clusters.