



HAL
open science

Neural coding of variable song structure in the songbird

Xavier Hinaut, Aurore Cazala, Catherine del Negro

► **To cite this version:**

Xavier Hinaut, Aurore Cazala, Catherine del Negro. Neural coding of variable song structure in the songbird. EBM 2017 - European Birdsong Meeting, May 2017, Bordeaux, France. pp.1. hal-01665824

HAL Id: hal-01665824

<https://inria.hal.science/hal-01665824>

Submitted on 17 Dec 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Neural coding of variable song structure in the songbird

Xavier Hinaut^{1,2,3,4}, Aurore Cazala¹, Catherine Del Negro¹


1 Université Paris Saclay, UMR 9197, Neuro-PSI, Orsay, France. 2 Inria Bordeaux Sud-Ouest, Talence, France. 3 LaBRI, UMR 5800, CNRS, Bordeaux INP, Université de Bordeaux, Talence, France. 4 Institut des Maladies Neurodégénératives, UMR 5293, CNRS, Université de Bordeaux, Bordeaux, France.

Introduction

Songbirds are an excellent model for exploring the neural coding of variable sequences of categorical acoustic elements. The domesticated canary, for instance, produces highly variable songs with complex transition rules between two consecutive acoustic elements. These transition rules are non-Markovian (i.e. the next acoustic element to be sung is dependent on several previous elements, not only the last one) [1]. In the HVC of Bengalese finches, Bouchard and Brainard [5] found that variations in responses to a given syllable could be explained by a positive linear dependence on the convergence probability of preceding sequences. Here, we reanalyse data from [3] to see if similar findings could be found for canaries.

Goal

Our aim is to understand how complex song sequences produced by canaries are coded in a sensorimotor area, the HVC nucleus.




The **HVC** (somehow analogous to vocal premotor cortical areas) participates in generating **temporal organization** of acoustic elements within songs [2]. In the HVC of canaries, most of cells show auditory responses and are selective for the **bird's own song (BOS)** over its temporal variants (with a different order of acoustic elements) [3], but how the sequential structure (ordering of acoustic elements) is coded is poorly understood.

HYPOTHESIS

HVC neurons encode all or a subset of **syntax features** of the songs heard.

Song structure and ways to explore it




Each canary has a **specific repertoire** composed of 15-35 distinct syllables (acoustic elements organized into sequences). According to [1], sequences of acoustic elements in canary songs show **long-distance dependencies** (e.g. "chunks").

No correlation found btw responses and prob. div./conv.

Z-score vs. Probability


We found no correlation between phrases neuronal responses (firing rate or Z-score) for the BOS or the randomized song and their probability of divergence and convergence. Below: the Z-score for 5 units of one bird (S62) given probability of divergence (left) and convergence (right).



References


- [1] Markovitz et al., PLoS Comp. Bio. (2013).
- [2] Hahnloser et al., Nature (2002).
- [3] Lehongre & Del Negro, Neurosci. (2011).
- [4] Fujimoto et al., J. Neuro (2011).
- [5] Bouchard & Brainard, J. Neuro. (2013).

Songs and song-evoked responses of a single unit (bird S62)



Current work: studying responses to chunk modifications

In a new study, we recorded auditory responses in HVC of anesthetized canaries using multi-electrode arrays (8 or 16). We used as auditory stimuli the bird's own song (BOS) and modified versions of the BOS, in which we manipulated either the local or global structure. We reversed the position of two consecutive elements (**local**) or reordered parts of the song (**global**).



Acknowledgments

The work was supported by Project IDEX NeuroSaclay "From Sequence Learning to Language Acquisition". We would like to thank Nicolas Giret for his help and advises. Poster template inspired from <http://quantum.opticsolomouc.org/archives/1026> under Creative Commons Attribution-ShareAlike 3.0 Unported License.

Contacts

xavier.hinaut@inria.fr, {aurore.cazala; catherine.del-negro}@u-psud.fr