



A multi-scale, multi-paradigm epidemiological modelling language for investigating Q Fever regional infection dynamics

Sébastien Picault, Yu-Lin Huang, Vianney Sicard, Thierry Hoch, Elisabeta Vergu, François Beaudeau, Pauline Ezanno

► To cite this version:

Sébastien Picault, Yu-Lin Huang, Vianney Sicard, Thierry Hoch, Elisabeta Vergu, et al.. A multi-scale, multi-paradigm epidemiological modelling language for investigating Q Fever regional infection dynamics. 6th International Conference on Infectious Disease Dynamics (Epidemics 6), Nov 2017, Sitges, Barcelona, Spain. , 2017, 6th International Conference on Infectious Disease Dynamics (Epidemics 6). hal-01705055

HAL Id: hal-01705055

<https://inria.hal.science/hal-01705055>

Submitted on 9 Feb 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A multi-scale, multi-paradigm epidemiological modelling language for investigating Q Fever regional infection dynamics

S. Picault^{1,2}, Y.-L. Huang¹, V. Sicard¹, T. Hoch¹, E. Vergu³, F. Beaudeau¹ and P. Ezanno¹

¹BIOEPAR, INRA Oniris, Nantes, France (firstname.lastname@oniris-nantes.fr) ²CRISTAL, Univ. Lille, France ³MaIAGE, INRA, Jouy-en-Josas, France

Understanding Q fever spread


The regional spread of Q fever, an endemic cattle zoonosis, relies upon airborne transmission and between-herd trade movements. Our goal was to better understand the contribution of each pathway to yearly herd incidence by coupling within- and between-herd dynamics.

To accelerate model design, explore multiple assumptions at each scale with a reduced amount of code to write, and enhance reliability, reusability and reproducibility, we used a generic framework, EMuLsion, which provides a domain-specific language (DSL) for explicitly describing all components of epidemiological models, and relies upon a multi-level agent-based architecture to encompass existing modelling paradigms [1].

Exploring model assumptions

Within-herd level

- simplification of previous model [2]
- intermittent shedding: **return to S** vs. **latent state**




Between-herd level



- heterogeneity in herd characteristics
- external risk at purchase
- Ermak-Stockie vs. Gaussian aerial dispersion

Within-herd Q fever model

Automated agents aggregation based on concerns: an hybrid approach




The EMuLsion approach: involving experts and users throughout model design




Model calibration at regional scale and influence of herd density

- within-herd model calibrated on previous model [2]
- between-herd model calibrated on observed annual herd incidence



Distribution of annual herd incidence as output of different models based on the combination of variants of three main features.



Map of Finistère with the density of initially infected herds (2012) and the location of herds infected in at least one run. Color shows the proportion of infections caused by wind vs. trade movements.

Future work

- extend DSL + framework to other diseases
- develop modelling standard in epidemiology
- automatize production of decision support tools

References

- [1] S. Picault, Y.-L. Huang, V. Sicard, and P. Ezanno (2017). Enhancing Sustainability of Complex Epidemiological Models through a Generic Multilevel Agent-based Approach. *26th Int. Joint Conf. on Artificial Intelligence*.
- [2] A. Courcoul, H. Monod, M. Nielen, D. Klinkenberg, L. Hogerwerf, F. Beaudeau, and E. Vergu (2011). Modelling the effect of heterogeneity of shedding on the within herd *Coxiella burnetii* spread and identification of key parameters by sensitivity analysis. *J. Theor. Biol.* **284**(1):130–141.

Funding

The work was funded by the French Research Agency (ANR) through projects MIHMES (ANR-10-BINF-07) and CADENCE (ANR-16-CE32-0007-01), the European fund for the Regional Development (FEDER) of Pays-de-la-Loire, and the Animal Health Division of the French National Institute for Agricultural Research (INRA).