



**HAL**  
open science

## Dynamics of an HBV infection model with cell-to-cell transmission and CTL immune response

Hajar Besbassi, Zineb Elrhoubari, Khalid Hattaf, Yousfi Noura

► **To cite this version:**

Hajar Besbassi, Zineb Elrhoubari, Khalid Hattaf, Yousfi Noura. Dynamics of an HBV infection model with cell-to-cell transmission and CTL immune response. 2018. hal-01719109v1

**HAL Id: hal-01719109**

**<https://inria.hal.science/hal-01719109v1>**

Preprint submitted on 28 Feb 2018 (v1), last revised 5 May 2019 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Dynamics of an HBV infection model with cell-to-cell transmission and CTL immune response

Hajar Besbassi<sup>\*,1</sup>, Zineb Elrhoubari<sup>1</sup>, Khalid Hattaf<sup>1,2</sup>, Yousfi Noura<sup>1</sup>

<sup>1</sup>Department of Mathematics and Computer Science LAMS, Faculty of Sciences Ben M'sik, Hassan II University, P.O Box 7955 Sidi Othman, Casablanca, Morocco

<sup>2</sup>Centre Régional des Métiers de l'Éducation et de la Formation (CRMEF), 20340 Derb Ghalef, Casablanca, Morocco

## Abstract

In this work, we propose a mathematical model to describe the dynamics of the hepatitis B virus (HBV) infection by taking into account the cure of infected cells, the export of precursor cytotoxic T lymphocytes (CTL) cells from the thymus and both modes of transmission that are the virus-to-cell infection and the cell-to-cell transmission. The local stability of the disease-free equilibrium and the chronic infection equilibrium is obtained via characteristic equations. Furthermore, the global stability of both equilibria is established by using two techniques, the direct Lyapunov method for the disease-free equilibrium and the geometrical approach for the chronic infection equilibrium.

**Keywords:** HBV infection, immune response, nonlinear incidence, global stability.

## Introduction

HBV infection is a major global health problem that can cause acute or chronic infection and puts people at high risk of death from cirrhosis and liver cancer. In 2015, hepatitis B resulted in 887000 deaths, mostly from complications (including cirrhosis and hepatocellular carcinoma) [1]. In addition, Cytotoxic T Lymphocytes (CTL) cells play an important role in antiviral defense by killing the infected cells. On the other hand, HBV can spread by two fundamental modes, one by virus-to-cell infection through the extracellular space and the other by cell-to-cell transfer involving direct cell-to-cell contact [2, 3]. For these

---

<sup>1</sup>Corresponding author, Email: hajar.besbassi@gmail.com

reasons, we propose the following model:

$$\begin{cases} \dot{x} &= \lambda - dx - f(x, y, v)v - g(x, y)y + \rho y, \\ \dot{y} &= f(x, y, v)v + g(x, y)y - (a + \rho)y - pyz, \\ \dot{v} &= ky - \mu v, \\ \dot{z} &= s + \frac{c y z}{\omega + y} - bz, \end{cases} \quad (1)$$

where  $x(t)$ ,  $y(t)$ ,  $v(t)$  and  $z(t)$  represent the concentrations of uninfected cells, infected cells, free virus and CTL cells at time  $t$ , respectively. Susceptible host (healthy hepatocytes) cells are produced at rate  $\lambda$ , die at rate  $dx$  and become infected either by free virus at rate  $f(x, y, v)v$  or by direct contact with an infected cell at rate  $g(x, y)y$ . Hence, the term  $f(x, y, v)v + g(x, y)y$  denotes the total infection rate of uninfected cells. Infected cells cured at rate  $\rho y$ , die at rate  $ay$  and are killed by the CTL immune response at rate  $pyz$ . Free virus is produced by an infected cell at rate  $ky$  and decays at rate  $\mu v$ . CTL cells expand in response to viral antigen derived from infected cells at rate  $\frac{c y z}{\omega + y}$ , where  $c$  is HBV-specific CTL stimulation rate and  $\omega$  represents virus load for half-maximal CTL cells stimulation [4] and decay in the absence of antigenic stimulation at the rate  $bz$ . The parameter  $s$  denotes the export of precursor CTL cells from the thymus [4].

As in [5], the incidence functions  $f(x, y, v)$  and  $g(x, y)$  for the two modes are continuously differentiable and satisfy the following hypotheses:

( $H_0$ )  $g(0, y) = 0$ , for all  $y \geq 0$ ;  $\frac{\partial g}{\partial x}(x, y) \geq 0$  (or  $g(x, y)$  is a strictly monotone increasing function with respect to  $x$  when  $f \equiv 0$ ) and  $\frac{\partial g}{\partial y}(x, y) \leq 0$ , for all  $x \geq 0$  and  $y \geq 0$ .

( $H_1$ )  $f(0, y, v) = 0$ , for all  $y \geq 0$  and  $v \geq 0$ ,

( $H_2$ )  $f(x, y, v)$  is a strictly monotone increasing function with respect to  $x$  (or  $\frac{\partial f}{\partial x}(x, v, y) \geq 0$  when  $g(x, y)$  is a strictly monotone increasing function with respect to  $x$ ), for any fixed  $y \geq 0$  and  $v \geq 0$ ,

( $H_3$ )  $f(x, y, v)$  is a monotone decreasing function with respect to  $y$  and  $v$ .

Biologically, the four hypotheses are reasonable and consistent with the reality. For more details on the biological significance of these four hypotheses, we refer the reader to the works [5, 6, 7]. Further, the general incidence functions  $f(x, y, v)$  and  $g(x, y)$  include various types of incidence rates existing in the literature.

The main objective of this work is to investigate the dynamical behavior of system (1). To do this end, we start with the existence, the positivity and

boundedness of solutions, which implies that our model is well-posed. After, we determine the basic reproduction number and steady states of the model. Finally, the local and global stabilities of the disease-free equilibrium and the chronic infection equilibrium are established.

## 1 Positivity and boundedness of solutions

The first important step is to validate our model as model that represents the evolution of cells. In the following result, we show that all cell numbers are non-negative and bounded.

**Theorem 1.1.** *All solutions of system (11) starting from positive initial value  $(x_0, y_0, v_0, z_0)$  remain bounded and positive for all  $t > 0$ . Moreover, we have*

$$(i) \quad T(t) \leq T(0) + \lambda/\delta,$$

$$(ii) \quad v(t) \leq v_0 + (k/u) \|y\|_\infty,$$

$$(iii) \quad z(t) \leq z_0 + s/b + (c/pw)(\lambda/b + x_0 + y_0 + \max(0, 1-d/b) \|x\|_\infty + \max(0, 1-a/b) \|y\|_\infty),$$

where  $T = x + y$  that represents the total cells of liver and  $\delta = \min(d, a)$ .

**Proof 1.2.** *First, we prove the positivity, that any solution starting in positive quadrant  $\mathbb{R}^4 = \{(x, y, v, z) \in \mathbb{R}^4 : x \geq 0, y \geq 0, v \geq 0, z \geq 0\}$  stays quadrant. In fact, for  $(x(t), y(t), v(t), z(t)) \in \mathbb{R}^4$  we have:*

$$\begin{aligned} \dot{x}|_{x=0} &= \lambda + \rho y \geq 0, & \dot{y}|_{y=0} &= f(x, 0, v)v \geq 0, \\ \dot{v}|_{v=0} &= ky \geq 0, & \dot{z}|_{z=0} &= s \geq 0, \end{aligned} \quad (2)$$

this immediately implies that all solutions of the system (1) with initial condition  $(x_0, y_0, v_0, z_0) \in \mathbb{R}^4$  stay in the first quadrant. Now, for the boundness of the solution of the system (1). We have that  $\dot{T} \leq \lambda - dx - ay$  we deduce that

$$T(t) \leq T(0)e^{-\delta t} + \frac{\lambda}{\delta}(1 - e^{-\delta t}), \quad (3)$$

since  $0 \leq e^{-\delta t} \leq 1$  and  $1 - e^{-\delta t} \leq 1$ .

From, the third equation of our model we have

$$\dot{v} = ky - uv,$$

which implies

$$v(t) = v(0)e^{-ut} + k \int_0^t y(s)e^{(s-t)u} ds, \quad (4)$$

thus

$$v(t) \leq v(0) + \frac{k}{u} \|y\|_\infty (1 - e^{(s-t)u}), \quad (5)$$

Since  $1 - e^{-\delta t} \leq 1$ , we conclude (ii).

Last, we get from the fourth equation of (1),

$$\dot{z} + bz \leq s + \frac{c}{w}yz, \quad (6)$$

Hence,

$$\dot{z} + bz \leq s + \frac{c}{pw}[\lambda - [\lambda - (\dot{x} + dx) - (\dot{y} + ay)]], \quad (7)$$

then,

$$\begin{aligned} z(t)e^{bt} - z_0 &\leq \left(\frac{s}{b} + \frac{c\lambda}{bpw}\right)(e^{bt} - 1) - \frac{c}{pw} \left[ \int_0^t e^{(b-d)s} \frac{d}{ds}(x(s)e^{ds})ds \right] \\ &\quad - \frac{c}{pw} \left[ \int_0^t e^{(b-a)s} \frac{d}{ds}(y(s)e^{ds})ds \right] \end{aligned} \quad (8)$$

By integration by parts, we have

$$\begin{aligned} \int_0^t e^{(b-d)s} \frac{d}{ds}(x(s)e^{ds})ds &= [x(s)e^{bs}]_0^t - (b-d) \int_0^t x(s)e^{bs}ds \\ \int_0^t e^{(b-a)s} \frac{d}{ds}(y(s)e^{ds})ds &= [y(s)e^{bs}]_0^t - (b-a) \int_0^t y(s)e^{bs}ds \end{aligned} \quad (9)$$

thus,

$$\begin{aligned} z(t) &\leq \left[ \frac{c}{pw}(x_0 + y_0) + z_0 \right] e^{-bt} + \left( \frac{s}{b} + \frac{c\lambda}{bpw} \right) (1 - e^{-bt}) \\ &\quad + \frac{c}{pw} \left[ \int_0^t [(b-d)x(s) + (b-a)y(s)]e^{b(s-t)ds} - x(t) - y(t) \right] \end{aligned} \quad (10)$$

In order to find an upper bound to this integral, we study

If  $b-d \leq 0$  and  $b-a \leq 0$ , we have

$$z(t) \leq z_0 + \frac{s}{b} + \frac{c}{pw} \left( \frac{\lambda}{b} + x_0 + y_0 \right) \quad (11)$$

If  $b-d \leq 0$  and  $b-a \geq 0$ , we have

$$z(t) \leq z_0 + \frac{s}{b} + \frac{c}{pw} \left[ \frac{\lambda}{b} + x_0 + y_0 + \left(1 - \frac{a}{b}\right) \|y\|_\infty \right] \quad (12)$$

If  $b-d \geq 0$  and  $b-a \geq 0$ , we have

$$z(t) \leq z_0 + \frac{s}{b} + \frac{c}{pw} \left[ \frac{\lambda}{b} + x_0 + y_0 + \left(1 - \frac{d}{b}\right) \|x\|_\infty \right] \quad (13)$$

If  $b - d \geq 0$  and  $b - a \geq 0$ , we have

$$z(t) \leq z_0 + \frac{s}{b} + \frac{c}{p\omega} \left[ \frac{\lambda}{b} + x_0 + y_0 + \left(1 - \frac{d}{b}\right) \|x\|_\infty + \left(1 - \frac{a}{b}\right) \|y\|_\infty \right]. \quad (14)$$

From (11)-(14), we can conclude (iii). ■

## 2 Basic reproduction number and Equilibria

Obviously, system (11) has always one infection free equilibrium  $E_f \left(\frac{\lambda}{d}, 0, 0, \frac{s}{b}\right)$ . Then we define the basic reproduction number of our model(11) as follows :

$$R_0 = \frac{kf\left(\frac{\lambda}{d}, 0, 0\right) + ug\left(\frac{\lambda}{d}, 0\right)}{u\left(a + \rho + p\frac{s}{b}\right)} \quad (15)$$

which can be rewritten as  $R_0 = R_{01} + R_{02}$  where

$$R_{01} = \frac{k}{a + \rho + p\frac{s}{b}} \times f\left(\frac{\lambda}{d}, 0, 0\right) \times \frac{1}{u}$$

and

$$R_{02} = g\left(\frac{\lambda}{d}, 0\right) \times \frac{1}{a + \rho + p\frac{s}{b}}$$

In the formula (15)  $\frac{1}{a + \rho + p\frac{s}{b}}$  denotes the average life expectancy of infected cells, which is less than  $\frac{1}{a}$  because of the role of immune cells ;  $\frac{k}{(a + \rho + p\frac{s}{b})}$  denotes the amount of virus generated from an infected during its survival period;  $\frac{1}{u}$  is the average life expectancy of viruses;  $\frac{\lambda}{d}$  denotes the number of susceptible cells at the beginning of the infectious process, which means that  $f\left(\frac{\lambda}{d}, 0, 0\right)$  and  $g\left(\frac{\lambda}{d}, 0\right)$  are the values of both incidence functions when all cells are uninfected. Hence,  $R_{01}$  is the basic reproduction number corresponding to virus-to-cell infection mode, whereas  $R_{02}$  is the basic reproduction number corresponding to cell-to-cell transmission mode. therefore,  $R_0$  describes the average number of newly infected cells generated from one infected cell at the beginning of the infectious process. To find the other equilibrium of the system

we solve the system :

$$\lambda - dx - f(x, y, v)v - g(x, y)y + \rho y = 0, \quad (16)$$

$$f(x, y, v)v + g(x, y)y - (a + \rho)y - pyz = 0, \quad (17)$$

$$ky - \mu v = 0, \quad (18)$$

$$s + \frac{cyz}{\omega + y} - bz = 0, \quad (19)$$

By (16) to (19), we have  $v = \frac{ky}{u}$ ,  $z = \frac{s(\omega + y)}{b\omega + (b - c)y}$ ,  $x = \psi(y)$  where

$$\psi(y) = \frac{\lambda}{d} - y \left( \frac{a}{d} - \frac{ps(\omega + y)}{db\omega + dy(b - c)} \right) \quad (20)$$

Now, we define on interval  $[0, +\infty) - \left\{ \frac{b\omega}{b-c} \right\}$  the function  $\Gamma$  by

$$\Gamma(y) = f \left( \psi(y), y, \frac{ky}{u} \right) + g(\psi(y), y) - u \left( a + \rho + \frac{ps(\omega + y)}{b\omega + y(b - c)} \right). \quad (21)$$

We have

$$\Gamma(0) = u(a + \rho + p\frac{s}{b})(R_0 - 1) \geq 0$$

and

$$\begin{aligned} \psi'(y) &= -\frac{a}{d} - \frac{ps(\omega + y)}{b\omega + y(b - c)} - y \frac{psd\omega}{(db\omega + dy(b - c))^2} < 0, \\ \Gamma'(y) &= \frac{\partial f}{\partial x} \psi'(y) + \frac{\partial f}{\partial y} + \frac{k}{u} \frac{\partial f}{\partial v} + \frac{\partial g}{\partial x} \psi'(y) + \frac{\partial g}{\partial y} - \frac{ucps\omega}{(b\omega + y(b - c))^2} < 0. \end{aligned} \quad (22)$$

Let  $\alpha = \frac{b\omega}{c-b}$  be a root of  $\Gamma$ , Then we need to consider two cases:

- (i) If  $c > b$ , then  $\alpha > 0$  and  $z = \frac{s(\omega + y)}{b\omega + y(b - c)} \geq 0$ . So we have that  $y < \alpha$ , which mean that there is no equilibrium point if  $y \geq \alpha$ . It is evident to show that

$$\lim_{y \rightarrow \alpha^-} \Gamma(y) = -\infty. \quad (23)$$

We deduce that  $0 < y^* < y$ . Hence, we get  $\psi(y) < \psi(y^*) < \psi(0)$  ( $\psi$  is decreasing on  $(0, y)$ ). thus  $0 < x^* < \lambda/d$ . Clearly  $v^*$  and  $z^*$  are positive.)

we deduce, for  $R_0 > 1$ , there exists a unique chronic infection equilibrium  $E^*(x^*, y^*, v^*, z^*)$  with  $y^* \in (0, y)$ ,  $x^* > 0$ ,  $v^* > 0$  and  $z^* > 0$ .

- (ii) If  $c < b$ , then  $\alpha < 0$  and  $\lim_{y \rightarrow \infty} \psi(y) = -\infty$ . As  $\psi(0) = \lambda/d > 0$ , consquently, there exists a unique  $\bar{y} \in (0, \alpha)$  such that  $\psi(\bar{y}) = 0$ . We have

$$\Gamma(\bar{y}) = -u \left( a + \rho + \frac{ps(\omega + \bar{y})}{b\omega + \bar{y}(b - c)} \right) < 0. \quad (24)$$

Following the same method, we can show that  $x^*, v^*$  and  $z^*$  are positive. Hence, there exists a unique  $E^*(x^*, y^*, v^*, z^*)$  with  $y^* \in (0, \bar{y}), x^* > 0, v^* > 0, z^* > 0$ .

The pervious discussions can be summarized in the following result.

**Theorem 2.1.**

- (i) When  $R_0 \leq 1$ , the model (1) has always one free infection equilibrium of the form  $E_f(\frac{\lambda}{d}, 0, 0, \frac{s}{b})$ .
- (ii) When  $R_0 > 1$ , the model (1) has a unique chronic infection equilibrium of the form  $E^*(x^*, y^*, v^*, z^*)$  with  $x^* > 0, y^* > 0, v^* > 0$ , and  $z^* > 0$ .

### 3 Local stability of equilibria

In this section, we discuss the local stability of both equilibria of system (1). Note that the Jacobian matrix of system (1) is given by

$$\begin{pmatrix} -d - v \frac{\partial f}{\partial x} - y \frac{\partial g}{\partial x} & -v \frac{\partial f}{\partial y} - y \frac{\partial g}{\partial y} - g(x, y) + \rho & -v \frac{\partial f}{\partial v} - f(x, y, v) & 0 \\ v \frac{\partial f}{\partial x} + y \frac{\partial g}{\partial x} & v \frac{\partial f}{\partial y} + y \frac{\partial g}{\partial y} + g(x, y) - (a + \rho + pz) & v \frac{\partial f}{\partial v} + f(x, y, v) & -\rho y \\ 0 & \frac{k}{c\omega} & -u & 0 \\ 0 & \frac{c\omega}{(\omega + y)^2} & 0 & \frac{cy}{\omega + y} - b \end{pmatrix} \quad (25)$$

Firstly, we have the following result.

**Theorem 3.1.** *The free infection equilibrium  $E_f$  is locally asymptotically stable if  $R_0 < 1$  and becomes unstable if  $R_0 > 1$ .*

**Proof 3.2.** *Evaluated (25) at  $E_f$  we get*

$$J_{E_f} = \begin{pmatrix} -d & -g(\frac{\lambda}{d}, 0) + \rho & -f(\frac{\lambda}{d}, 0, 0) & 0 \\ 0 & g(x, y) - (a + \rho + p\frac{s}{b}) & f(\frac{\lambda}{d}, 0, 0) & 0 \\ 0 & \frac{k}{c\omega} & -u & 0 \\ 0 & \frac{c\omega}{b\omega^2} & 0 & -b \end{pmatrix} \quad (26)$$

*the characteristic equation is given by*

$$(\xi + d)(\xi + b)(\xi^2 + [u + a + \rho + p\frac{s}{b} - g(\frac{\lambda}{d}, 0)]\xi + u(-g(d\frac{\lambda}{d}, 0) + a + \rho + p\frac{s}{b}) - kf(\frac{\lambda}{d}, 0, 0)) = 0 \quad (27)$$



Hence, the roots of (27) are

$$\begin{aligned}
\xi_1 &= -d, \\
\xi_2 &= -b, \\
\xi_3 &= \frac{-(u + a + \rho + p\frac{s}{b} - g(\frac{\lambda}{d}, 0)) - \sqrt{\Delta}}{2} \\
\xi_4 &= \frac{-(u + a + \rho + p\frac{s}{b} - g(\frac{\lambda}{d}, 0)) + \sqrt{\Delta}}{2}
\end{aligned} \tag{28}$$

where  $\Delta = (u + a + \rho + p\frac{s}{b} - g(\frac{\lambda}{d}, 0))^2 - 4u(a + \rho + p\frac{s}{b})(1 - R_0)$ . Clearly,  $\xi_1$ ,  $\xi_2$  and  $\xi_3$  are negative. However,  $\xi_4$  is negative if  $R_0 < 1$  and is positive if  $R_0 > 1$ . Therefore,  $E_f$  is locally asymptotically stable if  $R_0 < 1$  and unstable if  $R_0 > 1$ . ■

Now we focus on the local stability of the chronic infection equilibrium  $E^*$ .

**Theorem 3.3.** *The chronic infection equilibrium  $E^*$  is locally asymptotically stable if  $R_0 > 1$ .*

**Proof 3.4.** *Supposing that  $R_0 > 1$ . Evaluating (25) at  $E^*$  gives*

$$\xi^4 + a_1\xi^3 + a_2\xi^2 + a_3\xi + a_4 = 0, \tag{29}$$

where

$$\begin{aligned}
a_1 &= \alpha - N - u - \rho y w - Q + d + M, \\
a_2 &= u(N - \alpha - \rho y w) - \sigma k + (Q + d)(\alpha - N - u - \rho y w) - dQ + M(\alpha + \rho - u - \rho y w - Q), \\
a_3 &= u(Q + d)(N - \alpha - \rho y w) + \sigma k Q + d(Q(\alpha - N - u - \rho y w) - \sigma k) \\
&\quad + M[u(\rho - \alpha - \rho y w) + Q(\alpha + \rho - u - \rho y w)], \\
a_4 &= duQ(N - \alpha - \rho y w) - \sigma k Q + MuQ(\rho - \alpha - \rho y w),
\end{aligned}$$

with

$$\begin{aligned}
M &= v \frac{\partial f}{\partial x} - y \frac{\partial g}{\partial x}, \\
N &= v \frac{\partial f}{\partial y} + y \frac{\partial g}{\partial y} + g(x, y), \\
\sigma &= v \frac{\partial f}{\partial v} + f(x, y, v),
\end{aligned}$$

$$\alpha = a + \rho + pz, \quad w = \frac{cz\omega}{(\omega + y)^2}, \quad Q = \frac{cy}{\omega + y} - b,$$

When  $R_0 > 1$ , it is clear that  $a_1, a_2, a_3$  and  $a_4$  are positive. In addition,

$$\begin{vmatrix} a_1 & 1 \\ a_3 & a_2 \end{vmatrix} = a_1 a_2 - a_3 > 0 \quad (30)$$

In the same way, we have

$$\begin{vmatrix} a_1 & 1 & 0 \\ a_3 & a_2 & a_1 \\ 0 & a_4 & a_1 \end{vmatrix} = a_3 \begin{vmatrix} a_1 & 1 \\ a_2 & a_3 \end{vmatrix} - a_1^2 a_4 > 0. \quad (31)$$

From the Routh-Hurwitz theorem we know that all roots of (29) have negative real parts. Thus, the chronic infection equilibrium  $E^*$  is locally asymptotically stable for  $R_0 > 1$ . ■

## 4 Global Stability of Equilibria

In this section, we study the global stability of both equilibria. For the free infection equilibrium  $E_f$ , we assume that  $a \geq d$ . Therefore, we have the following result.

**Theorem 4.1.** *The free infection equilibrium  $E_f$  is globally asymptotically stable if  $R_0 \leq 1$ .*

**Proof 4.2.** *Consider*

$$\Omega = \left\{ (x, y, v, z) \in \mathbb{R}_+^4 : z \geq \frac{s}{b} \right\}$$

We see that any solution  $(x(t), y(t), v(t), z(t))$  starting in  $\Omega$  remains there forever. In fact, by theorem 1.1 we have  $(x(t), y(t), v(t), z(t)) \in \mathbb{R}_+^4$ . It remains to prove that  $z \geq \frac{s}{b}$  with  $z_0 \geq \frac{s}{b}$ . According to the fourth equation of (11), we have :

$$z(t) \geq \frac{s}{b} + \left( z_0 - \frac{s}{b} \right) e^{-bt} \Rightarrow z(t) \geq \frac{s}{b}, \quad (32)$$

Which implies that  $z \geq \frac{s}{b}$ . Then  $(x(t), y(t), v(t), z(t)) \in \Omega$ . We construct the Lyapunov functional  $L$  on  $\Omega$  as follow :

$$L(t) = y(t) + \frac{f(\frac{\lambda}{d}, 0, 0)}{u} v(t).$$

Calculating the time derivative of  $L$  along the positive solution of (1), we get

$$\frac{dL}{dt}(t) = \left( f(x, y, v) - f\left(\frac{\lambda}{d}, 0, 0\right) \right) v + (a + \rho + pz)y \left( \frac{kf\left(\frac{\lambda}{d}, 0, 0\right) + ug(x, y)}{u(a + \rho + pz)} - 1 \right)$$

. It is not hard to see that  $\limsup_{t \rightarrow \infty} x(t) \leq \frac{\lambda}{d}$  and  $\limsup_{t \rightarrow \infty} z(t) \leq \frac{s}{b}$ . This yields that all  $\Omega$  limit points satisfy  $x(t) \leq \frac{\lambda}{d}$  and  $z(t) \leq \frac{s}{b}$ . So it suffices to consider solutions for which  $x(t) \leq \frac{\lambda}{d}$  and  $z(t) \leq \frac{s}{b}$ . Using the expression of  $R_0$  given in (15), we obtain

$$\begin{aligned} \frac{dL}{dt}(t) &\leq \left( f(x, 0, 0) - f\left(\frac{\lambda}{d}, 0, 0\right) \right) v + (a + \rho + pz)(R_0 - 1)y \\ &\leq (a + \rho + pz)(R_0 - 1)y, \end{aligned}$$

Since  $R_0 \leq 1$ , we have  $\frac{dL}{dt}(t) \leq 0$ . In addition, it is easy to show that the largest compact invariant set in  $\{(x, y, v, z) \mid \frac{dL}{dt}(t) = 0\}$  is the singleton  $\{E_f\}$ . By the LaSalle invariance principle, the free infection equilibrium  $E_f$  is globally asymptotically stable for  $R_0 \leq 1$ . ■

Next, we will study the global dynamics of system (1) when  $R_0 > 1$ . Firstly, we need the following lemma.

**Lemma 4.3.** *The system (1) is uniformly persistent if  $R_0 > 1$ .*

**Proof 4.4.** *The maximal invariant set  $M$  on the boundary  $\partial\Omega$  is the singleton  $\{E_f\}$  and it is isolated. Further, Theorem 4.3 given in [8] ensures the equivalence between the uniform persistence of system (1) and the instability of the free infection equilibrium  $E_f$ . From theorem 3.1 we have  $E_f$  is unstable if  $R_0 > 1$ . Therefore, system (1) is uniformly persistent if  $R_0 > 1$ . ■*

Now, we focus on the global stability of  $E^*$  under the assumption  $R_0 > 1$  and the incidence function  $f$  satisfies the following property:

$$(H_4) \quad f(x, y, v) + v \frac{\partial f}{\partial v}(x, y, v) \geq 0 \text{ for all } x \geq 0, y \geq 0 \text{ and } v \geq 0.$$

**Theorem 4.5.** *Suppose  $R_0 < 1$  and (4) holds. Then the chronic infection equilibrium  $E^*$  is globally asymptotically stable.*

**Proof 4.6.** To study the global stability of  $E^*$ , we first consider the geometrical approach by Li and Muldowney[9].

Introduced by the following subsystem of (1):

$$\begin{cases} \dot{x} = \lambda - dx - f(x, y, v)v - g(x, y)y + \rho y, \\ \dot{y} = f(x, y, v)v + g(x, y)y - (a + \rho)y - pyz, \\ \dot{v} = ky - \mu v, \end{cases} \quad (33)$$

The Jacobian matrix of system (33) is

$$J = \begin{pmatrix} -d - v \frac{\partial f}{\partial x} - y \frac{\partial g}{\partial x} & -v \frac{\partial f}{\partial y} - y \frac{\partial g}{\partial y} - g(x, y) + \rho & -v \frac{\partial f}{\partial v} - f(x, y, v) \\ v \frac{\partial f}{\partial x} + y \frac{\partial g}{\partial x} & v \frac{\partial f}{\partial y} + y \frac{\partial g}{\partial y} + g(x, y) - (a + \rho + pz) & v \frac{\partial f}{\partial v} + f(x, y, v) \\ 0 & k & -u \end{pmatrix} \quad (34)$$

and its the second additive compound matrix is

$$J^{[2]} = \begin{pmatrix} j_{11} + j_{22} & j_{23} & -j_{13} \\ j_{32} & j_{11} + j_{33} & j_{12} \\ -j_{31} & j_{21} & j_{22} + j_{33} \end{pmatrix}, \quad (35)$$

where  $j_{kl}$  is the  $(k, l)$ the entry of the matrix  $J$ .

In this case, we choose  $P = \text{diag}(1, \frac{y}{v}, \frac{y}{v})$ . we get

$$P_f P^{-1} = \text{diag}(0, \frac{\dot{y}}{y} - \frac{\dot{v}}{v}, \frac{\dot{y}}{y} - \frac{\dot{v}}{v}),$$

where  $P_f$  is obtained by replacing each entry  $p_{ij}$  of  $P$  by its derivative in the direction of solution of (33). Also, we have

$$B = P_f P^{-1} + P J^{[2]} P^{-1} = \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix}, \quad (36)$$

where

$$\begin{aligned} B_{11} &= -(a + d + \rho + pz) - \frac{\partial f}{\partial x}v - \frac{\partial g}{\partial x}y + \frac{\partial f}{\partial y}v + \frac{\partial g}{\partial y}y + g(x, y), \\ B_{12} &= \left( \frac{v}{y} \left( \frac{\partial f}{\partial v}v + f \right) \quad \frac{v}{y} \left( \frac{\partial f}{\partial v}v + f \right) \right), \\ B_{21} &= \begin{pmatrix} \frac{ky}{v} \\ 0 \end{pmatrix}, \\ B_{22} &= \begin{pmatrix} \frac{\dot{y}}{y} - \frac{\dot{v}}{v} - u - d - \frac{\partial f}{\partial x}v - \frac{\partial g}{\partial x}y & \rho - \frac{\partial f}{\partial y}v - g(x, y) - \frac{\partial g}{\partial y}y, \\ \frac{\partial f}{\partial x}v + \frac{\partial g}{\partial x}y & \frac{\dot{y}}{y} - \frac{\dot{v}}{v} - a - \rho - pz - u + \frac{\partial f}{\partial y}v + g(x, y) + \frac{\partial g}{\partial y}y \end{pmatrix}, \end{aligned}$$

We choose a norm in  $\mathbb{R}^3$  as follow  $|\omega_1, \omega_2, \omega_3| = \max\{|\omega_1|, |\omega_2| + |\omega_3|\}$  for  $(\omega_1, \omega_2, \omega_3) \in \mathbb{R}^3$ . Thus the Lozinskii measure  $\mu$  with respect to this norm  $|\cdot|$  can be estimated as follows (see[10]):

$$\mu(B) \leq \sup\{g_1, g_2\}, \quad (37)$$

where  $g_1 = \mu_1(B_{11}) + |B_{12}|$  and  $g_2 = |B_{21}| + \mu_1(B_{22})$ .

Here  $\mu_1$  represents the Lozinskii measure with respect to  $l_1$  vector norm,  $|B_{12}|$  and  $|B_{21}|$  are matrix norms with respect to  $l_1$  norm. Furthermore, we have:

$$\begin{aligned} \mu_1(B_{11}) &= -(a + d + \rho + pz) - \frac{\partial f}{\partial x}v - \frac{\partial g}{\partial x}y + \frac{\partial f}{\partial y}v + \frac{\partial g}{\partial y}y + g(x, y), \\ |B_{12}| &= \frac{v}{y} \left( v \frac{\partial f}{\partial v} + f(x, y, v) \right) \\ &= \frac{\dot{y}}{y} + a + \rho + pz + \frac{v^2}{y} \frac{\partial f}{\partial v} - g(x, y), \\ |B_{21}| &= k \frac{y}{v} = \frac{\dot{v}}{v} + u, \\ \mu_1(B_{22}) &= \max \left\{ \frac{\dot{y}}{y} - \frac{\dot{v}}{v} - u - d, \frac{\dot{y}}{y} - \frac{\dot{v}}{v} - u - a - pz \right\} \\ &\leq \frac{\dot{y}}{y} - \frac{\dot{v}}{v} - u - \delta \end{aligned}$$

Consequently, we get

$$\begin{aligned} g_1 &= \frac{\dot{y}}{y} - d + \frac{v^2}{y} \frac{\partial f}{\partial v} - \frac{\partial f}{\partial x}v - \frac{\partial g}{\partial x}y + \frac{\partial f}{\partial y}v + \frac{\partial g}{\partial y}y \\ &\leq \frac{\dot{y}}{y} - \delta \end{aligned}$$

and

$$g_2 \leq \frac{\dot{y}}{y} - \delta$$

Therefore,

$$\mu(B) \leq \frac{\dot{y}}{y} - \delta$$

From Lemma 4.3, the system (1) is uniformly persistent for  $R_0 > 1$ . Then there exists a compact absorbing set  $K \subset \Omega$ [11]. Along each solution  $(x(t), y(t), v(t))$  of (1) with  $X_0 = (x(0), y(0), v(0)) \in K$ , we have

$$\frac{1}{t} \int_0^t \mu(B) ds \leq \frac{1}{t} \ln \left( \frac{y(t)}{y(0)} \right) - \delta,$$

which implies that

$$\bar{q}_2 = \limsup_{t \rightarrow \infty} \sup_{X_0 \in K} \frac{1}{t} \int_0^t \mu(B) ds \leq \frac{-\delta}{2} < 0.$$

Hence, the positive equilibrium  $(x^*, y^*, v^*)$  of the subsystem (33) is globally asymptotically stable. We consider the last equation of system(1)

$$\dot{z} = s + \frac{cyz}{\omega + y} - bz \quad (38)$$

and its limit system is

$$\dot{z} = s + \frac{cy^*z}{\omega + y^*} - bz \quad (39)$$

By  $b - \frac{cy^*}{\omega + y^*} = \frac{s}{z^*}$ , we obtain

$$\dot{z} = b \left( 1 - \frac{z}{z^*} \right) \quad (40)$$

Hence,

$$\lim_{t \rightarrow \infty} z(t) = z^* \quad (41)$$

Therefore, the chronic infection equilibrium  $E^*$  is globally asymptotically stable if  $R_0 > 1$ . ■

## 5 Discussion and conclusions

In this work, we have proposed an HBV infection model that takes into the account cell-to-cell transmission and CTL immune response. In the proposed model, the infection processes for the two modes of transmission are modeled by two general functions. We first proved the existence, positivity, and the boundedness of solutions of the problem which ensures that our model is well-posed. Under some assumptions about the general incidence function, the global dynamics of the model are fully characterized by a threshold parameter called the basic reproductive number  $R_0$ . More precisely, the infection-free equilibrium  $E_f$  is globally asymptotically stable if  $R_0 \leq 1$  which biologically means that the virus is cleared and the infection d-out. When  $R_0 > 1$ ,  $E_f$  become unstable and the chronic equilibrium  $E^*$  is globally asymptotically stable. Which means that the liver is hosting the virus. From the above analytical result, we deduce a strategy to control the HBV infection. This strategy is based on the reduction of  $R_0$  and makes its valeur lower or more equal 1. From explicit expression of  $R_0$  given in (15), the values of  $R_0$  can be reduced by increasing the export of the thymus. Therefore, we conclude that cellular humanity mediated by CTL cells plays an important role in the clearance of HBV from the liver.

## References

- [1] WHO, Hepatitis B, Fact sheet, April 2017, Available at: <http://www.who.int/mediacentre/factsheets/fs204/en/>.
- [2] W. Mothes, N. M Sherer, J. Jin, P. Zhong, Virus cell-to-cell transmission, *Journal of Virology* 84 (2010) 8360–8368.
- [3] P. Zhong, L.M. Agosto, J.B. Munro and W. Mothes, Cell-to-cell transmission of viruses, *Current Opinion in Virology* 3 (2013) 44–50.
- [4] G. Bocharov, B. Ludewig, A. Bertolotti et al., Underwhelming the immune response: effect of slow virus growth on CD 8<sup>+</sup> T Lymphocytes responses, *Journal of Virology* 78 (5) (2004) 2247–2254.
- [5] K. Hattaf, N. Yousfi, A generalized virus dynamics model with cell-to-cell transmission and cure rate, *Advances in Difference Equations* (2016) DOI 10.1186/s13662-016-0906-3.
- [6] K. Hattaf, N. Yousfi, A numerical method for a delayed viral infection model with general incidence rate, *J. King Saud Univ. Sci.* 28 (4) (2016) 368–374.
- [7] X.-Y. Wang, K. Hattaf, H.-F. Huo, H. Xiang, Stability analysis of a delayed social epidemics model with general contact rate and its optimal control, *Journal of Industrial and Management Optimization* 12 (4) (2016) 1267–1285.
- [8] Freedman, H, Ruan, S, Tang, M: Uniform persistence and flows near a closed positively invariant set. *J. Dyn. Differ. Equ.* 6, 583-600 (1994)
- [9] Li, MY, Muldowney, JS: A geometric approach to the global-stability problems. *SIAM J. Math. Anal.* 27, 1070-1083 (1996)
- [10] Martin, RH Jr.: Logarithmic norms and projections applied to linear differential systems. *J. Math. Anal. Appl.* 45, 432-454 (1974)
- [11] Butler, G, Waltman, P: Persistence in dynamical systems. *Proc. Am. Math. Soc.* 98, 425-430 (1986)