



**HAL**  
open science

## On the lower part of the lattice of partial clones

Miguel Couceiro, Lucien Haddad, Karsten Schölzel

► **To cite this version:**

Miguel Couceiro, Lucien Haddad, Karsten Schölzel. On the lower part of the lattice of partial clones. Journal of Multiple-Valued Logic and Soft Computing, 2019, 33 (3), pp.177-196. hal-01826870v4

**HAL Id: hal-01826870**

**<https://inria.hal.science/hal-01826870v4>**

Submitted on 31 Oct 2018

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# ON THE LOWER PART OF THE LATTICE OF PARTIAL CLONES

MIGUEL COUCEIRO, LUCIEN HADDAD, AND KARSTEN SCHÖLZEL

ABSTRACT. Let  $\mathbf{k}$  be a  $k$ -element set. We show that the lattice of all strong partial clones on  $\mathbf{k}$  has no minimal elements. Moreover, we show that if  $C$  is a strong partial clone, then the family of all partial subclones of  $C$  is of continuum cardinality. Finally we show that every non-trivial strong partial clone contains a family of continuum cardinality of strong partial subclones.

To the memory of Professor Ivan Stojmenovic

## 1. PRELIMINARIES

In this section we recall basic concepts and terminology needed throughout the paper. For further background, see [2, 5, 6, 7].

Let  $A$  be a finite set with  $|A| \geq 2$ . If  $|A| = k \geq 2$ , then we assume that  $A = \mathbf{k} := \{0, \dots, k-1\}$ . For a positive integer  $n$ , an  $n$ -ary *partial function* on  $\mathbf{k}$  is a map

$$f : \text{dom}(f) \rightarrow \mathbf{k}$$

where  $\text{dom}(f)$  is a subset of  $\mathbf{k}^n$  called the *domain* of  $f$ . We denote by  $\text{Par}^{(n)}(\mathbf{k})$  the set of all  $n$ -ary partial functions on  $\mathbf{k}$  and we set

$$\text{Par}(\mathbf{k}) := \bigcup_{n \geq 1} \text{Par}^{(n)}(\mathbf{k}).$$

For positive integers  $n, m \geq 1$ , and  $f \in \text{Par}^{(n)}(\mathbf{k})$  and  $g_1, \dots, g_n \in \text{Par}^{(m)}(\mathbf{k})$ , the *composition of  $f$  with  $g_1, \dots, g_n$*  is the partial function  $f[g_1, \dots, g_n] \in \text{Par}^{(m)}(\mathbf{k})$  whose domain  $\text{dom}(f[g_1, \dots, g_n])$  is the set of all

$$\vec{a} \in \bigcap_{i=1}^m \text{dom}(g_i) \quad \text{such that} \quad (g_1(\vec{a}), \dots, g_n(\vec{a})) \in \text{dom}(f),$$

and whose values are given by

$$f[g_1, \dots, g_n](\vec{a}) := f(g_1(\vec{a}), \dots, g_n(\vec{a}))$$

for all  $\vec{a} \in \text{dom}(f[g_1, \dots, g_n])$ . For every positive integer  $n$  and each  $1 \leq i \leq n$ , we denote by  $e_i^n$  the  $n$ -ary  $i$ -th projection function (or, simply, *projection*) defined by

$$e_i^n(a_1, \dots, a_n) = a_i$$

for all  $(a_1, \dots, a_n) \in \mathbf{k}^n$ . Furthermore, we use the notation

$$J_{\mathbf{k}} := \{e_i^n : 1 \leq i \leq n, n \in \mathbb{N}\}$$

for the set of all projections.

**Definition 1.** A partial clone on  $\mathbf{k}$  is a subset of  $\text{Par}(\mathbf{k})$  closed under composition and containing the set of all projection functions  $J_{\mathbf{k}}$ .

**Remark 2.** Partial clones can be equivalently defined in the formalism of Mal'cev; see, e.g., Chapter 20 of [7].

The set of partial clones on  $\mathbf{k}$  forms a lattice  $\mathcal{L}_{P_{\mathbf{k}}}$  under inclusion, in which the infimum is the set-theoretical intersection. Thus the intersection of arbitrarily many partial clones on  $\mathbf{k}$  is itself a partial clone on  $\mathbf{k}$ . Clearly, the set  $J_{\mathbf{k}}$  of all projections on  $\mathbf{k}$  is the smallest element in the lattice  $\mathcal{L}_{P_{\mathbf{k}}}$  of partial clones.

For  $f, g \in \text{Par}^{(n)}(\mathbf{k})$ , the partial function  $g$  is said to be a *subfunction* of  $f$ , fact that we denote by  $g \leq f$ , if

$$g = f|_{\text{dom}(g)},$$

i.e., if  $\text{dom}(g) \subseteq \text{dom}(f)$  and  $g(\vec{a}) = f(\vec{a})$  for all  $\vec{a} \in \text{dom}(g)$ .

**Definition 3.** A partial clone  $C$  on  $\mathbf{k}$  is said to be *strong* if it is closed under taking subfunctions, i.e., if the condition

$$g \leq f \implies g \in C$$

holds for every  $f \in C$  and  $g \in \text{Par}(\mathbf{k})$ . For  $F \subseteq \text{Par}(\mathbf{k})$ , we denote its strong closure by  $\text{Str}(F)$ , i.e.,

$$\text{Str}(F) := \{g \in \text{Par}(\mathbf{k}) : g \leq f \text{ for some } f \in F\}.$$

In what follows, by an  $n$ -ary *partial projection* we mean a subfunction of a projection function whose domain is a proper subset of  $\mathbf{k}^n$ .

**Example 4.** It is easy to see that the set

$$\{f \in \text{Par}(\mathbf{k}) : (0, \dots, 0) \in \text{dom}(f) \implies f(0, \dots, 0) = 0\}$$

is a strong partial clone on  $\mathbf{k}$ .

Strong partial clones have been widely studied in the literature, see, e.g., [2, 5, 6] and [7] for a long list of references. The recent paper [8] is dedicated to the study of strong partial clones of Boolean functions. It is well known that a partial clone  $C$  is strong if and only if  $\text{Str}(J_{\mathbf{k}}) \subseteq C$  (see, e.g., [2], Lemma 2.11). Moreover, it is easy to see that the intersection of an arbitrary family of strong partial clones on  $\mathbf{k}$  is also a strong partial clone. In fact, the set of all strong partial clones, ordered by inclusion, also forms a lattice  $\mathcal{L}_{\text{Str}(P_{\mathbf{k}})}$  whose smallest element is  $\text{Str}(J_{\mathbf{k}})$ .

For  $F \subseteq \text{Par}(\mathbf{k})$ , let  $\text{pclone}(F)$  denote the *strong partial clone generated by  $F$* , i.e., the intersection of all strong partial clones on  $\mathbf{k}$  containing  $F$ . As it is well-known, strong partial clones coincide exactly with the sets of polymorphisms of relations on  $\mathbf{k}$ , that we now briefly describe. For  $m \geq 1$ , let  $\rho$  be an  $m$ -ary relation on  $\mathbf{k}$  (i.e., a subset of  $\mathbf{k}^m$ ). Denote by  $\mathcal{M}(\rho, \text{dom}(f))$  the set of all  $m \times n$  matrices  $M$  on  $\mathbf{k}$  whose columns  $M_{*j} \in \rho$ , for  $j = 1, \dots, n$  and whose rows  $M_{i*} \in \text{dom}(f)$  for  $i = 1, \dots, m$ . A partial function  $f \in \text{Par}^{(n)}(\mathbf{k})$  is said to *preserve*  $\rho \subseteq \mathbf{k}^m$  if  $(f(M_{1*}), \dots, f(M_{m*})) \in \rho$  for all  $M \in \mathcal{M}(\rho, \text{dom}(f))$ . Set

$$\text{pPol } \rho := \{f \in \text{Par}(\mathbf{k}) : f \text{ preserves } \rho\}.$$

It is well known that  $\text{pPol } \rho$  is a strong partial clone on  $\mathbf{k}$  called the partial clone *determined by  $\rho$* . Note that  $f \in \text{pPol } \rho$  if there is no matrix  $M$  with columns in  $\rho$  and rows in  $\text{dom}(f)$ .

**Main achievements:** The main contributions of this paper are 3-fold. First, we show that

- (I) *minimal strong partial clones* (i.e., atoms of  $\mathcal{L}_{\text{Str}(P_{\mathbf{k}})}$ ) do not exist, and
- (II) the family  $\{D : D \text{ is a partial clone and } D \subseteq \text{Str}(J_{\mathbf{k}})\}$  is of continuum cardinality.

Since every strong partial clone contains  $\text{Str}(J_{\mathbf{k}})$ , it follows that any strong partial clone contains a family of partial subclones of continuum cardinality.

However, partial clones strictly included in  $\text{Str}(J_{\mathbf{k}})$  are not strong. Hence, in the last section of the paper, we show that

- (III) every strong partial clone  $C \neq \text{Str}(J_{\mathbf{k}})$  contains a family of strong partial subclones of continuum cardinality.

Some results established in this paper have been presented at the 47<sup>th</sup> IEEE *International Symposium on Multiple-Valued Logic* (Novi Sad, Serbia) and have already appeared in the conference paper [4].

## 2. MINIMAL STRONG PARTIAL CLONES

A *minimal partial clone* on  $\mathbf{k}$  is an atom of the lattice  $\mathcal{L}_{P_{\mathbf{k}}}$ . In other words, a partial clone  $C$  is minimal if there is no partial clone  $C_0$  such that  $J_{\mathbf{k}} \neq C_0 \subset C \subset \text{Par}(\mathbf{k})$ . Minimal partial clones consisting of partial functions have been completely described in [1], none of which is strong. The natural question is then to ask whether *strong minimal partial clones* exist, i.e., whether atoms of  $\mathcal{L}_{\text{Str}(P_{\mathbf{k}})}$  exist.

In this section we show that such partial clones do not exist. This result is not surprising, but to our knowledge it has never been published. The idea behind the proof we present here comes from a discussion between the second author and Ferdinand Börner just after the paper [1] has appeared. We will show that, given a partial function  $g$  that is not a partial projection function, we can construct a partial function  $h \in \text{pclone}(g)$ , with  $h \notin \text{Str}(J_{\mathbf{k}})$  and such that  $g \notin \text{pclone}(h)$ .

Let  $g \notin \text{Str}(J_{\mathbf{k}})$  be an  $n$ -ary partial function and  $C := \text{pclone}(g)$ . We distinguish two cases:

**Case 1.** For every  $(a_1, \dots, a_n) \in \text{dom}(g)$ ,  $g(a_1, \dots, a_n) \in \{a_1, \dots, a_n\}$ .

Let  $f$  be a restriction of  $g$  so that any further restriction of the domain of  $f$  results in a partial projection. Let  $m := |\text{dom}(f)|$  and let  $\alpha^1, \dots, \alpha^m \in \text{dom}(f) \subseteq \mathbf{k}^n$  with  $\alpha^i := (a_1^i, \dots, a_n^i)$  and such that

$$(1) \quad f \begin{pmatrix} a_1^1 & \dots & a_n^1 \\ a_1^2 & \dots & a_n^2 \\ \vdots & \vdots & \vdots \\ a_1^m & \dots & a_n^m \end{pmatrix} = \begin{pmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{pmatrix}.$$

Since  $f$  is not a partial projection, the right tuple  $(b_1, \dots, b_m)^T$  is not a column of the matrix on the left. Let  $M$  be the  $m \times n$  matrix in (1) and let  $\rho_M$  be the  $m$ -ary relation on  $\mathbf{k}$  consisting of all columns of  $M$ . Clearly  $f \notin \text{pPol}\rho_M$ , and thus  $g \notin \text{pPol}\rho_M$ .

Let  $\bar{f}$  be the  $(n+2)$ -ary partial function given by

$$\bar{f}(x_1, \dots, x_{n+2}) := f(x_1, \dots, x_n)$$

for all  $(x_1, \dots, x_{n+2}) \in \text{dom}(f) \times \mathbf{k}^2$ . Note that

$$\bar{f} \in \text{pclone}(f) \subseteq \text{pclone}(g).$$

Now take a  $c \in \mathbf{k}, c \neq b_1$ , and consider the subfunction  $h$  of  $\bar{f}$  whose domain is the set of all rows of the following matrix:

$$\begin{pmatrix} a_1^1 & \dots & a_n^1 & c & b_1 \\ a_1^1 & \dots & a_n^1 & b_1 & c \\ a_1^2 & \dots & a_n^2 & b_2 & b_2 \\ \vdots & & \vdots & \vdots & \vdots \\ a_1^m & \dots & a_n^m & b_m & b_m \end{pmatrix}$$

where  $b_1, b_2, \dots, b_m$  are the values of the partial function  $f$ . We therefore have

$$h \begin{pmatrix} a_1^1 & \dots & a_n^1 & c & b_1 \\ a_1^1 & \dots & a_n^1 & b_1 & c \\ a_1^2 & \dots & a_n^2 & b_2 & b_2 \\ \vdots & & \vdots & \vdots & \vdots \\ a_1^m & \dots & a_n^m & b_m & b_m \end{pmatrix} = \begin{pmatrix} b_1 \\ b_1 \\ b_2 \\ \vdots \\ b_m \end{pmatrix}.$$

Note that  $|\text{dom}(h)| = m + 1$ .

We show that any restriction of  $h$  to a set with at most  $m$  tuples is a partial projection. Let  $h'$  be a restriction of  $h$  with  $|\text{dom}(h')| \leq m$ . If  $(a_1^1, \dots, a_n^1, c, b_1) \notin \text{dom}(h')$ , then  $h' \leq e_{n+1}^{n+2}$ , i.e.,  $h'$  is a partial projection on its  $(n+1)$ -st coordinate. Similarly, if  $(a_1^1, \dots, a_n^1, b_1, c) \notin \text{dom}(h')$ , then  $h' \leq e_{n+2}^{n+2}$ . On the other hand if  $(a_1^i, \dots, a_n^i, b_i, b_i) \notin \text{dom}(h')$  for some  $i = 2, \dots, m$ , then  $h'$  is a partial projection function on the same coordinate as the subfunction of  $f$  whose domain is contained in the set of rows of  $M$  except the  $i$ -th row.

Now we show that  $h \in \text{pPol}\rho_M$ . Let  $N$  be an  $m \times (n+2)$  matrix with all columns in  $\rho_M$  and all rows in  $\text{dom}(h)$ . Then as  $N$  has  $m$  rows, the restriction of  $h$  on the rows of  $N$  is a partial projection function and so it preserves any relation, including the relation  $\rho_M$ . Thus  $h \in \text{pPol}\rho_M$ , i.e.,  $\text{pclone}(h) \subseteq \text{pPol}\rho_M$ . Since  $g \notin \text{pPol}\rho_M$  we conclude that  $g \notin \text{pclone}(h)$  and thus  $\text{pclone}(h) \subsetneq C$ .

**Case 2.** There is an  $n$ -tuple  $(a_1, \dots, a_n) \in \text{dom}(g)$  such that  $g(a_1, \dots, a_n) \notin \{a_1, \dots, a_n\}$ .

Let  $f$  be the restriction of  $g$  to the domain  $\{(a_1, \dots, a_n)\}$  and let  $f(a_1, \dots, a_n) = b$  with  $b \notin \{a_1, \dots, a_n\}$ . Choose  $c \neq b$  and define the  $(n+2)$ -ary partial function  $h$  by

$$\text{dom}(h) = \{(a_1, \dots, a_n, c, b), (a_1, \dots, a_n, b, c)\},$$

and  $h(a_1, \dots, a_n, c, b) = h(a_1, \dots, a_n, b, c) = b$  and proceed as in **Case 1**. Thus we have shown the following result.

**Theorem 5.** *Let  $C \neq \text{Str}(J_{\mathbf{k}})$  be a strong partial clone. Then there is a strong partial clone  $C'$  such that  $\text{Str}(J_{\mathbf{k}}) \subsetneq C' \subsetneq C$ . Hence there are no strong minimal partial clones on  $\mathbf{k}$ .*

### 3. CLONES OF PARTIAL PROJECTIONS

In this section we consider partial clones consisting only of projections and partial projection functions. For  $F \subseteq \text{Par}(\mathbf{k})$ , let  $\langle F \rangle$  denote the *partial clone generated by  $F$* , i.e., the intersection of all partial clones

on  $\mathbf{k}$  containing  $F$ . Notice that  $\langle F \rangle$  is not necessarily a strong partial clone.

Let  $R_n$  denote the set of all  $n$ -tuples on  $\mathbf{k}$  of weight 1, i.e.,

$$R_n := \{(x_1, \dots, x_n) \in \mathbf{k}^n : \sum_{i=1}^n x_i = 1\}.$$

Furthermore, let  $p_n$  be the  $n$ -ary partial projection given by

$$p_n(x_1, \dots, x_n) := \begin{cases} x_1 & \text{if } (x_1, \dots, x_n) \in R_n \\ \text{undefined} & \text{otherwise.} \end{cases}$$

Let  $\mathbf{P} := \{p_n : n \geq 2\}$ . We will show that for  $t \geq 2$ ,  $p_t \notin \langle \mathbf{P} \setminus \{p_t\} \rangle$ . Recall Lemma 2.10 in [2].

**Lemma 6.** *Let  $F \subseteq \text{Par}(\mathbf{k})$  and set  $\mathcal{D}_0 := F \cup J_{\mathbf{k}}$ . For  $\ell \geq 0$ , define*

$$\begin{aligned} \mathcal{D}_{\ell+1} &:= \mathcal{D}_\ell \cup \{f[g_1, \dots, g_n] \mid f \in \mathcal{D}_0^{(n)} \text{ and} \\ &\quad g_1, \dots, g_n \in \mathcal{D}_\ell^{(m)}, \text{ for some } n, m \geq 1\}. \end{aligned}$$

Then  $\langle F \rangle = \bigcup_{\ell \geq 0} \mathcal{D}_\ell$ . ■

Fix  $t \geq 2$  and  $F := \mathbf{P} \setminus \{p_t\}$ . Observe that, since  $F \subseteq \text{Str}(J_{\mathbf{k}})$ ,  $\langle F \rangle \subseteq \text{Str}(J_{\mathbf{k}})$ , i.e.,  $\langle F \rangle$  consists only of projections and partial projections.

**Lemma 7.** *Let  $\mathcal{D}_0, \mathcal{D}_1, \dots$  be as in Lemma 6. Then, for each  $\ell \geq 0$ ,  $\mathcal{D}_\ell$  does not contain any partial projection  $p$  of arity  $t$  with  $R_t \subseteq \text{dom}(p)$ .*

**Proof.** We prove Lemma 7 by induction on  $\ell \geq 0$ . For  $\ell = 0$ , we have  $\mathcal{D}_0 = (\mathbf{P} \setminus \{p_t\}) \cup J_{\mathbf{k}}$ , which does not contain any partial projection of arity  $t$ .

Now let  $\ell \geq 0$  and suppose that  $\mathcal{D}_\ell$  does not contain any partial projection  $p$  of arity  $t$  with  $R_t \subseteq \text{dom}(p)$ . We show that the same holds for  $\mathcal{D}_{\ell+1}$ . By the induction hypothesis and the definition of  $\mathcal{D}_{\ell+1}$ , we need to show that if  $f \in \mathcal{D}_0^{(n)}$ ,  $n \geq 1$ ,  $g_1, \dots, g_n \in \mathcal{D}_\ell^{(t)}$  and  $p = f[g_1, \dots, g_n]$ , then either  $p$  is a projection or a partial projection with  $R_t \not\subseteq \text{dom}(p)$ . Since  $g_1, \dots, g_n \in \mathcal{D}_\ell$ , it follows from the induction hypothesis that for every  $i = 1, \dots, n$ , either  $g_i$  is a total projection or  $g_i$  is a partial projection with  $R_t \not\subseteq \text{dom}(g_i)$ .



Suppose first that  $g_i$  is a partial projection for some  $i = 1, \dots, n$ . Then  $R_i \not\subseteq \text{dom}(g_i)$  and thus there is a  $\vec{v} \in R_i \setminus \text{dom}(g_i)$ . But then, by the definition of the composition of partial functions  $p := f[g_1, \dots, g_n]$  is not defined on the tuple  $\vec{v}$ , and thus  $\vec{v} \notin \text{dom}(p)$ . This shows that  $R_i \not\subseteq \text{dom}(p)$ .

Suppose now that each  $g_1, \dots, g_n$  is a total projection of arity  $t$ . If  $f \in J_{\mathbf{k}}$  is a projection, then  $p = f[g_1, \dots, g_n]$  is a total projection and we are done.

So suppose that  $f$  is a partial projection. Since  $f \in \mathcal{D}_0$ , we have  $f = p_n$  with  $n \neq t$ .

**Case 1.** Let  $n < t$ . Then there is a  $j \in \{1, \dots, t\}$  such  $e_j^t \notin \{g_1, \dots, g_n\}$ . Therefore

$$(2) \quad \vec{v}_j := (0, \dots, 0, \underbrace{1}_{j\text{-th position}}, 0, \dots, 0) \notin \text{dom}(p)$$

since  $g_1(\vec{v}_j) = \dots = g_n(\vec{v}_j) = 0$  and  $(0, \dots, 0) \notin \text{dom}(f)$ .

**Case 2.** Let  $t < n$ . Then there are  $j, k \in \{1, \dots, n\}$ ,  $j \neq k$ , such that  $g_j = g_k$ . Without loss of generality, suppose that  $g_j = g_k = e_j^t$ . In this case, the  $t$ -tuple  $\vec{v}_j$  defined in (2) does not belong to  $\text{dom}(p)$ , since the  $n$ -tuple  $(g_1(\vec{v}_j), \dots, g_n(\vec{v}_j))$  contains at least two entries equal to 1, and thus  $(g_1(\vec{v}_j), \dots, g_n(\vec{v}_j)) \notin \text{dom}(f)$ . This completes the proof of Lemma 7.  $\blacksquare$

From Lemma 7, it follows that  $p_t \notin \langle \mathbf{P} \setminus \{p_t\} \rangle$  for every  $t \geq 2$ , and thus we have the following corollary.

**Corollary 8.** *Let  $X, Y \subseteq \{2, 3, \dots\}$ . If  $X \neq Y$ , then*

$$\langle \{p_i : i \in X\} \rangle \neq \langle \{p_j : j \in Y\} \rangle.$$

Let  $\mathcal{F}_{J_{\mathbf{k}}}$  be the set of all partial clones contained in  $\text{Str}(J_{\mathbf{k}})$ .

**Corollary 9.** *The set  $\mathcal{F}_{J_{\mathbf{k}}}$  is of continuum cardinality on  $\mathbf{k}$ .*

Now as every strong partial clone on  $\mathbf{k}$  contains the set  $\text{Str}(J_{\mathbf{k}})$  we immediately have the following result.

**Theorem 10.** *Let  $C$  be a strong partial clone on  $\mathbf{k}$ . Then the interval of partial clones  $[J_{\mathbf{k}}, C]$  is of continuum cardinality.*

Note that none of the partial clones contained in  $\text{Str}(J_{\mathbf{k}})$  is strong. Therefore, Theorem 10 does not provide any insight into the interval of strong partial clones  $[\text{Str}(J_{\mathbf{k}}), C]$ . Hence, for a strong partial clone  $C$ , it is natural to ask what is the size of the interval

$$\text{Clone}(C) := \{D : D \text{ is a strong partial subclone of } C\}.$$

This is discussed in the next section.

#### 4. STRONG PARTIAL CLONES GENERATED BY A SINGLE PARTIAL FUNCTION

Let  $f$  be an  $n$ -ary partial function not in  $\text{Str}(J_{\mathbf{k}})$ . Then there are  $m$  tuples  $\alpha^1, \dots, \alpha^m \in \mathbf{k}^n$ , with  $\alpha^i := (a_1^i, \dots, a_n^i)$ , such that

$$f \begin{pmatrix} a_1^1 & \dots & a_n^1 \\ \vdots & \vdots & \vdots \\ a_1^i & \dots & a_n^i \\ \vdots & \vdots & \vdots \\ a_1^m & \dots & a_n^m \end{pmatrix} = \begin{pmatrix} b_1 \\ \vdots \\ b_i \\ \vdots \\ b_m \end{pmatrix}$$

and such that  $(b_1, \dots, b_m)^T$  is not a column of the matrix on the left hand side of the equation. Denote this  $m \times n$  matrix by  $A$ . In the sequel we assume that the  $n$ -ary partial function  $f$ , the matrix  $A$  and the number  $m$  of rows of  $A$  are fixed.

In this section we show that the strong partial clone  $\text{pclone}(f)$  contains a family of strong partial subclones of continuum cardinality.

We will distinguish two cases:

**Case 1.** The following condition is satisfied by the matrix  $A$ :

- (\*) there is  $\alpha \in \mathbf{k}$  such that the  $m$ -tuple  $(\alpha, \dots, \alpha)^T$  is not a column of  $A$ .

**Case 2.** The partial function  $f$  is not constant.

We will explain in next subsection why those two cases cover all possibilities.

4.1. **Matrix  $A$  satisfies condition (\*)**. For notational ease, we assume in this section that  $\alpha = 0$ , i.e., we assume that  $(0, \dots, 0)^T$  is not a column of  $A$ . Denote by  $I_r$  the *identity matrix* of size  $r \geq 2$  on  $\mathbf{k}$ , i.e.,  $I_r$  is the  $r \times r$  matrix with symbol 1 on its diagonal and with 0 elsewhere. Furthermore, consider the following infinite set of positive integers

$$E := \{2m^r + \sum_{1 \leq i \leq r} m^{i-1} : r \geq 0\}.$$

**Remark 11.** Notice that, for each  $r \geq 1$ , we have

$$2m^r + m^{r-1} + \dots + m + 1 = m(2m^{r-1} + \dots + m + 1) + 1.$$

Consequently, if  $t > \ell$  with  $t, \ell \in E$ , then  $t \geq m\ell + 1$ .

We will construct an infinite set of partial functions

$$\{f_\ell : \ell \in E\}$$

each of which is in  $\text{pclone}(f)$ , and with the property that for every  $\ell \in E$ ,  $f_\ell \notin \text{pclone}(\{f_k : k \neq \ell, k \in E\})$ .

For  $\ell \in E$ , let  $M_\ell$  be the matrix whose first  $n$  columns are

$$(a_i^1, \dots, a_i^1, a_i^2, \dots, a_i^2, \dots, a_i^m, \dots, a_i^m)^T, \quad i = 1, \dots, n,$$

and the last  $m \times \ell$  columns are the columns of the identity matrix  $I_{m \times \ell}$ . Thus,  $M_\ell$  is a matrix with  $m \times \ell$  rows and  $n + m \times \ell$  columns. Note also that each of the tuples  $(a_1^i, \dots, a_n^i)$  is repeated  $\ell$  times in  $M_\ell$ .

$$M_\ell = \left( \begin{array}{ccc|ccc} a_1^1 & \dots & a_n^1 & 1 & 0 & \dots & 0 \\ \vdots & & \vdots & \vdots & \vdots & & \vdots \\ a_1^1 & \dots & a_n^1 & 0 & \dots & \dots & 0 \\ a_1^2 & \dots & a_n^2 & 0 & \dots & \dots & 0 \\ \vdots & & \vdots & \vdots & \vdots & & \vdots \\ a_1^2 & \dots & a_n^2 & 0 & \dots & \dots & 0 \\ \vdots & & \vdots & \vdots & \vdots & & \vdots \\ a_1^m & \dots & a_n^m & 0 & \dots & \dots & 0 \\ \vdots & & \vdots & \vdots & \vdots & & \vdots \\ a_1^m & \dots & a_n^m & 0 & \dots & \dots & 1 \end{array} \right)$$

Let  $\rho_\ell$  be the relation that consists of all columns of  $M_\ell$ . Note that  $\rho_\ell$  is a relation of arity  $m \times \ell$  over  $\mathbf{k}$ .

Now let  $f_\ell$  be the partial function of arity  $n + m \times \ell$  whose domain is the set of all rows of the matrix  $M_\ell$ , and that is given by  $f_\ell(\vec{v}) = f(a_1^i, \dots, a_n^i)$ , where  $(a_1^i, \dots, a_n^i)$  are the first  $n$  coordinates of the tuple  $\vec{v}$ .

**Example 12.** Let  $\mathbf{k} = \{0, 1, 2\}$ , and let  $f$  be the ternary partial function defined by

$$f \begin{pmatrix} 0 & 0 & 1 \\ 0 & 1 & 1 \\ 2 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}.$$

Here  $m = n = 3$  and for  $7 = 2\dot{3} + 1 \in E$ , we get

$$f_7 \left( \begin{array}{ccc|ccc} 0 & 0 & 1 & 1 & 0 & \dots & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\ 0 & 0 & 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\ 0 & 1 & 1 & 0 & 0 & \dots & 0 & 0 \\ 2 & 0 & 1 & 0 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \dots & \vdots & \vdots \\ 2 & 0 & 1 & 0 & 0 & \dots & 1 & 0 \\ 2 & 0 & 1 & 0 & 0 & \dots & 0 & 1 \end{array} \right) = \begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \\ 0 \\ 0 \\ \vdots \\ 0 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}.$$

$\underbrace{\hspace{15em}}_{I_{21}}$

Note that this matrix has 7 rows starting with  $(0, 0, 1)$ , 7 rows starting with  $(0, 1, 1)$ , and 7 rows starting with  $(2, 0, 1)$ .

**Lemma 13.** For each  $\ell \in E$ , we have that  $f_\ell \in \text{pclone}(f)$  and that  $f_\ell$  is not a partial projection.

**Proof.** It is easy to see that  $f_\ell \in \text{pclone}(f)$ . Since  $f$  is not a partial projection, none of the first  $n$  columns of  $M_\ell$  is equal to

$$(b_1, \dots, b_1, b_2, \dots, b_2, \dots, b_n, \dots, b_n)^T.$$

Moreover, each of the last  $m \times \ell$  columns of  $M_\ell$  consists of one 1 and  $m\ell - 1$  0's, and thus none of those columns can be equal to

$$(b_1, \dots, b_1, b_2, \dots, b_2, \dots, b_n, \dots, b_n)^T.$$

■

**Lemma 14.** *If  $\ell, t \in E$ , then  $f_\ell \in \text{pPol}\rho_t$  if and only if  $\ell \neq t$ .*

**Proof.** Clearly,  $f_\ell \notin \text{pPol}\rho_\ell$ . We show that if  $t \neq \ell$ , then there is no matrix whose rows are rows of the matrix  $M_\ell$  and whose columns belong to the relation  $\rho_t$ .

For the sake of contradiction, suppose that such a matrix  $B$  exists. Then  $B$  has  $m \times t$  rows and  $n + m \times \ell$  columns. Let  $B_{[i,*]}$ ,  $i = 1 \dots m \times t$ , be all the rows of  $B$  and  $B_{[*],j}$ ,  $j = 1, \dots, n + m \times \ell$ , be all its columns.

**Case 1.**  $t < \ell$ .

For  $i = 1, \dots, m \times t$ , the  $i$ -th row of  $B$  has the form

$$B_{[i,*]} = (a_1^s, \dots, a_n^s, x_1, \dots, x_{m \times \ell})$$

where  $s \in \{1, \dots, m\}$  and exactly one of  $x_1, \dots, x_{m \times \ell}$  is equal to 1 and all other are 0. Given that  $B$  consists of  $m \times t$  such rows and given that  $mt < m \times \ell$ , there is a  $j \in \{n + 1, \dots, n + m \times \ell\}$  such that the column  $B_{[*],j}$  consists of only 0's. Hence, it does not belong to  $\rho_t$ , which yields the desired contradiction.

**Case 2.**  $\ell < t$ .

As  $\ell, t \in E$ , we must have  $t \geq m \times \ell + 1$ , and

$$M_{\ell[i,*]} = (a_1^s, \dots, a_n^s, 0, \dots, 0, \underbrace{1}_{i\text{-th position}}, 0, \dots, 0)$$

where  $1 \leq s \leq m$ . Therefore, if for some  $i \in \{1, \dots, m \times \ell\}$ , the  $i$ -th row  $M_{\ell[i,*]}$  of  $M_\ell$  does not appear in  $B$ , then  $B_{[*],n+i}$  will contain only 0's, and it will not belong to the relation  $\rho_t$ . Thus every row of  $M_\ell$

must appear in the matrix  $B$ . Since  $\ell < t$ , some of the rows of  $M_\ell$  are repeated in  $B$ .

Now if  $M_{\ell[i,*]}$  appears repeated exactly  $r$  times in  $B$ ,  $2 \leq r \leq m \times t$ , then  $B_{[*],n+i}$  will contain  $r$  1's. Given that this column belongs to  $\rho_t$  and given that each row of  $A$  occurs  $t$  times in  $M_t$ , we conclude that  $r$  is a multiple of  $t$ .

For  $i = 0, 1, \dots, s$  let  $r_i$  be the number of rows of  $M_\ell$  that are repeated  $i \times t$  times in  $B$ , with  $r_0$  being the number of rows in  $M_\ell$  that appear only once in  $B$ . Then counting the rows of  $B$  we get the equality

$$r_0 + t \times r_1 + 2t \times r_2 + 3t \times r_3 + \cdots + st \times r_s = m \times t,$$

i.e.,  $r_0$  is a multiple of  $t$ . As  $m \times \ell$  is the total number of rows in  $M_\ell$ , and since at least one row of  $M_\ell$  is repeated in  $B$ , we have that  $r_0 < m \times \ell$ . Furthermore, as  $t \geq m \times \ell + 1$ , we conclude that  $r_0 < m \times \ell < t$  and thus  $r_0 = 0$ . Hence, every row of  $M_\ell$  appears  $i \times t$  times in  $B$  with  $i \geq 1$ . It thus follows that  $B$  has no column with one 1 and  $m \times t - 1$  0's, i.e., every column of  $B$  is one of the first  $n$  columns of  $M_t$ .

Let  $B_R$  be the block matrix consisting of the last  $m \times \ell$  columns to the right of the matrix  $B$ . Let us count the number of 0's in  $B_R$ . Since the relation  $\rho_t$  contains no zero tuple, every column of  $B_R$  contains at least  $t$  1's, i.e., every column in  $B_R$  contains at most  $mt - t$  0's. Given that there are  $m \times \ell$  columns in  $B_R$ , there are at most  $m \times \ell \times (m \times t - t)$  0's in  $B_R$ . Furthermore, every row of  $B$  is a row of  $M_\ell$ , and thus every row of  $B_R$  contains exactly one 1 and  $m \times \ell - 1$  0's.

Hence, there are  $m \times t \times (m \times \ell - 1)$  0's in  $B_R$ . Since

$$m \times t \times (m \times \ell - 1) > m \times \ell \times (m \times t - t),$$

we conclude that at least one column of  $B_R$  must contain only 0's, which again yields a contradiction.  $\blacksquare$

**Corollary 15.** *If  $X \subseteq E$  and  $\ell \notin X$ , then  $f_\ell \notin \text{pclone}(\{f_t : t \in X\})$ .*

**Proof.** Since  $\ell \notin X$ , by Lemma 14 we have that  $f_t \in \text{pPol}\rho_\ell$  for every  $t \in X$ , and thus  $\text{pclone}(\{f_t : t \in X\}) \subseteq \text{pPol}\rho_\ell$ . Since  $f_\ell \notin \text{pPol}\rho_\ell$ , we have  $f_\ell \notin \text{pclone}(\{f_t : t \in X\})$ .  $\blacksquare$

Now let  $C$  be a strong partial clone and suppose that  $C$  contains an  $n$ -ary partial function  $f \notin \text{Str}(J_{\mathbf{k}})$  such that  $|\text{dom}(f)| = m$ . Suppose also that  $f$  satisfies condition (\*). Reconsider the sets  $E$  and  $\{f_\ell : \ell \in E\}$  defined above. By Corollary 15, for every  $X, Y \subseteq E$  with  $X \neq Y$ , the two strong partial clones  $\text{pclone}(\{f_t : t \in X\})$  and  $\text{pclone}(\{f_t : t \in Y\})$  are distinct and both are contained in  $\text{pclone}(f) \subseteq C$ . Therefore, we have the following corollary.

**Corollary 16.** *Let  $C$  be a strong partial clone on  $\mathbf{k}$  and suppose that  $C$  contains a partial function  $f \notin \text{Str}(J_{\mathbf{k}})$  whose matrix  $A$  satisfies (\*). Then the set of strong partial clones contained in  $C$  is of continuum cardinality.*

**Example 17.** *Let  $C$  be a strong partial clone on  $\mathbf{k}$  and suppose that  $C$  contains a unary partial function  $g$  such that  $1 \in \text{dom}(g)$  and  $g(1) = 0$ . Let  $f := g|_{\{1\}}$ , i.e.,  $f$  is given by  $\text{dom}(f) = \{1\}$  and  $f(1) = 0$ . Since  $C$  is a strong partial clone, we have  $f \in C$ .*

*Consider the set  $E$  defined above with  $m = 1$ , i.e.,*

$$E = \{2, 3, \dots\}.$$

*For  $\ell \in E$ , let  $f_\ell$  be the  $(\ell + 1)$ -ary partial function whose domain is the set of rows of the matrix*

$$M_\ell = \left( \begin{array}{c|cccc} 1 & 1 & 0 & 0 & \dots & 0 \\ 1 & 0 & 1 & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & & \vdots \\ 1 & 0 & 0 & 0 & \dots & 1 \end{array} \right)$$

*and let  $f_\ell(\vec{v}) := 0$  for all  $\vec{v} \in \text{dom}(f_\ell)$ . Then  $f_\ell$  is a subfunction of  $f[e_1^{\ell+1}]$  and so  $f_\ell \in C$ . By Corollary 15 we have that for every  $X, Y \subseteq E$ ,  $X \neq Y$ ,*

$$\text{pclone}(\{f_t : t \in X\}) \neq \text{pclone}(\{f_t : t \in Y\}).$$

*Hence the strong partial clone  $C$  contains each member of the family of strong partial clones  $\{\text{pclone}(\{f_t : t \in X\}) : X \subseteq E\}$ , which is of continuum cardinality.*

**Remark 18.** The construction presented in the above section cannot be used to show that  $\text{pclone}(f)$  contains a family of strong partial clones of continuum cardinality if the matrix  $A$  contains all constant columns. Indeed in such a case the proof of Lemma 14 fails. The following example illustrates this fact.

**Example 19.** Let  $\mathbf{k} = \{0, 1, 2\}$  and let  $f$  be the 4-ary partial function given by

$$f \begin{pmatrix} 0 & 1 & 2 & 0 \\ 0 & 1 & 2 & 1 \\ 0 & 1 & 2 & 2 \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \\ 0 \end{pmatrix}.$$

Note that here  $m = 3$  and  $n = 4$ . Thus  $E = \{2, 7, 22, \dots\}$ . For  $\ell = 2$  we have

$$M_2 = \left( \begin{array}{cccc|cccccc} 0 & 1 & 2 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & 1 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 2 & 1 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 2 & 2 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 2 & 2 & 0 & 0 & 0 & 0 & 0 & 1 \end{array} \right).$$

We want to show that  $f_7 \notin \text{pPol}_2$ , however the matrix  $M_7$  associated with the partial function  $f_7$  is of size  $21 \times 25$ . So for the sake of brevity we will use  $f_3$  instead of  $f_7$ . Here the partial function  $f_3$  is given by

$$f_3 \left( \begin{array}{cccc|cccccccc} 0 & 1 & 2 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 1 & 2 & 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 2 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 1 & 2 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 1 & 2 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{array} \right) = \begin{pmatrix} 1 \\ 1 \\ 1 \\ 2 \\ 2 \\ 2 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$



Let  $\rho_2$  be the relation of arity 6 that consists of all columns of  $M_2$ . The matrix below shows that  $f_3 \notin \text{pPol}\rho_2$ .

$$f_3 \left( \begin{array}{cccc|cccccccc} 0 & 1 & 2 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & 1 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 2 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 1 & 2 & 2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \end{array} \right) = \begin{pmatrix} 1 \\ 1 \\ 2 \\ 2 \\ 0 \\ 0 \end{pmatrix}$$

Therefore the construction presented in this subsection is not applicable if we start with an  $n$ -ary partial function  $f$  whose matrix  $A$  does not satisfy the condition (\*).

**4.2. The partial function  $f$  is non-constant.** Notice first that any partial function whose matrix does not satisfy the above condition (\*) is included in this case. Indeed, let  $f$  be such an  $n$ -ary partial function. Then each of the  $m$ -tuples  $(\alpha, \dots, \alpha)^T$ ,  $\alpha \in \mathbf{k}$ , is a column of  $A$ . If  $f$  was a constant function, then  $f$  would be a partial projection function, contradicting  $f \notin \text{Str}(J_{\mathbf{k}})$ .

Without loss of generality we may assume that the domain of the  $n$ -ary partial function  $f$  is repetition free. Indeed if there are  $1 \leq i < j \leq n$  such that  $x_i = x_j$  for all  $\vec{x} = (x_1, \dots, x_n) \in \text{dom}(f)$ , then we can define the  $(n-1)$ -ary partial function  $g$  by

$$g(x_1, \dots, x_i, \dots, x_{j-1}, x_{j+1}, x_n) := f(x_1, \dots, x_i, \dots, x_{j-1}, x_i, x_{j+1}, \dots, x_n)$$

and use  $g$  instead of  $f$ . Thus we assume that all  $n$  columns of  $A$  are distinct. We shall make use of the following definitions and notation.

Let  $\text{Col}(A)$  be the set of all the columns of  $A$ ,  $d_f := |\text{dom}(f)|$  and let  $\vec{v}_f := f(A) \in \mathbf{k}^{d_f}$ . For  $\vec{x} := (x_1, \dots, x_h) \in \mathbf{k}^h$  and  $\ell \geq 1$ , we define  $\vec{x}^{\times \ell} \in \mathbf{k}^{h\ell}$  by

$$\vec{x}^{\times \ell} = \left( \underbrace{x_1, \dots, x_1}_{\ell \text{ times}}, \underbrace{x_2, \dots, x_2}_{\ell \text{ times}}, \dots, \underbrace{x_h, \dots, x_h}_{\ell \text{ times}} \right),$$

and let  $[\vec{x}] := \{x_1, \dots, x_h\}$ . For a set  $X \subseteq \mathbf{k}$  with  $X = \{x_1 < \dots < x_{|X|}\}$  and  $a \in X$  we define  $\text{next}_X(a) \in X$  by

$$\text{next}_X(a) := \begin{cases} x_{i+1} & \text{if } a = x_i \text{ and } i < |X|, \\ x_1 & \text{if } a = x_{|X|}. \end{cases}$$

Furthermore, for  $\vec{x} = (x_1, \dots, x_h) \in [\vec{v}_f]^h$  and  $1 \leq i \leq h$  we define  $c_i(\vec{x})$  by

$$c_i(\vec{x}) := (x_1, \dots, x_{i-1}, \text{next}_{[\vec{v}_f]}(x_i), x_{i+1}, \dots, x_h).$$

Since the partial function  $f$  is non-constant, the set  $[\vec{v}_f]$  contains at least two different elements, and so  $c_i(\vec{x}) \neq \vec{x}$  for all  $\vec{x} \in [\vec{v}_f]^h$  and all  $i = 1, \dots, h$ .

Let  $\alpha \geq 0$  be the number of columns  $\vec{u}$  in the matrix  $A$  that satisfy  $[\vec{u}] = [\vec{v}_f]$ . Without loss of generality we may assume that all those  $\alpha$  columns (if any) are the first columns to the left of the matrix  $A$ .

Let  $\ell \geq 1$ . We define the relation  $\rho_\ell$  of arity  $\ell d_f$  by

$$\rho_\ell := \{c_i(\vec{v}_f^{\times \ell}) \mid 1 \leq i \leq \ell d_f\} \cup \{\vec{u}^{\times \ell} \mid \vec{u} \in \text{Col}(A)\}.$$

Notice that  $|\rho_\ell| = \ell d_f + n$ . We denote by  $M_\ell$  the matrix with  $\ell d_f$  rows and whose  $\ell d_f + n$  columns are the tuples of  $\rho_\ell$  written in the following order:

$$c_1(\vec{v}_f^{\times \ell}), \dots, c_{\ell d_f}(\vec{v}_f^{\times \ell}), \vec{u}_1^{\times \ell}, \dots, \vec{u}_n^{\times \ell},$$

where  $\vec{u}_1, \dots, \vec{u}_n$  are the columns of  $A$  written in the same order as they appear in  $A$ . We denote by  $f^{\times \ell}$  the  $(\ell d_f + n)$ -ary partial function whose domain is the set of all rows of  $M_\ell$  and defined by

$$f^{\times \ell}(M_\ell) := \vec{v}_f^{\times \ell}.$$

Notice that for every  $\mathbf{x} := (x_1, \dots, x_{\ell d_f + n}) \in \text{dom}(f^{\times \ell})$ , we have that  $x_1, \dots, x_{\ell d_f + \alpha} \in [\vec{v}_f]$ .

**Example 20.** Let  $\mathbf{k} = \{0, 1, 2\}$ ,  $\ell = 3$  and

$$f \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 1 \\ 0 & 0 & 2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}.$$

Here  $\vec{v}_f = (0, 0, 1)$ ,  $[\vec{v}_f] = \{0, 1\}$ ,  $\vec{v}_f^{\times 3} = (0, 0, 0, 0, 0, 0, 1, 1, 1)$ ,

$$A = \begin{pmatrix} 0 & 0 & 0 \\ 1 & 0 & 1 \\ 0 & 0 & 2 \end{pmatrix}$$

and  $\text{Col}(A) = \{(0, 1, 0)^T, (0, 0, 0)^T, (0, 1, 2)^T\}$ . In this case we have

$$f^{\times 3}(M_3) = f^{\times 3} \left( \begin{array}{cccccccccc|ccc} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 1 & 0 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 & 0 & 1 & 1 & 0 & 0 & 2 & 2 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 & 0 & 1 & 0 & 0 & 2 & 2 \\ 1 & 1 & 1 & 1 & 1 & 1 & 1 & 1 & 0 & 0 & 0 & 2 & 2 \end{array} \right) = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 1 \\ 1 \\ 1 \end{pmatrix}$$

**Remark 21.** Let  $\ell \geq 1$ . Then for  $i = 1, \dots, ld_f$ , the tuples  $\vec{v}_f^{\times \ell}$  and  $c_i(\vec{v}_f^{\times \ell})$  differ only by their  $i$ -th coordinate. Therefore, for  $i = 1, \dots, ld_f$ , the restriction of the partial function  $f^{\times \ell}$  to a set of tuples not containing the  $i$ -th row of  $M_\ell$  is the partial projection function on the  $i$ -th coordinate.

**Lemma 22.** Let  $\ell \geq 1$ . Then  $f^{\times \ell} \in \text{pclone}(f)$  and  $f^{\times \ell} \notin \text{pPol}\rho_\ell$ .

**Proof.** Let  $h$  be the  $(ld_f + n)$ -ary partial function such that  $\text{dom}(h) := \mathbf{k}^{ld_f} \times \text{dom}(f)$  and  $h(x_1, \dots, x_{ld_f}, y_1, \dots, y_n) := f(y_1, y_2, \dots, y_n)$ . Then  $f^{\times \ell}$  is the restriction of  $h$  on the rows of the matrix  $M_\ell$ , thus  $f^{\times \ell} \in \text{pclone}(f)$ .

Now since the set  $[\vec{v}_f]$  contains at least two different elements, we have that  $c_i(\vec{v}_f^{\times \ell}) \neq \vec{v}_f^{\times \ell}$  for all  $1 \leq i \leq ld_f$ . Moreover since  $f$  is not a partial projection, we have that  $\vec{u}^{\times \ell} \neq \vec{v}_f^{\times \ell}$  for all  $\vec{u} \in \text{Col}(A)$  and so  $\vec{v}_f^{\times \ell}$  is not a column of  $M_\ell$  proving that  $f^{\times \ell} \notin \text{pPol}\rho_\ell$ . ■

Next we want to show that  $f^{\times \ell} \in \text{pPol}\rho_m$  whenever  $\ell \neq m$ . In order to do so, we need the following definition. Let  $M_\ell^*$  be the submatrix of  $M_\ell$  whose columns are  $c_1(\vec{v}_f^{\times \ell}), \dots, c_{ld_f}(\vec{v}_f^{\times \ell})$  together with all columns  $\vec{u}^{\times \ell}$  such that  $[\vec{u}^{\times \ell}] = [\vec{v}_f]$ . Thus the matrix  $M_\ell^*$  has exactly  $ld_f + \alpha$

columns. Furthermore, let  $R_\ell$  be the set of all columns of  $M_\ell^*$ , thus  $R_\ell = \{\omega \in \rho_\ell \mid [\omega] = [v_f]\}$ .

Denote by  $F_\ell$  the  $(ld_f + \alpha)$ -ary partial function with domain the set of all rows of  $M_\ell^*$  and defined by  $F_\ell(M_\ell^*) := v_f^{\times \ell}$ . Now by construction of  $f^{\times \ell}$ , we have that the value of  $f^{\times \ell}(\mathbf{x})$  is uniquely determined by the first  $ld_f$  entries of  $\mathbf{x}$  for every row  $\mathbf{x}$  of the matrix  $M_\ell$ . Therefore the value of  $f^{\times \ell}$  is determined by the first  $ld_f$  columns of  $M_\ell$  and thus the partial function  $F_\ell$  is well defined.

Summing up, we have the following result.

**Lemma 23.** *For all  $\ell, m \geq 2$ , if  $F_\ell \in \text{pPol}R_m$  then  $f^{\times \ell} \in \text{pPol}\rho_m$ .*

**Proof.** Suppose that  $f^{\times \ell} \notin \text{pPol}\rho_m$ . So there is a matrix  $P$  whose  $ld_f + n$  columns are all from  $\rho_m$  and whose rows are from  $\text{dom}(f^{\times \ell})$ , and such that  $f^{\times \ell}(P) \notin \rho_m$ . Let  $P'$  be the submatrix of  $P$  consisting of the first  $ld_f + \alpha$  columns of  $P$ .

Since the first  $ld_f + \alpha$  entries of every tuple  $\mathbf{x} \in \text{dom}(f^{\times \ell})$  belong to  $[v_f]$ , we have that all columns of  $P'$  belong to  $R_m$ , and since  $F_\ell(P') = f^{\times \ell}(P) \notin \rho_m$  we have that  $F_\ell \notin \text{pPol}R_m$ . ■

**Lemma 24.** *Let  $\ell, m \geq 2$  with  $\ell \neq m$ . Then  $f^{\times \ell} \in \text{pPol}\rho_m$ .*

**Proof.** We distinguish two cases.

**Case 1.**  $m < \ell$ . Let  $M$  be a matrix whose rows are in  $\text{dom}(f^{\times \ell})$  and whose columns are in  $\rho_m$ . Thus  $M$  has  $md_f$  rows all of which are rows of  $M_\ell$  and  $M$  has  $ld_f + n$  columns all of which are columns of  $M_m$ . As  $m < \ell$  the matrix  $M$  has less rows than  $M_\ell$  and so by Remark 21,  $f^{\times \ell}$  restricted to the rows of  $M$  is a partial projection function. Thus  $f^{\times \ell}(M) \in \rho_m$  proving that  $f^{\times \ell} \in \text{pPol}\rho_m$ .

**Case 2.**  $\ell < m$ . We argue the contrapositive. Suppose that  $f^{\times \ell} \notin \text{pPol}\rho_m$ . Then there is a matrix  $M$  whose rows are in  $\text{dom}(f^{\times \ell})$  and whose columns are in  $\rho_m$  and such that  $f^{\times \ell}(M) \notin \rho_m$ . If some row of  $M_\ell$  is missing in  $M$ , then by Remark 21,  $f^{\times \ell}$  restricted to the rows of  $M$  is a partial projection function, but this contradicts our assumption that  $f^{\times \ell}(M) \notin \rho_m$ .

Therefore we assume that all rows of  $M_\ell$  appear in  $M$  and consequently the matrix  $M$  contains  $md_f - ld_f$  repeated rows. Also, as all columns of  $M_\ell$  are distinct, we have that all  $ld_f + n$  columns of  $M$  are distinct. Let  $M^*$  be the submatrix of  $M$  consisting of the first  $ld_f + \alpha$  columns of  $M$ . Since  $f^{\times \ell}(M) \notin \rho_m$ , we have, by Lemma 23, that  $F_\ell(M^*) \notin R_m$ . Since all columns of  $M$  are distinct, we have that all  $ld_f + \alpha$  columns of  $M^*$  are distinct. Furthermore, since the matrix  $M$  contains  $md_f - ld_f$  repeated rows we have that  $M^*$  contains at least  $md_f - ld_f$  repeated rows.

Set  $\vec{v}_f^{\times m} := (v_1, \dots, v_{md_f})$ . Let  $j_1 \neq j_2 \in \{1, \dots, md_f\}$  and suppose that the  $j_1$ -th row and the  $j_2$ -th row of  $M^*$  are identical.

Now if  $v_{j_1} \neq v_{j_2}$ , then none of the tuples  $c_i(\vec{v}_f^{\times m})$  ( $i = 1, \dots, ld_f$ ,  $i \neq j_1$  and  $i \neq j_2$ ) can appear as a column of  $M^*$  because the  $j_1$ -th ( $j_2$ -th) coordinate of  $c_i(\vec{v}_f^{\times m})$  is  $v_{j_1}$  (is  $v_{j_2}$ ). This contradicts the fact that all  $ld_f + \alpha$  columns of the matrix  $M^*$  are distinct.

Therefore we assume that if  $j_1, j_2 \in \{1, \dots, md_f\}$  are such that the  $j_1$ -th row and the  $j_2$ -th row of  $M^*$  are identical, then  $v_{j_1} = v_{j_2}$  and in such a case neither  $c_{j_1}(\vec{v}_f^{\times m})$  nor  $c_{j_2}(\vec{v}_f^{\times m})$  appear as columns in  $M^*$ . Since at least  $md_f - ld_f$  rows of  $M^*$  are repeated, we conclude that at least  $md_f - ld_f + 1$  tuples of the form  $c_i(\vec{v}_f^{\times m})$  cannot appear as columns of  $M^*$ . Consequently, the number of all possible distinct columns in  $M^*$  is at most

$$md_f + \alpha - (md_f - ld_f + 1) = ld_f - 1 < ld_f,$$

and thus the matrix  $M^*$  cannot have all its  $ld_f$  columns distinct. This contradiction shows that  $F_\ell \in \text{pPol}R_m$ , and by Lemma 23  $f^{\times \ell} \in \text{pPol}\rho_m$ .  $\blacksquare$

The proof of the following result is similar to that of Corollary 16.

**Corollary 25.** *Let  $C$  be a strong partial clone on  $\mathbf{k}$  and suppose that  $C$  contains a partial function  $f \notin \text{Str}(J_{\mathbf{k}})$  that is not a constant function. Then the set of strong partial clones contained in  $C$  is of continuum cardinality.*

By combining Corollaries 16 and 25, we obtain the third main result of this paper.

**Theorem 26.** *Let  $C \neq \text{Str}(J_{\mathbf{k}})$  be a strong partial clone over  $\mathbf{k}$ . Then the family of all strong partial clones contained in  $C$  is of continuum cardinality.*

#### REFERENCES

- [1] F. Börner, L. Haddad and R. Pöschel, Minimal Partial Clones. *Bulletin of the Australian Mathematical Society* **44** (1991), pp 405–415.
- [2] F. Börner and L. Haddad, Maximal Partial Clones with no finite basis, *Algebra Universalis* **40** (1998), pp 453–476.
- [3] F. Börner, Personal Communications.
- [4] M. Couceiro, L. Haddad and K. Schölzel. On the Nonexistence of Minimal Strong Partial Clones. *Proceedings of the 47th IEEE International Symposium on Multiple-Valued Logic*, 2017, Novi Sad, Serbia, pp 82 – 87.
- [5] M. Couceiro and L. Haddad. A Survey on Intersections of Maximal Partial Clones of Boolean Partial Functions. *Proceedings of the 42nd IEEE International Symposium on Multiple-Valued Logic*, 2012, Victoria, Canada, pp 287–292.
- [6] L. Haddad and K. Schölzel. Countable Intervals of Partial Clones. *Proceedings 44th IEEE International Symposium on Multiple-Valued Logic*, 2014, Bremen, Germany, pp 155–160.
- [7] D. Lau, Function algebras on finite sets. A basic course on many-valued logic and clone theory. *Springer Monographs in Mathematics* (2006).
- [8] K. Schölzel. Dichotomy on intervals of strong partial Boolean clones, *Algebra Universalis* **73** (2015), pp 347–368.