



**HAL**  
open science

# A Neuron-Glial Perspective for Computational Neuroscience

Maurizio de Pittà, Hugues Berry

► **To cite this version:**

Maurizio de Pittà, Hugues Berry. A Neuron-Glial Perspective for Computational Neuroscience. Maurizio de Pittà; Hugues Berry. Computational Glioscience, Spinger, pp.3-35, 2019, Springer Series in Computational Neuroscience, 978-3-030-00817-8. 10.1007/978-3-030-00817-8\_1. hal-01995849

**HAL Id: hal-01995849**

**<https://inria.hal.science/hal-01995849v1>**

Submitted on 27 Jan 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# A Neuron-Glial Perspective for Computational Neuroscience

Maurizio De Pittà

Basque Center for Applied Mathematics, Bilbao, Spain;  
EPI BEAGLE, INRIA Rhône-Alpes, Villeurbanne, France

Hugues Berry

EPI BEAGLE, INRIA Rhône-Alpes, Villeurbanne, France

June 30, 2018

## Abstract

There is growing excitement around glial cells, as compelling evidence point to new, previously unimaginable roles for these cells in information processing of the brain, with the potential to affect behavior and higher cognitive functions. Among their many possible functions, glial cells could be involved in practically every aspect of the brain physiology in health and disease. As a result, many investigators in the field welcome the notion of a Neuron-Glial paradigm of brain function, as opposed to Ramon y Cayal’s more classical neuronal doctrine which identifies neurons as the prominent, if not the only, cells capable of a signaling role in the brain. The demonstration of a brain-wide Neuron-Glial paradigm however remains elusive and so does the notion of what neuron-glial interactions could be functionally relevant for the brain computational tasks. In this perspective, we present a selection of arguments inspired by available experimental and modeling studies with the aim to provide a biophysical and conceptual platform to computational neuroscience no longer as a mere prerogative of neuronal signaling but rather as the outcome of a complex interaction between neurons and glial cells.

## 1 Introduction

A decade ago, Ben Barres began his famous review on “The Mystery and Magic of Glia” recalling how he first became interested in the function of glial cells as a young neurologist in training. As he looked at brain sections from various neurological diseases under the microscope, not only he came to realize that at least a half of the brain cells are glial cells – astrocytes, oligodendrocytes and microglial cells – but also that glial cell phenotypes are radically altered in many brain injury and disease (Barres, 2008). These early observations grew in him the wonder of what glial cells normally do, and what their role in the disease is – a question that still lingers, beyond his lifetime achievements and the many other findings by his peers.

---

**Abbreviations.** Ado: adenosine; AMPA (AMPA):  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (receptor); ANLS: astrocyte-to-neuron lactate shuttle; AP: action potential; AQP4: aquaporin channel type 4; cAMP: cyclic adenosine monophosphate; CICR: calcium-induced calcium release; CNS: central nervous system; ECS: extracellular space; GABA:  $\gamma$ -Aminobutyric acid; GGC: glutamate–glutamine cycle; GPCR: G protein-coupled receptor; IP<sub>3</sub>: inositol 1,4,5-trisphosphate; ISI: interspike interval; Kir: inwardly rectifying K<sup>+</sup> channel; LC: locus coeruleus; LTD: long-term depression; LTP: long-term potentiation; MCT: monocarboxylate transporter; NA: noradrenaline; NBC: Na<sup>+</sup>-HCO<sub>3</sub><sup>-</sup> cotransporter; NKP: Na<sup>+</sup>/K<sup>+</sup> ATPase pump; NMDA (NMDAR): *N*-Methyl-D-aspartate (receptor); SIC (SOC): slow inward (outward) current; SNR: signal-to-noise ratio; SON: supraoptic nucleus; SSR: steady-state synaptic release; TNF $\alpha$ : tumor necrosis factor alpha; V1: primary visual cortex.

It is almost thirty years since the seminal discovery by Anne Cornell-Bell and colleagues that glutamate evokes calcium concentration rises in astrocytes (Cornell-Bell et al., 1990) – a phenomenon later confirmed in slices (Dani et al., 1992; Porter and McCarthy, 1996; Newman and Zahs, 1997) and *in vivo* (Wang et al., 2006), which could also occur in other glia types (Biber et al., 1999; Alberdi et al., 2002; Butt et al., 2004) – and yet our understanding of the functional role of glial calcium signaling is still very limited. As  $\text{Ca}^{2+}$  signaling can propagate along astrocytic processes and even between glial cells as waves (Cornell-Bell et al., 1990; Dani et al., 1992; Newman and Zahs, 1997; Hirase et al., 2004; Nimmerjahn et al., 2004), the possibility that glial  $\text{Ca}^{2+}$  waves might constitute an extraneuronal signaling system in the CNS was raised (Newman and Zahs, 1997). The next step was reached with the demonstration that increases in cytosolic  $\text{Ca}^{2+}$  concentration of astrocytes could regulate the release of neuroactive molecules (Parpura et al., 1994; Nedergaard, 1994; Jeftinija et al., 1996; Coco et al., 2003; Benz et al., 2004) and hormones (Krzan et al., 2003) or influence ion homeostasis (Price et al., 2002). The possibility that these astrocytic  $\text{Ca}^{2+}$  transients could influence neurons (Parpura et al., 1994; Nedergaard, 1994) and vascular smooth muscles (Zonta et al., 2003; Mulligan and MacVicar, 2004; Gordon et al., 2007), led to the idea that astrocytes are powerful regulators of neuronal spiking, synaptic plasticity and brain blood flow (Nedergaard et al., 2003; Haydon and Carmignoto, 2006).

The current consensus is that neuronal spiking, synaptic plasticity and the coupling between neurons and the vasculature underpin higher brain functions (Laughlin and Sejnowski, 2003). The recognition that glial cells could be much more than passive bystanders of neurons and their synapses has therefore been welcomed as a revolution of the classical Neuron paradigm of the brain in favor of a more comprehensive Neuron-Glial paradigm (Haydon, 2001). Nonetheless few topics in neuroscience are as controversial as the idea that glia could be an active partner in information processing, and cognitive and behavioral tasks elaborated by neurons. In part, this is due to a lack of definite experimental evidence. But in our opinion, much of the controversy originates from the lack of a theoretical platform whereupon to elaborate the Neuron-Glial paradigm. This perhaps comes without surprise in a context where theoretical studies in neuroscience claim they are able to account for many of the computational tasks performed by our brain by the sole consideration of neurons (Abbott et al., 2016). However, understanding the activity patterns of neural activity that support sensory processing (Ganmor et al., 2015), memory (Buzsáki, 2010), decision-making (Beck et al., 2008) and cognition (Rigotti et al., 2013), remains a major challenge for neuroscience theories. In the following we argue that such theories should also take into account a glial component, as we point out to key mechanisms of computational value that are mediated by glia.

## 2 Glial codes

When talking about neurons, the term ‘neural code’ refers to the properties of a single sequence of action potentials (spike trains) or a spike train ensemble to encode, decode and process sensory and/or cognitive information (Perkel and Bullock, 1968). Experimental and theoretical approaches to investigate the neural code are often based on presenting different inputs (stimuli) to an animal, and quantifying various properties of the generated spike patterns (Bialek and Rieke, 1992). Is it possible to think of an analogous code for glial cells?

Because glial cells are notoriously electrically passive (Tasaki and Chang, 1958; Steinhäuser et al., 1992), their code (if any) must rely on mechanisms that are different from the membrane potential, and these mechanisms are likely to be found among the many intracellular signaling pathways occurring within glial cells. Supported by modeling arguments (Kummer et al., 2000; Violin et al., 2008; De Pittà et al., 2009), signals originating from receptors in the glial plasma

membrane could all be potential candidates, as long as they bear strong correlation between the triggering extracellular stimulus and the downstream intracellular events, so as not to lose information about the former. This could be the case for example of second messengers like inositol 1,4,5-trisphosphate (IP<sub>3</sub>), intracellular calcium or cyclic adenosine monophosphate (cAMP) which result from the activation of a rich panoply of G protein-coupled receptors (GPCRs) expressed by glial cells (Porter and McCarthy, 1997; Káradóttir and Attwell, 2007; Pocock and Kettenmann, 2007). However, the diversity of these receptors, and the fact that they are linked to different signaling cascades – G<sub>q</sub>-PCRs to IP<sub>3</sub>/Ca<sup>2+</sup> signaling, and G<sub>s</sub>- and G<sub>i/o</sub>-PCRs respectively to stimulation and inhibition of cAMP (Gilman, 1987) – suggest that there might be multiple glial codes at play.

Among the possible glial codes, calcium signaling is the most studied one, arguably because of the availability of indicators to monitor intracellular Ca<sup>2+</sup> (Rusakov, 2015), as well as for its recognized function as intracellular messenger in a multitude of other cells (Berridge et al., 2000). There are nonetheless further reasons why Ca<sup>2+</sup> signaling could be an effective code for glial cells. In particular, Ca<sup>2+</sup> signaling could serve all the four key functions originally proposed by Perkel and Bullock for a candidate neural code: stimulus representation, interpretation, transformation and transmission (Perkel and Bullock, 1968).

*Stimulus representation* is broadly defined as the capability to reflect (or encode) extracellular stimuli by detectable changes in the cell activity. Stimulus representation is clearly carried out by Ca<sup>2+</sup> signaling, which is at the core of our current notion of glial excitability (Zorec et al., 2012). By such notion, we refer to the ability of glial cells to respond to extracellular stimuli by transient increases of their intracellular Ca<sup>2+</sup> concentration (Butt, 2006; Agulhon et al., 2008; Kettenmann et al., 2011). And accordingly, we dub a glial cell as ‘activated’ when its intracellular Ca<sup>2+</sup> dynamics departs from a baseline concentration (Verkhatsky and Kettenmann, 1996; Hoffmann et al., 2003). Because intracellular Ca<sup>2+</sup> is a simple ion, one expects the associated signal should be encoded in temporal and spatial patterns of intracellular Ca<sup>2+</sup> concentration (Figure 1, *yellow pathways*), very similar to the use of electric voltage or current signals in information technology (Jaffe, 1993). The ways by which a stimulus could be represented by glial calcium are multiple. The arguably simplest possibly is that the concentration of the extracellular agonist is encoded in the frequency, or amplitude, or a combination thereof, of the sequence of Ca<sup>2+</sup> elevations ensuing from stimulation (Schuster et al., 2002; De Pittà et al., 2008).

*Interpretation* refers to the possibility to reliably decode the information on the stimulus carried by the candidate code. In the case of glial Ca<sup>2+</sup> signaling, decoding is a necessary readout step for any mechanism that uses intracellular Ca<sup>2+</sup> to perform a downstream action, be it for example, the release of neuroactive molecules (Liu et al., 2006; Frühbeis et al., 2013; Imura et al., 2013; Araque et al., 2014) and myelin-related proteins (Krämer-Albers et al., 2007), the regulation of the blood flow (Haydon and Carmignoto, 2006; Attwell et al., 2010) or the transcription of cellular factors (O’Neill and Kaltschmidt, 1997). All these processes seem to be strongly constrained by intracellular Ca<sup>2+</sup> dynamics: some of them require intracellular Ca<sup>2+</sup> concentration to overcome a threshold value in order to occur (Dolmetsch et al., 1997; Parpura and Haydon, 2000; Pasti et al., 2001; Liu et al., 2006), others depend, in magnitude, on intracellular Ca<sup>2+</sup> concentration (Zonta et al., 2003; Krämer-Albers et al., 2007). Gliotransmitter exocytosis from astrocytes appears to occur preferentially during rising phases of Ca<sup>2+</sup> elevations (Pasti et al., 1997; Marchaland et al., 2008), hinting a preferential dependence on the frequency rather than the duration of these elevations (De Pittà et al., 2011). Conversely, the activation of transcription factors positively correlates with the amplitude and duration of Ca<sup>2+</sup> increases (Dolmetsch et al., 1997). Taken together, these observations hint the existence of readout mechanisms that are finely tuned to extract selective features of the Ca<sup>2+</sup> signal.

*Transformation.* The  $\text{Ca}^{2+}$  signal that is generated within glial cells must be manipulated (i.e. transformed) in a controlled way to allow computations of some sort in order to be effectively regarded as a code. Consider for example the formation of a  $\text{Ca}^{2+}$  wave, that is a range of  $\text{Ca}^{2+}$  concentration transients traveling through the cytoplasm of a glial cell or a group of glial cells (Leybaert and Sanderson, 2012). One hypothesis states that these waves result from the spatiotemporal summation of short-lived, spatially-confined  $\text{Ca}^{2+}$  puffs (Marchant and Parker, 2001) whose coordination is possible by inherent cellular properties, like the characteristic distance between  $\text{Ca}^{2+}$  puff sites and their spatial coupling set by local  $\text{Ca}^{2+}$  buffers (Skupin et al., 2008). In this example, the generation of the wave is a transformation of the local  $\text{Ca}^{2+}$  signal. The growing recognition of a compartmentation of  $\text{Ca}^{2+}$  signals within glial cells, like astrocytes in the cortex (Bindocci et al., 2017) or Bergman cells in the cerebellum (Grosche et al., 1999), also supports the hypothesis of a transformation of  $\text{Ca}^{2+}$  signals by the cellular architecture, based on its control of the genesis, dynamics and extension of the waves.

*Transmission.* The candidate code must also enable transmission of information between different cells, allowing for the exchange and propagation of information across cell networks. The propagation of intercellular  $\text{Ca}^{2+}$  waves in astrocyte networks has been observed in the cortex (Nimmerjahn et al., 2004), the hippocampus (Kuga et al., 2011), the retina (Newman, 2001), or the cerebellum (Hoogland et al., 2009).  $\text{Ca}^{2+}$  waves could as well occur in microglia and oligodendrocytes either independently (Takeda et al., 1995; Wu et al., 2013; Jiang et al., 2017) or in interaction with astrocytes (Verderio and Matteoli, 2001; Schipke et al., 2002; Parys et al., 2010).

Taken together,  $\text{Ca}^{2+}$  signalling seems to match the four key functions above. There are however several obstacles on the path of the identification of  $\text{Ca}^{2+}$  signalling as a glial code. First of all,  $\text{Ca}^{2+}$  signalling is stochastic, a property that has mostly been studied with  $\text{Ca}^{2+}$  spikes in cultured astrocytes and microglia (Skupin et al., 2008; Thurley and Falcke, 2011; Thurley et al., 2012, 2014). This stochasticity largely complicates the analysis of  $\text{Ca}^{2+}$  signalling in glial cells. Moreover, at the molecular level, there might be more than a single  $\text{Ca}^{2+}$  signalling system at play in glial cells and those different systems might be segregated to distinct subparts of the cell (Volterra et al., 2014). The best known example of this anisotropy is provided by astrocytes, in which the main source of  $\text{Ca}^{2+}$  for the transients is believed to be  $\text{Ca}^{2+}$  exchange between the endoplasmic reticulum and the cytosol (Scemes and Giaume, 2006; Leybaert and Sanderson, 2012). However, knockout of  $\text{IP}_3\text{R}2$  receptors – often considered the main channels responsible for endoplasmic reticulum to cytosol  $\text{Ca}^{2+}$  transport in those cells (Zhang et al., 2014) – indeed abolished nearly all  $\text{Ca}^{2+}$  transients in the cell soma, but spared roughly half of the  $\text{Ca}^{2+}$  transients in the cell processes (Srinivasan et al., 2015). The molecular mechanisms for those  $\text{IP}_3\text{R}2$ -independent  $\text{Ca}^{2+}$  transients in the processes could rely on additional extracellular calcium fluxes from plasma membrane channels (Srinivasan et al., 2015) or on type-1 or type-3  $\text{IP}_3\text{R}$  (Sherwood et al., 2017), or a combination thereof.

The presence of different mechanisms of  $\text{Ca}^{2+}$  spiking has also been suggested to account for the variety of the spatiotemporal scales of the  $\text{Ca}^{2+}$  signals that include fast ( $\sim 0.3 - 0.93$  s)  $\text{Ca}^{2+}$  events, locally confined in astrocytic processes (Di Castro et al., 2011; Bindocci et al., 2017); as well as large, and longer ( $> 5$  s)  $\text{Ca}^{2+}$  signals spanning whole processes, and that can invade the whole cell (Volterra et al., 2014; Bindocci et al., 2017). This is reminiscent of the hierarchy of  $\text{Ca}^{2+}$ -induced- $\text{Ca}^{2+}$ -release (CICR)-mediated  $\text{Ca}^{2+}$  events that, from blips and puffs – respectively associated with the opening of individual  $\text{IP}_3\text{R}$  receptors or a cluster thereof – can ultimately lead to the generation of cell-wide  $\text{Ca}^{2+}$  spikes.

There is however evidence that the distribution of intervals between consecutive  $\text{Ca}^{2+}$  blips or puffs at the astrocyte periphery, maybe different from those reported for CICR-mediated  $\text{Ca}^{2+}$  events in primary processes and somata (Bindocci et al., 2017). Therefore, a major open

question in the field is to decide whether the different  $\text{Ca}^{2+}$  signals observed in astrocytes are the result of the spatiotemporal summation of a unique underlying microscopic mechanism or of different microscopic mechanisms, each possibly attached to a specific location in the cell.

### 3 Oligodendrocytes and regulation of axonal electric conduction

A unique specialization of glia in vertebrates is myelination, that is the ensheathing of axons by myelin (Waehnel et al., 1986; Zalc et al., 2008). Myelination provides at least two advantages. First, compact lipid-rich myelin provides high membrane electrical resistance and low capacitance, which prevents current loss and enables rapid and efficient action potential (AP) conduction. Second, increased conduction velocity allows for computational complexity within a compact nervous system such as that of vertebrates (Fields, 2008).

Oligodendrocytes are the specialized glial cells responsible both for axon myelination, and for the formation and maintenance of nodal structures and lifelong integrity of axons in the CNS (Nave, 2010). By controlling the structural properties that characterize myelinated axons, oligodendrocytes likely participate to the fine tuning of action potential shape and conduction velocity (Kimura and Itami, 2009; Tomassy et al., 2014; Ford et al., 2015; Hughes et al., 2018). This could be achieved either by changes of myelin thickness or internode length or node geometry, or by a combination thereof (Ullén, 2009; Fields, 2008; Arancibia-Carcamo et al., 2017). For instance, adjustments of conduction speed of myelinated axons can tune propagation times to mediate sound localization (Carr and Konishi, 1990; McAlpine and Grothe, 2003; Seidl et al., 2010; Ford et al., 2015) or promote synchronous neuronal firing (Sugihara et al., 1993; Lang and Rosenbluth, 2003). Together with delays induced by the AP-generation dynamics (Fourcaud-Trocmé et al., 2003) and those rising from synaptic processing (Markram et al., 1997), axonal conduction delays are an important property of neural interactions, which may induce a wealth of dynamical states with different spatiotemporal properties and domains of multistability which could serve different computational purposes (Roxin et al., 2005; Roxin and Montbrió, 2011).

The observation that the tuning of conduction delays by oligodendrocytes continues during adulthood (Dimou et al., 2008; Young et al., 2013) suggests the thought-provoking possibility that these different states could change in an experience-dependent way, i.e. in a manner that depends on neuronal electrical activity (Fields, 2013). The potential functional and computational implications of this two-way regulation of oligodendrocyte-mediated transmission delays and neuronal activity remain to be investigated but we note that the active control of conduction delays of axon ensembles by oligodendrocytes may lead to time-locked patterns of activation of the neurons – a concept known as polychronization which could be of great computational significance (Izhikevich, 2006).

### 4 Glia morphology and functional specialization

The factors underpinning morphogenesis of neuronal structures remain poorly understood, in particular when one includes neuron-glia interactions into the equation. Experimental studies have identified a combination of transcriptional programs and a battery of molecular signals that seem to be regulated both developmentally and regionally, in a cell-specific manner (Jadhav et al., 2009; Eroglu and Barres, 2010; Götz, 2013; Stassart et al., 2013). A paramount example of this complex combination of factors is provided by astrocytes, the glial cells recognized as key regulators of the neuronal connectome in the cortex and hippocampus (Fields et al., 2015). In the postnatal developing brain, astrocytes are found in spatially distinct domains and

express domain-specific genes that are needed to support formation of specific neural circuits and neuronal subtypes (Molofsky et al., 2014). Astrocytes also secrete a variety of molecules that regulate with spatiotemporal specificity all stages of the genesis of functional neural circuits (Allen, 2013; Clarke and Barres, 2013). This intimate relationship between astrocytes and synapses continues in the adult brain, with the processes of the astrocytes that wrap themselves around synapses (Reichenbach et al., 2010), engaging in a multitude of signals (Figure 1). Eventually, a single astrocyte contacts circa 20–150 thousand synapses in the rodent (Bushong et al., 2002), and  $\sim 300$  thousand to 2 million synapses in the human brain (Oberheim et al., 2009). What are the possible functional impacts of this ensheathing?

One obvious possibility is that astrocytic processes are strategically positioned at synaptic loci to act as physical barriers that constrain and regulate extracellular diffusion of neurotransmitters (Ventura and Harris, 1999). A realistic reconstruction of the extracellular space (ECS) between astrocytic processes and glutamatergic synapses of the hippocampus revealed a strong impact of astrocyte arrangements on glutamate diffusivity (Kinney et al., 2013). The degree of astrocytic ensheathing at glutamatergic synapses also directly dictates the rate of glutamate uptake by glial transporters – the main mechanism of glutamate uptake in the adult brain (Clements, 1996; Danbolt, 2001). The degree of astrocyte ensheathment seems inversely correlated with the size of dendritic spines (Medvedev et al., 2014). This suggests that, in agreement with independent theoretical arguments (Barbour, 2001), smaller synapses are strongly sealed by astrocytic processes whereas larger synapses might exhibit more glutamate spillover (Ventura and Harris, 1999).

An additional element of complexity is that astrocytic processes are highly motile and may undergo structural changes in response to neural activity (Heller and Rusakov, 2015). Activity-dependent plasticity of perisynaptic glial processes has been reported in the rat supraoptic nucleus in relation with parturition (Oliet et al., 2001; Piet et al., 2004b), in the hypothalamic suprachiasmatic nucleus in association with circadian light/dark cycles (Becquet et al., 2008), in the hippocampus (Haber et al., 2006; Verbich et al., 2012) and the cerebellum in correlation with neural activity (Lippman et al., 2008, 2010) and in the somatosensory cortex *in vivo* by sensory stimulation (Bernardinelli et al., 2014b; Perez-Alvarez et al., 2014). Those results suggest that activity-dependent plasticity of perisynaptic glial processes is a brain-wide phenomenon, although its functional consequences are likely region-specific (Khakh and Sofroniew, 2015). At hippocampal excitatory synapses for example, astrocyte processes preferentially colocalize with the postsynaptic element (Rusakov, 2001; Lehre and Rusakov, 2002). Simulations of glutamate diffusion at those synapses suggest that this arrangement favors glutamate spillover to the presynaptic elements (Rusakov, 2001; Lehre and Rusakov, 2002). This could promote activation of presynaptic metabotropic receptors, and account for feedback inhibition mediated by these receptors at mossy fiber-pyramidal cell synapses (Lehre and Rusakov, 2002; Nicoll and Schmitz, 2005; Omrani et al., 2009).

Glial transporters and the degree of glial coverage of synapses have also been implicated in the control of long-term plasticity at excitatory synapses. Astrocytic glutamate transporters gate spike-timing-dependent plasticity in the striatum *ex-vivo* (Valtcheva and Venance, 2016), as well as rate-coded long-term potentiation (LTP) in the amygdala (Tsvetkov et al., 2004) and in the hippocampus (Katagiri et al., 2001; Lushnikova et al., 2009). LTP induction in the hippocampus correlates with dynamical changes of astrocytic processes at the stimulated synapses (Lushnikova et al., 2009; Bernardinelli et al., 2014b; Perez-Alvarez et al., 2014), and these changes could associate with long-lasting spine formation and remodelling (Lushnikova et al., 2009; Verbich et al., 2012; Bernardinelli et al., 2014b). Taken together, those results pinpoint a key role for activity-dependent, dynamical reshaping of glial ensheathing in the control of synapse morphology, extracellular glutamate homeostasis, and the functionality of

neural circuits (Bernardinelli et al., 2014a).

## 5 Ion homeostasis and volume regulation

Glial cells express a vast cassette of ion channels, transporters and pumps whereby they can sense and regulate extracellular ion concentrations with important functional consequences on neural activity (Kettenmann and Ransom, 2013; Simard and Nedergaard, 2004). One of the first recognized possible functions of astrocytes was their importance in the maintenance of extracellular  $K^+$  (Figure 1, *blue pathway*) (Hertz, 1965; Orkand et al., 1966). Although traditionally considered in a pathological context including epilepsy and spreading depression (Somjen, 2002), electrophysiological recordings *in vivo* also implicate modulations of extracellular  $K^+$  by astrocytes in the regional control of neuronal excitability in the healthy brain (Amzica et al., 2002; Djukic et al., 2007; Chever et al., 2010). The resting membrane potential of neurons depends on intra- and extracellular concentrations of different ions (by the Goldman-Hodgkin-Katz equation) (Johnston and Wu, 1995), so changes of extracellular  $K^+$  concentration can alter neuronal excitability. Theoretical investigations suggest that the interplay of astrocytic  $Na^+/K^+$  pumps (NKPs) and Kir4.1 channels (Walz, 2000) can promote AP generation in conditions of low neuronal firing (Somjen et al., 2008), thus potentially contributing to the brain basal level of spontaneous neural activity (Deco et al., 2011). Alternatively, depending on the activity level, this could translate into a non-trivial positive feedback on neuronal firing, turning regular spiking dynamics into bursting (Somjen et al., 2008; Øyehaug et al., 2012; Cui et al., 2018).

Astrocyte-mediated regulation of extracellular  $K^+$  also affects neuronal excitability by modulating the time course of glutamate at excitatory synapses (Djukic et al., 2007). For each glutamate molecule transported, glial glutamate transporters co-transport 3  $Na^+$ , 1  $H^+$  and 1  $K^+$  (Levy et al., 1998). Therefore alteration of extracellular  $K^+$  concentration is expected to change the rate of glutamate uptake, thus modulating the excitatory drive of a neuron and its firing dynamics.

Astrocytes also express a number of transporters involved in pH regulation (Deitmer and Rose, 1996, see also Figure 1, *brown pathway*) that may be affected by alterations of  $K^+$  concentration and glutamate transport rate. The impact of these glial transport/uptake systems on neuronal computation would strongly depend on the stoichiometry of individual transporters and their spatial organisation on the glial plasma membrane along with that of ion channels and pumps. All this can vary depending on the brain region (Kofuji and Newman, 2004; Rose and Chatton, 2016). In the rat optic nerve, for example, extracellular  $K^+$  buffering seems more dependent on NKPs than on Kir channels (Ransom et al., 2000). Similarly, glial  $K^+$  spatial buffering, which ensues from the electrotonic propagation of a  $K^+$  current in glial cells from ECS regions where  $K^+$  is high to others where the concentration of this ion is low (Orkand et al., 1966), may be carried out differently in astrocytes vs. Müller glia, possibly because of the different expression and distribution of Kir channels in these cells (Kofuji and Newman, 2004).

Another specificity of astrocytes in connection with extracellular ion homeostasis is that their volume can change by activity-dependent modulations of extracellular  $K^+$  (Florence et al., 2012; Larsen et al., 2014). Although the molecular mechanisms of astrocytic swelling remain to be elucidated, colocalization of astrocytic Kir4.1 with aquaporin water channels AQP4 strongly supports a role for these channels in this phenomenon (Nagelhus et al., 2004). Compartmental modelling supports a mechanism whereby  $K^+$  influx into astrocytes through Kir4.1 channels, resulting from  $K^+$  buffering of neuronal activity, sets an osmolarity gradient across the astrocyte membrane which drives extracellular water into the astrocyte by AQP4 channels, thus increasing



the cell volume (Østby et al., 2009). Bicarbonate ( $\text{HCO}_3^-$ ) homeostasis also plays a crucial role in physiological astrocyte swelling (Florence et al., 2012), presumably by improving the steepness of the osmolarity gradient (Østby et al., 2009).

## 6 Gliotransmission

Gliotransmission, that is the active information transfer from astrocytes to neurons (Bezzi and Volterra, 2001) probably bears the widest implications in information processing. There is arguably no other concept that epitomizes the Neuron-Glia paradigm better than the “tripartite synapse”, whereby astrocyte is the third active element in synaptic information transfer besides the pre- and postsynaptic terminals, sensing synaptically-released neurotransmitters by a variety of mechanisms, and signaling back to synaptic terminals by gliotransmission (Figure 1, *red pathway*) (Araque et al., 1999, 2014). An emerging view is that gliotransmission is probably not a global, stereotyped On-Off phenomenon, but rather a multifaceted one, dependent on age, circuit, stimulation and the considered synapse (Savtchouk and Volterra, 2018). Indeed, not all synapses are tripartite:  $\sim 10\% - 50\%$  of cortical and hippocampal synapses are not adjacent to any astrocytic process (Ventura and Harris, 1999; Witcher et al., 2007; Kasthuri et al., 2015) and that proximity of glial processes to dendritic spines varies depending on the brain area (Ventura and Harris, 1999; Grosche et al., 2002) and on the considered synapse in a given area (Chao et al., 2002). The existence of such bidirectional communication between astrocytes and synapses in physiological conditions is therefore still lively debated (see e.g. Fiacco and McCarthy (2018) and Savtchouk and Volterra (2018)). However consideration of this possibility opens to profound functional implications as it puts astrocytes in the position to actively control synaptic transmission.

Although astrocytes can release a range of gliotransmitters, including glutamate, ATP, GABA and D-serine (Sahlender et al., 2014), the effects of these gliotransmitters can be generalised in terms of modifications of short- and long-term properties of synapses. Consider first short-term synaptic dynamics. An important consequence of this dynamics is that synapses can act as filters of the AP sequence they transmit (Fortune and Rose, 2001). However, the filtering characteristics of a given synapse is not fixed but can be adjusted through modulation of the initial release probability  $p_0$  (Dittman et al., 2000). Depending on receptor type, gliotransmitters may either increase or decrease the value of  $p_0$  at both excitatory and inhibitory synapses (Araque et al., 2014; Angulo et al., 2008), thereby turning high-pass filtering synapses into band-pass filtering ones, or band-pass filtering synapses into low-pass filtering ones, and vice versa (De Pittà et al., 2015). Low-pass filtering synaptic characteristics associates with derivative synaptic coding, that is the transmission of variations (i.e. the derivative) of AP sequences, as opposed to high-pass filtering, which results instead in integrative synaptic coding and carries information on the count of incoming APs (Tsodyks, 2005). Hence, modulation of synaptic filtering by gliotransmitters could make a synapse alternate between these different coding modes (De Pittà and Brunel, 2016).

The modulation of synaptic release probability by gliotransmitters may also occur on multiple time scales (reviewed by De Pittà et al., 2015). It may lasts for tens of seconds up to few minutes, thus affecting synaptic transmission and network computations merely for a transient period of time. But it can become persistent and last for tens of minutes, that is on time scales that could promote long-term plastic changes of the synapse (Bear and Malenka, 1994). This possibility was showed to account for long-term depression (LTD) at glutamatergic synapses between neurons in layers 4 and 2/3 of the barrel cortex (Min and Nevian, 2012). Long-term modulations of synaptic release are expected to substantially alter the recruitment of postsynaptic NMDA receptors, thereby modulating the influx of  $\text{Ca}^{2+}$  ions (Froemke and Dan, 2002;

Froemke et al., 2006, 2010). Because the time course of postsynaptic  $\text{Ca}^{2+}$  is believed to control the outcome of spike-timing-dependent plasticity (STDP) – namely whether LTP or LTD is observed (Ismailov et al., 2004; Nevian and Sakmann, 2006; Graupner and Brunel, 2010), the above arguments predict that persistent alterations of synaptic release by gliotransmitters modulate STDP. Computational arguments however hint that this modulation is not straight forward and depends on the timing of gliotransmitter release from perisynaptic astrocytic processes with respect to pre- and postsynaptic APs (De Pittà and Brunel, 2016).

Gliotransmitters may also modulate the probability of activation of postsynaptic receptors, either by the release of D-serine from astrocytes, which is a co-agonist for NMDA receptors with synaptically-released glutamate (Oliet and Mothet, 2009), or by slow depolarizing and hyperpolarizing currents (De Pittà and Brunel, 2016), respectively mediated by extrasynaptic NMDA and GABA receptors (Kozlov et al., 2006; Jiménez-González et al., 2011; Chen et al., 2012; Le Meur et al., 2012; Martín et al., 2015). How could gliotransmission affect learning via modulation of synaptic plasticity? Although this topic remains to be investigated, few theoretical studies offer some enticing insights into the question. Porto-Pazos and collaborators investigated the performance of an astrocyte-inspired learning rule to train deep learning networks in data classification and found that the trained neuron-glia networks were able to outperform identical networks without astrocytes in all discrimination tasks they implemented (Porto-Pazos et al., 2011; Alvarellos-González et al., 2012; Mesejo et al., 2015). Although those investigators do not provide any explanation for the possible mechanism whereby astrocyte-mediated plasticity could improve network learning and performance, a significant feature of their learning rule is that, for successful training, potentiation by astrocytes must be weaker than depression resulting from astrocyte inactivity (Alvarellos-González et al., 2012; Mesejo et al., 2015). In their model, this amounts to putting the threshold between LTP and LTD under the control of astrocyte activity. There is circumstantial evidence that astrocytes could modify the threshold for LTD vs. LTP induction, such as in the supraoptic nucleus – where astrocytic coverage of synapse is reduced during lactation (Panatier et al., 2006).

Philips et al. (2017) devised a modified version of the BCM rule (Bienenstock et al., 1982; Gerstner and Kistler, 2002) where the threshold rate of postsynaptic firing for induction of LTD vs. LTP varies proportionally with astrocyte activation, and investigated how this rule affects development of orientation preference maps (OPMs) in a self-organizing network model of the primary visual cortex (V1) (Stevens et al., 2013). This choice not only allows reproducing map orientation experimentally observed in V1, but also reveals that, upon reduction of astrocytic radius, the periodicity of OPMs increases while the width of individual hypercolumns decreases (Philips et al., 2017). Since astrocyte size varies across species (Oberheim et al., 2009; López-Hidalgo et al., 2016), these results predict a causal link between astrocytic radius and the different hypercolumn widths observed in different species (Kaschube et al., 2010).

## 7 Resource management

Glucose is the almost exclusive energetic fuel of the mammalian brain and can completely sustain neural activity acting as a substrate for the synthesis of lipids and amino acids, as well as of neurotransmitters like glutamate, GABA or acetylcholine (Dienel, 2012; Mergenthaler et al., 2013). Cerebral glucose fuels neural activity by glycolysis which converts glucose into pyruvate by the concomitant production of two molecules of ATP. Pyruvate may then either enter mitochondria, where it is oxidized by Krebs cycle, or be temporarily converted to lactate in the cytoplasm for later use. Moreover, lactate produced in one cell can also be released into the ECS and be used by other cells (Simpson et al., 2007). Glucose supply to the brain is mainly either through the interstitial fluid by blood circulation or by metabolism of intracellular glycogen

– the only glucose store. Multiple mechanisms ensure neurovascular coupling whereby local energy supply, together with the cerebral blood flow, is finely adjusted to variations of the neuronal activity to allow normal neuronal function (Magistretti and Allaman, 2015). Astrocytes have been implicated in all aspects of this coupling. They can regulate cerebral blood flow, by dilating or constricting blood vessels through multiple  $\text{Ca}^{2+}$ -dependent pathways (Figure 1, *magenta pathway*) (Iadecola and Nedergaard, 2007; Attwell et al., 2010). In parallel, they can actively supply glucose through glycogen and by production of lactate – possibly triggered by the energy demand of excitatory synaptic activity (Magistretti and Allaman, 2015).

The proposal of an astrocyte-to-neuron lactate shuttle (ANLS) was originally formulated by Pellerin and Magistretti (1994) and envisages the generation of lactate at excitatory synapses by astrocytes, and its export to neurons where it is converted to pyruvate for ATP generation in mitochondria (Pellerin and Magistretti, 2012). This scheme could regulate the energy supplied to neurons in response to their activity, since glutamate released by active neurons could promote lactate production in astrocytes by stimulating glycolytic ATP generation to power astrocytic uptake of glutamate in a positive feedback loop (Figure 1, *purple pathway*). However, it is not clear under what circumstances of neuronal activation the ANLS could occur, as well as if it could be a brain-wide signaling pathway, or rather depend on the architecture of local neuron-glia circuits (Vaishnavi et al., 2010; Bélanger et al., 2011; Magistretti and Allaman, 2015). Actually, the only evidence *in vivo* for the ANLS at present, sees it implicated in hippocampal LTP maintenance (Suzuki et al., 2011) and spatial working memory (Newman et al., 2011).

Astrocytes are also mediators of the glutamate–glutamine cycle (GGC) (Danbolt, 2001; Hertz, 2013), whereby synaptically released glutamate is taken up by astrocytic transporters and converted to glutamine by glutamine synthase – an enzyme exclusively found in astrocytes (Figure 1, *turquoise pathway*). Glutamine is then released to the ECS, where it is sequestered by neurons and reconverted to glutamate, and by this latter, possibly to GABA too (Rothman et al., 2003; Hertz, 2013). The GGC can continuously supply neurotransmitters to neurons to sustain synaptic transmission, both at excitatory and at inhibitory synapses. At cortical and hippocampal excitatory synapses however, neurotransmitter supply by GGC was proven necessary to sustain synaptic release for prolonged, high-frequency stimulations, but appeared dispensable in conditions of low synaptic activity (Tani et al., 2014).

## 8 Microglia in neuronal and astrocytic signaling

Microglial cells roughly constitute 5% to 15% of the brain’s cellular elements, representing the most numerous glial population in the brain after astrocytes and oligodendrocytes (Lawson et al., 1990; Pelvig et al., 2008). Although they are traditionally described to comprise the main component of the brain’s innate immune system by responding to any pathological insult, experimental observations over the past two decades also suggest an involvement of these cells in the genesis and function of neural circuits in the healthy brain (Kettenmann et al., 2013).

Microglial processes in the somatosensory and visual cortex make brief, repetitive contacts with synapses at a frequency of about once per hour (Nimmerjahn et al., 2005; Wake et al., 2009). The nature of these contacts is synapse-specific and varies with neuronal activity and brain region. Visual deprivation shifts the preferential association of V1 microglial processes to small dendritic spines that transiently grow, to a subset of bigger spines that persistently shrink (Tremblay et al., 2010). Reduction of spontaneous neuronal activity correlates with retraction of microglial processes in the visual cortex (Wake et al., 2009), but not in the somatosensory cortex (Nimmerjahn et al., 2005).

Besides a role in structural synaptic plasticity (Stevens et al., 2007; Schafer et al., 2012; Kettenmann et al., 2013), microglia could also participate in several forms of functional synaptic

plasticity (Kettenmann et al., 2013; Wake et al., 2013). This could be achieved either directly, by release from microglia of plasticity-inducing molecules, or indirectly, through the modulation of gliotransmission-mediated pathways of plasticity (Section 6). In the latter scenario in particular, microglia could either promote (Pascual et al., 2011) or amplify glutamate release from astrocytes (Bezzi et al., 2001) (Figure 2, schemes A and E), with the potential to modulate excitatory synaptic transmission and neuronal excitability in a variegated fashion.

The proinflammatory cytokine tumor necrosis factor alpha ( $\text{TNF}\alpha$ ) is tightly related to microglia. Brain  $\text{TNF}\alpha$  production can be stimulated by neuronal activity (Churchill et al., 2008), and the cytokine can be released into the ECS both by astrocytes and by microglia (Habbas et al., 2015; Lewitus et al., 2016).  $\text{TNF}\alpha$  is found in low concentrations in the ECS in physiological conditions (Santello and Volterra, 2012), where it seems necessary for normal glutamatergic gliotransmission (Bezzi et al., 2001; Stellwagen et al., 2005; Domercq et al., 2006; Santello et al., 2011; Pribiag and Stellwagen, 2013) and regulation of homeostatic synaptic plasticity mechanisms, like synaptic scaling (Steinmetz and Turrigiano, 2010). The implication of  $\text{TNF}\alpha$  in synaptic scaling may be important during competition with Hebbian plasticity, for instance for ocular dominance plasticity (Kaneko et al., 2008).

## 9 Glia in higher brain functions

A popular argument, often raised to wow the audience, is that Albert Einstein’s brain, when first inspected, did not look statistically significantly different from that of other individuals with normal cognitive ability, except for a higher glia/neuron ratio, leading to speculate that the larger number of glial cells could partly have accounted for Einstein’s exceptional intelligence (Diamond et al., 1985). The further observation that Einstein’s corpus callosum – the largest bundle of myelinated fibers in the brain that connects the two cerebral hemispheres – was thicker in several subregions than in elderly and younger individuals certainly added to the popularity of the argument (Men et al., 2014). Although the studies at the origin of those sensational results have been strongly opposed because of methodological issues or interpretation biases (Fields, 2009), the implication of glia in higher brain functions is still an hypothesis that cannot be rejected.

From an evolutionary point of view, glia emergence coincides with the appearance of the centralized nervous systems (Hartline, 2011), and later on, with the birth of radial glial cells which hallmark the development of nervous systems of chordates and vertebrates, and are precursors of astrocytes in the adult brain of these animals (Sild and Ruthazer, 2011). In parallel, the evolution of myelin reduced the energy required for neuronal communication and boosted the speed of impulse propagation, allowing complex nervous systems to operate quickly and efficiently (Zalc and Colman, 2000). Oligodendrocytes also provide trophic support for the axons they myelinate, thus allowing for longer axons and, ultimately, greater vertebrate size (Nave, 2010). As nervous systems increase in complexity, a trend in increased complexity is also observed for astrocytes (Verkhatsky and Nedergaard, 2016). Human and primate astrocytes are larger and more branched than rodent ones, with humans astrocytes generally being the largest (Oberheim et al., 2009). Moreover, human astrocytes morphologies are more divers, with some morphologies that are not observed in other species (Colombo and Reisin, 2004; Oberheim et al., 2009). A landmark study by Maiken Nedergaard’s group considered the injection of human glial progenitor cells into the ventricles of newborn mice (Han et al., 2013). As those progenitor cells differentiated into mature astrocytes and oligodendrocytes, the adult engrafted animals were found to outperform their littermates that did not receive human glial progenitor grafts, on multiple cognition tests, including novel object recognition and fear conditioning. Overall, these observations provide direct evidence that human astrocytes boost cognitive abilities of

mice, possibly by increasing neural plasticity (Han et al., 2013).

Several lines of evidence in humans and in other animal models link glia function with the development and maturation of multiple cognitive and motor skills (Fields, 2008; Oliveira et al., 2015). Beyond motor performance, sensory processing is of great importance for appropriate evaluation of behavior. The development and function of cortical maps seem coupled with astrocytic signaling (López-Hidalgo and Schummers, 2014). In the primary visual cortex for example, astrocytes are integral components of OPMs, as they are both visually responsive and capable of modulating visually driven responses in close register with receptive fields of individual neurons (Schummers et al., 2008). In addition, astrocytes, either individually or in association with microglia, could control OPM formation during developmental plasticity (Sections 6 and 8), as well as during adulthood (López-Hidalgo and Schummers, 2014). Gliotransmission has also been implicated in the sensory modulation of rhythmic activity of the central pattern generators responsible for breathing and chewing (Kadala et al., 2015; Del Negro et al., 2018). Stimulation of glutamatergic sensory fibers that project onto the trigeminal sensory-motor circuit for mastication can indeed activate astrocytes, triggering  $\text{Ca}^{2+}$ -dependent release of the astrocyte-specific  $\text{Ca}^{2+}$ -binding protein S100 $\beta$ . In turn, this results in a reduction of  $\text{Ca}^{2+}$  in the ECS which promotes rhythmic bursting at frequencies that are compatible with those observed for voluntary chewing (Morquette et al., 2015; Condamine et al., 2018). In addition, medial basal hypothalamic astrocytes can control feeding behavior bidirectionally by purinergic gliotransmission, regulating appetite under both favorable and unfavorable conditions (Yang et al., 2015). Thus, astrocytes critically regulate food intake which is a crucial behavior for energy homeostasis, preventing energy deficits or surfeit.

In the general framework of neuronal network theory, glia-mediated variations of network activity may ultimately be linked to modulations of the balance between excitation (E) and inhibition (I). This possibility arises by the analysis of the few currently available models of neuron-glia network dynamics which come in different flavors, as a result of the combination of different E-I network configurations with different choices of neuronal (and synaptic) models and astrocytic signaling pathways (Savin et al., 2009; Ullah et al., 2009; Volman et al., 2013; Savtchenko and Rusakov, 2014; Garnier et al., 2016). Nonetheless, all these models eventually envisage an effect of glial signaling in terms of a modulation of synaptic drive, either at excitatory (Savin et al., 2009; Savtchenko and Rusakov, 2014) or at excitatory and inhibitory synapses (Ullah et al., 2009; Volman et al., 2013; Garnier et al., 2016), which accounts for emergence of a variety of network activities. Similar observations may also be made by other models which consider different scenarios of gliotransmission, such as short-term modulation of E-to-I synaptic connections by glutamatergic or purinergic gliotransmission (Savtchenko and Rusakov, 2014), or homeostatic upregulation of excitation by glial  $\text{TNF}\alpha$ , although the latter scenario could also account for emergence of paroxysmal activity in various pathological conditions (Volman et al., 2013). Significantly, these models identify the spatial extent of gliotransmission as a key factor for the regulation of glia-mediated episodes of increased network activity (Volman et al., 2013; Savtchenko and Rusakov, 2014). In these models, the transient depression of synapses within an astrocytic anatomical domain correlates with a decrease of neuronal firing and synchronization, which is larger for larger astrocytic domains (Savtchenko and Rusakov, 2014). In the context of oscillatory network dynamics, this could then account for reductions of specific frequencies of oscillation with possibly multiple functional and behavioral consequences (Buzsáki and Draguhn, 2004). For example, selective, inducible inhibition of vesicular release from brain-wide astrocytic domains was shown to reduce the power spectrum of gamma frequency oscillations ( $\sim 25\text{--}80$  Hz) in living mice and, ultimately a performance deficit in novel object recognition tasks (Lee et al., 2014)(Lee et al., 2014). Another possible pathway that has indeed been linked with memory consolidation during fear conditioning is gliotransmitter

release by connexin 43 (Cx43) hemichannels (Stehberg et al., 2012).

Another illustration is the putative involvement of glia in sleep regulation. Many factors involved in the regulation of sleep are still unknown, yet mounting evidence indicate glia as integral component of the neurobiological substrate underpinning circadian and homeostatic mechanisms believed to regulate sleep (Halassa and Haydon, 2010; Jackson, 2011; Petit and Magistretti, 2016). In the suprachiasmatic nucleus of the hypothalamus where the main circadian clock resides, neuron-glia networks undergo rhythmic structural reorganization during the 24 h light/day cycle, which is deemed necessary for appropriate adjustment of the circadian clock to this cycle (Becquet et al., 2008). Gene transcription analysis revealed that several genes that control diverse aspects of glial morphology and function are differentially expressed during sleep vs. wakefulness, independently of the behavioral state (Cirelli et al., 2004; Bellesi et al., 2013, 2015). In parallel, compelling evidence also link astrocytes with homeostatic mechanisms of sleep. Extracellular accumulation during wakefulness of adenosine originating from astrocytic ATP, has been implicated in the progressive need of sleep (also dubbed “sleep pressure”) (Pascual et al., 2005; Schmitt et al., 2012; Bjorness et al., 2016), as well as in cognitive impairment associated with prolonged periods of sleep deprivation (Halassa et al., 2009; Florian et al., 2011). Spatially-confined extracellular glutamate increases that correlate with astrocytic  $\text{Ca}^{2+}$  activation could also promote putative sleep states at the level of individual brain regions, triggering there slow wave activity characteristic of non-random eye movement (NREM) sleep (Nir et al., 2011; Vyazovskiy et al., 2011; Poskanzer and Yuste, 2016).

## 10 Conclusions

The variety of experimental and theoretical arguments presented in the previous sections outlines a role for glial cells in the control of multiple mechanisms underpinning information processing by the brain. Although this role can in principle extend to the whole brain, in its many spatial and temporal scales of activity, the biophysical considerations made here also support the idea of a strong regional specialization for neuron-glia interactions and function, ensuing from evolutionary, developmental and environmental (activity-dependent) factors (Sections 2 and 4). The multiplicity of chemical signals that glial cells can sense and generate in response to neuronal activity also make these cells potential signaling hubs in the neuropil. Accordingly, glial cells could sense different aspects of neuronal activity, such as electrical signals, synaptic neurotransmitters, structural changes, energy demands and vascular metabolites, thanks to a rich cassette of membrane receptors, channels, transporters, and pumps (Sections 4 and 5). In turn, they can release multiple molecules with different, and potentially multiple signaling roles. Gliotransmitters like ATP and glutamate for example, could also be involved both in metabolic (Section 7) and in inflammatory processes (Section 8). Conversely, glial release of the proinflammatory cytokine  $\text{TNF}\alpha$  could be permissive towards gliotransmission and homeostatic mechanisms of plasticity (Section 8). While signaling multiplexing may be guaranteed by spatially and temporally distinct scales of action of different signals impinging on and mediated by glial cells, it should be emphasized that these different signals could also be interdependent, with potential for emergence of nontrivial correlations in the neural code (Section 9).

It may be objected that the majority of the arguments exposed here link glial signaling to neurobiological mechanisms that are generally regarded to underpin computations performed by the brain but for which a definitive experimental connection with higher brain functions is still missing (Section 10). This however does not weaken the original motivation of our hitherto discussion, that is the conceptualization of a Neuron-Glia paradigm of the brain; rather it should prompt readers to revise our knowledge of the literature from a different, potentially deeper perspective, inclusive of glial signaling. We indeed predict that many contradicting

results in neuroscience and poorly understood signaling pathways of the brain could benefit from novel investigations driven by a theoretical understanding of the possible functional relevance of neuron-glia interactions in brain physiology.


## Figure captions

**Figure 1.** Interaction of astrocytes with neuropil and vasculature. Simplified illustration of the main pathways of interaction of astrocytes with neurons at glutamatergic synapses. (*Yellow pathway*) *Calcium signaling.* Glutamate (Glu) spillover from the synaptic cleft targets metabotropic receptors (mGluR) on perisynaptic astrocytic processes which trigger inositol 1,4,5-trisphosphate (IP<sub>3</sub>)-mediated calcium (Ca<sup>2+</sup>) release from the cell’s endoplasmic reticulum (ER). IP<sub>3</sub> can also diffuse intra- and inter-cellularly by gap junction channels (GJCs), allowing regenerative propagation of Ca<sup>2+</sup> signaling to other regions of the cell or neighboring astrocytes. Intracellular Ca<sup>2+</sup> transients in astrocytic processes can also occur by spontaneous opening of transient receptor potential channels TRPA<sub>1</sub> (Shigetomi et al., 2013), or by neurotransmitter-gated ionotropic receptor channels (iRCs) such as AMPA and NMDA channels and purine-bound P2X channels (Newman, 2005; Suadicani et al., 2006; Hamilton et al., 2008; Palygin et al., 2010; Lind et al., 2013). Excess Ca<sup>2+</sup> is pumped out of the cell by the sodium Na<sup>+</sup>/Ca<sup>2+</sup> exchanger (NCX) or into the ER by (sarco)endoplasmic reticulum Ca<sup>2+</sup>/ATPase (SERCA) pumps. (*Red pathway*) *Gliotransmission.* Transient cytosolic Ca<sup>2+</sup> increases may trigger release of neuroactive molecules (or “gliotransmitters”) from astrocytes by multiple pathways. These pathways include glutamate release by exocytosis or by bestrophin-1 (Best-1) ion channels (Sahlender et al., 2014), and release of ATP by vesicular exocytosis (not shown) or by purine-permeable channels (PPCs) (Stout et al., 2002; Suadicani et al., 2006; Bowser and Khakh, 2007). Extracellular ATP rapidly degraded into adenosine (Ado) which, along with glutamate of astrocytic origin, can target different receptors on presynaptic and postsynaptic elements, thereby modulating synaptic transmission and neuronal excitability (Savtchouk and Volterra, 2018). Either with glutamate (not shown) or independently, astrocytes can also release D-serine (D-ser), which is the main co-agonist (with glutamate) of postsynaptic NMDA receptors at many synapses (Mothet et al., 2000; Henneberger et al., 2010; Papouin et al., 2012). (*Green pathway*) *Cytokine signaling* Among possible signals mediated by astrocytes is also the release of cytokine tumor necrosis factor alpha (TNF $\alpha$ ) by Ca<sup>2+</sup>-dependent TNF $\alpha$ -converting enzyme (TACE) (Bezzi et al., 2001). This cytokine – which is also present in the extracellular milieu at constitutive concentrations < 100 – 200 pM –, may target both astrocytic and synaptic TNF receptors (TNFR1). In the astrocyte this modulates glutamatergic gliotransmission (via a yet-unresolved pathway) (Santello and Volterra, 2012); at synapses it modulates instead insertion (endocytosis) of postsynaptic AMPA (GABA) receptors (not shown) (Stellwagen et al., 2005; Pribiag and Stellwagen, 2013; Lewitus et al., 2014). (*Turquoise pathway*) *Glutamate–glutamine cycle.* Astrocytic excitatory amino acid transporters (EAATs) are responsible for the uptake of a large fraction of glutamate at the synapse. Glutamate is converted into glutamine by glutamine synthetase (GS) and shuttled back to neurons by Na<sup>+</sup>-coupled neutral amino acid transporters (SNATs) (Mackenzie and Erickson, 2004). Once in the neuron, glutamine is reconverted to glutamate by phosphate-activate glutaminase (PAG), and the ensuing glutamate may either be (transiently) consumed by oxidative metabolism of mitochondria, or be used to refill synaptic vesicles. The whole process is coupled with ammonia (NH<sub>3</sub>) homeostasis and also accounts for GABA synthesis from glutamate at inhibitory synapses (not shown), possibly in association with astrocytic uptake by GABA transporter 3 (GAT3) (Rothman et al., 2003; Hertz, 2013). Significantly, expression of astrocytic EAATs and GATs may be modulated by intracellular Ca<sup>2+</sup> (*dashed yellow pathways*) (Mashimo et al., 2010; Shigetomi et al., 2012; Devaraju et al., 2013). (*Blue pathway*) *Potassium buffering.* Astrocytes buffer excess potassium (K<sup>+</sup>) released into the extracellular space as a result of neuronal activity. Excess extracellular K<sup>+</sup> is taken up mostly by inwardly rectifying K<sup>+</sup> channels (Kir4.1) in conjunction with Na<sup>+</sup>/K<sup>+</sup>-ATPase (NKP) pumps and Na<sup>+</sup>-K<sup>+</sup>-2Cl<sup>-</sup> cotransporters (NKCC1), all of which are richly expressed on perisynaptic



astrocytic processes. Potassium ions then travel down their concentration gradient through the astrocyte's cytoplasm, or through gap junction channels to other cells, where they are released at sites of lower extracellular  $K^+$  concentration or shunted to the vascular system (Kofuji and Newman, 2004; Simard and Nedergaard, 2004). The pathway is coupled with the cell's water homeostasis by aquaporin channels (AQP4), as well as with glutamate uptake, since intracellular  $K^+$  controls astrocytic transporters' stoichiometry together with extracellular  $Na^+$  and  $H^+$  (Section 5). (*Purple pathway*). *Lactate shuttle*. Glutamate uptake is accompanied by  $Na^+$  influx into the astrocyte which is counteracted by the action of NKPs. The resulting increase in ADP/ATP ratio triggers anaerobic glucose utilization in astrocytes by glycolysis which may be associated with glycogenolysis and/or with glucose uptake from the circulation by glucose transporter GLUT1. Lactate dehydrogenase (LDH) catalyzes the interconversion of pyruvate (Pyr) from glycolysis into lactate (and vice versa) with concomitant interconversion of NADH and  $NAD^+$ . Lactate (Lac) is then shuttled to neurons through monocarboxylate transporters (mainly MCT1 in astrocytes and MCT2 in neurons), where it can be used as an energy substrate after its reconversion to pyruvate (Allaman et al., 2011). (*Brown pathway*). *pH buffering*. Abundant carbonic anhydrase (CA) in astrocytes converts  $CO_2$  resulting from cellular metabolism (plus  $H_2O$ ) into  $H^+$  and  $HCO_3^-$ . Two  $HCO_3^-$  are transported into the extracellular space along with one  $Na^+$  via the  $Na^+$ -coupled bicarbonate transporter (NBC), thereby increasing the extracellular buffering power. The efflux of  $HCO_3^-$  induces extracellular alkalinization and also occurs by activation of  $GABA_A$  channels as well as by  $Cl^-/HCO_3^-$  exchanger (CIBX). Excess  $H^+$  in astrocytes is extruded via  $Na^+-H^+$  exchanger (NHX). Chloride and bicarbonate homeostasis are also intimately related to cell volume regulation (Section 5). (*Orange pathway*) *Glutathione metabolism*. Astrocytes release glutathione (GSH) in the extracellular space where it is cleaved by the astrocytic ectoenzyme  $\gamma$ -glutamyl transpeptidase ( $\gamma$ GT) to produce cysteinylglycine (CysGly). In turn, CysGly is cleaved by neuronal ectoaminopeptidase N (ApN), forming cysteine (Cys) and glycine (Gly), which serve as precursors for neuronal GSH synthesis (Aoyama et al., 2008). In the astrocyte the reduced form of glutathione (i.e. GSH) is obtained by multiple pathways including recycling of extracellular CysGly, *de novo* synthesis from intracellular glutamate, or precursors of it like glutamine for example (*dashed orange arrow*), or by oxidized glutathione (GSSG) via NADPH-dependent glutathione reductase (GR) (Dringen, 2000). Astrocytic GSSG ensues from oxidative metabolism but may also be linked with the ANLS via  $NADP^+$ -dependent ascorbic acid/vitamin C recycling (not shown, see Castro et al. (2009)). NADPH/ $NADP^+$  can instead be modulated by NADH/ $NAD^+$  either in the cytoplasm or in mitochondria (Ying, 2008), although astrocytic NADPH production seems strongly dependent on pentose phosphate pathway-mediated metabolism of glucose 6-phosphate (glucose-6P) (Dringen, 2000) – the main available form of intracellular glucose obtained from uptake from blood and by glycogenolysis. Shuttling of GSH between astrocytes and neurons is essential in providing precursors for neuronal GSH synthesis and makes astrocytes key players in neuroprotection against oxidative stress. (*Magenta pathway*) *Vascular coupling*.  $Ca^{2+}$  rise in the astrocyte may also stimulate generation of arachidonic acid (AA) from phospholipase  $A_2$  ( $PLA_2$ ), which is converted to prostaglandins (PGs) and epoxyeicosatrienoic acids (EETs) (by cyclooxygenases) to dilate blood vessels through astrocytic endfeet – which are specialized astrocytic processes contacting blood vessels. Alternatively, astrocytic AA passes via endfoot to the smooth muscle surrounding capillaries, and there it is converted into 20-hydroxy-eicosatetraenoic acid (20-HETE) (by  $\omega$ -hydroxylase) which constricts vessels (Attwell et al., 2010). ClC: chloride channel;  $NAD^+$  ( $NADH$ ): oxidized (reduced) nicotinamide adenine dinucleotide;  $NADP^+$  ( $NADPH$ ): oxidized (reduced) nicotinamide adenine dinucleotide phosphate; GPhos: glycogen phosphorylase; GGS/PTG: glycogen synthase/UTP–glycogen–phosphate uridylyltransferase; SACIC: swell-activated chloride channel; VGCC: voltage-gated  $Ca^{2+}$  channel; VGKC: voltage-gated  $K^+$  channel; VGNC: voltage-gated  $Na^+$  channel.

## Figures



**Figure 1.** Interaction of astrocytes with neuropil and vasculature.

## References

- Abbott, L. F., DePasquale, B., and Memmesheimer, R.-M. (2016). Building functional networks of spiking model neurons. *Nature Neuroscience*, 19(3):350.
- Abbott, L. F. and Nelson, S. B. (2000). Synaptic plasticity: taming the beast. *Nature*, 3:1178–1183.
- Aboitiz, F., Scheibel, A. B., Fisher, R. S., and Zaidel, E. (1992). Fiber composition of the human corpus callosum. *Brain Research*, 598(1-2):143–153.
- Agulhon, C., Petravicz, J., McMullen, A. B., Sweger, E. J., Minton, S. K., Taves, S. R., Casper, K. B., Fiacco, T. A., and McCarthy, K. D. (2008). What is the role of astrocyte calcium in neurophysiology? *Neuron*, 59:932–946.
- Alberdi, E., Sánchez-Gómez, M. V., Marino, A., and Matute, C. (2002).  $\text{Ca}^{2+}$  influx through AMPA or kainate receptors alone is sufficient to initiate excitotoxicity in cultured oligodendrocytes. *Neurobiology of Disease*, 9(2):234–243.
- Allam, S. L., Ghaderi, V. S., Bouteiller, J.-M. C., Legendre, A., Nicolas, A., Greget, R., Bischoff, S., Baudry, M., and Berger, T. W. (2012). A computational model to investigate astrocytic glutamate uptake influence on synaptic transmission and neuronal spiking. *Frontiers in Computational Neuroscience*, 6:70.
- Allaman, I., Belanger, M., and Magistretti, P. J. (2011). Astrocyte–neuron metabolic relationships: for better and for worse. *Trends in Neurosciences*, 34(2):76–87.
- Allen, N. J. (2013). Role of glia in developmental synapse formation. *Current Opinion in Neurobiology*, 23(6):1027–1033.
- Allen, N. J., Káradóttir, R., and Attwell, D. (2005). A preferential role for glycolysis in preventing the anoxic depolarization of rat hippocampal area CA1 pyramidal cells. *Journal of Neuroscience*, 25(4):848–859.
- Alvarellos-González, A., Pazos, A., and Porto-Pazos, A. B. (2012). Computational models of neuron-astrocyte interactions lead to improved efficacy in the performance of neural networks. *Computational and mathematical methods in medicine*, 2012:476324.
- Amzica, F., Massimini, M., and Manfredi, A. (2002). Spatial buffering during slow and paroxysmal sleep oscillations in cortical networks of glial cells in vivo. *Journal of Neuroscience*, 22(3):1042–1053.
- Angulo, M. C., Le Meur, K., Kozlov, A. S., Charpak, S., and Audinat, E. (2008). GABA, a forgotten gliotransmitter. *Progress in Neurobiology*, 86(3):297–303.
- Aoyama, K., Watabe, M., and Nakaki, T. (2008). Regulation of neuronal glutathione synthesis. *Journal of pharmacological sciences*, 108(3):227–238.
- Arancibia-Carcamo, I. L., Ford, M. C., Cossell, L., Ishida, K., Tohyama, K., and Attwell, D. (2017). Node of Ranvier length as a potential regulator of myelinated axon conduction speed. *eLife*, 6.
- Araque, A., Carmignoto, G., Haydon, P. G., Oliet, S. H. R., Robitaille, R., and Volterra, A. (2014). Gliotransmitters travel in time and space. *Neuron*, 81(4):728–739.

- Araque, A., Parpura, V., Sanzgiri, R. P., and Haydon, P. G. (1998). Glutamate-dependent astrocyte modulation of synaptic transmission between cultured hippocampal neurons. *Eur. J. Neurosci.*, 10:2129–2142.
- Araque, A., Parpura, V., Sanzgiri, R. P., and Haydon, P. G. (1999). Tripartite synapses: glia, the unacknowledged partner. *Trends Neurosci.*, 23(5):208–215.
- Assaf, Y. and Pasternak, O. (2008). Diffusion tensor imaging (DTI)-based white matter mapping in brain research: a review. *Journal of molecular neuroscience*, 34(1):51–61.
- Aston-Jones, G., Rajkowski, J., and Cohen, J. (1999). Role of locus coeruleus in attention and behavioral flexibility. *Biological Psychiatry*, 46(9):1309–1320.
- Attwell, D., Buchan, A. M., Charpak, S., Lauritzen, M., MacVicar, B. A., and Newman, E. A. (2010). Glial and neuronal control of brain blood flow. *Nature*, 468(7321):232–243.
- Attwell, D. and Gibb, A. (2005). Neuroenergetics and the kinetic design of excitatory synapses. *Nature Reviews Neuroscience*, 6(11):841.
- Attwell, D. and Laughlin, S. B. (2001). An energy budget for signaling in the grey matter of the brain. *Journal of Cerebral Blood Flow & Metabolism*, 21(10):1133–1145.
- Averbeck, B. B., Latham, P. E., and Pouget, A. (2006). Neural correlations, population coding and computation. *Nature reviews neuroscience*, 7(5):358.
- Bains, J. S. and Oliet, S. H. R. (2007). Glia: they make your memories stick! *Trends Neurosci.*, 30(8):417–424.
- Barbour, B. (2001). An evaluation of synapse independence. *J. Neurosci.*, 21(20):7969–7984.
- Barres, B. (2008). The mystery and magic of glia: a perspective on their roles in health and disease. *Neuron*, 60(3):430–440.
- Bear, M. F. and Malenka, R. C. (1994). Synaptic plasticity: LTP and LTD. *Current Opinion in Neurobiology*, 4(3):389–399.
- Beattie, E., Stellwagen, D., Morishita, W., Bresnahan, J., Ha, B., Von Zastrow, M., Beattie, M., and Malenka, R. (2002). Control of synaptic strength by glial TNF $\alpha$ . *Science*, 295(5563):2282–2285.
- Beck, J. M., Ma, W. J., Kiani, R., Hanks, T., Churchland, A. K., Roitman, J., Shadlen, M. N., Latham, P. E., and Pouget, A. (2008). Probabilistic population codes for Bayesian decision making. *Neuron*, 60(6):1142–1152.
- Becquet, D., Girardet, C., Guillaumond, F., François-Bellan, A.-M., and Bosler, O. (2008). Ultrastructural plasticity in the rat suprachiasmatic nucleus. Possible involvement in clock entrainment. *Glia*, 56(3):294–305.
- Bekar, L. K., He, W., and Nedergaard, M. (2008). Locus coeruleus  $\alpha$ -adrenergic-mediated activation of cortical astrocytes in vivo. *Cerebral Cortex*, 18(12):2789–2795.
- Bélanger, M., Allaman, I., and Magistretti, P. J. (2011). Brain energy metabolism: focus on astrocyte-neuron metabolic cooperation. *Cell Metabolism*, 14(6):724–738.
- Bellesi, M., de Vivo, L., Tononi, G., and Cirelli, C. (2015). Effects of sleep and wake on astrocytes: clues from molecular and ultrastructural studies. *BMC Biology*, 13(1):66.

- Bellesi, M., Pfister-Genskow, M., Maret, S., Keles, S., Tononi, G., and Cirelli, C. (2013). Effects of sleep and wake on oligodendrocytes and their precursors. *Journal of Neuroscience*, 33(36):14288–14300.
- Bengtsson, S. L., Nagy, Z., Skare, S., Forsman, L., Forssberg, H., and Ullén, F. (2005). Extensive piano practicing has regionally specific effects on white matter development. *Nature Neuroscience*, 8(9):1148.
- Bennett, M., Farnell, L., and Gibson, W. (2008). A quantitative model of cortical spreading depression due to purinergic and gap-junction transmission in astrocyte networks. *Biophys. J.*, 95(12):5648–5660.
- Benz, B., Grima, G., and Do, K. Q. (2004). Glutamate-induced homocysteic acid release from astrocytes: possible implication in glia-neuron signaling. *Neuroscience*, 124(2):377–386.
- Bernardinelli, Y., Muller, D., and Nikonenko, I. (2014a). Astrocyte-synapse structural plasticity. *Neural Plasticity*, 2014:23215.
- Bernardinelli, Y., Randall, J., Janett, E., Nikonenko, I., König, S., Jones, E. V., Flores, C. E., Murai, K. K., Bochet, C. G., Holtmaat, A., and Muller, D. (2014b). Activity-dependent structural plasticity of perisynaptic astrocytic domains promotes excitatory synapse stability. *Current Biology*, 24(15):1679–1688.
- Berridge, C. W. (2008). Noradrenergic modulation of arousal. *Brain Research Reviews*, 58(1):1–17.
- Berridge, C. W. and Waterhouse, B. D. (2003). The locus coeruleus–noradrenergic system: modulation of behavioral state and state-dependent cognitive processes. *Brain Research Reviews*, 42(1):33–84.
- Berridge, M. J., Lipp, P., and Bootman, M. D. (2000). The versatility and universality of calcium signalling. *Nat. Rev. Mol. Cell. Biol.*, 1:11–21.
- Berry, M., Ibrahim, M., Carlile, J., Ruge, F., Duncan, A., and Butt, A. M. (1995). Axon-glia relationships in the anterior medullary velum of the adult rat. *Journal of Neurocytology*, 24(12):965–983.
- Bezzi, P., Domercq, M., Brambilla, L., Galli, R., Schols, D., De Clercq, E., Vescovi, A., Bagetta, G., Kollias, G., Meldolesi, J., and Volterra, A. (2001). CXCR4-activated astrocyte glutamate release via TNF $\alpha$ : amplification by microglia triggers neurotoxicity. *Nat. Neurosci.*, 4(7):702–710.
- Bezzi, P. and Volterra, A. (2001). A neuron-glia signalling network in the active brain. *Curr. Opinion Neurobiol.*, 11:387–394.
- Bialek, W. and Rieke, F. (1992). Reliability and information transmission in spiking neurons. *Trends in Neurosciences*, 15(11):428–434.
- Biber, K., Laurie, D. J., Berthele, A., Sommer, B., Tölle, T. R., Gebicke-Härter, P.-J., Van Calker, D., and Boddeke, H. W. G. M. (1999). Expression and signaling of group I metabotropic glutamate receptors in astrocytes and microglia. *Journal of Neurochemistry*, 72(4):1671–1680.

- Bienenstock, E., Cooper, L., and Munro, P. (1982). Theory for the development of neuron selectivity: orientation specificity and binocular interaction in visual cortex. *J. Neurosci.*, 2(1):32–48.
- Bindocci, E., Savtchouk, I., Liaudet, N., Becker, D., and Carriero, G. and Volterra, A. (2017). Three-dimensional  $\text{Ca}^{2+}$  imaging advances understanding of astrocyte biology. *Science*, 356:6339.
- Bjorness, T. E., Dale, N., Mettlach, G., Sonneborn, A., Sahin, B., Fienberg, A. A., Yanagisawa, M., Bibb, J. A., and Greene, R. W. (2016). An adenosine-mediated glial-neuronal circuit for homeostatic sleep. *Journal of Neuroscience*, 36(13):3709–3721.
- Bonansco, C., Couve, A., Perea, G., Ferradas, C. A., Roncagliolo, M., and Fuenzalida, M. (2011). Glutamate released spontaneously from astrocytes sets the threshold for synaptic plasticity. *Eur. J. Neurosci.*, 33:1483–1492.
- Borst, J. G. G. and Sakmann, B. (1999). Effect of changes in action potential shape on calcium currents and transmitter release in a calyx-type synapse of the rat auditory brainstem. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*, 354(1381):347–355.
- Boudkkazi, S., Fronzaroli-Molinieres, L., and Debanne, D. (2011). Presynaptic action potential waveform determines cortical synaptic latency. *The Journal of Physiology*, 589(5):1117–1131.
- Bourque, C. W. and Renaud, L. P. (1990). Electrophysiology of mammalian magnocellular vasopressin and oxytocin neurosecretory neurons. *Frontiers in Neuroendocrinology*, 11(3):183–212.
- Bouzier-Sore, A.-K. and Pellerin, L. (2013). Unraveling the complex metabolic nature of astrocytes. *Frontiers in Cellular Neuroscience*, 7:179.
- Bowser, D. N. and Khakh, B. S. (2007). Vesicular ATP is the predominant cause of intercellular calcium waves in astrocytes. *J. Gen. Physiol.*, 129(6):485–491.
- Brea, J., Senn, W., and Pfister, J.-P. (2013). Matching recall and storage in sequence learning with spiking neural networks. *The Journal of Neuroscience*, 33(23):9565–9575.
- Buonomano, D. V. and Merzenich, M. M. (1998). Cortical plasticity: from synapses to maps. *Annual review of neuroscience*, 21(1):149–186.
- Bushong, E. A., Martone, M. E., Jones, Y. Z., and Ellisman, M. H. (2002). Protoplasmic astrocyte in CA1 stratum radiatum occupy separate anatomical domains. *J. Neurosci.*, 22(1):183–192.
- Butt, A. M. (2006). Neurotransmitter-mediated calcium signalling in oligodendrocyte physiology and pathology. *Glia*, 54(7):666–675.
- Butt, A. M. (2013). *Neuroglia*, chapter Structure and function of oligodendrocytes, pages 62–73. Oxford University Press.
- Butt, A. M., Ibrahim, M., and Berry, M. (1998). Axon-myelin sheath relations of oligodendrocyte unit phenotypes in the adult rat anterior medullary velum. *J. Neurocytol.*, 27(4):259–269.
- Butt, A. M., Pugh, M., Hubbard, P., and James, G. (2004). Functions of optic nerve glia: axoglial signalling in physiology and pathology. *Eye*, 18(11):1110.

- Buzsáki, G. (2010). Neural syntax: cell assemblies, synapsembles, and readers. *Neuron*, 68(3):362–385.
- Buzsáki, G. and Draguhn, A. (2004). Neuronal oscillations in cortical networks. *Science*, 304(5679):1926–1929.
- Calvetti, D. and Somersalo, E. (2011). Dynamic activation model for a glutamatergic neurovascular unit. *Journal of Theoretical Biology*, 274(1):12–29.
- Calvetti, D. and Somersalo, E. (2012). Ménage à trois: the role of neurotransmitters in the energy metabolism of astrocytes, glutamatergic, and GABAergic neurons. *Journal of Cerebral Blood Flow & Metabolism*, 32(8):1472–1483.
- Carmona, M. A., Murai, K. K., Wang, L., Roberts, A. J., and Pasquale, E. B. (2009). Glial ephrin-A3 regulates hippocampal dendritic spine morphology and glutamate transport. *Proceedings of the National Academy of Sciences*, 106(30):12524–12529.
- Carr, C. E. and Konishi, M. (1990). A circuit for detection of interaural time differences in the brain stem of the barn owl. *Journal of Neuroscience*, 10(10):3227–3246.
- Carr, C. E. and Soares, D. (2002). Evolutionary convergence and shared computational principles in the auditory system. *Brain, Behavior and Evolution*, 59(5-6):294–311.
- Carr, C. E., Soares, D., Parameshwaran, S., and Perney, T. (2001). Evolution and development of time coding systems. *Current Opinion in Neurobiology*, 11(6):727–733.
- Castro, M. A., Beltrán, F. A., Brauchi, S., and Concha, I. I. (2009). A metabolic switch in brain: glucose and lactate metabolism modulation by ascorbic acid. *Journal of neurochemistry*, 110(2):423–440.
- Chao, T. I., Rickmann, M., and Wolff, J. R. (2002). The synapse-astrocyte boundary: an anatomical basis for an integrative role of glia in synaptic transmission. In Volterra, A., Magistretti, P. J., and Haydon, P. G., editors, *The Tripartite Synapse: Glia in Synaptic Transmission*, chapter 1, pages 3–23. Oxford University Press, New York.
- Chen, N., Sugihara, H., Sharma, J., Perea, G., Petravicz, J., Le, C., and Sur, M. (2012). Nucleus basalis enabled stimulus specific plasticity in the visual cortex is mediated by astrocytes. *Proc. Natl. Acad. Sci. USA*, 109(41):E2832E2841.
- Chesler, M. (2003). Regulation and modulation of pH in the brain. *Physiological Reviews*, 83(4):1183–1221.
- Chever, O., Djukic, B., McCarthy, K. D., and Amzica, F. (2010). Implication of Kir4.1 channel in excess potassium clearance: an in vivo study on anesthetized glial-conditional Kir4.1 knockout mice. *Journal of Neuroscience*, 30(47):15769–15777.
- Chever, O., Pannasch, U., Ezan, P., and Rouach, N. (2014). Astroglial connexin 43 sustains glutamatergic synaptic efficacy. *Phil. Trans. R. Soc. B*, 369(1654):20130596.
- Churchill, L., Rector, D. M., Yasuda, K., Fix, C., Rojas, M. J., Yasuda, T., and Krueger, J. M. (2008). Tumor necrosis factor  $\alpha$ : activity dependent expression and promotion of cortical column sleep in rats. *Neuroscience*, 156(1):71–80.
- Cirelli, C., Gutierrez, C. M., and Tononi, G. (2004). Extensive and divergent effects of sleep and wakefulness on brain gene expression. *Neuron*, 41(1):35–43.



- Clarke, L. E. and Barres, B. A. (2013). Emerging roles of astrocytes in neural circuit development. *Nature Reviews Neuroscience*, 14(5):311–321.
- Clements, J. D. (1996). Transmitter timecourse in the synaptic cleft: its role in central synaptic function. *Trends Neurosci.*, 19(5):163–171.
- Coco, S., Calegari, F., Pravettoni, E., Pozzi, D., Taverna, E., Rosa, P., Matteoli, M., and Verderio, C. (2003). Storage and release of ATP from astrocytes in culture. *J. Biol. Chem.*, 278(2):1354–1362.
- Colombo, J. A., Härtig, W., Lipina, S., and Bons, N. (1998). Astroglial interlaminar processes in the cerebral cortex of prosimians and Old World monkeys. *Anatomy and embryology*, 197(5):369–376.
- Colombo, J. A. and Reisin, H. D. (2004). Interlaminar astroglia of the cerebral cortex: a marker of the primate brain. *Brain research*, 1006(1):126–131.
- Condamine, S., Lavoie, R., Verdier, D., and Kolta, A. (2018). Functional rhythmogenic domains defined by astrocytic networks in the trigeminal main sensory nucleus. *Glia*, 66(2):311–326.
- Cornell-Bell, A. H., Finkbeiner, S. M., Cooper, M. S., and Smith, S. J. (1990). Glutamate induces calcium waves in cultured astrocytes: long-range glial signaling. *Science*, 247(4941):470–473.
- Covelo, A. and Araque, A. (2018). Neuronal activity determines distinct gliotransmitter release from a single astrocyte. *eLife*, 7:e32237.
- Cui, Y., Yang, Y., Ni, Z., Dong, Y., Cai, G., Foncelle, A., Ma, S., Sang, K., Tang, S., Li, Y., Berry, H., Shengzi, W., and Hailan, H. (2018). Astroglial Kir4.1 in the lateral habenula drives neuronal bursts in depression. *Nature*, 554(7692):323.
- Cunha, R. and Ribeiro, J. (2000). ATP as a presynaptic modulator. *Life Sciences*, 68:119–137.
- D’Ambrosio, R., Gordon, D. S., and Winn, H. R. (2002). Differential role of KIR channel and Na<sup>+</sup>/K<sup>+</sup>-pump in the regulation of extracellular K<sup>+</sup> in rat hippocampus. *Journal of Neurophysiology*, 87(1):87–102.
- Danbolt, N. C. (2001). Glutamate uptake. *Progress Neurobiol.*, 65:1–105.
- Danbolt, N. C., Lehre, K. P., Dehnes, Y., and Ullensvang, K. (2002). Transporters for synaptic transmitter on the glial cell plasma membrane. In Volterra, A., Magistretti, P. J., and Haydon, P. G., editors, *The Tripartite Synapse: Glia in Synaptic Transmission*, chapter 1, pages 47–61. Oxford University Press, New York.
- Dani, J. W., Chernjavsky, A., and Smith, S. J. (1992). Neuronal activity triggers calcium waves in hippocampal astrocyte networks. *Neuron*, 8:429–440.
- De Pittà, M. and Brunel, N. (2016). Modulation of synaptic plasticity by glutamatergic gliotransmission: A modeling study. *Neural Plasticity*, page 7607924.
- De Pittà, M., Brunel, N., and Volterra, A. (2015). Astrocytes: orchestrating synaptic plasticity? *Neuroscience*.
- De Pittà, M., Goldberg, M., Volman, V., Berry, H., and Ben-Jacob, E. (2009). Glutamate-dependent intracellular calcium and IP<sub>3</sub> oscillating and pulsating dynamics in astrocytes. *J. Biol. Phys.*, 35:383–411.

- De Pittà, M., Volman, V., Berry, H., and Ben-Jacob, E. (2011). A tale of two stories: astrocyte regulation of synaptic depression and facilitation. *PLoS Comput. Biol.*, 7(12):e1002293.
- De Pittà, M., Volman, V., Levine, H., Pioggia, G., De Rossi, D., and Ben-Jacob, E. (2008). Coexistence of amplitude and frequency modulations in intracellular calcium dynamics. *Phys. Rev. E*, 77(3):030903(R).
- Debanne, D., Campanac, E., Bialowas, A., Carlier, E., and Alcaraz, G. (2011). Axon physiology. *Physiological Reviews*, 91(2):555–602.
- Deco, G., Jirsa, V. K., and McIntosh, A. R. (2011). Emerging concepts for the dynamical organization of resting-state activity in the brain. *Nature Reviews Neuroscience*, 12(1):43.
- Deitmer, J. W. and Rose, C. R. (1996). pH regulation and proton signalling by glial cells. *Progress in Neurobiology*, 48(2):73–103.
- Del Castillo, J. and Katz, B. (1954). Quantal components of the end-plate potential. *J. Physiol.*, 124:560–573.
- Del Negro, C. A., Funk, G. D., and Feldman, J. L. (2018). Breathing matters. *Nature Reviews Neuroscience*, 19(6):351–367.
- Deng, Q., Terunuma, M., Fellin, T., Moss, S. J., and Haydon, P. (2011). Astrocytic activation of A1 receptors regulates the surface expression of NMDA receptors through a Src kinase dependent pathway. *Glia*, 59:1084–1093.
- Dere, E., De Souza-Silva, M. A., Frisch, C., Teubner, B., Söhl, G., Willecke, K., and Huston, J. P. (2003). Connexin30-deficient mice show increased emotionality and decreased rearing activity in the open-field along with neurochemical changes. *European Journal of Neuroscience*, 18(3):629–638.
- Devaraju, P., Sun, M.-Y., Myers, T. L., Lauderdale, K., and Fiacco, T. A. (2013). Astrocytic group I mGluR-dependent potentiation of astrocytic glutamate and potassium uptake. *Journal of Neurophysiology*, 109(9):2404–2414.
- Di Castro, M., Chuquet, J., Liaudet, N., Bhaukaurally, K., Santello, M., Bouvier, D., Tiret, P., and Volterra, A. (2011). Local  $\text{Ca}^{2+}$  detection and modulation of synaptic release by astrocytes. *Nat. Neurosci.*, 14:12761284.
- Diamond, M. C., Scheibel, A. B., Murphy Jr., G. M., and Harvey, T. (1985). On the brain of a scientist: Albert Einstein. *Experimental neurology*, 88(1):198–204.
- Dienel, G. A. (2012). Fueling and imaging brain activation. *ASN neuro*, 4(5):AN20120021.
- Diering, G. H., Nirujogi, R. S., Roth, R. H., Worley, P. F., Pandey, A., and Huganir, R. L. (2017). Homer1a drives homeostatic scaling-down of excitatory synapses during sleep. *Science*, 355(6324):511–515.
- Dimou, L., Simon, C., Kirchhoff, F., Takebayashi, H., and Götz, M. (2008). Progeny of Olig2-expressing progenitors in the gray and white matter of the adult mouse cerebral cortex. *Journal of Neuroscience*, 28(41):10434–10442.
- Ding, F., O'Donnell, J., Thrane, A. S., Zeppenfeld, D., Kang, H., Xie, L., Wang, F., and Nedergaard, M. (2013).  $\alpha$ 1-Adrenergic receptors mediate coordinated  $\text{Ca}^{2+}$  signaling of cortical astrocytes in awake, behaving mice. *Cell calcium*, 54(6):387–394.

- Dittman, J. S., Kreitzer, A. C., and Regehr, W. G. (2000). Interplay between facilitation, depression, and residual calcium at three presynaptic terminals. *J. Neurosci.*, 20(4):1374–1385.
- Dityatev, A. and Rusakov, D. A. (2011). Molecular signals of plasticity at the tetrapartite synapse. *Current Opinion in Neurobiology*, 21(2):353–359.
- Djukic, B., Casper, K. B., Philpot, B. D., Chin, L.-S., and McCarthy, K. D. (2007). Conditional knock-out of Kir4.1 leads to glial membrane depolarization, inhibition of potassium and glutamate uptake, and enhanced short-term synaptic potentiation. *Journal of Neuroscience*, 27(42):11354–11365.
- Doiron, B., Litwin-Kumar, A., Rosenbaum, R., Ocker, G. K., and Josić, K. (2016). The mechanics of state-dependent neural correlations. *Nature Neuroscience*, 19(3):383.
- Dolmetsch, R. E., Lewis, R. S., Goodnow, C. C., and Healy, J. I. (1997). Differential activation of transcription factors induced by  $\text{Ca}^{2+}$  response amplitude and duration. *Nature*, 386(24):855–858.
- Domercq, M., Brambilla, L., Pilati, E., Marchaland, J., Volterra, A., and Bezzi, P. (2006). P2Y1 receptor-evoked glutamate exocytosis from astrocytes: control by tumor necrosis factor- $\alpha$  and prostaglandins. *J. Biol. Chem.*, 281:30684–30696.
- Dringen, R. (2000). Metabolism and functions of glutathione in brain. *Progress in Neurobiology*, 62(6):649–671.
- Dringen, R. (2005). Oxidative and antioxidative potential of brain microglial cells. *Antioxidants & Redox Signaling*, 7(9-10):1223–1233.
- Ecker, A. S., Berens, P., Tolias, A. S., and Bethge, M. (2011). The effect of noise correlations in populations of diversely tuned neurons. *Journal of Neuroscience*, 31(40):14272–14283.
- Eroglu, C. and Barres, B. A. (2010). Regulation of synaptic connectivity by glia. *Nature*, 468(7321):223–231.
- Falk, D., Lepore, F. E., and Noe, A. (2012). The cerebral cortex of Albert Einstein: a description and preliminary analysis of unpublished photographs. *Brain*, 136(4):1304–1327.
- Feldman, D. E. and Brecht, M. (2005). Map plasticity in somatosensory cortex. *Science*, 310(5749):810–815.
- Fellin, T., Ellenbogen, J. M., De Pittà, M., Ben-Jacob, E., and Halassa, M. (2012). Astrocyte regulation of sleep circuits: experimental and modeling perspectives. *Front. Comput. Neurosci.*, 6(65).
- Fellin, T., Pascual, O., Gobbo, S., Pozzan, T., Haydon, P. G., and Carmignoto, G. (2004). Neuronal synchrony mediated by astrocytic glutamate through activation of extrasynaptic NMDA receptors. *Neuron*, 43:729–743.
- Fiacco, T. A. and McCarthy, K. D. (2018). Multiple lines of evidence indicate that gliotransmission does not occur under physiological conditions. *Journal of Neuroscience*, 38(1):3–13.
- Fields, R. D. (2008). White matter in learning, cognition and psychiatric disorders. *Trends in Neurosciences*, 31(7):361–370.

- Fields, R. D. (2009). *The other brain: From dementia to schizophrenia, how new discoveries about the brain are revolutionizing medicine and science*. Simon and Schuster.
- Fields, R. D. (2013). *Neuroglia*, chapter Regulation of myelination by functional activity, pages 573–586. Oxford Univ Press.
- Fields, R. D., Woo, D. H., and Basser, P. J. (2015). Glial regulation of the neuronal connectome through local and long-distant communication. *Neuron*, 86(2):374–386.
- Filosa, A., Paixão, S., Honsek, S. D., Carmona, M. A., Becker, L., Feddersen, B., Gaitanos, L., Rudhard, Y., Schoepfer, R., Klopstock, T., Kullander, K., Rose, C. R., Pasquale, E. B., and Klein, R. (2009). Neuron-glia communication via EphA4/ephrin-A3 modulates LTP through glial glutamate transport. *Nature Neuroscience*, 12(10):1285.
- Filosa, J. A., Morrison, H. W., Iddings, J. A., Du, W., and Kim, K. J. (2016). Beyond neurovascular coupling, role of astrocytes in the regulation of vascular tone. *Neuroscience*, 323:96–109.
- Firth, S. I., Wang, C., and Feller, M. B. (2005). Retinal waves: mechanisms and function in visual system development. *Cell Calcium*, 37(5):425–432.
- FitzHugh, R. (1962). Computation of impulse initiation and saltatory conduction in a myelinated nerve fiber. *Biophysical Journal*, 2(1):11–21.
- Florence, C. M., Baillie, L. D., and Mulligan, S. J. (2012). Dynamic volume changes in astrocytes are an intrinsic phenomenon mediated by bicarbonate ion flux. *PLoS One*, 7(11):e51124.
- Florian, C., Vecsey, C. G., Halassa, M. M., Haydon, P. G., and Abel, T. (2011). Astrocyte-derived adenosine and A1 receptor activity contribute to sleep loss-induced deficits in hippocampal synaptic plasticity and memory in mice. *Journal of Neuroscience*, 31(19):6956–6962.
- Fontainhas, A. M., Wang, M., Liang, K. J., Chen, S., Mettu, P., Damani, M., Fariss, R. N., Li, W., and Wong, W. T. (2011). Microglial morphology and dynamic behavior is regulated by ionotropic glutamatergic and GABAergic neurotransmission. *PLoS One*, 6(1):e15973.
- Ford, M. C., Alexandrova, O., Cossell, L., Stange-Marten, A., Sinclair, J., Kopp-Scheinflug, C., Pecka, M., Attwell, D., and Grothe, B. (2015). Tuning of ranvier node and internode properties in myelinated axons to adjust action potential timing. *Nature Communications*, 6:8073.
- Fortune, E. S. and Rose, G. J. (2001). Short-term synaptic plasticity as a temporal filter. *Trends Neurosci.*, 24(7):381–385.
- Fossat, P., Turpin, F. R., Sacchi, S., Dulong, J., Shi, T., Rivet, J., Sweedler, J. V., Pollegioni, L., Millan, M. J., Oliet, S. H. R., and Mothet, J. (2012). Glial D-serine gates NMDA receptors at excitatory synapses in prefrontal cortex. *Cerebral Cortex*, 22(3):595–606.
- Fourcaud-Trocmé, N., Hansel, D., van Vreeswijk, C., and Brunel, N. (2003). How spike generation mechanisms determine the neuronal response to fluctuating inputs. *J. Neurosci.*, 23(37):11628–11640.
- Franze, K., Grosche, J., Skatchkov, S. N., Schinkinger, S., Foja, C., Schild, D., Uckermann, O., Travis, K., Reichenbach, A., and Guck, J. (2007). Müller cells are living optical fibers in the vertebrate retina. *Proceedings of the National Academy of Sciences*, 104(20):8287–8292.

- Frisch, C., Theis, M., De Souza-Silva, M. A., Dere, E., Söhl, G., Teubner, B., Namestkova, K., Willecke, K., and Huston, J. P. (2003). Mice with astrocyte-directed inactivation of connexin43 exhibit increased exploratory behaviour, impaired motor capacities, and changes in brain acetylcholine levels. *European Journal of Neuroscience*, 18(8):2313–2318.
- Froemke, R. C. and Dan, Y. (2002). Spike-timing-dependent synaptic modification induced by natural spike trains. *Nature*, 416(6879):433–438.
- Froemke, R. C., Debanne, D., and Bi, G. (2010). Temporal modulation of spike-timing-dependent plasticity. *Frontiers in synaptic neuroscience*, 2(19):1–16.
- Froemke, R. C., Tsay, I. A., Raad, M., Long, J. D., and Dan, Y. (2006). Contribution of individual spikes in burst-induced long-term synaptic modification. *Journal of Neurophysiology*, 95(3):1620–1629.
- Frühbeis, C., Fröhlich, D., Kuo, W. P., Amphornrat, J., Thilemann, S., Saab, A. S., Kirchhoff, F., Möbius, W., Goebbels, S., Nave, K., et al. (2013). Neurotransmitter-triggered transfer of exosomes mediates oligodendrocyte–neuron communication. *PLoS Biology*, 11(7):e1001604.
- Funahashi, S., Bruce, C. J., and Goldman-Rakic, P. S. (1989). Mnemonic coding of visual space in the monkey’s dorsolateral prefrontal cortex. *Journal of Neurophysiology*, 61(2):331–349.
- Ganmor, E., Segev, R., and Schneidman, E. (2015). A thesaurus for a neural population code. *eLife*, 4:e06134.
- Garnier, A., Vidal, A., and Benali, H. (2016). A theoretical study on the role of astrocytic activity in neuronal hyperexcitability by a novel neuron-glia mass model. *The Journal of Mathematical Neuroscience*, 6(1):10.
- Gerstner, W. and Kistler, W. M. (2002). Mathematical formulations of Hebbian learning. *Biological Cybernetics*, 87(5-6):404–415.
- Giaume, C., Koulakoff, A., Roux, L., Holcman, D., and Rouach, N. (2010). Astroglial networks: a step further in neuroglial and gliovascular interactions. *Nat. Rev. Neurosci.*, 11(2):87–99.
- Gibbs, M. E., O’dowd, B. S., Hertz, L., Robinson, S. R., Sedman, G. L., and Ng, K. T. (1996). Inhibition of glutamine synthetase activity prevents memory consolidation. *Cognitive Brain Research*, 4(1):57–64.
- Gilman, A. G. (1987). G proteins: transducers of receptor-generated signals. *Annual Review of Biochemistry*, 56(1):615–649.
- Goldman, L. and Albus, J. S. (1968). Computation of impulse conduction in myelinated fibers; theoretical basis of the velocity-diameter relation. *Biophysical Journal*, 8(5):596–607.
- Goldman-Rakic, P. S. (1995). Cellular basis of working memory. *Neuron*, 14(3):477–485.
- Gordon, G., Baimoukhametova, D., Hewitt, S., Rajapaksha, W., Fisher, T., and Bains, J. (2005). Norepinephrine triggers release of glial ATP to increase postsynaptic efficacy. *Nat. Neurosci.*, 8(8):1078–1086.
- Gordon, G. R. J., Choi, H. B., Rungta, R. L., Ellis-Davies, G. C. R., and MacVicar, B. A. (2008). Brain metabolism dictates the polarity of astrocyte control over arterioles. *Nature*, 456(7223):745.

- Gordon, G. R. J., Mulligan, S. J., and MacVicar, B. A. (2007). Astrocyte control of the cerebrovasculature. *Glia*, 55(12):1214–1221.
- Götz, M. (2013). *Neuroglia*, chapter Radial glial cells, pages 50–61. Oxford University Press.
- Gourine, A. V., Kasymov, V., Marina, N., Tang, F., Figueiredo, M. F., Lane, S., Teschemacher, A. G., Spyer, M. K., Deisseroth, K., and Kasparov, S. (2010). Astrocytes control breathing through ph-dependent release of ATP. *Science*, 329(5991):571–575.
- Graupner, M. and Brunel, N. (2010). Mechanisms of induction and maintenance of spike-timing dependent plasticity in biophysical synapse models. *Frontiers in Computational Neuroscience*, 4(136):1–19.
- Grosche, J., Kettenmann, H., and Reichenbach, A. (2002). Bergmann glial cells form distinct morphological structures to interact with cerebellar neurons. *J. Neurosci. Res.*, 68:128–149.
- Grosche, J., Matyash, V., Möller, T., Verkhratsky, A., Reichenbach, A., and Kettenmann, H. (1999). Microdomains for neuron-glia interaction: parallel fiber signaling to bergmann glial cells. *Nature*, 2(2):139–143.
- Gruetter, R. (2003). Glycogen: the forgotten cerebral energy store. *Journal of Neuroscience Research*, 74(2):179–183.
- Guinan, J. J., Norris, B. E., and Guinan, S. S. (1972). Single auditory units in the superior olivary complex: II: locations of unit categories and tonotopic organization. *International Journal of Neuroscience*, 4(4):147–166.
- Habbas, S., Santello, M., Becker, D., Stubbe, H., Zappia, G., Liaudet, N., Klaus, F. R., Kollias, G., Fontana, A., Pryce, C. R., Suter, T., and Volterra, A. (2015). Neuroinflammatory TNF $\alpha$  impairs memory via astrocyte signaling. *Cell*, 163(7):1730–1741.
- Haber, M., Zhou, L., and Murai, K. K. (2006). Cooperative astrocyte and dendritic spine dynamics at hippocampal excitatory synapses. *J. Neurosci.*, 26(35):8881–8891.
- Halassa, M. and Haydon, P. (2010). Integrated brain circuits: astrocytic networks modulate neuronal activity and behavior. *Annu. Rev. Physiol.*, 72:335–355.
- Halassa, M. M., Fellin, T., Takano, H., Dong, J.-H., and Haydon, P. G. (2007). Synaptic islands defined by the territory of a single astrocyte. *J. Neurosci.*, 27(24):6473–6477.
- Halassa, M. M., Florian, C., Fellin, T., Munoz, J. R., Lee, S. Y., Abel, T., Haydon, P. G., and Frank, M. G. (2009). Astrocytic modulation of sleep homeostasis and cognitive consequences of sleep loss. *Neuron*, 61(2):213–219.
- Halter, J. A. and Clark Jr., J. W. (1991). A distributed-parameter model of the myelinated nerve fiber. *Journal of Theoretical Biology*, 148(3):345–382.
- Hamilton, N., Vayro, S., Kirchhoff, F., Verkhratsky, A., Robbins, J., Gorecki, D. C., and Butt, A. M. (2008). Mechanisms of ATP- and glutamate-mediated calcium signaling in white matter astrocytes. *Glia*, 56(7):734–749.
- Han, J., Kesner, P., Metna-Laurent, M., Duan, T., Xu, L., Georges, F., Koehl, M., Abrous, D., Mendizabal-Zubiaga, J., Grandes, P., Liu, Q., Bai, G., Wang, W., Xiong, L., Ren, W., Marsicano, G., and Zhang, X. (2012). Acute cannabinoids impair working memory through astroglial CB<sub>1</sub> receptor modulation of hippocampal LTD. *Cell*, 148(5):1039–1050.

- Han, X., Chen, M., Wang, F., Windrem, M., Wang, S., Shanz, S., Xu, Q., Oberheim, N. A., Bekar, L., Betstadt, S., Silva, A. J., Takano, T., Goldman, S. A., and Nedergaard, M. (2013). Forebrain engraftment by human glial progenitor cells enhances synaptic plasticity and learning in adult mice. *Cell Stem Cell*, 12(3):342–353.
- Hansen, B. J., Chelaru, M. I., and Dragoi, V. (2012). Correlated variability in laminar cortical circuits. *Neuron*, 76(3):590–602.
- Harris, J. J., Jolivet, R., and Attwell, D. (2012). Synaptic energy use and supply. *Neuron*, 75(5):762–777.
- Harrison, J. K., Jiang, Y., Chen, S., Xia, Y., Maciejewski, D., McNamara, R. K., Streit, W. J., Salafranca, M. N., Adhikari, S., Thompson, D. A., Botti, P., Bacon, K. B., and Feng, L. (1998). Role for neuronally derived fractalkine in mediating interactions between neurons and CX3CR1-expressing microglia. *Proceedings of the National Academy of Sciences*, 95(18):10896–10901.
- Hartline, D. K. (2011). The evolutionary origins of glia. *Glia*, 59(9):1215–1236.
- Haydon, P. G. (2001). Glia: listening and talking to the synapse. *Nat. Rev. Neurosci.*, 2:185–193.
- Haydon, P. G. and Carmignoto, G. (2006). Astrocyte control of synaptic transmission and neurovascular coupling. *Physiol. Rev.*, 86:1009–1031.
- Héja, L., Barabás, P., Nyitrai, G., Kékesi, K. A., Lasztóczy, B., TH<sup>+</sup>oke, O., Tárkányi, G., Madsen, K., Schousboe, A., Dobolyi, Á., Palkovits, M., and Kardos, J. (2009). Glutamate uptake triggers transporter-mediated GABA release from astrocytes. *PloS One*, 4(9):e7153.
- Héja, L., Nyitrai, G., Kékesi, O., Dobolyi, Á., Szabó, P., Fiáth, R., Ulbert, I., Pál-Szenthe, B., Palkovits, M., and Kardos, J. (2012). Astrocytes convert network excitation to tonic inhibition of neurons. *BMC Biology*, 10(1):26.
- Heller, J. P. and Rusakov, D. A. (2015). Morphological plasticity of astroglia: understanding synaptic microenvironment. *Glia*, 63(12):2133–2151.
- Henneberger, C., Papouin, T., Oliet, S. H. R., and Rusakov, D. A. (2010). Long-term potentiation depends on release of D-serine from astrocytes. *Nature*, 463:232–237.
- Herculano-Houzel, S. (2011). Not all brains are made the same: new views on brain scaling in evolution. *Brain, Behavior and Evolution*, 78(1):22–36.
- Herculano-Houzel, S. (2014). The glia/neuron ratio: how it varies uniformly across brain structures and species and what that means for brain physiology and evolution. *Glia*, 62(9):1377–1391.
- Herrero, J. L., Roberts, M. J., Delicato, L. S., Gieselmann, M. A., Dayan, P., and Thiele, A. (2008). Acetylcholine contributes through muscarinic receptors to attentional modulation in v1. *Nature*, 454(7208):1110.
- Herrmann, C. S., Fründ, I., and Lenz, D. (2010). Human gamma-band activity: a review on cognitive and behavioral correlates and network models. *Neuroscience & Biobehavioral Reviews*, 34(7):981–992.
- Herrmann, C. S., Munk, M. H. J., and Engel, A. K. (2004). Cognitive functions of gamma-band activity: memory match and utilization. *Trends in Cognitive Sciences*, 8(8):347–355.

- Hertz, L. (1965). Possible role of neuroglia: a potassium-mediated neuronal–neuroglial–neuronal impulse transmission system. *Nature*, 206(4989):1091.
- Hertz, L. (2004). Intercellular metabolic compartmentation in the brain: past, present and future. *Neurochemistry International*, 45(2-3):285–296.
- Hertz, L. (2013). The glutamate–glutamine (GABA) cycle: importance of late postnatal development and potential reciprocal interactions between biosynthesis and degradation. *Frontiers in Endocrinology*, 4:59.
- Hertz, L., Peng, L., and Dienel, G. A. (2007). Energy metabolism in astrocytes: high rate of oxidative metabolism and spatiotemporal dependence on glycolysis/glycogenolysis. *Journal of Cerebral Blood Flow & Metabolism*, 27(2):219–249.
- Hirase, H., Qian, L., Barthó, P., and Buzsáki, G. (2004). Calcium dynamics of cortical astrocytic networks *in vivo*. *PLoS Biol.*, 2(4):0494–0496.
- Hoffmann, A., Kann, O., Ohlemeyer, C., Hanisch, U.-K., and Kettenmann, H. (2003). Elevation of basal intracellular calcium as a central element in the activation of brain macrophages (microglia): suppression of receptor-evoked calcium signaling and control of release function. *Journal of Neuroscience*, 23(11):4410–4419.
- Hoogland, T. M., Kuhn, B., Göbel, W., Huang, W., Nakai, J., Helmchen, F., Flint, J., and Wang, S. S. (2009). Radially expanding transglial calcium waves in the intact cerebellum. *Proceedings of the National Academy of Sciences*, 106(9):3496–3501.
- Hooks, B. M. and Chen, C. (2006). Distinct roles for spontaneous and visual activity in remodeling of the retinogeniculate synapse. *Neuron*, 52(2):281–291.
- Hoshiko, M., Arnoux, I., Avignone, E., Yamamoto, N., and Audinat, E. (2012). Deficiency of the microglial receptor CX3CR1 impairs postnatal functional development of thalamocortical synapses in the barrel cortex. *Journal of Neuroscience*, 32(43):15106–15111.
- Howarth, C., Gleeson, P., and Attwell, D. (2012). Updated energy budgets for neural computation in the neocortex and cerebellum. *Journal of Cerebral Blood Flow & Metabolism*, 32(7):1222–1232.
- Hu, Y., Zylberberg, J., and Shea-Brown, E. (2014). The sign rule and beyond: boundary effects, flexibility, and noise correlations in neural population codes. *PLoS Computational Biology*, 10(2):e1003469.
- Hua, J. Y. and Smith, S. J. (2004). Neural activity and the dynamics of central nervous system development. *Nature Neuroscience*, 7(4):327.
- Huang, Y. H. and Bergles, D. E. (2004). Glutamate transporters bring competition to the synapse. *Current Opinion in Neurobiology*, 14(3):346–352.
- Hughes, E. G., Orthmann-Murphy, J. L., Langseth, A. J., and Bergles, D. E. (2018). Myelin remodeling through experience-dependent oligodendrogenesis in the adult somatosensory cortex. *Nature Neuroscience*, 21:696–706.
- Iadecola, C. and Nedergaard, M. (2007). Glial regulation of the cerebral microvasculature. *Nature Neuroscience*, 10(11):1369–1376.



- Imeri, L. and Opp, M. R. (2009). How (and why) the immune system makes us sleep. *Nature Reviews Neuroscience*, 10(3):199.
- Imura, Y., Morizawa, Y., Komatsu, R., Shibata, K., Shinozaki, Y., Kasai, H., Moriishi, K., Moriyama, Y., and Koizumi, S. (2013). Microglia release ATP by exocytosis. *Glia*, 61(8):1320–1330.
- Ingiosi, A. M., Opp, M. R., and Krueger, J. M. (2013). Sleep and immune function: glial contributions and consequences of aging. *Current Opinion in Neurobiology*, 23(5):806–811.
- Ishibashi, T., Dakin, K. A., Stevens, B., Lee, P. R., Kozlov, S. V., Stewart, C. L., and Fields, R. D. (2006). Astrocytes promote myelination in response to electrical impulses. *Neuron*, 49(6):823–832.
- Ismailov, I., Kalikulov, D., Inoue, T., and Friedlander, M. J. (2004). The kinetic profile of intracellular calcium predicts long-term potentiation and long-term depression. *The Journal of Neuroscience*, 24(44):9847–9861.
- Israel, J.-M., Le Masson, G., Theodosis, D. T., and Poulain, D. A. (2003). Glutamatergic input governs periodicity and synchronization of bursting activity in oxytocin neurons in hypothalamic organotypic cultures. *European Journal of Neuroscience*, 17(12):2619–2629.
- Izhikevich, E. M. (2006). Polychronization: computation with spikes. *Neural Computation*, 18(2):245–282.
- Jackson, F. R. (2011). Glial cell modulation of circadian rhythms. *Glia*, 59(9):1341–1350.
- Jadhav, A. P., Roesch, K., and Cepko, C. L. (2009). Development and neurogenic potential of müller glial cells in the vertebrate retina. *Progress in retinal and eye research*, 28(4):249–262.
- Jaffe, L. F. (1993). Classes and mechanisms of calcium waves. *Cell Calcium*, 14(10):736–745.
- Jeftinija, S. D., Jeftinija, K. V., Stefanovic, G., and Liu, F. (1996). Neuroligand-evoked calcium-dependent release of excitatory amino acids from cultured astrocytes. *J. Neurochem.*, 66:676–684.
- Jiang, P., Xing, F., Guo, B., Yang, J., Li, Z., Wei, W., Hu, F., Lee, I., Zhang, X., Pan, L., and Xu, J. (2017). Nucleotide transmitters ATP and ADP mediate intercellular calcium wave communication via P2Y<sub>12/13</sub> receptors among BV-2 microglia. *PloS one*, 12(8):e0183114.
- Jiménez-González, C., Pirttimaki, T., Cope, D. W., and Parri, H. R. (2011). Non-neuronal, slow GABA signalling in the ventrobasal thalamus targets  $\delta$ -subunit-containing GABA<sub>A</sub> receptors. *European Journal of Neuroscience*, 33(8):1471–1482.
- Jinno, S., Fleischer, F., Eckel, S., Schmidt, V., and Kosaka, T. (2007). Spatial arrangement of microglia in the mouse hippocampus: a stereological study in comparison with astrocytes. *Glia*, 55(13):1334–1347.
- Johnston, D. and Wu, S. M.-S. (1995). *Foundations of cellular neurophysiology*. MIT Press, Cambridge, MA.
- Jolivet, R., Allaman, I., Pellerin, L., Magistretti, P. J., and Weber, B. (2010). Comment on recent modeling studies of astrocyte–neuron metabolic interactions. *Journal of Cerebral Blood Flow & Metabolism*, 30(12):1982–1986.

- Joris, P. X., Carney, L. H., Smith, P. H., and Yin, T. C. (1994). Enhancement of neural synchronization in the anteroventral cochlear nucleus. I. Responses to tones at the characteristic frequency. *Journal of neurophysiology*, 71(3):1022–1036.
- Jourdain, P., Bergersen, L. H., Bhaukaurally, K., Bezzi, P., Santello, M., Domercq, M., Matute, C., Tonello, F., Gundersen, V., and Volterra, A. (2007). Glutamate exocytosis from astrocytes controls synaptic strength. *Nat. Neurosci.*, 10(3):331–339.
- Kadala, A., Verdier, D., Morquette, P., and Kolta, A. (2015). Ion homeostasis in rhythmogenesis: the interplay between neurons and astroglia. *Physiology*, 30(5):371–388.
- Kaila, K., Voipio, J., Paalasmaa, P., Pasternack, M., and Deisz, R. A. (1993). The role of bicarbonate in GABA<sub>A</sub> receptor-mediated IPSPs of rat neocortical neurones. *The Journal of Physiology*, 464(1):273–289.
- Kakegawa, W., Miyoshi, Y., Hamase, K., Matsuda, S., Matsuda, K., Kohda, K., Emi, K., Motohashi, J., Konno, R., Zaitso, K., and Yuzaki, M. (2011). D-Serine regulates cerebellar LTD and motor coordination through the  $\delta 2$  glutamate receptor. *Nature Neuroscience*, 14(5):603–611.
- Kaneko, M., Stellwagen, D., Malenka, R. C., and Stryker, M. P. (2008). Tumor necrosis factor- $\alpha$  mediates one component of competitive, experience-dependent plasticity in developing visual cortex. *Neuron*, 58:673–680.
- Káradóttir, R. and Attwell, D. (2007). Neurotransmitter receptors in the life and death of oligodendrocytes. *Neuroscience*, 145(4):1426–1438.
- Kaschube, M., Schnabel, M., Löwel, S., Coppola, D. M., White, L. E., and Wolf, F. (2010). Universality in the evolution of orientation columns in the visual cortex. *Science*, 330(6007):1113–1116.
- Kasthuri, N., Hayworth, K. ., Berger, D. R., Schalek, R. L., Conchello, J. A., Knowles-Barley, S., Lee, D., Vázquez-Reina, A., Kaynig, V., Jones, T. R., Roberts, M., Lyskowski, J. M., Tapia, J. C. Seung, H. S., Roncal, W. G., Vogelstein, J. T., Burns, R., Sussman, D. L., Priebe, C. E., Pfister, H., and Lichtman, J. W. (2015). Saturated reconstruction of a volume of neocortex. *Cell*, 162(3):648–661.
- Katagiri, H., Tanaka, K., and Manabe, T. (2001). Requirement of appropriate glutamate concentrations in the synaptic cleft for hippocampal LTP induction. *European Journal of Neuroscience*, 14(3):547–553.
- Katz, B. (1969). *The release of neural transmitter substances*, volume 10. Liverpool University Press.
- Keane, A. and Gong, P. (2015). Propagating waves can explain irregular neural dynamics. *Journal of Neuroscience*, 35(4):1591–1605.
- Kettenmann, H., Hanisch, U.-K., Noda, M., and Verkhratsky, A. (2011). Physiology of microglia. *Physiological Reviews*, 91(2):461–553.
- Kettenmann, H., Kirchhoff, F., and Verkhratsky, A. (2013). Microglia: new roles for the synaptic stripper. *Neuron*, 77(1):10–18.
- Kettenmann, H. and Ransom, B. R. (2004). *Neuroglia, The concept of neuroglia: A historical perspective*. Oxford University Press, New York, 1st edition.

- Kettenmann, H. and Ransom, B. R. (2013). *Neuroglia*. Oxford University Press, 3rd edition.
- Khakh, B. S. and Sofroniew, M. V. (2015). Diversity of astrocyte functions and phenotypes in neural circuits. *Nature Neuroscience*, 18(7):942.
- Khandelwal, P. J., Herman, A. M., and Moussa, C. E.-H. (2011). Inflammation in the early stages of neurodegenerative pathology. *Journal of neuroimmunology*, 238(1):1–11.
- Kimura, F. and Itami, C. (2009). Myelination and isochronicity in neural networks. *Frontiers in Neuroanatomy*, 3:12.
- Kinney, G. A. and Spain, W. J. (2002). Synaptically evoked GABA transporter currents in neocortical glia. *Journal of Neurophysiology*, 88(6):2899–2908.
- Kinney, J. P., Spacek, J., Bartol, T. M., Bajaj, C. L., Harris, K. M., and Sejnowski, T. J. (2013). Extracellular sheets and tunnels modulate glutamate diffusion in hippocampal neuropil. *Journal of Comparative Neurology*, 521(2):448–464.
- Kisvárdy, Z. F., Toth, E., Rausch, M., and Eysel, U. T. (1997). Orientation-specific relationship between populations of excitatory and inhibitory lateral connections in the visual cortex of the cat. *Cerebral cortex*, 7(7):605–618.
- Klinkenberg, I., Sambeth, A., and Blokland, A. (2011). Acetylcholine and attention. *Behavioural Brain Research*, 221(2):430–442.
- Ko, H., Hofer, S. B., Pichler, B., Buchanan, K. A., Sjöström, P. J., and Mrsic-Flogel, T. D. (2011). Functional specificity of local synaptic connections in neocortical networks. *Nature*, 473(7345):87.
- Kofuji, P. and Newman, E. A. (2004). Potassium buffering in the central nervous system. *Neuroscience*, 129(4):1043–1054.
- Kozlov, A. S., Angulo, M. C., Audinat, E., and Charpak, S. (2006). Target cell-specific modulation of neuronal activity by astrocytes. *Proc. Natl. Acad. Sci. USA*, 103(26):10058–10063.
- Krämer-Albers, E.-M., Bretz, N., Tenzer, S., Winterstein, C., Möbius, W., Berger, H., Nave, K.-A., Schild, H., and Trotter, J. (2007). Oligodendrocytes secrete exosomes containing major myelin and stress-protective proteins: Trophic support for axons? *Proteomics-clinical applications*, 1(11):1446–1461.
- Kreutzberg, G. W. (1996). Microglia: a sensor for pathological events in the CNS. *Trends in Neurosciences*, 19(8):312–318.
- Kriener, B., Helias, M., Aertsen, A., and Rotter, S. (2009). Correlations in spiking neuronal networks with distance dependent connections. *Journal of Computational Neuroscience*, 27(2):177–200.
- Krzan, M., Stenovec, N., Kreft, M., Pangrsic, T., Grilc, S., Haydon, P. G., and Zorec, R. (2003). Calcium-dependent exocytosis of atrial natriuretic peptide from astrocytes. *J. Neurosci.*, 23:1580–1583.
- Kuga, N., Sasaki, T., Takahara, Y., Matsuki, N., and Ikegaya, Y. (2011). Large-scale calcium waves traveling through astrocytic networks *in vivo*. *J. Neurosci.*, 31(7):2607–2614.

- Kummer, U., Olsen, L. F., Green, A. K., Bomberg-Bauer, E., and Baier, G. (2000). Switching from simple to complex oscillations in calcium signaling. *Biophys. J.*, 79:1188–1199.
- Labin, A. M. and Ribak, E. N. (2010). Retinal glial cells enhance human vision acuity. *Physical Review Letters*, 104(15):158102.
- Labin, A. M., Safuri, S. K., Ribak, E. N., and Perlman, I. (2014). Müller cells separate between wavelengths to improve day vision with minimal effect upon night vision. *Nature Communications*, 5:4319.
- Lang, E. J. and Rosenbluth, J. (2003). Role of myelination in the development of a uniform olivocerebellar conduction time. *Journal of Neurophysiology*, 89(4):2259–2270.
- Larsen, B. R., Assentoft, M., Cotrina, M. L., Hua, S. Z., Nedergaard, M., Kaila, K., Voipio, J., and MacAulay, N. (2014). Contributions of the Na<sup>+</sup>/K<sup>+</sup>-ATPase, NKCC1, and Kir4.1 to hippocampal K<sup>+</sup> clearance and volume responses. *Glia*, 62(4):608–622.
- Laughlin, S. B. (2001). Energy as a constraint on the coding and processing of sensory information. *Current Opinion in Neurobiology*, 11(4):475–480.
- Laughlin, S. B. and Sejnowski, T. J. (2003). Communication in neuronal networks. *Science*, 301(5641):1870–1874.
- Lawson, L. J., Perry, V. H., Dri, P., and Gordon, S. (1990). Heterogeneity in the distribution and morphology of microglia in the normal adult mouse brain. *Neuroscience*, 39(1):151–170.
- Le Meur, K., Mendizabal-Zubiaga, J., Grandes, P., and Audinat, E. (2012). GABA release by hippocampal astrocytes. *Front. Comp. Neurosci.*, 6.
- Lee, H. S., Ghetti, A., Pinto-Duarte, A., Wang, X., Dziewczapolski, G., Galimi, F., Huitron-Resendiz, S., Piña-Crespo, J. C., Roberts, A. J., Verma, I. M., Sejnowski, T. J., and Heinemann, S. F. (2014). Astrocytes contribute to gamma oscillations and recognition memory. *Proc. Nat. Acad. Sci. USA*, pages E3343–E3352.
- Lee, M., McGeer, E. G., and McGeer, P. L. (2011). Mechanisms of GABA release from human astrocytes. *Glia*, 59:1600–1611.
- Lee, S., Yoon, B., Berglund, K., Oh, S., Park, H., Shin, H., Augustine, G., and Lee, C. (2010). Channel-mediated tonic GABA release from glia. *Sci. STKE*, 330(6005):790–796.
- Lehre, K. P. and Rusakov, D. A. (2002). Asymmetry of glia near central synapses favors presynaptically directed glutamate escape. *Biophys. J.*, 83(1):125–134.
- Levy, L. M., Warr, O., and Attwell, D. (1998). Stoichiometry of the glial glutamate transporter GLT-1 expressed inducibly in a chinese hamster ovary cell line selected for low endogenous Na<sup>+</sup>-dependent glutamate uptake. *Journal of Neuroscience*, 18(23):9620–9628.
- Lewitus, G. M., Konefal, S. C., Greenhalgh, A. D., Pribiag, H., Augereau, K., and Stellwagen, D. (2016). Microglial TNF- $\alpha$  suppresses cocaine-induced plasticity and behavioral sensitization. *Neuron*, 90(3):483–491.
- Lewitus, G. M., Pribiag, H., Duseja, R., St-Hilaire, M., and Stellwagen, D. (2014). An adaptive role of TNF $\alpha$  in the regulation of striatal synapses. *Journal of Neuroscience*, 34(18):6146–6155.

- Leybaert, L. and Sanderson, M. J. (2012). Intercellular  $\text{Ca}^{2+}$  waves: mechanisms and function. *Physiological reviews*, 92(3):1359–1392.
- Lind, B. L., Brazhe, A. R., Jessen, S. B., Tan, F. C. C., and Lauritzen, M. J. (2013). Rapid stimulus-evoked astrocyte  $\text{Ca}^{2+}$  elevations and hemodynamic responses in mouse somatosensory cortex *in vivo*. *Proceedings of the National Academy of Sciences*, 110(48):E4678–E4687.
- Lippman, J. J., Lordkipanidze, T., Buell, M. E., Yoon, S. O., and Dunaevsky, A. (2008). Morphogenesis and regulation of Bergmann glial processes during Purkinje cell dendritic spine ensheathment and synaptogenesis. *Glia*, 56(13):1463–1477.
- Lippman, J. J., Lordkipanidze, T., Cobb, N., and Dunaevsky, A. (2010). Bergmann glial ensheathment of dendritic spines regulates synapse number without affecting spine motility. *Neuron Glia Biology*, 6(3):193–200.
- Litwin-Kumar, A. and Doiron, B. (2012). Slow dynamics and high variability in balanced cortical networks with clustered connections. *Nature Neuroscience*, 15(11):1498.
- Liu, G. J., Kalous, A., Werry, E. L., and Bennett, M. R. (2006). Purine release from spinal cord microglia after elevation of calcium by glutamate. *Molecular Pharmacology*, 70(3):851–859.
- López-Hidalgo, M., Hoover, W. B., and Schummers, J. (2016). Spatial organization of astrocytes in ferret visual cortex. *Journal of Comparative Neurology*, 524(17):3561–3576.
- López-Hidalgo, M., Salgado-Puga, K., Alvarado-Martínez, R., Medina, A. C., Prado-Alcalá, R. A., and García-Colunga, J. (2012). Nicotine uses neuron-glia communication to enhance hippocampal synaptic transmission and long-term memory. *PloS one*, 7(11):e49998.
- López-Hidalgo, M. and Schummers, J. (2014). Cortical maps: a role for astrocytes? *Current Opinion in Neurobiology*, 24:176–189.
- Losi, G., Mariotti, L., and Carmignoto, G. (2014). GABAergic interneuron to astrocyte signalling: a neglected form of cell communication in the brain. *Phil. Trans. Royal Soc. B: Biological Sciences*, 369(1654):20130609.
- Lowery, R. L., Tremblay, M.-È., Hopkins, B. E., and Majewska, A. K. (2017). The microglial fractalkine receptor is not required for activity-dependent plasticity in the mouse visual system. *Glia*, 65(11):1744–1761.
- Luders, E., Thompson, P. M., and Toga, A. W. (2010). The development of the corpus callosum in the healthy human brain. *Journal of Neuroscience*, 30(33):10985–10990.
- Lundgaard, I., Lu, M. L., Yang, E., Peng, W., Mestre, H., Hitomi, E., Deane, R., and Nedergaard, M. (2017). Glymphatic clearance controls state-dependent changes in brain lactate concentration. *Journal of Cerebral Blood Flow & Metabolism*, 37(6):2112–2124.
- Lushnikova, I., Skibo, G., Muller, D., and Nikonenko, I. (2009). Synaptic potentiation induces increased glial coverage of excitatory synapses in ca1 hippocampus. *Hippocampus*, 19(8):753–762.
- Lutz, S. E., Zhao, Y., Gulinello, M., Lee, S. C., Raine, C. S., and Brosnan, C. F. (2009). Deletion of astrocyte connexins 43 and 30 leads to a dysmyelinating phenotype and hippocampal CA1 vacuolation. *Journal of Neuroscience*, 29(24):7743–7752.

- Ma, B.-F., Xie, M.-J., and Zhou, M. (2012). Bicarbonate efflux via GABA<sub>A</sub> receptors depolarizes membrane potential and inhibits two-pore domain potassium channels of astrocytes in rat hippocampal slices. *Glia*, 60(11):1761–1772.
- Mackenzie, B. and Erickson, J. D. (2004). Sodium-coupled neutral amino acid (System N/A) transporters of the SLC38 gene family. *Pflügers Archiv. – Eur. J. Physiol.*, 447(5):784–795.
- Magistretti, P. J. and Allaman, I. (2015). A cellular perspective on brain energy metabolism and functional imaging. *Neuron*, 86(4):883–901.
- Magistretti, P. J. and Chatton, J.-Y. (2005). Relationship between L-glutamate-regulated intracellular Na<sup>+</sup> dynamics and ATP hydrolysis in astrocytes. *Journal of Neural Transmission*, 112(1):77–85.
- Magistretti, P. J., Morrison, J. H., Shoemaker, W. J., Sapin, V., and Bloom, F. E. (1981). Vasoactive intestinal polypeptide induces glycogenolysis in mouse cortical slices: a possible regulatory mechanism for the local control of energy metabolism. *Proceedings of the National Academy of Sciences*, 78(10):6535–6539.
- Mangia, S., DiNuzzo, M., Giove, F., Carruthers, A., Simpson, I. A., and Vannucci, S. J. (2011). Response to comment on recent modeling studies of astrocytaneuron metabolic interactions: Much ado about nothing. *Journal of Cerebral Blood Flow & Metabolism*, 31(6):1346–1353.
- Marchaland, J., Calì, C., Voglmaier, S. M., Li, H., Regazzi, R., Edwards, R. H., and Bezzi, P. (2008). Fast subplasma membrane Ca<sup>2+</sup> transients control exo-endocytosis of synaptic-like microvesicles in astrocytes. *J. Neurosci.*, 28(37):9122–9132.
- Marchant, J. S. and Parker, I. (2001). Role of elementary Ca<sup>2+</sup> puffs in generating repetitive Ca<sup>2+</sup> oscillations. *The EMBO Journal*, 20(1-2):65–76.
- Markram, H., Lübke, J., Frotscher, M., Roth, A., and Sakmann, B. (1997). Physiology and anatomy of synaptic connections between thick tufted pyramidal neurones in the developing rat neocortex. *The Journal of Physiology*, 500(2):409–440.
- Markram, H., Wang, Y., and Tsodyks, M. (1998). Differential signaling via the same axon of neocortical pyramidal neurons. *Proc. Natl. Acad. Sci. USA*, 95:5323–5328.
- Martín, R., Bajo-Grañeras, R., Moratalla, R., Perea, G., and Araque, A. (2015). Circuit-specific signaling in astrocyte-neuron networks in basal ganglia pathways. *Science*, 349(6249):730–734.
- Mashimo, M., Okubo, Y., Yamazawa, T., Yamasaki, M., Watanabe, M., Murayama, T., and Iino, M. (2010). Inositol 1,4,5-trisphosphate signaling maintains the activity of glutamate uptake in Bergmann glia. *European Journal of Neuroscience*, 32(10):1668–1677.
- Masland, R. H. (2001). The fundamental plan of the retina. *Nature Neuroscience*, 4(9):877.
- Matsuzaki, M., Ellis-Davies, G., Nemoto, T., Miyashita, Y., Iino, M., and Kasai, H. (2001). Dendritic spine geometry is critical for AMPA receptor expression in hippocampal CA1 pyramidal neurons. *Nat. Neurosci.*, 4(11):1086–1092.
- Matyash, V. and Kettenmann, H. (2010). Heterogeneity in astrocyte morphology and physiology. *Brain Res. Rev.*, 63(1-2):2–10.
- McAlpine, D. and Grothe, B. (2003). Sound localization and delay lines—do mammals fit the model? *Trends in Neurosciences*, 26(7):347–350.

- McCormick, D. A., Shu, Y., Hasenstaub, A., Sanchez-Vives, M., Badoual, M., and Bal, T. (2003). Persistent cortical activity: mechanisms of generation and effects on neuronal excitability. *Cerebral Cortex*, 13(11):1219–1231.
- McCoy, M. K. and Tansey, M. G. (2008). TNF signaling inhibition in the CNS: implications for normal brain function and neurodegenerative disease. *Journal of Neuroinflammation*, 5(1):45.
- Medvedev, N., Popov, V., Henneberger, C., Kraev, I., Rusakov, D. A., and Stewart, M. G. (2014). Glia selectively approach synapses on thin dendritic spines. *Phil. Trans. R. Soc. B*, 369(1654):20140047.
- Men, W., Falk, D., Sun, T., Chen, W., Li, J., Yin, D., Zang, L., and Fan, M. (2014). The corpus callosum of Albert Einsteins brain: another clue to his high intelligence? *Brain*, 137(4):e268–e268.
- Mergenthaler, P., Lindauer, U., Dienel, G. A., and Meisel, A. (2013). Sugar for the brain: the role of glucose in physiological and pathological brain function. *Trends in Neurosciences*, 36(10):587–597.
- Mesejo, P., Ibáñez, O., Fernández-Blanco, E., Cedrón, F., Pazos, A., and Porto-Pazos, A. B. (2015). Artificial neuron–glia networks learning approach based on cooperative coevolution. *International Journal of Neural Systems*, 25(04):1550012.
- Min, R. and Nevian, T. (2012). Astrocyte signaling controls spike timing-dependent depression at neocortical synapses. *Nat. Neurosci.*, 15(5):746–753.
- Molofsky, A. V., Kelley, K. W., Tsai, H.-H., Redmond, S. A., Chang, S. M., Madireddy, L., Chan, J. R., Baranzini, S. E., Ullian, E. M., and Rowitch, D. H. (2014). Astrocyte-encoded positional cues maintain sensorimotor circuit integrity. *Nature*, 509(7499):189.
- Montero, T. D. and Orellana, J. A. (2015). Hemichannels: new pathways for gliotransmitter release. *Neuroscience*, 286:45–59.
- Moore, J. W., Joyner, R. W., Brill, M. H., Waxman, S. D., and Najar-Joa, M. (1978). Simulations of conduction in uniform myelinated fibers. Relative sensitivity to changes in nodal and internodal parameters. *Biophysical Journal*, 21(2):147–160.
- Moreno-Bote, R., Beck, J., Kanitscheider, I., Pitkow, X., Latham, P., and Pouget, A. (2014). Information-limiting correlations. *Nature Neuroscience*, 17(10):1410.
- Morquette, P., Verdier, D., Kadala, A., Féthière, J., Philippe, A. G., Robitaille, R., and Kolta, A. (2015). An astrocyte-dependent mechanism for neuronal rhythmogenesis. *Nature Neuroscience*, 18(6):844.
- Mota, B. and Herculano-Houzel, S. (2014). All brains are made of this: a fundamental building block of brain matter with matching neuronal and glial masses. *Frontiers in Neuroanatomy*, 8:127.
- Mothet, J.-P., Parent, A. T., Wolosker, H., Brady, R. O., Linden, D. J., Ferris, C. D., Rogawski, M. A., and Snyder, S. H. (2000). D-serine is an endogenous ligand for the glycine site of the N-methyl-D-aspartate receptor. *Proceedings of the National Academy of Sciences*, 97(9):4926–4931.
- Müller, T., Möller, T., Neuhaus, J., and Kettenmann, H. (1996). Electrical coupling among Bergmann glial cells and its modulation by glutamate receptor activation. *Glia*, 17:274–284.

- Mulligan, S. and MacVicar, B. (2004). Calcium transients in astrocyte endfeet cause cerebrovascular constrictions. *Nature*, 431(7005):195–199.
- Murai, K. K., Nguyen, L. N., Irie, F., Yamaguchi, Y., and Pasquale, E. B. (2003). Control of hippocampal dendritic spine morphology through ephrin-A3/EphA4 signaling. *Nature Neuroscience*, 6(2):153.
- Nagelhus, E. A., Mathiisen, T. M., and Ottersen, O. P. (2004). Aquaporin-4 in the central nervous system: cellular and subcellular distribution and coexpression with KIR4.1. *Neuroscience*, 129(4):905–913.
- Navarrete, M. and Araque, A. (2008). Endocannabinoids mediate neuron-astrocyte communication. *Neuron*, 57(6):883–893.
- Navarrete, M., Díez, A., and Araque, A. (2014). Astrocytes in endocannabinoid signalling. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 369(1654):20130599.
- Navarrete, M., Perea, G., de Sevilla, D., Gómez-Gonzalo, M., Núñez, A., Martín, E., and Araque, A. (2012). Astrocytes mediate in vivo cholinergic-induced synaptic plasticity. *PLoS Biol.*, 10(2):e1001259.
- Nave, K.-A. (2010). Myelination and support of axonal integrity by glia. *Nature*, 468(7321):244.
- Nedergaard, M. (1994). Direct signaling from astrocytes to neurons in cultures of mammalian brain cells. *Science*, 263(5154):1768.
- Nedergaard, M., Ransom, B. R., and Goldman, S. A. (2003). New roles for astrocytes: redefining the functional architecture of the brain. *Trends Neurosci.*, 26(10):523–530.
- Nevian, T. and Sakmann, B. (2006). Spine  $\text{Ca}^{2+}$  signaling in spike-timing-dependent plasticity. *Journal of Neuroscience*, 26(43):11001–11013.
- Newman, E. (2005). Calcium increases in retinal glial cells evoked by light-induced neuronal activity. *J. Neurosci.*, 25(23):5502–5510.
- Newman, E. A. (2001). Propagation of intercellular calcium waves in retinal astrocytes and Müller cells. *J. Neurosci.*, 21:2215–2223.
- Newman, E. A. and Zahs, K. R. (1997). Calcium waves in retinal glial cells. *Science*, 275:844–847.
- Newman, L. A., Korol, D. L., and Gold, P. E. (2011). Lactate produced by glycogenolysis in astrocytes regulates memory processing. *PLoS One*, 6(12):e28427.
- Nicoll, R. A. and Schmitz, D. (2005). Synaptic plasticity at hippocampal mossy fibre synapses. *Nature Reviews Neuroscience*, 6(11):863.
- Nie, H., Zhang, H., and Weng, H. (2010). Bidirectional neuron–glia interactions triggered by deficiency of glutamate uptake at spinal sensory synapses. *J. Neurophysiol.*, 104(2):713–725.
- Nikola, G., Andon, V., and Kaynak, O. (2003). Sliding mode algorithm for online learning in analog multilayer feedforward neural networks. In *Artificial Neural Networks and Neural Information Processing–ICANN/ICONIP 2003*, pages 1064–1072. Springer.
- Nimmerjahn, A., Kirchhoff, F., and Helmchen, F. (2005). Resting microglial cells are highly dynamic surveillants of brain parenchyma in vivo. *Science*, 308(5726):1314–1318.



- Nimmerjahn, A., Kirchhoff, F., Kerr, J. N. D., and Helmchen, F. (2004). Sulforhodamine 101 as a specific marker of astroglia in the neocortex *in vivo*. *Nat. Methods*, 1:31–37.
- Nir, Y., Staba, R. J., Andrillon, T., Vyazovskiy, V. V., Cirelli, C., Fried, I., and Tononi, G. (2011). Regional slow waves and spindles in human sleep. *Neuron*, 70(1):153–169.
- Oberheim, N. A., Takano, T., Han, X., He, W., Lin, J. H. C., Wang, F., Xu, Q., Wyatt, J. D., Pilcher, W., Ojemann, J. G., Ransom, B. R., Goldman, S. A., and Nedergaard, M. (2009). Uniquely hominid features of adult human astrocytes. *J. Neurosci.*, 29(10):3276–3287.
- O’Connor, D. H., Wittenberg, G. M., and Wang, S. S.-H. (2005). Graded bidirectional synaptic plasticity is composed of switch-like unitary events. *Proceedings of the National Academy of Sciences*, 102(27):9679–9684.
- Ohki, K., Chung, S., Ch’ng, Y. H., Kara, P., and Reid, R. C. (2005). Functional imaging with cellular resolution reveals precise micro-architecture in visual cortex. *Nature*, 433(7026):597.
- Olah, M., Biber, K., Vinet, J., and Boddeke, H. W. G. M. (2011). Microglia phenotype diversity. *CNS & Neurological Disorders-Drug Targets*, 10(1):108–118.
- Oliet, S. H. R. and Mothet, J. (2009). Regulation of  $\text{i}_\text{h}$ / $\text{i}_\text{h}$ -methyl-D-aspartate receptors by astrocytic D-serine. *Neuroscience*, 158(1):275–283.
- Oliet, S. H. R., Piet, R., and Poulain, D. A. (2001). Control of glutamate clearance and synaptic efficacy by glial coverage of neurons. *Science*, 292:923–926.
- Oliveira, J. F., Sardinha, V. M., Guerra-Gomes, S., Araque, A., and Sousa, N. (2015). Do stars govern our actions? Astrocyte involvement in rodent behavior. *Trends in neurosciences*, 38(9):535–549.
- Omrani, A., Melone, M., Bellesi, M., Safiulina, V., Aida, T., Tanaka, K., Cherubini, E., and Conti, F. (2009). Up-regulation of GLT-1 severely impairs LTD at mossy fibre–CA3 synapses. *The Journal of Physiology*, 587(19):4575–4588.
- O’Neill, L. A. J. and Kaltschmidt, C. (1997). NF- $\kappa$ B: a crucial transcription factor for glial and neuronal cell function. *Trends in Neurosciences*, 20(6):252–258.
- Orkand, R. K., Nicholls, J. G., and Kuffler, S. W. (1966). Effect of nerve impulses on the membrane potential of glial cells in the central nervous system of amphibia. *Journal of Neurophysiology*, 29(4):788–806.
- Østby, I., Øyehaug, L., Einevoll, G. T., Nagelhus, E. A., Plahte, E., Zeuthen, T., Lloyd, C. M., Ottersen, O. P., and Omholt, S. W. (2009). Astrocytic mechanisms explaining neural-activity-induced shrinkage of extraneuronal space. *PLoS Computational Biology*, 5(1):e1000272.
- Otsu, Y., Couchman, K., Lyons, D. G., Collot, M., Agarwal, A., Mallet, J.-M., Pfrieger, F. W., Bergles, D. E., and Charpak, S. (2015). Calcium dynamics in astrocyte processes during neurovascular coupling. *Nature Neuroscience*, 18(2):210.
- Øyehaug, L., Østby, I., Lloyd, C. M., Omholt, S. W., and Einevoll, G. T. (2012). Dependence of spontaneous neuronal firing and depolarisation block on astroglial membrane transport mechanisms. *Journal of Computational Neuroscience*, 32(1):147–165.

- Palygin, O., Lalo, U., Verkhratsky, A., and Pankratov, Y. (2010). Ionotropic NMDA and P2X<sub>1/5</sub> receptors mediate synaptically induced Ca<sup>2+</sup> signalling in cortical astrocytes. *Cell calcium*, 48(4):225–231.
- Panatier, A., Theodosis, D. T., Mothet, J. P., Touquet, B., Pollegioni, L., Poulain, D. A., and Oliet, S. H. (2006). Glia-derived D-serine controls nmda receptor activity and synaptic memory. *Cell*, 125:775–784.
- Pannasch, U., Freche, D., Dallérac, G., Ghézali, G., Escartin, C., Ezan, P., Cohen-Salmon, M., Benchenane, K., Abudara, V., Dufour, A., Lübke, J. H. R., Déglon, N., Knott, G., Holcman, D., and Rouach, N. (2014). Connexin 30 sets synaptic strength by controlling astroglial synapse invasion. *Nature neuroscience*, 17(4):549.
- Paolicelli, R. C., Bolasco, G., Pagani, F., Maggi, L., Scianni, M., Panzanelli, P., Giustetto, M., Ferreira, T. A., Guiducci, E., Dumas, L., Ragozzino, D., and Gross, C. T. (2011). Synaptic pruning by microglia is necessary for normal brain development. *Science*, 333(6048):1456–1458.
- Papouin, T., Ladépêche, L., Ruel, J., Sacchi, S., Labasque, M., Hanini, M., Groc, L., Pollegioni, L., Mothet, J., and Oliet, S. (2012). Synaptic and extrasynaptic NMDA receptors are gated by different endogenous coagonists. *Cell*, 150(3):633–646.
- Parpura, V., Basarsky, T. A., Liu, F., Jeftinija, K., Jeftinija, S., and Haydon, P. G. (1994). Glutamate-mediated astrocyte-neuron signalling. *Nature*, 369:744–747.
- Parpura, V. and Haydon, P. G. (2000). Physiological astrocytic calcium levels stimulate glutamate release to modulate adjacent neurons. *Proc. Natl. Acad. Sci. USA*, 97(15):8629–8634.
- Parys, B., Côté, A., Gallo, V., De Koninck, P., and Sík, A. (2010). Intercellular calcium signaling between astrocytes and oligodendrocytes via gap junctions in culture. *Neuroscience*, 167(4):1032–1043.
- Pascual, O., Ben Achour, S., Rostaing, P., Triller, A., and Bessis, A. (2011). Microglia activation triggers astrocyte-mediated modulation of excitatory neurotransmission. *Proceedings of the National Academy of Sciences*, pages 1–9.
- Pascual, O., Casper, K. B., Kubera, C., Zhang, J., Revilla-Sanchez, R., Sul, J. Y., Takano, H., Moss, S. J., McCarthy, K., and Haydon, P. G. (2005). Astrocytic purinergic signaling coordinates synaptic networks. *Science*, 310:113–116.
- Pasquale, E. B. (2008). Eph-ephrin bidirectional signaling in physiology and disease. *Cell*, 133(1):38–52.
- Pasti, L., Volterra, A., Pozzan, T., and Carmignoto, G. (1997). Intracellular calcium oscillations in astrocytes: a highly plastic, bidirectional form of communication between neurons and astrocytes *in situ*. *J. Neurosci.*, 17(20):7817–7830.
- Pasti, L., Zonta, M., Pozzan, T., Vicini, S., and Carmignoto, G. (2001). Cytosolic calcium oscillations in astrocytes may regulate exocytotic release of glutamate. *J. Neurosci.*, 21(2):477–484.
- Paukert, M., Agarwal, A., Cha, J., Doze, V. A., Kang, J. U., and Bergles, D. E. (2014). Norepinephrine controls astroglial responsiveness to local circuit activity. *Neuron*, 82(6):1263–1270.

- Pellerin, L. and Magistretti, P. J. (1994). Glutamate uptake into astrocytes stimulates aerobic glycolysis: a mechanism coupling neuronal activity to glucose utilization. *Proceedings of the National Academy of Sciences*, 91(22):10625–10629.
- Pellerin, L. and Magistretti, P. J. (2012). Sweet sixteen for ANLS. *Journal of Cerebral Blood Flow & Metabolism*, 32(7):1152–1166.
- Pelvig, D. P., Pakkenberg, H., Stark, A. K., and Pakkenberg, B. (2008). Neocortical glial cell numbers in human brains. *Neurobiology of Aging*, 29(11):1754–1762.
- Penfield, W. (1932). *Cytology and Cellular Pathology of the Nervous System*, volume 2, chapter Neuroglia, normal and pathological, pages 421–479. Hafner, New York.
- Perea, G. and Araque, A. (2005). Properties of synaptically evoked astrocyte calcium signal reveal synaptic information processing by astrocyte. *J. Neurosci.*, 25(9):2192–2203.
- Perea, G. and Araque, A. (2007). Astrocytes potentiate transmitter release at single hippocampal synapses. *Science*, 317:1083–1086.
- Perea, G., Yang, A., Boyden, E. S., and Sur, M. (2014). Optogenetic astrocyte activation modulates response selectivity of visual cortex neurons in vivo. *Nature Communications*, 5:ncomms4262.
- Perez-Alvarez, A., Navarrete, M., Covelo, A., Martin, E. D., and Araque, A. (2014). Structural and functional plasticity of astrocyte processes and dendritic spine interactions. *The Journal of Neuroscience*, 34(38):12738–12744.
- Perkel, D. H. and Bullock, T. H. (1968). Neural coding: a report based on an nrp work session. *Neurosci. Res. Program Bull.*, 6:219349.
- Pernice, V., Staude, B., Cardanobile, S., and Rotter, S. (2011). How structure determines correlations in neuronal networks. *PLoS Computational Biology*, 7(5):e1002059.
- Petit, J.-M., Gyger, J., Burette-Godinot, S., Fiumelli, H., Martin, J.-L., and Magistretti, P. J. (2013). Genes involved in the astrocyte-neuron lactate shuttle (ANLS) are specifically regulated in cortical astrocytes following sleep deprivation in mice. *Sleep*, 36(10):1445–1458.
- Petit, J.-M. and Magistretti, P. J. (2016). Regulation of neuron–astrocyte metabolic coupling across the sleep–wake cycle. *Neuroscience*, 323:135–156.
- Philips, R. T., Sur, M., and Chakravarthy, V. S. (2017). The influence of astrocytes on the width of orientation hypercolumns in visual cortex: A computational perspective. *PLoS Computational Biology*, 13(10):e1005785.
- Piet, R., Poulain, D. A., and Oliet, S. H. (2004a). Contribution of astrocytes to synaptic transmission in the rat supraoptic nucleus. *Neurochemistry International*, 45(2-3):251–257.
- Piet, R., Vargová, L., Syková, E., Poulain, D., and Oliet, S. (2004b). Physiological contribution of the astrocytic environment of neurons to intersynaptic crosstalk. *Proc. Natl. Acad. Sci. USA*, 101(7):2151–2155.
- Pinheiro, P. S. and Mulle, C. (2008). Presynaptic glutamate receptors: physiological functions and mechanisms of action. *Nature Reviews Neuroscience*, 9:423–436.

- Pocock, J. M. and Kettenmann, H. (2007). Neurotransmitter receptors on microglia. *Trends in Neurosciences*, 30(10):527–535.
- Porter, J. T., Cauli, B., Staiger, J. F., Lambolez, B., Rossier, J., and Audinat, E. (1998). Properties of bipolar VIPergic interneurons and their excitation by pyramidal neurons in the rat neocortex. *European Journal of Neuroscience*, 10(12):3617–3628.
- Porter, J. T. and McCarthy, K. D. (1996). Hippocampal astrocytes *in situ* respond to glutamate released from synaptic terminals. *J. Neurosci.*, 16(16):5073–5081.
- Porter, J. T. and McCarthy, K. D. (1997). Astrocytic neurotransmitter receptors *in situ* and *in vivo*. *Progress in Neurobiology*, 51(4):439–455.
- Porto-Pazos, A. B., Veiguela, N., Mesejo, P., Navarrete, M., Alvarellos, A., Ibáñez, O., Pazos, A., and Araque, A. (2011). Artificial astrocytes improve neural network performance. *PLoS one*, 6(4):e19109.
- Poskanzer, K. E. and Yuste, R. (2016). Astrocytes regulate cortical state switching *in vivo*. *Proceedings of the National Academy of Sciences*, 113(19):E2675–E2684.
- Pribrag, H. and Stellwagen, D. (2013). TNF- $\alpha$  downregulates inhibitory neurotransmission through protein phosphatase 1-dependent trafficking of GABA<sub>A</sub> receptors. *The Journal of Neuroscience*, 33(40):15879–15893.
- Price, D. L., Ludwig, J. W., Mi, H., Schwarz, T. L., and Ellisman, M. H. (2002). Distribution of rSlo Ca<sup>2+</sup>-activated K<sup>+</sup> channels in rat astrocyte perivascular endfeet. *Brain Research*, 956(2):183–193.
- Rajani, V., Zhang, Y., Jalubula, V., Rancic, V., Sheikhabaehi, S., Zwicker, J. D., Pagliardini, S., Dickson, C. T., Ballanyi, K., Kasparov, S., Gourine, A. V., and Funk, G. D. (2017). Release of ATP by preBötzing complex astrocytes contributes to the hypoxic ventilatory response via a Ca<sup>2+</sup>-dependent P2Y<sub>1</sub> receptor mechanism. *The Journal of physiology*, 589:4583–4600.
- Ransom, C. B., Ransom, B. R., and Sontheimer, H. (2000). Activity-dependent extracellular K<sup>+</sup> accumulation in rat optic nerve: the role of glial and axonal Na<sup>+</sup> pumps. *The Journal of physiology*, 522(3):427–442.
- Reichenbach, A., Derouiche, A., and Kirchhoff, F. (2010). Morphology and dynamics of perisynaptic glia. *Brain Research Reviews*, 63(1):11–25.
- Reichenbach, A. and Robinson, S. R. (1995). Phylogenetic constraints on retinal organisation and development. *Progress in Retinal and Eye Research*, 15(1):139–171.
- Richardson, A. G., McIntyre, C. C., and Grill, W. M. (2000). Modelling the effects of electric fields on nerve fibres: influence of the myelin sheath. *Medical and Biological Engineering and Computing*, 38(4):438–446.
- Rigotti, M., Barak, O., Warden, M. R., Wang, X.-J., Daw, N. D., Miller, E. K., and Fusi, S. (2013). The importance of mixed selectivity in complex cognitive tasks. *Nature*, 497(7451):585.
- Roberts, M. T., Seeman, S. C., and Golding, N. L. (2013). A mechanistic understanding of the role of feedforward inhibition in the mammalian sound localization circuitry. *Neuron*, 78(5):923–935.

- Rose, C. R. and Chatton, J.-Y. (2016). Astrocyte sodium signaling and neuro-metabolic coupling in the brain. *Neuroscience*, 323:121–134.
- Rothman, D. L., Behar, K. L., Hyder, F., and Shulman, R. G. (2003). In vivo NMR studies of the glutamate neurotransmitter flux and neuroenergetics: implications for brain function. *Annual Review of Physiology*, 65(1):401–427.
- Rothstein, J. D., Dykes-Hoberg, M., Pardo, C. A., Bristol, L. A., Jin, L., Kuncl, R. W., Kanai, Y., Hediger, M. A., Wang, Y., Schielke, J. P., , and Welty, D. F. (1996). Knockout of glutamate transporters reveals a major role for astroglial transport in excitotoxicity and clearance of glutamate. *Neuron*, 16(3):675–686.
- Roxin, A., Brunel, N., and Hansel, D. (2005). Role of delays in shaping spatiotemporal dynamics of neuronal activity in large networks. *Physical Review Letters*, 94(23):238103.
- Roxin, A. and Montbrió, E. (2011). How effective delays shape oscillatory dynamics in neuronal networks. *Physica D: Nonlinear Phenomena*, 240(3):323–345.
- Rusakov, D. A. (2001). The role of perisynaptic glial sheaths in glutamate spillover and extracellular  $\text{Ca}^{2+}$  depletion. *Biophysical Journal*, 81(4):1947–1959.
- Rusakov, D. A. (2015). Disentangling calcium-driven astrocyte physiology. *Nature Reviews Neuroscience*, 16:226–233.
- Rushton, W. A. H. (1951). A theory of the effects of fibre size in medullated nerve. *The Journal of Physiology*, 115(1):101–122.
- Sabatini, B. L. and Regehr, W. G. (1999). Timing of synaptic transmission. *Annual Review of Physiology*, 61(1):521–542.
- Sahlender, D. A., Savtchouk, I., and Volterra, A. (2014). What do we know about gliotransmitter release from astrocytes? *Phil. Tran. R. Soc. B*, 369:20130592.
- Salami, M., Itami, C., Tsumoto, T., and Kimura, F. (2003). Change of conduction velocity by regional myelination yields constant latency irrespective of distance between thalamus and cortex. *Proceedings of the National Academy of Sciences*, 100(10):6174–6179.
- Sanchez-Vives, M. V. and McCormick, D. A. (2000). Cellular and network mechanisms of rhythmic recurrent activity in neocortex. *Nature Neuroscience*, 3(10):1027.
- Santello, M., Bezzi, P., and Volterra, A. (2011).  $\text{TNF}\alpha$  controls glutamatergic gliotransmission in the hippocampal dentate gyrus. *Neuron*, 69:988–1001.
- Santello, M. and Volterra, A. (2012).  $\text{TNF}\alpha$  in synaptic function: switching gears. *Trends in Neurosci.*, 35(10):638–647.
- Sarpeshkar, R. (1998). Analog versus digital: extrapolating from electronics to neurobiology. *Neural Computation*, 10(7):1601–1638.
- Sasaki, T., Matsuki, N., and Ikegaya, Y. (2011). Action-potential modulation during axonal conduction. *Science*, 331(6017):599–601.
- Savin, C., Triesch, J., and Meyer-Hermann, M. (2009). Epileptogenesis due to glia-mediated synaptic scaling. *Journal of The Royal Society Interface*, 6(37):655–668.

- Savtchenko, L. P. and Rusakov, D. A. (2014). Regulation of rhythm genesis by volume-limited, astroglia-like signals in neural networks. *Phil. Trans. Royal Soc. B: Biological Sciences*, 369(1654):20130614.
- Savtchouk, I. and Volterra, A. (2018). Gliotransmission: Beyond black-and-white. *Journal of Neuroscience*, 38(1):14–25.
- Scemes, E. and Giaume, C. (2006). Astrocyte calcium waves: What they are and what they do. *Glia*, 54:716–725.
- Schafer, D. P., Lehrman, E. K., Kautzman, A. G., Koyama, R., Mardinly, A. R., Yamasaki, R., Ransohoff, R. M., Greenberg, M. E., Barres, B. A., and Stevens, B. (2012). Microglia sculpt postnatal neural circuits in an activity and complement-dependent manner. *Neuron*, 74(4):691–705.
- Schipke, C. G., Boucsein, C., Ohlemeyer, C., Kirchhoff, F., and Kettenmann, H. (2002). Astrocyte  $\text{Ca}^{2+}$  waves trigger responses in microglial cells in brain slices. *The FASEB Journal*, 16(2):255–257.
- Schmithorst, V. J. and Wilke, M. (2002). Differences in white matter architecture between musicians and non-musicians: a diffusion tensor imaging study. *Neuroscience letters*, 321(1-2):57–60.
- Schmitt, L., Sims, R., Dale, N., and Haydon, P. (2012). Wakefulness affects synaptic and network activity by increasing extracellular astrocyte-derived adenosine. *J. Neurosci.*, 32(13):4417–4425.
- Scholz, J., Klein, M. C., Behrens, T. E. J., and Johansen-Berg, H. (2009). Training induces changes in white-matter architecture. *Nature Neuroscience*, 12(11):1370.
- Schummers, J., Yu, H., and Sur, M. (2008). Tuned responses of astrocytes and their influence on hemodynamic signals in the visual cortex. *Sci. STKE*, 320(5883):1638.
- Schuster, S., Marhl, M., and Höfer, T. (2002). Modelling of simple and complex calcium oscillations from single-cell responses to intercellular signalling. *Eur. J. Biochem.*, 269:1333–1355.
- Seidl, A. H., Rubel, E. W., and Harris, D. M. (2010). Mechanisms for adjusting interaural time differences to achieve binaural coincidence detection. *Journal of Neuroscience*, 30(1):70–80.
- Seifert, G. and Steinhäuser, C. (2017). Heterogeneity and function of hippocampal macroglia. *Cell and Tissue Research*, pages 1–18.
- Seung, H. S., Lee, D. D., Reis, B. Y., and Tank, D. W. (2000). The autapse: a simple illustration of short-term analog memory storage by tuned synaptic feedback. *Journal of Computational Neuroscience*, 9(2):171–185.
- Sheikhhahaei, S., Turovsky, E. A., Hosford, P. S., Hadjihambi, A., Theparambil, S. M., Liu, B., Nephtali, M., Teschemacher, A. G., Kasparov, S., Smith, J. C., and Gourine, A. V. (2018). Astrocytes modulate brainstem respiratory rhythm-generating circuits and determine exercise capacity. *Nature communications*, 9(1):370.
- Sherwood, M. W., Arizono, M., Hisatsune, C., Bannai, H., Ebisui, E., Sherwood, J. L., Panatier, A., Oliet, S. H. R., and Mikoshiba, K. (2017). Astrocytic ip3rs: Contribution to  $\text{ca}^{2+}$  signalling and hippocampal ltp. *Glia*, 65(3):502–513.

- Shigetomi, E., Jackson-Weaver, O., Huckstepp, R. T., O'Dell, T. J., and Khakh, B. S. (2013). TRPA1 channels are regulators of astrocyte basal calcium levels and long-term potentiation via constitutive D-serine release. *The Journal of Neuroscience*, 33(24):10143–10153.
- Shigetomi, E., Tong, X., Kwan, K., Corey, D., and Khakh, B. (2012). TRPA1 channels regulate astrocyte resting calcium levels and inhibitory synapse efficacy via GAT-3. *Nat. Neurosci.*, 15(1):70–80.
- Siegel, J. M. (2005). Clues to the functions of mammalian sleep. *Nature*, 437(7063):1264.
- Sild, M. and Ruthazer, E. S. (2011). Radial glia: progenitor, pathway, and partner. *The Neuroscientist*, 17(3):288–302.
- Simard, M. and Nedergaard, M. (2004). The neurobiology of glia in the context of water and ion homeostasis. *Neuroscience*, 129(4):877–896.
- Simpson, I. A., Carruthers, A., and Vannucci, S. J. (2007). Supply and demand in cerebral energy metabolism: the role of nutrient transporters. *Journal of Cerebral Blood Flow & Metabolism*, 27(11):1766–1791.
- Sipe, G. O., Lowery, R. L., Tremblay, M.-È., Kelly, E. A., Lamantia, C. E., and Majewska, A. K. (2016). Microglial P2Y<sub>12</sub> is necessary for synaptic plasticity in mouse visual cortex. *Nature Communications*, 7:10905.
- Sira-Ramirez, H. and Colina-Morles, E. (1995). A sliding mode strategy for adaptive learning in adalines. *IEEE Transactions on Circuits and Systems I: Fundamental Theory and Applications*, 42(12):1001–1012.
- Skupin, A., Kettenmann, H., Winkler, U., Wartenberg, M., Sauer, H., Tovey, S. C., Taylor, C. W., and Falcke, M. (2008). How does intracellular Ca<sup>2+</sup> oscillate: by chance or by clock? *Biophys. J.*, 94:2404–2411.
- Somersalo, E., Cheng, Y., and Calvetti, D. (2012). The metabolism of neurons and astrocytes through mathematical models. *Annals of Biomedical Engineering*, 40(11):2328–2344.
- Somjen, G. G. (2002). Ion regulation in the brain: implications for pathophysiology. *The Neuroscientist*, 8(3):254–267.
- Somjen, G. G., Kager, H., and Wadman, W. J. (2008). Computer simulations of neuron-glia interactions mediated by ion flux. *Journal of Computational Neuroscience*, 25(2):349–365.
- Sorg, O. and Magistretti, P. J. (1991). Characterization of the glycogenolysis elicited by vasoactive intestinal peptide, noradrenaline and adenosine in primary cultures of mouse cerebral cortical astrocytes. *Brain Research*, 563(1-2):227–233.
- Sorg, O. and Magistretti, P. J. (1992). Vasoactive intestinal peptide and noradrenaline exert long-term control on glycogen levels in astrocytes: blockade by protein synthesis inhibition. *Journal of Neuroscience*, 12(12):4923–4931.
- Srinivasan, R., Huang, B. S., Venugopal, S., Johnston, A. D., Chai, H., Zeng, H., Golshani, P., and Khakh, B. S. (2015). Ca<sup>2+</sup> signaling in astrocytes from Ip3r2<sup>-/-</sup> mice in brain slice and during startle responses *in vivo*. *Nature Neuroscience*, 18:708–717.
- Stassart, R., Goebbels, S., and Nave, K.-A. (2013). *Neuroglia*, chapter Factors controlling myelin formation, pages 555–572. Oxford University Press.

- Stehberg, J., Moraga-Amaro, R., Salazar, C., Becerra, A., Echeverría, C., Orellana, J. A., Bultynck, G., Ponsaerts, R., Leybaert, L., Simon, F., Simon, F., Sáez, J. C., and Retamal, M. A. (2012). Release of gliotransmitters through astroglial connexin 43 hemichannels is necessary for fear memory consolidation in the basolateral amygdala. *The FASEB Journal*, 26(9):3649–3657.
- Steinhäuser, C., Berger, T., Frotscher, M., and Kettenmann, H. (1992). Heterogeneity in the membrane current pattern of identified glial cells in the hippocampal slice. *European Journal of Neuroscience*, 4(6):472–484.
- Steinmetz, C. C. and Turrigiano, G. G. (2010). Tumor necrosis factor- $\alpha$  signaling maintains the ability of cortical synapses to express synaptic scaling. *Journal of Neuroscience*, 30(44):14685–14690.
- Stellwagen, D., Beattie, E. C., Seo, J. Y., and Malenka, R. C. (2005). Differential regulation of AMPA receptor and GABA receptor trafficking by tumor necrosis factor- $\alpha$ . *J. Neurosci.*, 25(12):3219–3228.
- Stellwagen, D. and Malenka, R. C. (2006). Synaptic scaling mediated by glial TNF- $\alpha$ . *Nature*, 440(7087):1054–1059.
- Stevens, B., Allen, N. J., Vazquez, L. E., Howell, G. R., Christopherson, K. S., Nouri, N., Micheva, K. D., Mehalow, A. K., Huberman, A. D., Stafford, B., Sher, A., Litke, A. M., Lambiris, J. D., Smith, S. J., John, S. W. M., and Barres, B. A. (2007). The classical complement cascade mediates CNS synapse elimination. *Cell*, 131(6):1164–1178.
- Stevens, J.-L. R., Law, J. S., Antolík, J., and Bednar, J. A. (2013). Mechanisms for stable, robust, and adaptive development of orientation maps in the primary visual cortex. *Journal of Neuroscience*, 33(40):15747–15766.
- Stobart, J. L., Ferrari, K. D., Barrett, M. J. P., Glück, C., Stobart, M. J., Zuend, M., and Weber, B. (2018). Cortical circuit activity evokes rapid astrocyte calcium signals on a similar timescale to neurons. *Neuron*, 98:726–735.
- Stout, C. E., Costantin, J. L., Naus, C. C. G., and Charles, A. C. (2002). Intercellular calcium signaling in astrocytes via ATP release through connexin hemichannels. *J. Biol. Chem.*, 277(12):10482–10488.
- Strohschein, S., Hüttmann, K., Gabriel, S., Binder, D. K., Heinemann, U., and Steinhäuser, C. (2011). Impact of aquaporin-4 channels on K<sup>+</sup> buffering and gap junction coupling in the hippocampus. *Glia*, 59(6):973–980.
- Suadicani, S. O., Brosnan, C. F., and Scemes, E. (2006). P2X<sub>7</sub> receptors mediate ATP release and amplification of astrocytic intercellular Ca<sup>2+</sup> signaling. *Journal of Neuroscience*, 26(5):1378–1385.
- Südhof, T. C. (2004). The synaptic vesicle cycle. *Annu. Rev. Neurosci.*, 27:509–547.
- Sugihara, I., Lang, E. J., and Llinás, R. (1993). Uniform olivocerebellar conduction time underlies purkinje cell complex spike synchronicity in the rat cerebellum. *The Journal of Physiology*, 470(1):243–271.
- Suzuki, A., Stern, S. A., O., B., Huntely, G., Wlaker, R. H., Magistretti, P. J., and Alberini, C. M. (2011). Astrocyte-neuron lactate transport is required for long-term memory formation. *Cell*, 144:810–823.



- Swadlow, H. A. (1985). Physiological properties of individual cerebral axons studied in vivo for as long as one year. *Journal of Neurophysiology*, 54(5):1346–1362.
- Takata, N., Mishima, T., Hisatsune, C., Nagai, T., Ebisui, E., Mikoshiba, K., and Hirase, H. (2011). Astrocyte calcium signaling transforms cholinergic modulation to cortical plasticity *in vivo*. *J. Neurosci.*, 31(49):18155–18165.
- Takeda, M., Nelson, D. J., and Soliven, B. (1995). Calcium signaling in cultured rat oligodendrocytes. *Glia*, 14(3):225–236.
- Tang, F., Lane, S., Korsak, A., Paton, J. F. R., Gourine, A. V., Kasparov, S., and Teschemacher, A. G. (2014). Lactate-mediated glia-neuronal signalling in the mammalian brain. *Nature Communications*, 5:3284.
- Tani, H., Dulla, C. G., Farzampour, Z., Taylor-Weiner, A., Huguenard, J. R., and Reimer, R. J. (2014). A local glutamate-glutamine cycle sustains synaptic excitatory transmitter release. *Neuron*, 81(4):888–900.
- Tasaki, I. and Chang, J. J. (1958). Electric response of glia cells in cat brain. *Science*, 128(3333):1209–1210.
- Taschenberger, H. and von Gersdorff, H. (2000). Fine-tuning an auditory synapse for speed and fidelity: developmental changes in presynaptic waveform, EPSC kinetics, and synaptic plasticity. *Journal of Neuroscience*, 20(24):9162–9173.
- Theodosis, D. T. and Poulain, D. A. (1984). Evidence for structural plasticity in the supraoptic nucleus of the rat hypothalamus in relation to gestation and lactation. *Neuroscience*, 11(1):183–193.
- Thurley, K. and Falcke, M. (2011). Derivation of  $\text{Ca}^{2+}$  signals from puff properties reveals that pathway function is robust against cell variability but sensitive for control. *Proc. Nat. Acad. Sci. USA*, 108(1):427–432.
- Thurley, K., Skupin, A., Thul, R., and Falcke, M. (2012). Fundamental properties of  $\text{Ca}^{2+}$  signals. *Biochimica et Biophysica Acta*, 1820:11851194.
- Thurley, K., Tovey, S. C., Moenke, G., Prince, V. L., Meena, A., Thomas, A. P., Skupin, A., Taylor, C. W., and Falcke, M. (2014). Reliable encoding of stimulus intensities within random sequences of intracellular  $\text{Ca}^{2+}$  spikes. *Science Signaling*, 7(331):ra59.
- Tomassy, G. S., Berger, D. R., Chen, H., Kasthuri, N., Hayworth, K. J., Vercelli, A., Seung, H. S., Lichtman, J. W., and Arlotta, P. (2014). Distinct profiles of myelin distribution along single axons of pyramidal neurons in the neocortex. *Science*, 344(6181):319–324.
- Tomassy, G. S., Dershowitz, L. B., and Arlotta, P. (2016). Diversity matters: a revised guide to myelination. *Trends in Cell Biology*, 26(2):135–147.
- Tononi, G. and Cirelli, C. (2014). Sleep and the price of plasticity: from synaptic and cellular homeostasis to memory consolidation and integration. *Neuron*, 81(1):12–34.
- Toyoizumi, T., Kaneko, M., Stryker, M. P., and Miller, K. D. (2014). Modeling the dynamic interaction of Hebbian and homeostatic plasticity. *Neuron*, 84(2):497–510.
- Tremblay, M.-È., Lowery, R. L., and Majewska, A. K. (2010). Microglial interactions with synapses are modulated by visual experience. *PLoS biology*, 8(11):e1000527.

- Tsodyks, M. (2005). Activity-dependent transmission in neocortical synapses. In Chow, C. C., Gutkin, B., D., H., C., M., and Dalibard, J., editors, *Methods and Models in Neurophysics*, pages 245–265. Elsevier.
- Tsvetkov, E., Shin, R.-M., and Bolshakov, V. Y. (2004). Glutamate uptake determines pathway specificity of long-term potentiation in the neural circuitry of fear conditioning. *Neuron*, 41(1):139–151.
- Tuchin, V. (2000). *Tissue Optics*. SPIE Press, Bellingham, WA.
- Ullah, G., Cressman Jr., J. R., Barreto, E., and Schiff, S. J. (2009). The influence of sodium and potassium dynamics on excitability, seizures, and the stability of persistent states: II. Network and glial dynamics. *Journal of computational neuroscience*, 26(2):171–183.
- Ullén, F. (2009). Is activity regulation of late myelination a plastic mechanism in the human nervous system? *Neuron Glia Biology*, 5(1-2):29–34.
- Vaishnavi, S. N., Vlassenko, A. G., Rundle, M. M., Snyder, A. Z., Mintun, M. A., and Raichle, M. E. (2010). Regional aerobic glycolysis in the human brain. *Proceedings of the National Academy of Sciences*, 107(41):17757–17762.
- Valtcheva, S. and Venance, L. (2016). Astrocytes gate Hebbian synaptic plasticity in the striatum. *Nature Communications*, 7:13845.
- Ventura, R. and Harris, K. M. (1999). Three-dimensional relationships between hippocampal synapses and astrocytes. *J. Neurosci.*, 19(16):6897–6906.
- Verbich, D., Prenosil, G. A., Chang, P. K.-Y., Murai, K. K., and McKinney, R. A. (2012). Glial glutamate transport modulates dendritic spine head protrusions in the hippocampus. *Glia*, 60(7):1067–1077.
- Verderio, C. and Matteoli, M. (2001). ATP mediates calcium signaling between astrocytes and microglial cells: modulation by IFN- $\gamma$ . *The Journal of Immunology*, 166(10):6383–6391.
- Verkhratsky, A. and Kettenmann, H. (1996). Calcium signaling in glial cells. *Trends Neurosci.*, 19:346–352.
- Verkhratsky, A. and Nedergaard, M. (2016). The homeostatic astroglia emerges from evolutionary specialization of neural cells. *Phil. Trans. R. Soc. B*, 371(1700):20150428.
- Violin, J. D., DiPilato, L. M., Yildirim, N., Elston, T. C., Zhang, J., and Lefkowitz, R. J. (2008).  $\beta$ 2-adrenergic receptor signaling and desensitization elucidated by quantitative modeling of real time cAMP dynamics. *Journal of Biological Chemistry*, 283(5):2949–2961.
- Volman, V., Bazhenov, M., and Sejnowski, T. J. (2013). Divide and conquer: functional segregation of synaptic inputs by astrocytic microdomains could alleviate paroxysmal activity following brain trauma. *PLoS Computational Biology*, 9(1):e1002856.
- Volman, V., Ben-Jacob, E., and Levine, H. (2007). The astrocyte as a gatekeeper of synaptic information transfer. *Neural Computation*, 19:303–326.
- Volterra, A., Liaudet, N., and Savtchouk, I. (2014). Astrocyte  $\text{Ca}^{2+}$  signalling: an unexpected complexity. *Nature Reviews Neuroscience*, 15:327–334.

- Vyazovskiy, V. V., Cirelli, C., Pfister-Genskow, M., Faraguna, U., and Tononi, G. (2008). Molecular and electrophysiological evidence for net synaptic potentiation in wake and depression in sleep. *Nature Neuroscience*, 11(2):200.
- Vyazovskiy, V. V. and Harris, K. D. (2013). Sleep and the single neuron: the role of global slow oscillations in individual cell rest. *Nature Reviews Neuroscience*, 14(6):443.
- Vyazovskiy, V. V., Olcese, U., Hanlon, E. C., Nir, Y., Cirelli, C., and Tononi, G. (2011). Local sleep in awake rats. *Nature*, 472(7344):443.
- Waehneltdt, T. V., Matthieu, J.-M., and Jeserich, G. (1986). Appearance of myelin proteins during vertebrate evolution. *Neurochemistry International*, 9(4):463–474.
- Wake, H., Moorhouse, A. J., Jinno, S., Kohsaka, S., and Nabekura, J. (2009). Resting microglia directly monitor the functional state of synapses *in vivo* and determine the fate of ischemic terminals. *Journal of Neuroscience*, 29(13):3974–3980.
- Wake, H., Moorhouse, A. J., Miyamoto, A., and Nabekura, J. (2013). Microglia: actively surveying and shaping neuronal circuit structure and function. *Trends in Neurosciences*, 36(4):209–217.
- Walz, W. (2000). Role of astrocytes in the clearance of excess extracellular potassium. *Neurochemistry international*, 36(4-5):291–300.
- Wang, F., Xu, Q., Wang, W., Takano, T., and Nedergaard, M. (2012). Bergmann glia modulate cerebellar purkinje cell bistability via  $\text{Ca}^{2+}$ -dependent  $\text{K}^+$  uptake. *Proc. Natl. Acad. Sci. USA*, 109(20):7911–7916.
- Wang, X., Lou, N., Xu, Q., Tian, G.-F., Peng, W. G., Han, X., Kang, J., Takano, T., and Nedergaard, M. (2006). Astrocytic  $\text{Ca}^{2+}$  signaling evoked by sensory stimulation *in vivo*. *Nat. Neurosci.*, 9(6):816–823.
- Waxman, S. G. (1980). Determinants of conduction velocity in myelinated nerve fibers. *Muscle & Nerve*, 3(2):141–150.
- Waxman, S. G. and Bennett, M. V. L. (1972). Relative conduction velocities of small myelinated and non-myelinated fibres in the central nervous system. *Nature New Biology*, 238(85):217.
- Wheeler, D. B., Randall, A., and Tsien, R. W. (1996). Changes in action potential duration alter reliance of excitatory synaptic transmission on multiple types of  $\text{Ca}^{2+}$  channels in rat hippocampus. *Journal of Neuroscience*, 16(7):2226–2237.
- Wiesel, T. N. (1982). Postnatal development of the visual cortex and the influence of environment. *Nature*, 299(5884):583.
- Winship, I., Plaa, N., and Murphy, T. (2007). Rapid astrocyte calcium signals correlate with neuronal activity and onset of the hemodynamic response *in vivo*. *J. Neurosci.*, 27(23):6268–6272.
- Witcher, M., Kirov, S., and Harris, K. (2007). Plasticity of perisynaptic astroglia during synaptogenesis in the mature rat hippocampus. *Glia*, 55(1):13–23.
- Witelson, S. F., Kigar, D. L., and Harvey, T. (1999). The exceptional brain of Albert Einstein. *The Lancet*, 353(9170):2149–2153.

- Woo, D. H., Han, K., Shim, J. W., Yoon, B., Kim, E., Bae, J. Y., Oh, S., Hwang, E. M., Marmorstein, A. D., Bae, Y. C., Park, J., and Lee, C. J. (2012). TREK-1 and Best1 channels mediate fast and slow glutamate release in astrocytes upon GPCR activation. *Cell*, 151(1):25–40.
- Wu, X., Pan, L., Liu, Y., Jiang, P., Lee, I., Drevensek-Olenik, I., Zhang, X., and Xu, J. (2013). Cell–cell communication induces random spikes of spontaneous calcium oscillations in multi-BV-2 microglial cells. *Biochemical and Biophysical Research Communications*, 431(4):664–669.
- Wu, Y., Dissing-Olesen, L., MacVicar, B. A., and Stevens, B. (2015). Microglia: dynamic mediators of synapse development and plasticity. *Trends in Immunology*, 36(10):605–613.
- Xiang, J., Ennis, S. R., Abdelkarim, G. E., Fujisawa, M., Kawai, N., and Keep, R. F. (2003). Glutamine transport at the blood–brain and blood–cerebrospinal fluid barriers. *Neurochemistry International*, 43(4-5):279–288.
- Xie, L., Kang, H., Xu, Q., Chen, M. J., Liao, Y., Thiyagarajan, M., ODonnell, J., Christensen, D. J., Nicholson, C., Iliff, J. J., Takahiro, T., Deane, R., and Nedergaard, M. (2013). Sleep drives metabolite clearance from the adult brain. *science*, 342(6156):373–377.
- Yamazaki, Y., Hozumi, Y., Kaneko, K., Sugihara, T., Fujii, S., Goto, K., and Kato, H. (2007). Modulatory effects of oligodendrocytes on the conduction velocity of action potentials along axons in the alveus of the rat hippocampal CA1 region. *Neuron Glia Biology*, 3(4):325–334.
- Yang, L., Qi, Y., and Yang, Y. (2015). Astrocytes control food intake by inhibiting AGRP neuron activity via adenosine a1 receptors. *Cell Reports*, 11(5):798–807.
- Yang, S., Qiao, H., Wen, L., Zhou, W., and Zhang, Y. (2005). D-serine enhances impaired long-term potentiation in CA1 subfield of hippocampal slices from aged senescence-accelerated mouse prone/8. *Neuroscience Letters*, 379(1):7–12.
- Yang, Y., Ge, W., Chen, Y., Zhang, Z., Shen, W., Wu, C., Poo, M., and Duan, S. (2003). Contribution of astrocytes to hippocampal long-term potentiation through release of d-serine. *Proc. Natl. Acad. Sci. USA*, 100(25):15194–15199.
- Ying, W. (2008). NAD<sup>+</sup>/NADH and NADP<sup>+</sup>/NADPH in cellular functions and cell death: regulation and biological consequences. *Antioxidants & Redox Signaling*, 10(2):179–206.
- Young, K. M., Psachoulia, K., Tripathi, R. B., Dunn, S., Cossell, L., Attwell, D., Tohyama, K., and Richardson, W. D. (2013). Oligodendrocyte dynamics in the healthy adult CNS: evidence for myelin remodeling. *Neuron*, 77(5):873–885.
- Yu, X., Zhihong, M., and Rahman, S. M. M. (1998). Adaptive sliding mode approach for learning in a feedforward neural network. *Neural Computing & Applications*, 7(4):289–294.
- Zalc, B. and Colman, D. R. (2000). Origins of vertebrate success. *Science*, 288(5464):271–271.
- Zalc, B., Goujet, D., and Colman, D. (2008). The origin of the myelination program in vertebrates. *Current Biology*, 18(12):R511–R512.
- Zenke, F., Gerstner, W., and Ganguli, S. (2017). The temporal paradox of Hebbian learning and homeostatic plasticity. *Current Opinion in Neurobiology*, 43:166–176.

- Zhan, Y., Paolicelli, R. C., Sforzini, F., Weinhard, L., Bolasco, G., Pagani, F., Vyssotski, A. L., Bifone, A., Gozzi, A., Ragozzino, D., and Gross, C. T. (2014). Deficient neuron-microglia signaling results in impaired functional brain connectivity and social behavior. *Nature Neuroscience*, 17(3):400.
- Zhang, Y., Chen, K., Sloan, S. A., Bennett, M. L., Scholze, A. R., O’Keeffe, S., Phatnani, H. P., Guarnieri, P., Caneda, C., Ruderisch, N., Deng, S., Liddelow, S. A., Zhang, C., Daneman, R., Maniatis, T., Barres, B. A., and Wu, J. Q. (2014). An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells of the cerebral cortex. *Journal of Neuroscience*, 34(36):11929–11947.
- Zhang, Z., Gong, N., Wang, W., Xu, L., and Xu, T. (2008). Bell-shaped D-serine actions on hippocampal long-term depression and spatial memory retrieval. *Cerebral Cortex*, 18(10):2391–2401.
- Zohary, E., Shadlen, M. N., and Newsome, W. T. (1994). Correlated neuronal discharge rate and its implications for psychophysical performance. *Nature*, 370(6485):140.
- Zonta, M., Angulo, M. C., Gobbo, S., Rosengarten, B., Hossmann, K. A., Pozzan, T., and Carmignoto, G. (2003). Neuron-to-astrocyte signaling is central to the dynamic control of brain microcirculation. *Nat. Neurosci.*, 6:43–40.
- Zorec, R., Araque, A., Carmignoto, G., Haydon, P., Verkhratsky, A., and Parpura, V. (2012). Astroglial excitability and gliotransmission: An appraisal of  $\text{Ca}^{2+}$  as a signaling route. *ASN Neuro*, 4(2):e00080.
- Zucker, R. S. and Regehr, W. G. (2002). Short-term synaptic plasticity. *Annu. Rev. Physiol.*, 64:355–405.