



Anticipation in the retina and the primary visual cortex: towards an integrated retino-cortical model for motion processing

Selma Souihel, Bruno Cessac, Matteo Di Volo, Alain Destexhe, Frederic Chavane, Sandrine Chemla, Olivier Marre

► To cite this version:

Selma Souihel, Bruno Cessac, Matteo Di Volo, Alain Destexhe, Frederic Chavane, et al.. Anticipation in the retina and the primary visual cortex: towards an integrated retino-cortical model for motion processing. Waves Côte d'Azur, Jun 2019, Nice, France. hal-02172010

HAL Id: hal-02172010

<https://inria.hal.science/hal-02172010>

Submitted on 3 Jul 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



AGENCE NATIONALE DE LA RECHERCHE



UNIVERSITÉ
CÔTE D'AZUR

Anticipation in the retina and the primary visual cortex : towards an integrated retino-cortical model for motion processing

PhD student : Selma Souihel
Adviser : Dr Bruno Cessac
Team : Biovision



In collaboration with :

Matteo Di Volo
Alain Destexhe




Frédéric Chavane
Sandrine Chemla





Olivier Marre



The visual flow




The visual flow



Source : Ryskamp et al. 2014

Stating the problem : Visual Anticipation

Anticipation is carried out by the primary visual cortex (V1) through an activation wave



Source :
Benvenutti et
al. 2015

But the retina is not a mere transmitter, it is able to perform many computations such as :

- Orientation sensitivity
- Contrast gain control
- Sensitivity to differential motion
- and « Motion Anticipation »


➤ What does retinal anticipation add to the cortical one ?

I) Anticipation in the retina

The Hubel-Wiesel view of vision



Nobel prize 1981

Ganglion cells response is the convolution of the stimulus with a spatio-temporal receptive field followed by a non linearity



The Hubel-Wiesel view of vision


Nobel prize 1981



Source : Berry et al. 1999

The Hubel-Wiesel view of vision

Nobel prize 1981





Source : Berry et al. 1999

- Which mechanism can account for motion anticipation in the retina ?

Building a 2D retina model for motion anticipation

1) Gain control

How does it work ?




$$\mathcal{G}_B(A) = \begin{cases} 0, & \text{if } A \leq 0; \\ \frac{1}{1+A^6}, & \text{else.} \end{cases}$$

$$R_{B_i} = \mathcal{N}_B(V_{B_i}) \mathcal{G}_B(A_{B_i}).$$

Building a 2D retina model for motion anticipation

1) Gain control (Chen et al. 2013)



- Bipolar voltage :

$$V_{B_i}(t) = V_{i_{drive}}(t) + P_{B_i}(t).$$

- Non-linear function :

$$\mathcal{N}_B(V_{B_i}) = \begin{cases} 0, & \text{if } V_{B_i} \leq \theta_B; \\ V_{B_i} - \theta_B, & \text{else.} \end{cases}$$

- Activation function :

$$\frac{dA_{B_i}}{dt} = -\frac{A_{B_i}}{\tau_a} + h\mathcal{N}(V_{B_i}(t)).$$

- Gain Control function :


$$\mathcal{G}_B(A) = \begin{cases} 0, & \text{if } A \leq 0; \\ \frac{1}{1+A^6}, & \text{else.} \end{cases}$$

- Output :

$$R_{B_i} = \mathcal{N}_B(V_{B_i}) \mathcal{G}_B(A_{B_i}).$$

Building a 2D retina model for motion anticipation

1) Gain control (Chen et al. 2013)



- Ganglion voltage

$$V_{G_k} = \sum_i W_{G_k}^{B_i} R_{B_i}$$

- Non-linear function :

$$\mathcal{N}_{G_F}(V) = \begin{cases} 0, & \text{if } V \leq 0; \\ \alpha_{G_F}(V - \theta_{G_F}), & \text{if } \theta_{G_F} \leq V \leq N_{G_F}^{\max}/\alpha_{G_F} + \theta_{G_F} \\ N_{G_F}^{\max}, & \text{else.} \end{cases}$$

- Activation function :

$$\frac{dA_{G_F k_F}}{dt} = -\frac{A_{G_F k_F}}{\tau_{G_F}} + h_{G_F} \mathcal{N}_{G_F} \left(V_{G_F k_F} \right)$$

- Gain Control function :


$$\mathcal{G}_{G_F}(A) = \begin{cases} 0, & \text{if } A \leq 0; \\ \frac{1}{1+A}, & \text{else.} \end{cases}$$

- Output :


$$R_{G_F k_F} \left(V_{G_F k_F}, A_{G_F k_F} \right) = \mathcal{N}_{G_F}(V_{G_F k_F}) \mathcal{G}_{G_F}(A_{G_F k_F}).$$

1D results : smooth motion anticipation with gain control

Bipolar layer






Ganglion layer



1D results : smooth motion anticipation with gain control



Anticipation variability with stimulus parameters



Building a 2D retina model for motion anticipation




2) Gap junctions connectivity

- A class of direction selective RGCs are connected through gap junctions
- Their activity comprises the activity pooled from bipolar cells and the activity coming from the downstream RGCs, in the direction of motion




$$R_{G_D k_D} = V_{G_D k_D} + \beta R_{G_D k_D - 1}$$

1D results : smooth motion anticipation with gap junctions




1D results : smooth motion anticipation with gap junctions



1D results : smooth motion anticipation with gap junctions

Anticipation variability with stimulus parameters



Building a 2D retina model for motion anticipation

3) Amacrine cells connectivity

- A class of RGCs are selective to differential motion



Building a 2D retina model for motion anticipation

3) Amacrine cells connectivity


- A class of RGCs are selective to differential motion



- The circuitry involves amacrine cells connectivity upstream of ganglion cells

Connectivity pathways

2) Amacrine cells connectivity



- Bipolar voltage :

$$\frac{dV_{B_i}}{dt} = -\frac{1}{\tau_B} V_{B_i} + \sum_{j=1}^{N_A} W_{B_i}^{A_j} V_{A_j} + F_{B_i}(t).$$

- External drive :

$$F_{B_i}(t) = \left[K_i *^S \left(\frac{\mathcal{S}}{\tau_B} + \frac{d\mathcal{S}}{dt} \right) \right] (t)$$

- Amacrine voltage :


$$\frac{dV_{A_j}}{dt} = -\frac{1}{\tau_A} V_{A_j}(t) + \sum_{i=1}^{N_A} W_{A_j}^{B_i} R_{B_i}(t).$$

- Coupled dynamics :


$$\begin{cases} \frac{dV_{B_i}}{dt} &= -\frac{1}{\tau_B} V_{B_i} + \sum_{j=1}^{N_A} W_{B_i}^{A_j} V_{A_j} + F_{B_i}(t) \\ \frac{dA_{B_i}}{dt} &= -\frac{A_{B_i}}{\tau_a} + h \mathcal{N}(V_{B_i}(t)), \\ \frac{dV_{A_j}}{dt} &= -\frac{1}{\tau_A} V_{A_j}(t) + \sum_{i=1}^{N_A} W_{A_j}^{B_i} R_{B_i}(t). \end{cases}$$

1D results : smooth motion anticipation with amacrine connectivity

Bipolar layer




Ganglion layer




1D results : smooth motion anticipation with amacrine connectivity

Anticipation variability with stimulus parameters




Comparing the performance of the three layers




Suggesting new experiments : 2D results


1) Angular anticipation



B)




C)




Suggesting new experiments : 2D results

1) Angular anticipation




Suggesting new experiments : 2D results

2) Anticipation and shape




Suggesting new experiments : 2D results

2) Anticipation on a noisy background




II) Anticipation in V1

Anticipation in V1







Source : Benvenutti et al. 2015




Anticipation in the cortex : VSDI data analysis

(Data courtesy of F. Chavane et S. Chemla)




A mean field model to reproduce VSDI recordings


Zerlaut et al 2016
Chemla et al 2018



Response of the cortical model to a LN retina drive




Response of the cortical model to a retina drive with gain control




Comparing simulation results to VSDI recordings

Cortex experimental recordings



Simulation results
Response to an LN model of the retina



Simulation results
Response to a gain control model of the retina

Conclusions

- We developed a 2D retina with three ganglion cell layers, implementing gain control and connectivity.
- We use the output of our model as an input to a mean field model of V1, and were able to reproduce anticipation as observed in VSDI

Questions :

- How to improve object identification 1) exploring the model's parameters and 2) using connectivity ?
- Is our model able to anticipate more complex trajectories, with accelerations for instance ?
- How to calibrate connectivity using biology ?
- How does anticipation affect higher order correlations ?
- Would it be possible to design psycho-physical tests clearly showing the role of the retina in visual anticipation ?


Thank you for your attention !

Supplementary material


Stimulus integration

$$\left[K_i \stackrel{S,t}{*} \mathcal{S} \right] (t) = \int_{-\infty}^t K_T(t-u) \left[\int_{\mathbb{R}^2} K_{i,S}(x,y) \mathcal{S}(x,y,u) dx dy \right] du \equiv V_{i_{drive}}(t).$$

A)



B)




$$K_{i,S}(x,y) = \frac{A_1}{2\pi\sqrt{\det C_1}} e^{-\frac{1}{2}\tilde{X}_i \cdot C_1^{-1} \cdot \tilde{X}_i} - \frac{A_2}{2\pi\sqrt{\det C_2}} e^{-\frac{1}{2}\tilde{X}_i \cdot C_2^{-1} \cdot \tilde{X}_i},$$

$$K_T(t) = \left(\frac{K_1}{\sqrt{2\pi}\sigma_1} e^{-\frac{(t-\mu_1)^2}{2\sigma_1^2}} - \frac{K_2}{\sqrt{2\pi}\sigma_2} e^{-\frac{(t-\mu_2)^2}{2\sigma_2^2}} \right) H(t)$$

(Data courtesy of O.
Marre)

Stimulus integration : anisotropy




$$\sigma_{x'} = \frac{\sigma_x \sigma_y}{\sqrt{\sigma_x^2 \cos^2 \theta + \sigma_y^2 \sin^2 \theta}}$$

$$\sigma_\phi = \frac{\sqrt{\sigma_y^2 \cos^2 \theta + \sigma_x^2 \sin^2 \theta}}{\sin \phi}$$


$$\tan(\phi) = \frac{\sigma_y^2 \cos^2 \theta + \sigma_x^2 \sin^2 \theta}{(\sigma_x^2 - \sigma_y^2) \cos \theta \sin \theta}$$

$$I = \sigma_x \sqrt{\frac{\pi}{2}} \sum_{(i;j) \in [0, s_x] \times [0, s_y]} \int_y^{(y+1)\frac{\delta}{\sin(\phi)}} C_{ij} e^{-\frac{(y' - y'_0)^2}{2\sigma_\phi^2}}$$

$$[erf(\frac{(-\cos(\phi)y' + x + 1)\delta - x'_0}{\sqrt{2}\sigma_{x'}}) - erf(\frac{(-\cos(\phi)y' + x)\delta - x'_0}{\sqrt{2}\sigma_{x'}})]dy'$$



Connectivity graph



- Variables of the model :
 - Number of branches N
 - Branch length X
 - Branch angle α

$$F_X(x) = \int_0^1 \frac{1}{\pi\sqrt{1-\delta^2}} \frac{x}{x+\delta} d\delta$$

$$\sin(\beta + \frac{\arcsin |y_j - y_i|}{\sqrt{(x_j - y_i)^2 + (y_j - x_j)^2}})$$

Cortex mean field model

Zerlaut et al 2016
Chemla et al 2018

Single neuron model (The adaptative exponential integrate and fire model Brette and Gerstner, 2005)

$$\begin{cases} C_m \frac{dV}{dt} = g_L (E_L - V) + I_{syn}(V, t) + k_a e^{\frac{V - V_{thre}}{k_a}} - I_w \\ \tau_w \frac{dI_w}{dt} = -I_w + a \cdot (V - E_L) + \sum_{t_s \in \{t_{spike}\}} b \delta(t - t_s) \end{cases}$$

The conductance-based exponential synapse

$$I_{syn}(V, t) = \sum_{s \in \{e, i\}} \sum_{t_s \in \{t_s\}} Q_s (E_s - V) e^{-\frac{t - t_s}{\tau_s}} \mathcal{H}(t - t_s)$$

Semi analytical transfer function :

$$\nu_{out} = \mathcal{F}(\nu_e, \nu_i) = \frac{1}{2\tau_V} \cdot Erfc\left(\frac{V_{thre}^{eff} - \mu_V}{\sqrt{2}\sigma_V}\right) \quad \text{with} \quad V_{thre}^{eff}(\mu_V, \sigma_V, \tau_V^N) = P_0 + \sum_{x \in \{\mu_V, \sigma_V, \tau_V^N\}} P_x \cdot \left(\frac{x - x^0}{\delta x^0}\right) + P_{\mu_G} \log\left(\frac{\mu_G}{g_L}\right) + \sum_{x, y \in \{\mu_V, \sigma_V, \tau_V^N\}} P_{xy} \cdot \left(\frac{x - x^0}{\delta x^0}\right) \left(\frac{y - y^0}{\delta y^0}\right)$$

Cortex mean field model

Zerlaut et al 2016
Chemla et al 2018

The mean, standard deviation and auto-correlation time of the excitatory and inhibitory conductance read :

$$\begin{array}{ll}
 \mu_{Ge}(\nu_e, \nu_i) = \nu_e K_e \tau_e Q_e & \rightarrow \mu_G(\nu_e, \nu_i) = \mu_{Ge} + \mu_{Gi} + g_L \\
 \sigma_{Ge}(\nu_e, \nu_i) = \sqrt{\frac{\nu_e K_e \tau_e}{2}} Q_e & \tau_m(\nu_e, \nu_i) = \frac{C_m}{\mu_G} \\
 \mu_{Gi}(\nu_e, \nu_i) = \nu_i K_i \tau_i Q_i & \downarrow \\
 \sigma_{Gi}(\nu_e, \nu_i) = \sqrt{\frac{\nu_i K_i \tau_i}{2}} Q_i & \mu_V(\nu_e, \nu_i) = \frac{\mu_{Ge} E_e + \mu_{Gi} E_i + g_L E_L}{\mu_G} \\
 & \sigma_V(\nu_e, \nu_i) = \sqrt{\sum_s K_s \nu_s \frac{(U_s \cdot \tau_s)^2}{2(\tau_m^{\text{eff}} + \tau_s)}} \\
 & \tau_V(\nu_e, \nu_i) = \left(\frac{\sum_s (K_s \nu_s (U_s \cdot \tau_s)^2)}{\sum_s (K_s \nu_s (U_s \cdot \tau_s)^2 / (\tau_m^{\text{eff}} + \tau_s))} \right)
 \end{array}$$

Finally, the transfer function reads :

$$\nu_{out} = \mathcal{F}(\nu_e, \nu_i) = \frac{1}{2 \tau_V} \cdot \operatorname{Erfc}\left(\frac{V_{thre}^{eff} - \mu_V}{\sqrt{2} \sigma_V}\right)$$

Cortex mean field model

Zerlaut et al 2016
Chemla et al 2018

Master equation for first and second moments local population dynamics (El Boustani and Destexhe, 2009) read :

$$\left\{ \begin{array}{l} T \frac{\partial \nu_\mu}{\partial t} = (\mathcal{F}_\mu - \nu_\mu) + \frac{1}{2} c_{\lambda\eta} \frac{\partial^2 \mathcal{F}_\mu}{\partial \nu_\lambda \partial \nu_\eta} \\ T \frac{\partial c_{\lambda\eta}}{\partial t} = A_{\lambda\eta} + (\mathcal{F}_\lambda - \nu_\lambda)(\mathcal{F}_\eta - \nu_\eta) + \\ \quad c_{\lambda\mu} \frac{\partial \mathcal{F}_\mu}{\partial \nu_\lambda} + c_{\mu\eta} \frac{\partial \mathcal{F}_\mu}{\partial \nu_\eta} - 2c_{\lambda\eta} \end{array} \right. \longrightarrow T \frac{\partial \nu_\mu}{\partial t} = \mathcal{F}_\mu - \nu_\mu$$

$$A_{\lambda\eta} = \begin{cases} \frac{\mathcal{F}_\lambda (1/T - \mathcal{F}_\lambda)}{N_\lambda} & \text{if } \lambda = \eta \\ 0 & \text{otherwise} \end{cases}$$