



**HAL**  
open science

## Crooked functions

Pascale Charpin

► **To cite this version:**

| Pascale Charpin. Crooked functions. 2019. hal-02337711

**HAL Id: hal-02337711**

**<https://inria.hal.science/hal-02337711>**

Preprint submitted on 29 Oct 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# *Crooked* functions

Pascale Charpin\*

October 29, 2019

## Abstract

Crooked permutations were defined twenty years ago. It was firstly shown that they can be used to construct interesting objects in graph theory. The field of applications was extended later, since crooked functions, bijective or not, correspond to APN functions and to some optimal codes. We adopt an unified presentation, of crooked functions, explaining the connexion with partially-bent functions. We then complete some known results and propose new properties. For instance, crooked functions allow to construct sets of bent functions, or simply define some permutations.

**Keywords:** Vectorial function, Boolean function, derivative, differential set, plateaued function, partially-bent functions, bent functions, APN function, AB function, permutation.

## 1 Introduction

The *crooked functions* have been introduced by Bending and Fon-Der-Flaass in 1998, as combinatorial objects of great interest [1]. Such a function has been defined from  $V$  to  $W$ , two  $n$ -dimensional vector spaces over  $\mathbb{F}_2$ , by the following property: the image set of any of its derivatives is the complement of a hyperplane. This characterization implies that a crooked function is bijective, and allows, in particular, to construct distance regular graphs. Later, several authors have developed this work, and have generalised the

---

\*INRIA, 2 rue Simone Iff, Paris 75012

previous definition. They notably related the crooked functions with several optimal objects, which have applications both to cryptography and coding theory [12, 13].

In this paper, we want to recall what is known about crooked functions today. In that sense, this paper is a survey, including several known results. However, we propose another approach, starting from the so-called *partially-bent* functions, and present some new results.

After preliminaries, we propose a brief survey on the (few) papers concerning crooked functions. As we can know, the list of references includes all this papers. Section 4 is devoted to the structure of crooked functions. We begin by proving the link between partially-bent functions and crooked functions. We later differentiate the two cases: odd and even number of variables. The odd case was mainly treated in the first papers, since in this case, crooked functions could be permutations. In Sections 5 and 6, we show how to construct, respectively, a set of bent functions and a set of permutations, since the nice structure of any crooked function. We conclude by the main conjecture about the existence of crooked functions.

## 2 Definitions, basic properties

Throughout this paper,  $|E|$  denotes the cardinality of the set  $E$ , and  $E^* = E \setminus \{0\}$ . Let  $F$  be a mapping, from the finite field  $\mathbb{F}_{2^n}$  to itself. Such a function is called a *vectorial function*, while a function  $f$ , from  $\mathbb{F}_{2^n}$  to  $\mathbb{F}_2$  is, as usually, a *Boolean function*. We denote by  $\mathcal{I}m(\xi)$  the image set of any function  $\xi$ .

A vectorial function  $F$ , from  $\mathbb{F}_{2^n}$  to itself, is said to be an *almost perfect nonlinear (APN) function* if and only if all the equations,

$$F(x) + F(x + a) = c, \quad a, c \in \mathbb{F}_{2^n}, \quad a \neq 0, \quad (1)$$

have zero or two solutions in  $\mathbb{F}_{2^n}$ , say  $x$  and  $x + a$ . Throughout this paper, we call *derivative of  $F$* , with respect to  $a$ , the function from  $\mathbb{F}_{2^n}$  to  $V$ :

$$x \mapsto D_a F(x) = F(x) + F(x + a), \quad a \in \mathbb{F}_{2^n}^*.$$

where  $V$  will be a subfield of  $\mathbb{F}_{2^n}$ , usually  $\mathbb{F}_{2^n}$  or  $\mathbb{F}_2$ . We call *differential set*, in point  $a$ , the image set of  $D_a F$ :

$$\mathcal{I}m(D_a F) = \{ F(x) + F(x + a) \mid x \in \mathbb{F}_{2^n} \}. \quad (2)$$

Clearly, when  $F$  is APN, we have for any  $a \in \mathbb{F}_{2^n}^*$ :

$$D_a F(x) = D_a F(x + a) = c, \text{ for some } c \text{ and for only one } x.$$

This means that  $D_a F$  is a 2-to-1 function. Thus, one can formulate the APN property as follows.

**Proposition 1** *Any vectorial function  $F$ , over  $\mathbb{F}_{2^n}$ , is APN if and only if all their differential sets have cardinality  $2^{n-1}$ .*

The  $2^n$ , so-called, *components of  $F$*  are the Boolean functions

$$f_\lambda : x \mapsto \text{Tr}(\lambda F(x)), \quad \lambda \in \mathbb{F}_{2^n},$$

where  $f_0$  is the null function, by convention. They are linearly defined by means of the absolute trace on  $\mathbb{F}_{2^n}$ :

$$x \mapsto \text{Tr}(x) = x + x^2 + \dots + x^{2^{n-1}}.$$

The dual  $V^\perp$ , of any subspace  $V$  of  $\mathbb{F}_{2^n}$ , is the subspace of those  $y$  such that  $\text{Tr}(yx) = 0$ , for all  $x \in V$ . The *Walsh transform* of a Boolean function  $f$ , is defined as

$$a \in \mathbb{F}_{2^n} \mapsto \mathcal{W}_f(a) = \sum_{x \in \mathbb{F}_{2^n}} (-1)^{f(x) + \text{Tr}(ax)}.$$

Recall the *Parseval's relation*:

$$\sum_{a \in \mathbb{F}_{2^n}} (\mathcal{W}_f(a))^2 = 2^{2n}.$$

We will need the following result:

**Lemma 1** [6, Lemma V.2] *Let  $f$  be a Boolean function over  $\mathbb{F}_{2^n}$ , and let  $V$  be a subspace of  $\mathbb{F}_{2^n}$  of dimension  $k$ . Then*

$$\sum_{v \in V} (\mathcal{W}_f(v))^2 = 2^k \sum_{u \in V^\perp} \sum_{x \in \mathbb{F}_{2^n}} (-1)^{f(x) + f(x+u)}.$$

We now define particular APN functions, which exist for odd  $n$  only.

**Definition 1** *The function  $F$  is said to be an almost bent (AB) function if the numbers*

$$\mu_F(a, \lambda) = \sum_{x \in \mathbb{F}_{2^n}} (-1)^{\text{Tr}(\lambda F(x) + ax)}, \quad (3)$$

*are equal to 0 or  $\pm 2^{\frac{n+1}{2}}$  only, when  $a \in \mathbb{F}_{2^n}$  and  $\lambda \in \mathbb{F}_{2^n}^*$ .*

$\mathcal{W}_f(u)$	number of $u \in \mathbb{F}_{2^n}$
0	$2^n - 2^{n-s}$
$2^{(n+s)/2}$	$2^{n-s-1} + (-1)^{f(0)} 2^{(n-s)/2-1}$
$-2^{(n+s)/2}$	$2^{n-s-1} - (-1)^{f(0)} 2^{(n-s)/2-1}$

Table 1: Walsh spectrum of the Boolean  $s$ -plateaued function  $f$

Note that  $\mu_F(a, \lambda) = \mathcal{W}_{f_\lambda}(a)$  for any fixed  $\lambda$ .

A Boolean function  $f$ , over  $\mathbb{F}_{2^n}$ , is said to be *bent* when  $\mathcal{W}_f$  takes two values  $\{\pm 2^{n/2}\}$  only, so that  $n$  must be even. It is said to be *s-plateaued* when either it is bent or  $\mathcal{W}_f$  takes three values,

$$\{0, \pm 2^{(n+s)/2}\}, \text{ with } 1 \leq s \leq n-2 \text{ and } n+s \text{ even.}$$

By convention, a bent function is 0-plateaued. The value  $2^{(n+s)/2}$  is the *amplitude* of  $f$ . A *plateaued vectorial function* is a vectorial function whose components are plateaued Boolean functions. It is said that  $F$  is *plateaued with single amplitude*, when all components of  $F$  have the same amplitude.

The *sum-of-square* indicator of  $f$  is defined by

$$\nu(f) = \sum_{a \in \mathbb{F}_{2^n}} \mathcal{W}_{D_a f}^2(0) = 2^{-n} \sum_{b \in \mathbb{F}_{2^n}} \mathcal{W}_f^4(b). \quad (4)$$

If  $f$  is  $s$ -plateaued then  $\nu(f) = 2^{2n+s}$ . Moreover, the vectorial function  $F$  is APN if and only if

$$\sum_{\lambda \in \mathbb{F}_{2^n}^*} \nu(f) = 2^{2n+1}(2^n - 1) \quad (5)$$

(see [2, Corollary 1]). A Boolean function  $f$  is said to be *balanced* if it takes the values 0 and 1 the same number of times. Recall a well-known result:

**Theorem 1** *A vectorial function  $F$  is a permutation if and only if all its components  $f_\lambda$ ,  $\lambda \in \mathbb{F}_{2^n}^*$ , are balanced.*

### 3 Brief record

We use our terminology, of the previous section, rather than of the initial works on crooked functions. The next definition was proposed by Bending and Fon-Der-Flaass in [1], twenty years ago.

**Definition 2** Let  $F$  be a function from  $\mathbb{F}_{2^n}$  to itself. This function is called *crooked* if it satisfies the following three properties:

- (i)  $F(0) = 0$ ;
- (ii)  $F(x) + F(y) + F(z) + F(x + y + z) \neq 0$ , for any three distinct  $x, y, z$ ;
- (iii)  $D_a F(x) + D_a F(y) + D_a F(z) \neq 0$ , for arbitrary  $x, y, z$  and any  $a \neq 0$ .

This definition implies that such a function  $F$  is a bijection over  $\mathbb{F}_{2^n}$ , where  $n$  must be odd. Note that the condition (ii) means that  $F$  is APN (see (1)). Also,  $F$  is crooked if and only if any of its differential sets is a complement of a hyperplane. Further, other properties are studied, in [1], such as some relations of crooked permutations with bent functions of dimension  $n - 1$ , and with the so-called *Kerdock sets*.

Crooked permutations allow to construct some *distance regular graphs*. This was shown in [1], generalizing previous constructions. Later, van Dam and Fon-Der-Flaass proposed another construction, and then another generalisation (see, in particular, Theorem 3 of [12]). Conversely, Godsil and Roy have shown that crooked permutations can be fully characterized by Preparata codes of minimum distance 5. Similarly, some distance-regular graphs provide crooked permutations [15, Theorems 3,5].

Since the high interest for APN functions in cryptography and coding theory [10], the existence of crooked functions was later the core of the research on crooked functions. Kyureghyan proposed another definition of crooked functions, identifying all APN functions which are such that their differential sets are affine hyperplanes. She established the basic properties of such functions. She notably proved that the *monomial crooked* functions are quadratic [16, 17].

The APN quadratic functions are crooked. We do not know if crooked functions of higher degree do exist. This is a recurring question, about which only negative results have been obtained. An important result was obtained by Bierbrauer and Kyureghyan: *binomial crooked* functions are quadratic [4].

## 4 Structure of crooked functions

A hyperplane of  $\mathbb{F}_{2^n}$  can be seen as a subspace of  $\mathbb{F}_{2^n}$ , as follows defined:

$$H_\lambda = \{ y \mid \text{Tr}(\lambda y) = 0 \}, \lambda \in \mathbb{F}_{2^n}^*. \quad (6)$$

We will denote by  $\overline{H_\lambda}$  the complement of  $H_\lambda$ . The dual of  $H_\lambda$  is obviously  $H_\lambda^\perp = \{0, \lambda\}$ .

The next definition of *crooked* functions is due to Kyureghan [17]. The corpus of such functions includes the crooked permutations, but also a large variety of non bijective crooked functions, especially in even dimension.

**Definition 3** *A function  $F$ , from  $\mathbb{F}_{2^n}$  to itself, is called crooked when it is such that, for every  $a$ , its differential set*

$$S_a = \{ F(x) + F(x + a) \mid x \in \mathbb{F}_{2^n} \},$$

*is an affine hyperplane.*

We directly deduce from Proposition 1:

**Claim 1** *Any crooked function is an APN function.*

Assuming that  $S_a = H_\lambda$ , for some  $\lambda$ , we get

$$\text{Tr}(\lambda(D_a F(x))) = c, \text{ for all } x, \text{ where } c \in \mathbb{F}_2.$$

This means that the derivative of the component  $f_\lambda$  of  $F$ , in point  $a$ , is a constant function. In this case,  $a$  is said to be a *linear structure* of  $f_\lambda$ . The *linear space* of  $f_\lambda$  is the subspace, including  $a = 0$ , of their linear structures. We will see that crooked functions have always components with nonzero linear structures.

## 4.1 Crooked and partially-bent functions

Let  $f$  be any Boolean function over  $\mathbb{F}_{2^n}$ . Denote by  $N_d$ , the number of balanced derivatives of  $f$ , and by  $N_f$ , the size of the set  $\{a \mid \mathcal{W}_f(a) = 0\}$ . Partially-bent functions were introduced in [9], as functions satisfying

$$(2^n - N_d)(2^n - N_f) = 2^n. \tag{7}$$

There is another characterization of partially-bent functions, which allows to determine their Walsh spectrum.

**Theorem 2** [9, Theorem] *A Boolean function  $f$  is partially-bent if and only if its derivatives are either constant or balanced.*

This leads to the precise description of the Walsh spectrum of any partially-bent function. Note that bent functions are particular partially-bent functions, with  $N_d = 2^n - 1$  and  $N_f = 0$ . Moreover, a partially-bent function is, in a certain sense, obtained by concatenating the same bent function, several times. The next result is partly given by [9, Proposition 2]. See also [17, Theorem 1], which concerns crooked functions but, actually, holds for any partially-bent function. For the Walsh spectrum, see a proof in [7, Proposition 4].

**Corollary 1** *Let  $f$  be a partially-bent Boolean function. Assume that  $f$  has a linear space  $V$ , of dimension  $s > 0$ .*

*Then  $f$  is a  $s$ -plateaued function, such that  $n + s$  is even, and  $\mathcal{W}_f$  takes three values,  $\{0\}$  and  $\{\pm 2^{(n+s)/2}\}$ . The Walsh spectrum of  $f$  is given by the Table 1 above.*

The definition of vectorial partially-bent functions  $F$  is simply derived.

**Definition 4** *Let  $F : \mathbb{F}_{2^n} \rightarrow \mathbb{F}_{2^n}$ . The function  $F$  is said to be a partially-bent vectorial function if every component of  $F$  is partially-bent. When  $n$  is even, some components can be bent.*

**Theorem 3** *Let  $F$  be a vectorial function over  $\mathbb{F}_{2^n}$  with components  $f_\lambda$ . Set, for  $a \in \mathbb{F}_{2^n}^*$ ,*

$$\Lambda_a = \{ \lambda \in \mathbb{F}_{2^n}^* \mid D_a f_\lambda \text{ is constant} \} \cup \{0\},$$

*and denote by  $\ell(a)$  the dimension of  $\Lambda_a$ . Then we have:*

- *Assume that  $F$  is partially-bent. Then the differential sets of  $F$  are affine subspaces. For any  $a \in \mathbb{F}_{2^n}^*$ , this subspace is of codimension  $\ell(a)$ , with  $\ell(a) \geq 1$  and  $D_a F$  is a  $2^{\ell(a)}$ -to-1 function.*
- *If  $F$  is a crooked function then  $F$  is partially-bent.*

*Proof.* Recall that  $D_a f_\lambda(x) = \text{Tr}(\lambda D_a F(x))$ , for all  $x$ . Obviously,  $\Lambda_a$  is a subspace of  $\mathbb{F}_{2^n}$ . Now, fixing  $a$  and  $x$ , we compute

$$\begin{aligned} B(a, x) &= \sum_{\lambda \in \mathbb{F}_{2^n}^*} \sum_{y \in \mathbb{F}_{2^n}} (-1)^{D_a f_\lambda(x) + D_a f_\lambda(y)} \\ &= \sum_{y \in \mathbb{F}_{2^n}} \sum_{\lambda \in \mathbb{F}_{2^n}^*} (-1)^{\text{Tr}(\lambda(D_a F(x) + D_a F(y)))} \\ &= 2^n \times |\{ y \mid D_a F(x) = D_a F(y) \}|. \end{aligned}$$



On the other hand, we set for any  $\lambda \in \mathbb{F}_{2^n}^*$ :

$$B(\lambda) = \sum_{y \in \mathbb{F}_{2^n}} (-1)^{D_a f_\lambda(x) + D_a f_\lambda(y)} = (-1)^{D_a f_\lambda(x)} \sum_{y \in \mathbb{F}_{2^n}} (-1)^{D_a f_\lambda(y)}.$$

We first assume that  $F$  is partially bent. Note that  $\ell(a) \geq 1$ , since otherwise we would have that the Boolean function  $D_a f_\lambda$  is balanced, for any  $\lambda \in \mathbb{F}_{2^n}^*$ , from Theorem 2. This is impossible since  $D_a F$  cannot be a permutation (see Theorem 1).

Clearly,  $B(\lambda) = 0$  if and only if the function  $D_a f_\lambda$  is balanced. If it is not balanced, then  $\lambda \in \Lambda_a$ , and this function is constantly equal to either 0 or 1. In both cases, we get  $B(\lambda) = 2^n$ . Hence, we get

$$B(a, x) = \sum_{\lambda \in \Lambda_a} B(\lambda) = 2^n 2^{\ell(a)}, \text{ for all } x,$$

implying that the number of  $y$  such that  $D_a F(x) = D_a F(y)$  equals  $2^{\ell(a)}$ , i.e.,  $D_a F$  is  $2^{\ell(a)}$ -to-1. Since  $\mathcal{I}m(D_a F)$  is contained in an affine subspace of codimension  $\ell(a)$ , according to the definition of  $\Lambda_a$ ,  $\mathcal{I}m(D_a F)$  is equal to this affine subspace.

Now, we suppose that  $F$  is a crooked function. Consider any component  $f_\lambda$  of  $F$ . Let  $a \in \mathbb{F}_{2^n}^*$  such that  $D_a f_\lambda$  is not constant. Set  $V = \mathcal{I}m(D_a F)$ . Then we have that any  $x$  satisfies:

$$\text{Tr}(\lambda D_a F(x)) = 0 \text{ if and only if } x \in V \cap H_\lambda.$$

There are  $2^{n-1}$  such  $x$ , since  $D_a F$  is 2-to-1 and  $V$  is an affine hyperplane, which is neither  $H_\lambda$  nor its complement. Hence  $D_a f_\lambda$  is balanced. We have proved that  $f_\lambda$  is partially-bent, completing the proof.  $\diamond$

Let  $F$  be any quadratic function:

$$F(x) = \sum_{0 \leq i < j < n} u_{i,j} x^{2^i + 2^j}, \quad u_{i,j} \in \mathbb{F}_{2^n}.$$

The derivatives of  $F$  are affine functions, say  $L_a$  for any  $a \in \mathbb{F}_{2^n}^*$ . Thus,  $F$  is partially-bent; it is crooked if and only if every  $L_a$  is an affine function with kernel of dimension 1.

**Corollary 2** *Any quadratic vectorial function is partially-bent. It is crooked if and only if it is APN.*

## 4.2 Crooked of odd dimension

In this section, we study crooked functions of odd dimension, bijective or not. Theorem 4 (below) is the main result, describing the exceptional properties of such functions. These results are quite known, but were partially presented in several papers [13, 16, 17]. First, it is easy to describe the set of crooked permutations.

**Lemma 2** *Let  $F$  be a crooked function such that  $F(0) = 0$ , with differential sets  $S_a$ . Let  $a$  and  $\lambda$  be such that  $S_a$  equals either  $H_\lambda$  or  $\overline{H_\lambda}$ , where  $H_\lambda$  is defined by (6). Then we have:*

$$S_a = \overline{H_\lambda} \iff \text{Tr}(\lambda F(a)) = 1.$$

*Besides,  $F$  is a permutation if and only if  $S_a = \overline{H_\lambda}$ , for any such pair  $(\lambda, a)$ . In this case,  $n$  is odd.*

*Proof.* By hypothesis, we have  $\text{Tr}(\lambda D_a F(x)) = c$  for all  $x$ , where  $c \in \{0, 1\}$ . In particular,  $\text{Tr}(\lambda F(a)) = c$ ; further,  $c = 0$  means that  $\mathcal{I}m(D_a F) = H_\lambda$ .

The function  $F$  is not bijective if and only if  $F(x) = F(x + a)$ , for some pair  $(x, a)$ . Equivalently, there is  $(\lambda, a)$  such that  $\mathcal{I}m(D_a F) = H_\lambda$ , since 0 belongs to  $\mathcal{I}m(D_a F)$ .

When  $n$  is even,  $F$  cannot be a permutation, since it is an APN function, which is plateaued (see [2, Theorem 3]).  $\diamond$

When  $n$  is odd, a crooked function may not always be bijective, as we show by the next example.

**Example 1** *Assume that  $n$  is odd. It is well-known that*

$$F : x \mapsto x^{2^t+1}, \quad \text{with } \gcd(t, n) = 1,$$

*is an AB permutation. So, it is a crooked permutation. Consider now the function  $x \mapsto G(x) = x^{2^t+1} + x$ , which is AB too, and then crooked. Since  $G(0) = G(1) = 0$ ,  $G$  is not a permutation.*

**Definition 5** *A Boolean function is said to be near-bent if it is 1-plateaued, i.e., its Walsh transform takes the values 0 and  $\pm 2^{(n+1)/2}$  only.*

**Theorem 4** *Let  $F$  be a crooked function over  $\mathbb{F}_{2^n}$ ,  $n$  odd, such that  $F(0) = 0$ . Every differential set  $S_a$ ,  $a \in \mathbb{F}_{2^n}^*$ , is an affine hyperplane:*

$$S_a = \beta + H_\lambda, \quad \beta \in \mathbb{F}_{2^n}, \quad \text{for some } \lambda \in \mathbb{F}_{2^n}^*.$$

*Then the following properties hold:*

- (i) *The differential sets  $S_a$  are two by two distinct, corresponding to two by two distinct hyperplanes  $H_\lambda$ . More precisely, to every  $a \in \mathbb{F}_{2^n}^*$  corresponds one and only one  $\lambda \in \mathbb{F}_{2^n}^*$ .*
- (ii) *Any component  $f_\lambda$  of  $F$  is a near-bent Boolean function with linear space of dimension 1, say  $\{0, a\}$ . When  $f_\lambda$  is balanced, its derivative in point  $a$  is equal to 1. This holds for any  $f_\lambda$ , when  $F$  is a permutation,*
- (iii)  *$F$  is an AB function.*

*Proof.* Since  $F$  is crooked, it is partially-bent. Let  $f_\lambda$ ,  $\lambda \in \mathbb{F}_{2^n}^*$ , be the components of  $F$ . Thus, for any  $\lambda$  and for any  $a$ , the derivative of  $f_\lambda$ , in point  $a$ , is either constant or balanced. Hence

$$\sum_{x \in \mathbb{F}_{2^n}} (-1)^{f_\lambda(x) + f_\lambda(x+a)} \in \{0, \pm 2^n\}.$$

For any fixed  $\lambda$ , there is at least one  $a$ , say  $a(\lambda)$ , such that  $D_a f_\lambda$  is constant, since otherwise the function  $f_\lambda$  would be bent, which is impossible when  $n$  is odd. Thus, we get the set of the  $a(\lambda)$ , whose size is at most  $2^n - 1$ . However,  $a(\lambda) = a(\mu) = b$ , for some  $\mu \neq \lambda$ , would mean the following: the Boolean functions

$$x \mapsto \text{Tr}(\lambda D_b F(x)) \quad \text{and} \quad x \mapsto \text{Tr}(\mu D_b F(x))$$

are both constant. This would imply that  $S_b$  is of codimension at least 2, a contradiction. To each  $\lambda$  corresponds one and only one  $a$ , completing the proof of (i).

We deduce that any component  $f_\lambda$  has only one nonzero linear structure, say  $a$ , *i.e.*, its linear space has dimension 1. Obviously, if  $f_\lambda$  is balanced, its derivative in point  $a$  is constantly equal to 1. When  $F$  is a permutation, all  $f_\lambda$  are balanced, completing the proof of (ii).

From Corollary 1, every  $f_\lambda$  is near-bent, providing

$$\mathcal{W}_{f_\lambda}(a) \in \{0, \pm 2^{(n+1)/2}\}, \quad \text{for all } a \in \mathbb{F}_{2^n}.$$

According to Definition 1,  $F$  is an AB function, completing the proof.  $\diamond$

From Table 1, we know that the support of the Walsh spectrum of any near-bent Boolean function on  $\mathbb{F}_{2^n}$  has size  $2^{n-1}$ . It could be an affine subspace of codimension 1, as it holds for components of some crooked functions.

**Proposition 2** *Let  $F$  be a vectorial function over  $\mathbb{F}_{2^n}$ ,  $n$  odd, such that  $F(0) = 0$ . Assuming that  $F$  is crooked, the set*

$$W_\lambda = \{ a \in \mathbb{F}_{2^n} \mid \mathcal{W}_{f_\lambda}(a) = 0 \}$$

*is an affine subspace of codimension 1, for all  $\lambda \in \mathbb{F}_{2^n}^*$ .*

*Conversely, assume that the sets  $W_\lambda$  are affine hyperplanes. In this case, if  $F$  is APN then  $F$  is crooked.*

*Proof.* Assume that  $F$  is crooked and let  $\lambda \in \mathbb{F}_{2^n}^*$ . Since  $F$  is an AB function,  $f_\lambda$  is partially bent, with linear space  $\{0, b\}$ , for some nonzero  $b$ . Obviously, the function  $g_a : x \mapsto f_\lambda(x) + \text{Tr}(ax)$  is partially bent too, with linear space  $\{0, b\}$ , for any  $a \in \mathbb{F}_{2^n}$ . We know that any partially-bent function is balanced if and only if it is not constant on its linear space (see [9, Proposition 2]). Thus,  $g_a$  is balanced if and only if

$$D_b g_a(x) = 1, \quad \text{i.e., } D_b f_\lambda(x) + \text{Tr}(ab) = 1, \text{ for all } x.$$

Hence, we get  $\text{Tr}(ab) = 1 + c$ ,  $c \in \mathbb{F}_2$ , since  $D_b f_\lambda$  is a constant function. We can suppose that  $c = 0$ . Thus,  $g_a$  is balanced if and only if  $\text{Tr}(ab) = 1$ , that is  $a \in \overline{H_b}$ . Thus

$$W_\lambda = \{ a \in \mathbb{F}_{2^n} \mid \text{Tr}(ab) = 1 \} = \overline{H_b}.$$

Similarly, with  $c = 1$  we obtain  $W_\lambda = H_b$ .

Conversely, suppose that  $W_\lambda$  is an affine subspace of codimension 1, say  $u + H_b$ ,  $u \in \mathbb{F}_{2^n}$ , for some  $b$ . So,  $H_b^\perp = \{0, b\}$  and we have from Lemma 1:

$$\sum_{a \in H_b} \mathcal{W}_{f_\lambda}(a)^2 = 2^{n-1} \left( \sum_{x \in \mathbb{F}_{2^n}} (-1)^{\text{Tr}(\lambda D_0 F(x))} + \sum_{x \in \mathbb{F}_{2^n}} (-1)^{\text{Tr}(\lambda D_b F(x))} \right)$$

Since Perceval's relation, the sum above on the left is equal either to 0 or to  $2^{2n}$ . We deduce that  $D_b f_\lambda$  is a constant function. So, if  $D_b F$  is 2-to1, then  $\text{Im}(D_b F)$  is equal to either  $H_\lambda$  or  $\overline{H_\lambda}$ .  $\diamond$

### 4.3 Crooked of even dimension

When  $n$  is even, crooked functions are partially-bent functions which have bent components. We recall this property below, by Theorem 5.

**Theorem 5** *Let  $F$  be a plateaued function over  $\mathbb{F}_{2^n}$ , with  $n$  even and  $n > 4$ . Let  $2^{(t_\lambda+n)/2}$  be the amplitude of any component of  $F$ , namely of any  $f_\lambda$ ,  $\lambda \in \mathbb{F}_{2^n}^*$ . Denote by  $B$  the number of bent components of  $F$ . Then  $F$  is APN if and only if*

$$B = \sum_{\lambda \in \mathbb{F}_{2^n}^*, t_\lambda > 0} (2^{t_\lambda} - 2). \quad (8)$$

*This property holds, in particular, for crooked functions.*

*Consequently,  $B$  satisfies*

$$\frac{2(2^n - 1)}{3} \leq B < 2^n - 2^{n/2}, \quad (9)$$

*where the lower bound is reached if and only if  $t_\lambda = 2$  for all non zero  $t_\lambda$ .*

*Proof.* The equality (8) has been proved by [11, Proposition 6], but may be computed by using (5). Indeed, since each  $f_\lambda$  is plateaued, we have  $\nu(f_\lambda) = 2^{t_\lambda+2n}$ , for any  $\lambda$ , and then:

$$A = \sum_{\lambda \in \mathbb{F}_{2^n}^*} \nu(f_\lambda) = 2^{2n} \sum_{\lambda \in \mathbb{F}_{2^n}^*} 2^{t_\lambda}.$$

According to (5),  $F$  is APN if and only if  $A = 2^{2n+1}(2^n - 1)$ , providing

$$B + \sum_{\lambda \in \mathbb{F}_{2^n}^*, t_\lambda > 0} 2^{t_\lambda} = 2(2^n - 1) = 2(B + N), \quad (10)$$

since  $2^n - 1 = B + N$  with  $N = |\{\lambda \mid t_\lambda > 0\}|$ . Hence, the equality above is equivalent to (8). Note that  $t_\lambda > 0$  implies  $t_\lambda \geq 2$ , since  $n + t_\lambda$  must be even.

The lower bound of  $B$ , in (9), is known (see [2, Corollary 3]). The upper bound has been proved by [19, Theorem 3]. It cannot be reached here, since we must have  $B \equiv 2 \pmod{4}$ , from (10).  $\diamond$

By the study of bent components, one understand precisely the difference between odd and even cases. Let  $F$  be a crooked function on  $\mathbb{F}_{2^n}$ , with components  $f_\lambda$ . We have seen that, when  $n$  is odd, there is a one to one

correspondence  $\lambda \mapsto a(\lambda)$ , where  $a(\lambda)$  is the unique linear structure of  $f_\lambda$ . When  $n$  is even, the linear space of  $f_\lambda$  has dimension  $s$ , where  $s$  is even. So, either  $f_\lambda$  is bent ( $s = 0$ ) or  $s \geq 2$ . The number of hyperplanes involved in the structure of  $F$  is at most  $(2^n - 1)/3$ . For any  $a$ , there is one and only one  $\lambda$  such that  $\mathcal{I}m(D_a F)$  is equal to  $H_\lambda$  or to  $\overline{H_\lambda}$ . But, the function  $D_a f_\lambda$  is constant for any  $a$  belonging to the linear space of  $f_\lambda$ . There are at least three such nonzero  $a$ , when  $f_\lambda$  is not bent.

**Proposition 3** *Let  $F$  be a crooked function on  $\mathbb{F}_{2^n}$ ,  $n$  even, with components  $f_\lambda$ . For any  $a \in \mathbb{F}_{2^n}^*$ , there is a unique  $\lambda$  such that the derivative of  $f_\lambda$ , in point  $a$ , is a constant function.*

*Conversely, any function  $f_\lambda$  is either bent or with linear space  $V$  of dimension  $k \geq 2$ .*

*Proof.* By hypothesis,  $\mathcal{I}m(D_a F)$  is an hyperplane  $H_\lambda$ , or the complement of  $H_\lambda$ , for any  $a \in \mathbb{F}_{2^n}^*$ . Hence  $Tr(\lambda D_a F(x)) = c$ , for all  $x$ , where  $c \in \mathbb{F}_2$ . Such a  $\lambda$  is unique, because otherwise  $\mathcal{I}m(D_a F)$  would be an affine subspace of codimension 2.

Since  $F$  is a plateaued APN function, at least  $2(2^n - 1)/3$  components of  $F$  are bent. Let  $f_\lambda$  be a non bent component. Its linear set has an even dimension, so this dimension must be greater than or equal to 2.  $\diamond$

## 5 Bent from crooked functions

In this section, we consider crooked functions over  $\mathbb{F}_{2^n}$ , where  $n$  is odd. The components of such a function are *near-bent* Boolean functions. Moreover, every component has (only) one derivative which is a constant function. (see Theorem 4). We will show that a set of  $2^n - 1$  bent functions of  $n + 1$  variables can be derived, using this strong property. Our main reference is the construction of Leander and McGuire [18], on the near-bent Boolean functions, that we apply to vectorial functions which are crooked. We first recall some facts which are more or less known, maybe not in this form.

**Lemma 3** *Let  $f$  be a near-bent Boolean function over  $\mathbb{F}_{2^n}$ , where  $n$  is odd. Assume that  $f(0) = 0$ . Then (i) and (ii) are equivalent:*

(i)  *$f$  has a constant derivative in point  $a \in \mathbb{F}_{2^n}^*$ .*

(ii) *There exists a such that*

$$\{ u \in \mathbb{F}_{2^n} \mid \mathcal{W}_f(u) = 0 \} = \{ u \in \mathbb{F}_{2^n} \mid \text{Tr}(ua) = 1 + f(a) \}.$$

*Proof.* Note that,  $f$  cannot have more than one linear structure, since it is near-bent. Assume that there is (a unique)  $a$  such that  $D_a f(x) = c$ , for all  $x$ , where  $c \in \mathbb{F}_2$ . Thus,  $c = f(0) + f(a) = f(a)$ . Now, we compute  $\mathcal{W}_f(u)$ :

$$\begin{aligned} \mathcal{W}_f(u) &= \sum_{x \in \mathbb{F}_{2^n}} (-1)^{f(x) + \text{Tr}(ux)} = \sum_{x \in \mathbb{F}_{2^n}} (-1)^{f(x+a) + \text{Tr}(u(x+a))} \\ &= \sum_{x \in \mathbb{F}_{2^n}} (-1)^{f(x) + f(a) + \text{Tr}(u(x+a))} \\ &= (-1)^{f(a) + \text{Tr}(ua)} \mathcal{W}_f(u), \end{aligned}$$

since  $f(x) + f(x+a) = f(a)$ . Clearly, if  $u$  is such that  $\text{Tr}(ua) + f(a) = 1$  then  $\mathcal{W}_f(u) = 0$ . But, there are  $2^{n-1}$  such  $u$ , proving that (ii) holds.

Now suppose that (ii) holds, for some  $a$ . Recall that

$$H_a = \{ u \in \mathbb{F}_{2^n} \mid \text{Tr}(ua) = 0 \} \text{ so that } H_a^\perp = \{0, a\}.$$

Thus, the set of those  $u$  such that  $\mathcal{W}_f(u) = 0$  is either equal to the hyperplane  $H_a$ , or equal to its complement, according to either  $f(a) = 1$  or  $f(a) = 0$ . So, we can apply Lemma 1:

$$\sum_{v \in H_a} \mathcal{W}_f(v)^2 = 2^{n-1} \left( \sum_{x \in \mathbb{F}_{2^n}} (-1)^0 + \sum_{x \in \mathbb{F}_{2^n}} (-1)^{f(x) + f(x+a)} \right).$$

By hypothesis, this sum equals 0 if  $f(a) = 1$  and  $2^{2n}$  if  $f(a) = 0$ , since the Parseval's relation. This is possible only if  $D_a f$  is a constant derivative of  $f$ . It is the only one such derivative, since  $f$  is near-bent.  $\diamond$

**Lemma 4** *Let  $f$  be a near-bent Boolean function on  $\mathbb{F}_{2^n}$  ( $n$  odd) such that  $f(0) = 0$ . Assume that  $f$  has a linear structure  $a$ . Let  $g$  be the function from  $G = \mathbb{F}_{2^n} \times \mathbb{F}_2$  to  $\mathbb{F}_2$ :*

$$g(x, y) = (y + 1)f(x) + y(f(x) + \text{Tr}(a^{-1}x)) \quad (11)$$

*Then  $g$  is a bent function of  $n + 1$  variables.*

*Proof.* For any  $u$ , we denote by  $f_u$  the Boolean function  $x \mapsto f(x) + \text{Tr}(ux)$ . Let  $b = a^{-1}$ . The restriction of  $g$  to  $\mathbb{F}_{2^n}$  ( $y = 0$ ) and to its complement in  $G$  ( $y = 1$ ) are respectively  $f$  and  $f_b$  which are both near-bent Boolean functions. Moreover

$$D_a f(x) = f(a) \quad \text{and} \quad D_a f_b(x) = f(a) + \text{Tr}(ba) = f(a) + 1 = f_b(a),$$

since  $\text{Tr}(ba) = \text{Tr}(1) = 1$ . Let  $u \in \mathbb{F}_{2^n}^*$ . Applying Lemma 3,  $f_u$  is balanced if and only if  $\text{Tr}(ua) = 1 + f(a)$  and  $f_{b+u}$  is balanced if and only if

$$\text{Tr}(a(u + b)) = 1 + f(a), \quad \text{providing} \quad \text{Tr}(ua) = 1 + 1 + f(a) = f(a).$$

Thus  $f_u$  is balanced if and only if  $f_{b+u}$  is not balanced. This is equivalent to say that  $g$  is bent (see [6, Theorem V.3] or [18, Theorem 2]).  $\diamond$

Now, we consider a crooked function  $F$ , from  $\mathbb{F}_{2^n}$  to itself. According to Theorem 4, we know that all components  $f_\lambda$  of  $F$  are near-bent with a (unique) constant derivative. This allows to derive a specific set of bent functions from any crooked function, in odd dimension.

**Theorem 6** *Let  $F$  be a crooked function over  $\mathbb{F}_{2^n}$  where  $n$  is odd, such that  $F(0) = 0$ . Denote by  $a_\lambda$  the linear structure of the component  $f_\lambda$  of  $F$ . Then we get a set  $B(F)$  of  $2^n - 1$  bent functions  $g_\lambda$ , each from  $G = \mathbb{F}_{2^n} \times \mathbb{F}_2$  to  $\mathbb{F}_2$ :*

$$B(F) = \{ g_\lambda(x, y) = (y + 1)f_\lambda(x) + y (f_\lambda(x) + \text{Tr}(a_\lambda^{-1}x)) \mid \lambda \in \mathbb{F}_{2^n}^* \}.$$

*Proof.* Since  $F$  is a crooked function, there is a bijective correspondence between the  $\lambda \in \mathbb{F}_{2^n}^*$ , and then the functions  $f_\lambda$ , and the linear structures  $a_\lambda$  of every  $f_\lambda$ . Thus, Lemma 4 applies for every  $\lambda$ , using  $a_\lambda^{-1}$ , providing  $2^n - 1$  distinct bent functions.  $\diamond$

**Remark 1** *The construction, given by Theorem 6, could be of interest regarding the problem of the existence of crooked functions. It would be useful to exhibit some properties of  $B(F)$ , or to construct other sets of bent functions. For instance,  $g_\lambda$  could have an higher degree, by replacing  $f_\lambda$  by another Boolean function, affinely equivalent to  $f_\lambda$ . The question is to know if such a set of bent functions does exist, for some degree greater than 2.*

*Note that a set like  $B(F)$  is not a subspace, but could generate a linear code with few weights.*

*To derive effectively bent functions from a given crooked function, it is necessary to have an expression of the linear structures  $a_\lambda$ . This is generally an open problem, even when  $F$  is a quadratic APN function.*



**Example 2** Let  $F(x) = x^{2^k+1}$ ,  $F : \mathbb{F}_{2^n} \rightarrow \mathbb{F}_{2^n}$ ,  $n$  odd. It is well-known that  $F$  is crooked if and only if  $\gcd(k, n) = 1$ . For any  $\lambda \in \mathbb{F}_{2^n}^*$ , the unique linear structure of  $f_\lambda$  is  $a_\lambda = \lambda^{-1/(2^k+1)}$ . Thus, for any  $\lambda \in \mathbb{F}_{2^n}^*$ ,

$$g_\lambda(x, y) = (y + 1)\text{Tr}\left(\lambda x^{2^k+1}\right) + y\left(\text{Tr}(\lambda x^{2^k+1}) + \text{Tr}(\lambda^{1/(2^k+1)}x)\right),$$

is a bent function,  $(x, y) \mapsto g_\lambda(x, y)$ , from  $\mathbb{F}_{2^n} \times \mathbb{F}_2$  to  $\mathbb{F}_2$ .

## 6 Permutations from crooked functions

Any crooked function allows to construct a set of permutations, via their derivatives.

**Theorem 7** Let  $F : \mathbb{F}_{2^n} \rightarrow \mathbb{F}_{2^n}$ , such that  $F(0) = 0$ . Assume that  $F$  is a crooked function. Define, for any  $a \in \mathbb{F}_{2^n}^*$ :

$$x \mapsto G_a(x) = F(x) + F(x + a) + F(a).$$

Let us define:

- $\lambda \in \mathbb{F}_{2^n}^*$ , defining the hyperplane  $H_\lambda$ , which is the image set of  $G_a$ ;
- $\beta$  be such that  $\text{Tr}(\lambda\beta) = 1$ , i.e.  $\beta \notin H_\lambda$ ;
- $H_\mu$  be any hyperplane such that  $\text{Tr}(\mu a) = 1$ , i.e.,  $a \notin H_\mu$ .

Then, the function

$$x \mapsto R_a(x) = G_a(x) + \beta\text{Tr}(\mu x)$$

is a permutation, such that  $R_a(H_\mu) = H_\lambda$ .

*Proof.* Let  $x \neq y$  such that  $R_a(x) = R_a(y)$ . We get

$$G_a(x) + G_a(y) = \beta(\text{Tr}(\mu(x + y))).$$

If  $\text{Tr}(\mu(x + y)) = 0$  then  $G_a(x) + G_a(y) = 0$ , which implies  $y = x + a$ , since  $G_a$  is 2-to-1 and  $G_a(x) = G_a(x + a)$ . In this case, we get  $\text{Tr}(\mu(x + y)) = \text{Tr}(\mu a) = 0$ , a contradiction.

Now suppose that  $Tr(\mu(x+y)) = 1$ . Hence  $G_a(x) = G_a(y) + \beta$ . But this is impossible, since  $G_a(x)$  and  $G_a(y)$  belong to  $H_\lambda$  while  $\beta \notin H_\lambda$ . Thus, we get again a contradiction and can conclude that  $R_a$  is a permutation.

By construction,  $R_a(H_\mu) = H_\lambda$ , since  $Tr(\mu x) = 0$  for all  $x \in H_\mu$  and  $G_a(H_\mu) = H_\lambda$ . Indeed,  $G_a$  is 2-to-1 and its image set is  $H_\lambda$ . For any pair  $(x, x+a)$ , we have  $x \in H_\mu$  if and only if  $x+a \in \overline{H_\mu}$  (with  $G_a(x) = G_a(x+a)$ ).

◇

## 7 Comments to conclude

As a prelude of this conclusion, we want to recall the main conjecture concerning crooked functions:

**Conjecture:** Let  $F : \mathbb{F}_{2^n} \rightarrow \mathbb{F}_{2^n}$  be a crooked function, as defined by Definition 3. Then  $F$  is quadratic or, in other terms, has algebraic degree 2.

By the two previous sections, we want to exhibit specific constructions derived from crooked functions (only). Our purpose is to open new ways to study the existence of crooked functions. We are convinced that other properties have to be found, increasing the knowledge on crooked functions, especially on the existence of such functions.

On the other hand, the quadratic functions form a corpus of great interest about which many problems remain open. In particular, we are still far from understanding the different structures of quadratic APN functions of even dimension. The determination of the number of bent components of such a function would be of great interest. Negative answers, concerning the APN property of quadratic vectorial functions, have been obtained, for instance in [2][5]. In [3], a description of the corpus of binomial crooked functions is presented. As a conclusion, Bierbrauer has conjectured that, up to equivalence, no other binomial crooked functions exist.

*Crooked functions of codimension  $k$ ,  $1 \leq k \leq n - 1$* , appeared during the cryptanalysis of a hash function, called MARACA [8]. Here, the differential sets are affine subspaces of same codimension. The authors of this cryptanalysis, Canteaut and Naya-Placienta, introduced the *crooked property*. They have shown that this property could make a cryptographic primitive very weak. We will explore, in a more general context, the *functions having the crooked property*, in a forthcoming work.

## References

- [1] T. Bending and D. Fon-Der-Flaass. Crooked functions, bent functions, and distance regular graphs. *Electronic Journal of Combinatorics*, 5(1), 1998. R34.
- [2] T.P. Berger, A. Canteaut, P. Charpin, and Y. Laigle-Chapuy. On almost perfect nonlinear functions. *IEEE Trans. Inform. Theory*, 52(9):4160–4170, September 2006.
- [3] J. Bierbrauer, A family of crooked functions, *Designs, Codes and Cryptography*, (2009) 50:235-241.
- [4] J. Bierbrauer and G. Kyureghyan, Crooked binomials, *Des. Codes Cryptogr.* 46 (2008) 269-301.
- [5] E. Byrne and G. McGuire, On the non-existence of quadratic APN and crooked functions on finite fields, *Proc. of the Workshop on Coding and Cryptography*, WCC 2004, 316-324.
- [6] A. Canteaut, C. Carlet, P. Charpin, and C. Fontaine. On cryptographic properties of the cosets of  $R(1, m)$ . *IEEE Transactions on Information Theory*, 47(4):1494–1513, 2001.
- [7] A. Canteaut, P. Charpin. Decomposing bent functions. *IEEE Trans. Inform. Theory*, 49(8), pp. 2004-19, August 2003.
- [8] A. Canteaut and M. Naya-Plasencia. Structural weaknesses of permutations with a low differential uniformity and generalized crooked functions . In *Finite Fields: Theory and Applications - FQ9 - Contemporary Mathematics*, AMS, number 518, pp. 55-71, 2010.
- [9] C. Carlet. Partially-bent functions. *Designs, Codes and Cryptography*, (3):135–145, 1993.
- [10] C. Carlet, P. Charpin, and V. Zinoviev. Codes, bent functions and permutations suitable for DES-like cryptosystems. *Designs, Codes and Cryptography*, 15(2):125–156, 1998.
- [11] P. Charpin and J. Peng. Differential uniformity and the associated codes of cryptographic functions. *Advances in Mathematics of Communications*, AIMS, Special issue on *Applications of discrete mathematics in*

*secure communication*, Subhamoy Maitra Ed., November 2019, 13(4): 579-600.

- [12] E.R. van Dam and D. Fon-Der-Flaass. Uniformly packed codes and more distance regular graphs from crooked functions. *J. Algebraic Combin.* 12 (2000), no. 2, 115–121. 94B05
- [13] E.R. van Dam and D. Fon-Der-Flaass. Codes, graphs, and schemes from nonlinear functions. *European J. Combin.* 24 (1) (2003) 85-98.
- [14] J.F. Dillon and G. McGuire, Near bent functions on a hyperplane, *Finite Fields Appl.*, 14 (2008), no. 3, 715-720.
- [15] C. Godsil and A. Roy. Two characterization of crooked functions, *IEEE Trans. Inform. Theory* 54 (2008), no. 2, 864-866.
- [16] G. Kyureghyan, The only crooked power functions are  $x^{2^k+2^l}$ , *European J. Combin.* 28 (2007) 1345–1350.
- [17] G. Kyureghyan, Crooked maps in  $F_{2^n}$ , *Finite Fields Appl.* 13(3), pp. 713-726 (2007).
- [18] G. Leander and G. McGuire, Construction of bent functions from near-bent functions, *Journal of Combinatorial Theory, Series A* 116 (2009) 960-970.
- [19] A. Pott, E. Pasalic, A. Muratovic-Ribic and S. Bajric, On the maximum number of bent component of vectorial functions, *IEEE Transactions on Information Theory*, Vol. 64, No. 1, January 2018, pp. 403-411.
- [20] Y. Zheng and X. M. Zhang, On plateaued functions, *IEEE Trans. Inform. Theory*, 47(2001), No. 3, pp. 1215–1223.