



HAL
open science

The circuit - hypergraph of a tournament

Jean-Claude Bermond

► **To cite this version:**

Jean-Claude Bermond. The circuit - hypergraph of a tournament. Infinite and finite sets, Colloq Kezsthely 1973, deidcated to P.Erdös on his 60th birthday Colloq. Math Soc Janos Bolyai, 1975, Keszthely, Hungary. pp.165-180. hal-02366962

HAL Id: hal-02366962

<https://inria.hal.science/hal-02366962>

Submitted on 16 Nov 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

THE CIRCUIT – HYPERGRAPH OF A TOURNAMENT

J-C. BERMOND*

Let $\tau(T_n)$ be the minimal number of arcs which must be deleted (or reversed) in a tournament T_n in order to obtain a graph without circuits. In this article we give results on the function $f(n) = \max_{T_n} \tau(T_n)$ (maximum taken over all the tournaments with n vertices), answering a conjecture of Kotzig [7]. Other known results and references can be found in [2].

1. DEFINITIONS AND PRELIMINARY RESULTS

Throughout this paper $G = (X, U)$ will denote a finite oriented graph, where X is the set of vertices and U the set of arcs. T_n will denote a tournament (graph (X, U) in which each pair of vertices x, y is joined by one and only one of the arcs \overrightarrow{xy} and \overrightarrow{yx}), with $|X| = n$, and so $|U| = \frac{n(n-1)}{2}$. A circuit of G is a sequence of *different* vertices $x_1 x_2 \dots x_i x_{i+1} \dots x_p$ such that $\overrightarrow{x_i x_{i+1}}$ ($i = 1, 2, \dots, p-1$) and $\overrightarrow{x_p x_1}$ are arcs of G .

*Chemin de Jaumeron 91190-GIF sur Yvette France

Definition 1. To a graph $G = (X, U)$ we associate the hypergraph $C(G) = (U, C)$, called the *circuit-hypergraph of G* , whose set of vertices is U (set of arcs of G) and whose edges are the circuits of G . This hypergraph has been studied by Chaty [4].

Definition 2 (see Berge [1]). A *transversal set* of a hypergraph H is a set of vertices of H that meet every edge of the hypergraph and the *transversal number* $\tau(H)$ is the minimum number of elements of a transversal set.

Let $\tau(G)$ denote the transversal number of the circuit-hypergraph $C(G)$ associated to G , that is, the minimum number of arcs that we must delete from G in order to obtain a graph without circuits. It has been proved by Grindberg and Dambit, Remage and Thompson (see [2], p. 9) that $\tau(G)$ is also the minimum number of arcs whose orientation must be reversed in order to obtain a graph without circuits.

Example. For the tournament T_5 of Fig. 1 14, 15, 25 is a transversal set, so $\tau(T_5) \geq 3$. We shall see later that $\tau(T_5) = 3$.




Fig. 1

Instead of $\tau(G)$, we can consider the number $w(G)$, the stability number of the circuit-hypergraph of G , which is the maximum cardinality of a set of arcs such that the partial graph generated by these arcs is without circuits (such a set of arcs is called a set of consistent arcs by Erdős and Moon [5]). We have the relation $\tau(G) + w(G) = |U|$. $\tau(G)$ has been introduced by Slater [9] and $w(G)$ by Erdős and Moon [5]. The number $\tau(G)$ has importance in "paired comparisons" (see [2])

for further details and references). In this case G is a tournament and one of the problems is to determine the function $f(n) = \max_{T_n} \tau(T_n)$, where the maximum is taken over all the tournaments with n vertices; as has remarked Kotzig [7], $f(n)$ is also the maximum of $\tau(G)$ taken over all the antisymmetric graphs G with n vertices.

Various results have been obtained about the function $f(n)$ (see [2]), essentially upper bounds and that $\frac{f(n)}{\binom{n}{2}} \rightarrow \frac{1}{2}$ if $n \rightarrow \infty$ by Erdős and Moon [15]. Here we are interested in lower bounds of $f(n)$.

Definition 3. H being a hypergraph, the *matching number* of H : $\nu(H)$ is the maximum number of disjoint edges of H .

Let $\nu(G)$ denote the matching number of the circuit-hypergraph $C(G)$, that is the maximum number of arc-disjoint circuits of G .

By a theorem on hypergraph ([1], p. 405) we have:

Proposition 1. For every graph G : $\tau(G) \geq \nu(G)$.

Example. In the tournament T_5 of Fig. 1 we have 3 arc-disjoint circuits: 124, 135, 2345. Thus $\nu(T_5) \geq 3$ and by Proposition 1 $\tau(T_5) = \nu(T_5) = 3$.

Definition 4. The maximum number of edge-disjoint cycles of the non-oriented complete graph K_n will be denoted by $\pi(n)$. It has been proved by Chartrand, Geller and Hedetniemi [3] that $\pi(n) = \left\lceil \frac{n}{3} \left\lfloor \frac{n-1}{2} \right\rfloor \right\rceil$ (where $\lfloor x \rfloor$ is the greatest integer not exceeding x).

Proposition 2. $f(n) \geq \pi(n) = \left\lceil \frac{n}{3} \left\lfloor \frac{n-1}{2} \right\rfloor \right\rceil$: (For a proof see [2]).

By using the upper bound of Jung [6] (see also [2]) we have:

Proposition 3. For $n \leq 9$: $f(n) = \pi(n)$.

In [2] p. 21 and 22 we have determined the values of $f(10)$ and $f(11)$.

Proposition 4. $f(10) = 15$ ($\pi(10) = 13$); $f(11) = 20$ ($\pi(11) = 18$).

Kotzig [7] has also found independently Propositions 1, 2, 3 and the value of $f(10)$. He has also conjectured that:

Conjecture 1 (Kotzig [7]). For $n \geq 10$ $f(n) > \pi(n)$.

In his article Kotzig has proved that if $f(n) > \pi(n)$ then $f(3n) > \pi(3n)$. Here we shall prove similar results for $f(3n + 1)$ and $f(3n + 2)$ and so to prove the conjecture it would be sufficient to prove it for a small number of values of n . For this purpose we need results on the transversal number of the circuit-hypergraph associated to the lexicographic product (and generalized product) of two graphs.

2. LEXICOGRAPHIC PRODUCT OF GRAPHS

Definition. Let $G = (X, U)$ and $H = (Y, V)$ be two (oriented) graphs the *lexicographic product of G by H* , noted $G \otimes H$, is the graph which has the Cartesian product $X \times Y$ as set of vertices and in which (x_1, y_1) is joined to (x_2, y_2) if and only if either $\overrightarrow{x_1 x_2}$ is an arc of G or $x_1 = x_2$ and $\overrightarrow{y_1 y_2}$ is an arc of H . $G \otimes H$ is called composition of G with H in Moon [8] p. 78. (We can consider this graph as obtained by replacing each vertex of G by a copy of H).

Example: $G = C_3$, $H = K_2$.




Fig. 2

Proposition 5. Let $G = (X, U)$ and $H = (Y, V)$ be two graphs with $|X| = p$ and $|Y| = q$, then $\tau(G \otimes H) \leq p\tau(H) + q^2\tau(G)$.

Proof. First we prove that if G and H are two graphs without circuits then $G \otimes H$ is without circuits. Suppose, on the contrary, that $G \otimes H$ contains a circuit $(x_1, y_1)(x_2, y_2) \dots (x_k, y_k)$. If all the first indices x_i of the vertices of the circuit were equal, then $y_1 y_2 \dots y_k$ would be a circuit of H contrary to the hypothesis. Thus, two of the x_i 's are different; let us consider the different x_i 's in the order in which they appear in the circuit: $x_{i_1}, x_{i_2}, \dots, x_{i_p}$, then by definition of the lexicographic product $x_{i_1} x_{i_2} \dots x_{i_p}$ contains a circuit of G , contrary to the hypothesis.

According to the definition of $\tau(G)$ and $\tau(H)$ we can obtain a graph G' without circuits, by deleting $\tau(G)$ arcs from G and a graph H' without circuits by deleting $\tau(H)$ arcs from H . By the first part of the proof $G' \otimes H'$ is a graph without circuits and is obtained from $G \otimes H$ by deleting the $q^2\tau(G)$ arcs of the form $(x_1, y)(x_2, y')$, $\overrightarrow{x_1 x_2}$ being an arc of $G - G'$ and the $p\tau(H)$ arcs of the form $(x, y_1)(x, y_2)$ where $\overrightarrow{y_1 y_2}$ is an arc of $H - H'$ (the x is the same).

Conjecture 2. $\tau(G \otimes H) = p\tau(H) + q^2\tau(G)$.

We are able to prove the conjecture, when $\tau(G) = \nu(G)$, by considering $\nu(G \otimes H)$.

Proposition 6. Let S_q be the graph with q vertices and no arcs, then $\nu(G \otimes S_q) \geq q^2\nu(G)$.

Proof. Let the vertices of S_p be labelled: $1, 2, \dots, q$. To the circuit $x_1 x_2 \dots x_i \dots x_k$ of G , we associate the q^2 arc-disjoint circuits $(x_1, j)(x_2, j + \alpha) \dots (x_i, j + (i - 1)\alpha) \dots (x_{k-1}, j + (k - 2)\alpha)(x_k, \alpha)$, where $j = 1, 2, \dots, q$, $\alpha = 1, 2, \dots, q$ and where the second indices of the vertices are to be taken modulo q . Then to the $\nu(G)$ arc-disjoint circuits of G , we associate $q^2\nu(G)$ arc-disjoint circuits in $G \otimes S_q$.

Conjecture 3. $\nu(G \otimes S_q) = q^2\nu(G)$.





Fig. 3

Lemma. $\tau(S_p \otimes H) = p\tau(H)$ and $\nu(S_p \otimes H) = p\nu(H)$.

Proof. This results from the decomposition of $S_p \otimes H$ into the union of p disjoint copies of H .

Proposition 7. $\nu(G \otimes H) \geq p\nu(H) + q^2\nu(G)$.

Proof. This results from the decomposition of $G \otimes H$ into the union of the two arc-disjoint graphs $S_p \otimes H$ and $G \otimes S_q$.

Proposition 8. $\tau(G \otimes H) \geq p\tau(H) + q^2\nu(G)$.

Proof. This results from the decomposition of $G \otimes H$ into $S_p \otimes H$

and $G \otimes S_q$, the lemma applied to $\tau(S_p \otimes H)$ and the fact that:
 $\tau(G \otimes S_q) \geq \nu(G \otimes S_q) \geq q^2 \nu(G)$ (Propositions 1 and 6).

Corollary. *When $\tau(G) = \nu(G)$, then Conjectures 2 and 3 are true.*

Theorem 1. $f(pq) \geq pf(q) + q^2 \pi(p)$.

Proof. The theorem can be deduced from Proposition 8, by choosing as graph G a tournament T_p , with $\nu(T_p) = \pi(p)$, and as graph H a tournament T_q , with $\tau(T_q) = f(q)$.

Examples. $p = 3, q = 10: f(30) \geq 145, (\pi(30) = 140); p = 3, q = 11, f(33) \geq 181, (\pi(33) = 176); p = 7, q = 10, f(70) \geq 805, (\pi(70) = 793), p = 7, q = 11: f(77) \geq 987, (\pi(77) = 975)$.

In the case $p = 3$ Theorem 1 leads to $f(3n) \geq n^2 + 3f(n)$. This result has been proved by Kotzig ([7]); from this result Kotzig has deduced the following theorem:

Theorem 2 (Kotzig [7]). *If $f(n) > \pi(n)$, then $f(3n) > \pi(3n) + 1$.*

Proof. From the value of $\pi(n) = \left\lfloor \frac{n}{3} \left[\frac{n-1}{2} \right] \right\rfloor$, we obtain the values of the following table, which proves that $n^2 + 3\pi(n) \geq \pi(3n) - 1$.

n	$\pi(n)$	$n^2 + 3\pi(n)$	$\pi(3n)$
$6k$	$6k^2 - 2k$	$54k^2 - 6k$	$54k^2 - 6k$
$6k + 1$	$6k^2 + k$	$54k^2 + 15k + 1$	$54k^2 + 15k + 1$
$6k + 2$	$6k^2 + 2k$	$54k^2 + 30k + 4$	$54k^2 + 30k + 4$
$6k + 3$	$6k^2 + 5k + 1$	$54k^2 + 51k + 12$	$54k^2 + 51k + 12$
$6k + 4$	$6k^2 + 6k + 1$	$54k^2 + 66k + 19$	$54k^2 + 66k + 20$
$6k + 5$	$6k^2 + 9k + 3$	$54k^2 + 87k + 34$	$54k^2 + 87k + 35$

If $f(n) > \pi(n)$, then $f(n) \geq \pi(n) + 1$ and so by Theorem 1:

$$f(3n) \geq n^2 + 3f(n) \geq n^2 + 3\pi(n) + 3 \geq \pi(3n) + 2 > \pi(3n) + 1.$$

3. GENERALIZATION AND APPLICATIONS TO $f(n)$

Definition. Let p, q, h be three integers; let $G = (X, U)$ be a graph with $|X| = p$; let $H = (Y \cup Z, V)$ be a graph with vertex set the union of the disjoint sets Y and Z , where $|Y| = q$ and $|Z| = h$ (the vertices of Y are labelled from 1 to q and those of Z from $q + 1$ to $q + h$); and let H' be the subgraph of H generated by Z , and H'' the subgraph of H generated by Y . We define a new graph $G^\pm(H, H')$ as follows: its vertex set is $(X \times Y) \cup Z$ and the vertices are labelled (i, j) with $i = 1, 2, \dots, p$ and $j = 1, 2, \dots, q$ (if they belong to $X \times Y$ and (k) with $k = q + 1, \dots, q + h$ (if they belong to Z); in $G^\pm(H, H')$ the subgraph generated by $X \times Y$ is isomorphic to $G \otimes H''$, the subgraphs H_{i_0} (p in number) generated by the vertices (i_0, j) with $j = 1, \dots, q$ and (k) with $k = q + 1, \dots, q + h$ are isomorphic to H for all i_0 . Thus two vertices of $G^\pm(H, H')$ are joined if and only if they are of the form (i_1, j_1) and (i_2, j_2) if either $\overrightarrow{i_1 i_2}$ is an arc of G or $i_1 = i_2$ and $\overrightarrow{j_1 j_2}$ is an arc of H ; (i, j) and $(k) - \overrightarrow{j k}$ being an arc of H ; (k) and $(k') - \overrightarrow{k k'}$ belong an arc of H .

In the case $Z = \phi$, we obtain the lexicographic product $G \otimes H$.

Example.




Fig. 4

We want to find $\tau(G^\pm(H, H'))$ or at least a lower bound; this can be

done by decomposing $G^\pm(H, H')$ into the union of arc-disjoint graphs. We shall give here a lower bound which is good when $\frac{h}{q}$ is small.

Proposition 9. *Let $m(H')$ denote the number of arcs of H' , then:*

$$\tau(G^\pm(H, H')) \geq q^2 \nu(G) + p\tau(H) - (p-1)(m(H') - \tau(H')).$$

Proof. $G^\pm(H, H')$ can be considered as the arc-disjoint union of $G \otimes S_q$, H' , and the p copies of $H - H': H_{i_0} - H'$ (formed by the partial graphs of H_{i_0} obtained by deleting the arcs of H'). Let S be a transversal set of $G^\pm(H, H')$:

$$(1) |S \cap (G \otimes S_q)| \geq \tau(G \otimes S_q) \geq q^2 \nu(G) \text{ by Propositions 1 and 6.}$$

(2) $|S \cap H'| \leq m(H') - \tau(H')$. Indeed let us consider the graph \bar{H}' obtained by reversing the orientation of all the arcs of H' (in number $m(H')$), $\tau(\bar{H}') = \tau(H')$ and so if we reverse in H' strictly more than $m(H') - \tau(H')$ arcs, the obtained graph can be deduced from \bar{H}' by reversing the orientation of strictly less than $\tau(\bar{H}')$ and so it contains a circuit; thus the cardinality of a transversal set is less than or equal to $m(H') - \tau(H')$; $S \cap H'$ being a transversal set of H' we have (2).

(3) $|S \cap H_{i_0}| \geq \tau(H)$ for all i_0 because $S \cap H_{i_0}$ is a transversal set of H_{i_0} which is isomorphic to H . Using the decomposition of $G^\pm(H, H')$ we have: $|S \cap G^\pm(H, H')| = |S \cap (G \otimes S_q)| + \sum_{i_0=1}^p |S \cap (H_{i_0} - H')| + |S \cap H'|$. From $S \cap (H_{i_0} - H') = S \cap H_{i_0} - S \cap H'$ and from (1), (2), (3) we deduce: $|S \cap G^\pm(H, H')| \geq q^2 \nu(G) + p\tau(G) - (p-1)(m(H') - \tau(H'))$ and since S was an arbitrary transversal set, the same lower bound hold for $\tau(G^\pm(H, H')) = \min |S|$.

Proposition 9 implies the following theorem for $f(n)$:

Theorem 3.

$$f(qp + h) \geq q^2 \pi(p) + \max_{H, H'} \{p\tau(H) - (p-1)(m(H') - \tau(H'))\},$$

where the maximum is taken over all graphs H with $q + h$ vertices, H'

being any subgraph of H with h vertices.

Remark. In the case $h = 0$, Proposition 9 and Theorem 3 reduce to Proposition 8 and Theorem 1 respectively. We shall give now lower bounds of $f(pq + h)$ for $h = 1, 2, 3$ (Theorems 4, 6, 8).

Theorem 4. $f(qp + 1) \geq q^2 \pi(p) + pf(q + 1)$.

Proof. We choose a graph H with $\tau(H) = f(q + 1)$; H' being a subgraph with one vertex $m(H') = 0$ and $\tau(H') = 0$.

Examples.

$$\begin{array}{llll} p = 3, q = 9 & f(28) \geq 126 & \pi(28) = 121 \\ p = 3, q = 10 & f(31) \geq 160 & \pi(31) = 155 \\ p = 7, q = 9 & f(64) \geq 672 & \pi(64) = 661 \\ p = 7, q = 10 & f(71) \geq 840 & \pi(71) = 828 \end{array}$$

Theorem 5. If $f(n + 1) > \pi(n + 1)$, then $f(3n + 1) > \pi(3n + 1) + 1$.

Proof. The following table shows that $n^2 + 3\pi(n + 1) \geq \pi(3n + 1) - 1$.

n	$\pi(n + 1)$	$n^2 + 3\pi(n + 1)$	$\pi(3n + 1)$
$6k$	$6k^2 + k$	$54k^2 + 3k$	$54k^2 + 3k$
$6k + 1$	$6k^2 + 2k$	$54k^2 + 18k + 1$	$54k^2 + 18k + 1$
$6k + 2$	$6k^2 + 5k + 1$	$54k^2 + 39k + 7$	$54k^2 + 39k + 7$
$6k + 3$	$6k^2 + 6k + 1$	$54k^2 + 54k + 12$	$54k^2 + 54k + 13$
$6k + 4$	$6k^2 + 9k + 3$	$54k^2 + 75k + 25$	$54k^2 + 75k + 26$
$6k + 5$	$6k^2 + 10k + 4$	$54k^2 + 90k + 37$	$54k^2 + 90k + 37$

Thus, if $f(n + 1) \geq \pi(n + 1) + 1$, we have by Theorem 4:

$$f(3n + 1) \geq n^2 + 3f(n + 1) \geq n^2 + 3\pi(n + 1) + 3 \geq \pi(3n + 1) + 2.$$

Theorem 6. $f(qp + 2) \geq q^2 \pi(p) + pf(q + 2) - (p - 1)$.

Proof. We choose a graph H with $\tau(H) = f(q + 2)$; H' having two vertices $m(H') = 1$ and $\tau(H') = 0$.

Examples.

$$\begin{array}{lll}
 p = 3, q = 8 & f(26) \geq 107 & \pi(26) = 104 \\
 p = 3, q = 9 & f(29) \geq 139 & \pi(29) = 135 \\
 p = 7, q = 8 & f(58) \geq 547 & \pi(58) = 541 \\
 p = 7, q = 9 & f(65) \geq 701 & \pi(65) = 693
 \end{array}$$

Theorem 7. *If $f(n+2) > \pi(n+2)$ and $n \not\equiv 2 \pmod{6}$ then $f(3n+2) > \pi(3n+2)$ and if $f(n+2) > \pi(n+2) + 1$ and $n \equiv 2 \pmod{6}$ then $f(3n+2) > \pi(3n+2) + 2$.*

Proof. The following table shows that if $n \equiv 2 \pmod{6}$, we have: $\pi(3n+2) = n^2 + 3\pi(n+2) + 1$, and if $n \not\equiv 2 \pmod{6}$: $n^2 + 3\pi(n+2) \geq \pi(3n+2)$.

n	$\pi(n+2)$	$n^2 + 3\pi(n+2)$	$\pi(3n+2)$
$6k$	$6k^2 + 2k$	$54k^2 + 6k$	$54k^2 + 6k$
$6k+1$	$6k^2 + 5k + 1$	$54k^2 + 27k + 4$	$54k^2 + 27k + 3$
$6k+2$	$6k^2 + 6k + 1$	$54k^2 + 42k + 7$	$54k^2 + 42k + 8$
$6k+3$	$6k^2 + 9k + 3$	$54k^2 + 63k + 18$	$54k^2 + 63k + 18$
$6k+4$	$6k^2 + 10k + 4$	$54k^2 + 78k + 28$	$54k^2 + 78k + 28$
$6k+5$	$6k^2 + 13k + 7$	$54k^2 + 99k + 46$	$54k^2 + 99k + 45$

Thus if $n \not\equiv 2 \pmod{6}$, by the hypothesis and by Theorem 6 $f(3n+2) \geq n^2 + 3f(n+2) - 2 \geq n^2 + 3\pi(n+2) + 1 \geq \pi(3n+2) + 1$. If $n \equiv 2 \pmod{6}$ by the hypothesis $f(n+2) \geq \pi(n+2) + 2$, and by Theorem 6 $f(3n+2) \geq n^2 + 3f(n+2) - 2 \geq n^2 + 3\pi(n+2) + 4 \geq \pi(3n+2) + 3$.

Proposition 10. *If, in an interval $[a, 3a-1]$, $f(n) > \pi(n)$ and for $n \equiv 4 \pmod{6}$ $f(n) > \pi(n) + 1$, then in the interval $[3a, 9a-1]$, $f(n) > \pi(n)$ and for $n \equiv 4 \pmod{6}$ $f(n) > \pi(n) + 1$.*

Proof. Let n belong to the interval $[3a, 9a-1]$.

Case 1. $n = 3p$, then p belongs to $[a, 3a-1]$; thus, by the hypothesis $f(p) > \pi(p)$ and by Theorem 2 $f(3p) > \pi(3p) + 1$.

Case 2. $n = 3p + 1$ (in particular this case contains $n \equiv 4 \pmod{6}$). Then $p + 1$ belongs to $[a, 3a]$; thus by the hypothesis and for $p = 3a$ by Case 1 $f(p + 1) > \pi(p + 1)$ and by Theorem 5 $f(3p + 1) > \pi(3p + 1) + 1$.

Case 3. $n = 3p + 2$, then $p + 2$ belongs to $[a, 3a + 1]$; more if $p + 2 \equiv 4 \pmod{6}$, $p \equiv 2 \pmod{6}$; thus by the hypothesis and for $p = 3a$ and $p = 3a + 1$ by Case 1 and 2, $f(p + 2) > \pi(p + 2)$ and for $p \equiv 2 \pmod{6}$, $f(p + 2) > \pi(p + 2) + 1$ and so, by Theorem 7, $f(3p + 2) > \pi(3p + 2)$.

Corollary. To prove Conjecture 1 of Kotzig: $f(n) > \pi(n)$ for $n \geq 10$ it suffices to prove that for n belonging to $[10], [29]$: $f(n) > \pi(n)$ and for $n \equiv 4 \pmod{6}$ in this interval $f(n) > \pi(n) + 1$.

In Proposition 4 we have proved that $f(10) > \pi(10) + 1$ and that $f(11) > \pi(11)$. In the example of Theorem 4 we have $f(28) > \pi(28) + 1$ and in the examples of Theorem 6 $f(26) > \pi(26)$ and $f(29) > \pi(29)$. We shall decide new values in the following propositions.

Theorem 8. $f(qp + 3) \geq q^2 \pi(p) + pf(q + 3) - 2(p - 1)$.

Proof. Let us choose as graph H in Theorem 3 a tournament with $\tau(H) = f(q + 3)$ and, as subgraph H' , a 3-circuit (this is possible otherwise H would be a tournament without circuit), then $m(H') = 3$ and $\tau(H') = 1$.

Corollary. $f(24) > \pi(24)$ and $f(27) > \pi(27)$.

Proof. We apply Theorem 8 with $p = 3$ and $q = 7, 8$; we deduce $f(24) \geq 90$ $\pi(24) = 88$; $f(27) \geq 120$ $\pi(27) = 117$.

Proposition 11. $f(12) \geq 21$ (Conjecture 1 is true for 12).

Proof. Let us consider the tournament T_{11} defined in [2] p. 21, where \overrightarrow{ij} is an arc of T_{11} , what we note by $i > j$, if and only if $j - i$ is congruent to a square modulo 11. Let T_{12} be the tournament obtained by adding to T_{11} a vertex, denoted by 12, and the arcs $\overrightarrow{12i}$ for $i = 1, 2, 4, 5, 6, 10$ and $\overrightarrow{j12}$ for $j = 3, 7, 8, 9, 11$. T_{12} can be written

$1 > 2, 4, 5, 6, 10,$	$2 > 3, 5, 6, 7, 11,$	$3 > 1, 4, 6, 7, 8, 12,$
$4 > 2, 5, 7, 8, 9,$	$5 > 3, 6, 8, 9, 10,$	$6 > 4, 7, 9, 10, 11,$
$7 > 1, 5, 8, 10, 11, 12,$	$8 > 1, 2, 6, 9, 11, 12,$	$9 > 1, 2, 3, 7, 10, 12,$
$10 > 2, 3, 4, 8, 11,$	$11 > 1, 3, 4, 5, 9, 12,$	$12 > 1, 2, 4, 5, 6, 10.$

To prove Conjecture 1, as $\pi(12) = 20$, we have to prove that $\tau(T_{12}) \geq 21$. Let S be a transversal set with $\tau(T_{12})$ elements.

Case 1. Let us suppose that S contains no arc with end-vertex 12, and let T_i be the subtournament generated by the vertices 1, 2, 4, 5, 6, 10 and T_j by the vertices 3, 7, 8, 9, 11; the subtournament generated by 2, 4, 5, 6, 10 and T_j being isomorphic to the tournament of the example of Fig. 1 we have $\tau(T_i) = 3$ and $\tau(T_j) = 3$. $S \cap T_i$ must be a transversal set of T_i , thus $|S \cap T_i| \geq 3$ and similarly $|S \cap T_j| \geq 3$; further S must contain all the 15 arcs \vec{ij} with $i = 1, 2, 4, 5, 6, 10$; $j = 3, 7, 8, 9, 11$ otherwise $T_{12} - S$ would contain the circuit $12ij$. Thus $|S| \geq 21$.

Case 2. Let us suppose that S contains at least an arc with end-vertex 12; $S \cap T_{11}$ must be a transversal set of T_{11} and $\tau(T_{11}) = 20$ (see [2]), thus $|S| \geq 21$. In both cases $|S| \geq 21$ and so $\tau(T_{12}) \geq 21$.

Proposition 12. $f(18) > \pi(18)$ and $f(19) > \pi(19)$.

Proof. We have $\pi(18) = 48$ and $\pi(19) = 57$. Let us consider the tournament T_{19} in which \vec{ij} is an arc if and only if $j - i$ is congruent to a square modulo 19 (quadratic residue). This tournament is vertex transitive and so $\tau(T_{19}) = 9 + \tau(T_{18})$ where T_{18} is a tournament obtained by deleting any vertex, for example 19. In T_{18} all the vertices of score (out-degree) 9 are equivalent and all the vertices of score 8 are equivalent: it suffices to consider the automorphisms associated to the powers of the permutation $1 \rightarrow 5 \rightarrow 6 \rightarrow 11 \rightarrow 17 \rightarrow 9 \rightarrow 7 \rightarrow 16 \rightarrow 4 \rightarrow 1$ and $2 \rightarrow 10 \rightarrow 12 \rightarrow 3 \rightarrow 15 \rightarrow 18 \rightarrow 14 \rightarrow 13 \rightarrow 8 \rightarrow 2$. Thus $\tau(T_{18}) = 8 + \tau(T_{17})$ where T_{17} is obtained by deleting any one vertex, for example 18. We shall prove that $\pi(T_{17}) \geq 41$, so $\tau(T_{17}) \geq 41$ and thus $\tau(T_{18}) \geq 49 > \pi(18)$ and $\tau(T_{19}) \geq 58 > \pi(19)$. T_{17} can be written:

$1 > 2, 5, 6, 7, 8, 10, 12, 17;$	$2 > 3, 6, 7, 8, 9, 11, 13;$
$3 > 1, 4, 7, 8, 9, 10, 12, 14;$	$4 > 1, 2, 5, 8, 9, 10, 11, 13, 15;$
$5 > 2, 3, 6, 9, 10, 11, 12, 14, 16;$	$6 > 3, 4, 7, 10, 11, 12, 13, 15, 17;$
$7 > 4, 5, 8, 11, 12, 13, 14, 16;$	$8 > 5, 6, 9, 12, 13, 14, 15, 17;$
$9 > 1, 6, 7, 10, 13, 14, 15, 16;$	$10 > 2, 7, 8, 11, 14, 15, 16, 17;$
$11 > 1, 3, 8, 9, 12, 15, 16, 17;$	$12 > 2, 4, 9, 10, 13, 16, 17;$
$13 > 1, 3, 5, 10, 11, 14, 17;$	$14 > 1, 2, 4, 6, 11, 12, 15;$
$15 > 1, 2, 3, 5, 7, 12, 13, 16;$	$16 > 1, 2, 3, 4, 6, 8, 13, 14, 17;$
$17 > 2, 3, 4, 5, 7, 9, 14, 15;$	

T_{17} contains the following 41 circuits:

1 2 3; 1 5 16; 1 6 11; 1 7 13; 1 8 15; 1 10 14; 1 12 4; 1 17 9; 2 6 17;
 2 7 14; 2 8 12; 2 11 15; 2 13 10; 3 4 13; 3 7 5; 3 8 17; 3 9 15; 3 10 11;
 3 12 16; 3 14 6; 4 8 14; 4 9 6; 4 10 17; 4 11 16; 4 15 7; 5 6 15; 5 9 13;
 5 10 8; 5 11 17; 6 7 8; 6 10 16; 7 11 9; 7 16 17; 8 9 16; 8 13 11;
 10 15 12; 12 17 14; 13 17 15; 14 15 16; 7 12 9 10; 11 12 13 14.

By the aid of a computer program Y. Kodratoff (C.N.R.S. Paris) has found that for this tournament $\tau(T_{17}) = 47$, and for the subtournament T_{16} , obtained by deleting the vertex 4, $\tau(T_{16}) = 40$; thus we have:

Proposition 13. $f(19) \geq 64$; $f(18) \geq 55$; $f(17) \geq 47 > 45 = \pi(17)$
 and $f(16) \geq 40 > 38 = \pi(16) + 1$.

The program applied to other tournaments has given:

Proposition 14. $f(13) \geq 27 > 26 = \pi(13)$.

The program was applied to the following tournament T_{13} :

$1 > 2, 4, 5, 6, 10, 12;$	$2 > 3, 5, 6, 7, 11, 13;$	$3 > 1, 4, 6, 7, 8, 12;$
$4 > 2, 5, 7, 8, 9, 13;$	$5 > 3, 6, 8, 9, 10, 13;$	$6 > 4, 7, 9, 10, 11, 13;$
$7 > 1, 5, 8, 10, 11, 12;$	$8 > 1, 2, 6, 9, 11, 12;$	$9 > 1, 2, 3, 7, 10, 12;$
$10 > 2, 3, 4, 8, 11, 13;$	$11 > 1, 3, 4, 5, 9, 12;$	$12 > 2, 4, 5, 6, 10, 13;$
$13 > 1, 3, 7, 8, 9, 11.$		

The tournament T_{13} being regular, we have Proposition 11 as corollary.

Proposition 15. $f(15) \geq 36 > \pi(15)$ and $f(14) \geq 29 > \pi(14)$.

Let us consider the tournament T_{15} isomorphic to the subtournament of the tournament T_{31} of the quadratic residues module 31, generated by the vertices i such that $\overrightarrow{i31}$ is an arc of T_{31} . T_{15} can be written:

$$\begin{array}{ll}
 1 > 2, 4, 6, 7, 8, 12, 13; & 2 > 3, 5, 7, 8, 10, 11, 13; \\
 3 > 1, 4, 6, 7, 9, 11, 13; & 4 > 2, 5, 7, 8, 9, 12, 14; \\
 5 > 1, 3, 6, 7, 9, 10, 14; & 6 > 2, 4, 7, 8, 10, 11, 15; \\
 7 > 8, 9, 10, 11, 12, 14, 15; & 8 > 3, 5, 9, 11, 12, 13, 15; \\
 9 > 1, 2, 6, 10, 11, 12, 15; & 10 > 1, 3, 4, 8, 11, 13, 14; \\
 11 > 1, 4, 5, 12, 13, 14, 15; & 12 > 2, 3, 5, 6, 10, 13, 15; \\
 13 > 4, 5, 6, 7, 9, 14, 15; & 14 > 1, 2, 3, 6, 8, 9, 12; \\
 15 > 1, 2, 3, 4, 5, 10, 14.
 \end{array}$$

T_{15} is vertex-transitive; therefore the program was applied to a subtournament T_{14} and gave $\tau(T_{14}) = 29$ and therefore $\tau(T_{15}) = 36$.

Remarks.

1. It is enough to prove Conjecture 1 for the odd values of n , because if $f(2p+1) > \pi(2p+1)$ then $f(2p) > \pi(2p)$: suppose that $f(2p) \leq \pi(2p)$, from $\pi(2p+1) = \pi(2p) + p$ and $f(2p+1) \leq f(2p) + p$ (see [2] p. 11) we deduce $f(2p+1) \leq \pi(2p+1)$ contrary to the hypothesis.

2. The result of Erdős and Moon [5] shows that if $n \rightarrow \infty$, $f(n) \sim \frac{n(n-1)}{4}$ but $\pi(n) \sim \frac{n(n-1)}{6}$; thus Conjecture 1 is true if n is sufficiently large.

In summary to prove Conjecture 1 of Kotzig it remains to prove that $f(n) > \pi(n)$ for $n = 21, 23, 25$ and that $f(22) > \pi(22) + 1$. (We hope to verify the conjecture for these values by computer search).

REFERENCES

- [1] C. Berge, *Graphes et Hypergraphes*, Paris, Dunod, 1970.
- [2] J-C. Bermond, Ordres a distance minimum d'un tournoi et graphes partiels sans circuits maximaux, *Math. Sci. Hum.*, 37 (1972), 5-25.
- [3] G. Chartrand – D. Geller – S. Hedetniemi, Graphs with forbidden subgraphs, *J. Comb. Theory*, Vol. 10. No. 1, Ser. B, (1971), 12-41.
- [4] G. Chaty, Cheminements remarquables dans les graphes: Existence, obtention, conservation, Thèse, Université Paris VI, Juin 1971.
- [5] P. Erdős – J.W. Moon, On sets of consistent arcs in tournaments, *Canadian Math. Bull.*, 8 (1965), 269-271.
- [6] H.A. Jung, On subgraphs without cycles in tournament, *Combinatorial theory and its applications II*, Balatonfüred, (eds. P. Erdős, A. Rényi and V.T. Sós) North-Holland, 1970, 675-677.
- [7] A. Kotzig, On maximal order of cyclicity of antisymmetric directed graphs, *preprint of Centre de Recherches Mathematiques*, Montreal.
- [8] J.W. Moon, *Topics on tournaments*, Holt, New York, 1968.
- [9] P. Slater, Inconsistencies in a schedule of paired comparisons, *Biometrika*, 48 (1961), 303-312.