



Toll-like receptor 3 downregulation is an escape mechanism from apoptosis during hepatocarcinogenesis

Marc Bonnin, Nadim Fares, Barbara Testoni, Yann Estornes, Kathrin Weber, Béatrice Vanbervliet, Lydie Lefrancois, Amandine Garcia, Alain Kfoury, Floriane Pez, et al.

► To cite this version:

Marc Bonnin, Nadim Fares, Barbara Testoni, Yann Estornes, Kathrin Weber, et al.. Toll-like receptor 3 downregulation is an escape mechanism from apoptosis during hepatocarcinogenesis. *Journal of Hepatology*, 2019, 71 (4), pp.763-772. 10.1016/j.jhep.2019.05.031 . hal-02367101

HAL Id: hal-02367101

<https://inria.hal.science/hal-02367101>

Submitted on 20 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

KfouryToll-like receptor 3 downregulation is an escape mechanism from apoptosis during hepatocarcinogenesis

Short Title: Poor prognosis of TLR3-negative hepatocarcinoma resistant to TLR3-triggered apoptosis

Marc Bonnin^{1*}, Nadim Fares^{1*}, Barbara Testoni¹, Yann Estornes¹, Kathrin Weber¹, Béatrice Vanbervliet¹, Lydie Lefrançois¹, Amandine Garcia¹, Alain Kfoury¹, Floriane Pez¹, Isabelle Coste¹, Pierre Saintigny^{1,2}, Alain Viari³, Kévin Lang¹, Baptiste Guey¹, Valérie Hervieu⁴, Brigitte Bancel⁴, Birke Bartoch¹, David Durantel¹, Toufic Renno¹, Philippe Merle^{1,5} and Serge Lebecque^{1,6}.

Affiliations of authors:

1. INSERM, U1052, Cancer Research Center of Lyon (CRCL), Université de Lyon (UCBL1), CNRS UMR_5286, Centre Léon Bérard, Lyon, France
2. Department of Translational Research and Innovation and Department of Medicine, Centre Léon Bérard, Lyon, France
3. Synergie Lyon Cancer, Plateforme de Bioinformatique 'Gilles Thomas' Centre Léon Bérard, Lyon, France.
4. Service d'Anatomopathologie, Groupement Hospitalier Est, Hospices Civils de Lyon, Lyon, France.
5. Groupement Hospitalier Lyon Nord, Hepatology Unit, Lyon, France
6. Hospices Civils de Lyon, Laboratoire d'Anatomie et Cytologie Pathologiques, Centre Hospitalier Lyon Sud, Pierre Bénite, France

* Both authors contributed equally to this work.

Correspondence should be addressed to S. Lebecque, CRCL, Centre Léon Bérard, 28, rue de Laennec, 69008 Lyon Cedex 08, France (serge.lebecque@univ-lyon1.fr) (Tél: 33619042629) or to P. Merle, CRCL, 151 Cours Albert Thomas, 69424 Lyon Cedex 03, France (philippe.merle@inserm.fr)

Keywords: TLR3, hepatocarcinoma; apoptosis; pattern recognition receptor; prognostic factor

Words count: 4808 (abstract, lay summary, highlights, introduction, results, discussion, bibliography, M&M)

6 Figures, 1 table (4 supplemental figures, 2 supplemental tables)

List of abbreviations:

TLR3:	Toll-like receptor 3
HCC:	Hepatocarcinoma
IHC:	Immunohistochemistry
PHH:	Primary hepatocytes
HR:	Hazard ratio
CI:	Confidence interval

Disclosures: the authors declare no conflict of interest

Financial support: This research was funded by the French National Cancer Institute (PAIR2009-140) and by the Fondation ARC pour la Recherche sur le Cancer (DOC20121206071). The study sponsors had no role in the design of the study; the collection, analysis, and interpretation of the data; the writing of the manuscript; or the decision to submit the manuscript for publication.

Author Contributions: study concept and design, analysis and interpretation of data: Marc Bonnin, Nadim Fares, Philippe Merle and Serge Lebecque; acquisition of data: Marc Bonnin, Nadim Fares, Floriane Pez, Yann Estornes, Lydie Lefrançois, Amandine Garcia, Alain Kfoury, Béatrice Vanbervliet, Isabelle Coste and Baptiste Guey; design and production of lentiviral vectors: Birke Bartoch; immunohistology: Brigitte Bancel and Valérie Hervieu; statistical analysis: Kévin Lang, Alain Viari, Pierre Saintigny, Nadim Fares and Philippe Merle; drafting of the manuscript: Philippe Merle and Serge Lebecque; critical revision of the manuscript for important intellectual content: Yann Estornes, Kathrin Weber, David Durantel, Toufic Renno, Marc Bonnin and Nadim Fares.

Abstract

Background and aims: Low levels of TLR3 in patients with hepatocarcinoma (HCC) is of poor prognosis owing primarily to the loss of inflammatory signaling for the recruitment of immune cells to the liver. We herein explore the role of TLR3-triggered apoptosis in HCC cells.

Methods: iQRT-PCR, Western blotting, immunohistochemistry and comparative genomic hybridization were used to analyze human and mouse HCC cell lines, and human surgically-resected primary HCC tumors, and to correlate TLR3 expression with HCC patient outcome. Functional analyses were performed following lentiviral transduction for restoration of TLR3 in HCC cells. The role of TLR3-triggered apoptosis in HCC was analyzed *in vivo* in a transgenic mouse model of HCC.

Results: Lower expression of TLR3 in tumor vs. the non-tumor matched tissue was observed at both mRNA and protein levels in primary HCC, and was predictive of shorter recurrence-free survival after surgical resection in both univariate (hazard ratio [HR], 1.79; 95% confidence interval [CI]: 1.04-3.06; $P = 0.03$) and multivariate analyses (HR = 1.73; CI: 1.01-2.97; $P = 0.04$). Immunohistochemistry confirmed frequent downregulation of TLR3 in human and mouse primary HCC cells. None of the 6 human HCC cell lines analyzed expressed TLR3, and ectopic expression of TLR3 following lentiviral transduction not only restored the inflammatory response but also sensitized cells to TLR3-triggered apoptosis. Lastly, in the transgenic mouse model of HCC, absence of TLR3 expression was accompanied by a lower rate of preneoplastic hepatocyte apoptosis and accelerated hepatocarcinogenesis without altering the tumor immune infiltrate.

Conclusion: Downregulation of TLR3 protects transforming hepatocytes from direct TLR3-triggered apoptosis, thereby contributing to hepatocarcinogenesis and poor patient outcome.

Lay summary

HCC is a heterogeneous disease with poor prognosis and limited methods are currently available for predicting patient survival. TLR3 downregulation is associated with shorter survival in HCC patients due to a lack of TLR3-dependent recruitment and activation of immune cells in the liver. We show that following their transformation, hepatocytes become sensitive to TLR3-triggered apoptosis. We also demonstrate that in transgenic mice developing spontaneous HCC, the absence of TLR3 accelerates hepatocarcinogenesis and is associated with a lower rate of apoptosis in the liver, without altering the liver immune

infiltrate. Therefore, the poor prognosis of TLR3^{low} HCC is likely linked to their resistance to TLR3-triggered apoptosis and, in TLR3⁺ HCC, TLR3 may become a promising therapeutic target.

Highlights

- Downregulation of TLR3 in HCC is associated with poor prognosis and with resistance to TLR3-triggered apoptosis.
- Downregulation of TLR3 is an escape mechanism for HCC cells which prevents their apoptosis and enhances tumor progression, independently of the immune response.
- HCC expressing TLR3 may represent suitable targets for therapies based on the use of TLR3 ligands.

Introduction

Hepatocytes are constantly exposed to chemicals, activated by bacterial products, and frequently infected by viruses. These assaults can lead to chronic hepatitis or cirrhosis, two conditions associated with hepatocarcinogenesis. Similarly to other cells, hepatocytes defend themselves against viruses by initiating an innate immune response resulting in the production of type I interferons endowed with anti-viral activities¹. Activation of the innate immune system depends on the detection by pattern recognition receptors (PRRs) of molecular cues signaling the presence of microbes². Among TLRs, which represent the largest family of human PRRs, TLR3 is expressed in the lysosomes of various types of cells, including immune cells, epithelial cells, and hepatocytes. TLR3 recognizes dsRNA³ of viral⁴ origin but also self dsRNA⁵, and activates NF- κ B-, JNK-, AP-1-, and IRF3-dependent signaling pathways in normal cells. In the liver, recognition of HCV dsRNA replicative intermediates by TLR3 protects human hepatocytes *in vitro*^{6,7}, but there is no direct evidence of its involvement in viral hepatitis. Aside from the inflammatory response, TLR3 can also trigger apoptosis in cancer cells⁸⁻¹². We previously described the signaling complex that physically recruits caspase 8 to TLR3 and drives cancer cell death while sparing normal epithelial cells¹³, and a former publication suggests that TLR3 signaling could be “skewed to apoptosis” in hepatocarcinoma¹⁴. In this context, poor prognosis associated with low TLR3 mRNA in HCC and decreased expression of the protein by human transformed hepatocytes has previously been reported. Indeed, Chew et al. proposed a model wherein TLR3 expressed by parenchymal and non-parenchymal liver cells, including NK cells, drives the intratumoral recruitment of T cells that consequently kill cancer cells¹⁵. In the present study, we investigate the expression of TLR3 in cancer hepatocytes and its involvement in apoptotic signaling, and describe TLR3 suppression as an escape mechanism from apoptosis leading to an enhanced hepatocarcinogenesis.

Results

TLR3 is frequently downregulated in human primary HCC

Quantitative comparison of TLR3 protein expression by WB in non-tumoral vs. corresponding tumoral liver samples showed highly significant suppression of TLR3 in 30 tumors (Wilcoxon matched-pairs signed rank test, $p < 0.0001$) (**Figure 1.A and Supplemental Figure 1.A**). We then used a reliable protocol¹⁶ to determine by IHC which cells had lost their TLR3 expression in HCC. In normal liver, all hepatocytes showed a homogeneous and punctuated pattern of cytoplasmic staining compatible with TLR3 localization in endolysosomes (**Supplemental Figure 1.B, upper pictures**). A strong TLR3 signal was also observed in biliary epithelial cells (**Supplemental Figure 1.B, lower left picture**). However, no other cell type expressed TLR3. In HCC (**Figure 1.B**), staining of transformed hepatocytes was heterogeneous, displaying the punctuated pattern seen in normal liver or an enhanced signal close to the nucleus. Similarly to the normal liver, biliary epithelial cells were strongly TLR3⁺ in all tumors, while inflammatory cells were negative (**Supplemental Figure 1.B, lower right picture**). In agreement with our WB data, with Human Protein Atlas data (see supplemental M&M) and with previous reports^{17–19}, semi-quantitative assessment by IHC in tumor vs. matched non-tumor tissues also revealed that TLR3 expression decreased in transformed compared to non-transformed peritumoral hepatocytes in 11/20 HCC samples (**Supplemental Table 1**). Online data from microarray (EBI²⁰, $n = 75$) and RNAseq (TCGA, $n = 50$) analyses demonstrated that TLR3 mRNA expression is also significantly lower ($p < 0.0001$) in human HCC compared to normal liver (**Supplemental Figures 1.C and D, respectively**). Interestingly, such a downregulation is observed neither for TRIF (the adapter shared by TLR3 and TLR4), nor for the two cytoplasmic dsRNA receptors RIG-I and MDA-5. These data prompted us to analyze by RT-qPCR the expression of TLR3 mRNA in 126 primary HCC from patients who had undergone tumor resection (French Cohort features are shown in **Supplemental Table 2**). Compared to normal liver, TLR3 mRNA was expressed at similar or higher level in 80% of the samples (hereafter referred to as the “TLR3 non low” group), but significantly downregulated (cut-off value $< [\text{mean value} - 2\text{SEM}]$ of 8 normal livers) in 26/126 tumors (20%). The downregulation of TLR3 mRNA was not associated with etiology (χ^2 test, $p = 0.70$, data not shown). Patients with a low TLR3 mRNA expression had a significantly shorter recurrence-free survival (RFS) (**Figure 1.C**) with a median RFS for TLR3-low cases of 16 months versus 33 months in the TLR3 non low group (Log-rank test, $p = 0.01$). Moreover, low TLR3 expression was a prognostic factor for early tumor recurrence in univariate analysis (HR =

1.79; CI: 1.04-3.06; $P = 0.03$) and remained independent in multivariate analysis (HR = 1.73; CI: 1.01-2.97; $P = 0.04$). (**Table 1**). Next, we analyzed the correlation between the TLR3 mRNA level and RFS for 306 HCC patients from the TCGA series (see supplemental M&M) who had undergone tumor resection and for which follow-up data was available. By analogy with results described above, we confirmed that the 20% HCC patients ($n = 60$) with the lowest TLR3 mRNA level had a significantly shorter RFS (Log-rank test, $p = 0.006$) (**Supplemental Figure 1.D**). A similar result was observed using a cut-off of 50% to separate TLR3^{high} from TLR3^{low} HCC (Log-rank test, $p = 0.002$, data not shown). The transcription factor, p53, known to enhance TLR3 transcription²¹, is the second most frequently mutated gene in HCC, occurring in about 30% of cases²², and *TP53* mutations are associated with poor prognosis in HCC²³. Although the *TP53* mutation and low level of TLR3 mRNA statuses were correlated (Fischer's exact test, $p = 0.027$), these two parameters remained independent prognostic factors of RFS in multivariate analysis once tumor size had been omitted, owing to its prevailing influence (p53: HR = 1.55, CI = 1.12-2.15, $p = 0.009$; TLR3 mRNA: HR = 1.55, CI = 1.13-2.13, $p = 0.007$). Lastly, closer analysis of the EBI data²⁰ indicated that the loss of TLR3 mRNA expression occurred only after transformation and, as previously reported¹⁵, increased during disease progression (**Figure 1.D**). Altogether, our results confirm that TLR3 downregulation occurs frequently in HCC cells and show that low TLR3 mRNA expression in HCC is an independent prognostic factor for shorter RFS after surgery. The progressive loss of TLR3 mRNA during hepatocarcinogenesis led us to investigate the functions of the receptor in normal vs. transformed hepatocytes.

TLR3 triggers an inflammatory signal in normal human hepatocytes

We used primary human hepatocytes (PHH) and the non-transformed hepatocyte progenitor cell line HepaRG²⁴, which had been shown to respond to Poly(I:C)²⁵, to study the expression and function of TLR3 in normal hepatocytes. In agreement with Luangsay et al.²⁵, PHH and HepaRG expressed similar levels of TLR3 mRNA (**Figure 2A**) and protein by WB (**Figure 2.B**) and flow cytometry (**Figure 2.C**). Western blot analysis revealed that while they both expressed the 72 kDa C-terminal fragment of the cleaved-functional receptor²⁶, PHH alone expressed the 130 kDa full length TLR3 (**Figure 2.B**). In response to the synthetic TLR3 ligand Poly(I:C), PHH and HepaRG both secreted comparable amounts of RANTES and IP10, two chemokines mostly dependent on the transcription factors NF- κ B and IRF, respectively (**Figure 2.D, E**), while they were completely resistant to TLR3-triggered apoptosis (**Figure 2.F**).

TLR3 is absent from human HCC cell lines

In contrast, TLR3 mRNA expression was very low compared to normal hepatocytes in 6/6 HCC cell lines analyzed (**Figure 3.A**), and the protein could not be detected either by flow cytometry (**Figure 3.B**) or by WB in all HCC cell lines (**Figure 3.C**). Of note, the absence of TLR3 was not correlated with the p53 status of the cell lines, as SK-Hep-1 and HepG2 cells express WT p53 while the HuH7 and PLC/PRF/5 cells harbor mutated p53 and Hep3B and FOCUS cells have both alleles deleted^{27,28}. In agreement with the absence of TLR3, Poly(I:C) had no detectable effect on the secretion of RANTES and IP10 (**Figure 3.D, E**) and on the survival (**Figure 3.F**) of Hep3B and HepG2 cells, which we then selected for further studies on the functions of TLR3 in transformed hepatocytes.

Re-expression of exogenous TLR3 sensitizes transformed human hepatocytes to TLR3-triggered apoptosis and inflammatory response

To study the functions of TLR3 in HCC cells, we transduced Hep3B and HepG2 cells with either empty (E-LV) or TLR3-HA-encoding (TLR3-LV) doxycycline-inducible lentiviruses, and confirmed the expression of TLR3 by flow cytometry (**Figure 4.A**) and WB analysis (**Figure 4.B**). Of note, in agreement with a previous publication showing that retroviruses can activate TLR3 and IFN-I production²⁹, transduction with E-LV or with TLR3-LV without induction was by itself sufficient to drive detectable expression of endogenous TLR3 in approximately 15 % of Hep3B cells, corroborating findings by Khvalevsky et al.¹⁴. Re-expression of either endogenous TLR3 by E-LV in Hep3B cells and/or exogenous TLR3 in both cell lines had no effect on cell proliferation in standard culture conditions (data not shown), but restored a strong dose- and cell line-dependent secretion of RANTES and a weaker production of IP-10 in response to dsRNA (**Figure 4.C, D**). Furthermore, and in contrast to wild-type cells, the forced exogenous (HepG2 cells) and/or endogenous (Hep3B cells) expression of TLR3 in both HCC cell resulted in a dose-dependent cell death in response to Poly(I:C) (**Figure 4.E**). This death was an apoptotic cell death, as it was accompanied by caspase 3 activation in the case of HepG2 cells¹⁴, and Hep3B cells were entirely protected by the pancaspase-inhibitor Z-VAD (**Supplemental Figure 2**).

TLR3 inhibits hepatocarcinogenesis in transgenic mice

Similarly to humans, TLR3 expression was undetectable by Western blot in two mouse HCC cell lines (LL-11 and MH22a), and was very weak in Hepa1-6 cells compared to normal

mouse spleen and liver (**Supplemental Figure 3.A**). Moreover, IHC revealed that the expression of TLR3 was strongly repressed in HCC developing spontaneously in transgenic AsVB male mice³⁰ expressing the SV40-TAg in hepatocytes (**Figure 5.A**), as well as in HCC occurring after treatment of WT mice with DEN (**Figure 5.B**). To address the role of TLR3 in hepatocarcinogenesis *in vivo*, we crossed AsVB mice with TLR3 WT and TLR3 KO mice and compared the development of liver tumors in their descendants. At 10 weeks of age, the number of macroscopic tumors was higher (Two-tailed Mann-Whitney test, $p = 0.047$) in TLR3^{-/-} mice than in their TLR3^{+/+} littermates (**Figure 5.C, D**). The number of microscopic tumors was also higher at 8 ($p = 0.012$) and 12 weeks ($p < 0.001$) in TLR3^{-/-} than in wild type mice (**Figure 5.E** and **Supplemental Figure 3.B**). Strikingly, the average size of tumors was comparable in both groups of mice at 8, 10 and 12 weeks, (**Supplemental Figure 3.C**) as was the frequency of Ki67⁺ proliferating cells (**Figure 5.F**). Altogether, those data indicate that the absence of TLR3 was associated with an increased rate of successful hepatocyte transformation without alteration in cancer cell proliferation after their transformation.

Loss of TLR3 is correlated with decreased apoptosis in HCC without alteration of immune cell infiltration

IHC analysis showed no significant decrease in the density of CD3⁺ (**Figure 6.A**) and CD8⁺ T cells (**Figure 6.B**) infiltrating the tumors in TLR3^{-/-} vs. TLR3^{+/+} mice. Accordingly, analysis by flow cytometry showed no difference in the numbers of CD3⁺ and CD4⁺ T lymphocytes and NK cells infiltrating non-tumoral as well as tumoral liver from either TLR3^{+/+} or TLR3^{-/-} mice (**Figure 6.C-E**). Conversely, the frequency of cleaved-caspase 3-positive hepatocytes decreased in the liver of TLR3^{-/-} mice compared to WT mice (**Figure 6.F and G**), and was inversely correlated with the density of microscopic tumor nodules in male ASV-B mice (Pearson correlation, $r = -0.042$, $p = 0.002$) (**Supplemental Figure 4**). In summary, in AsVB transgenic mice, TLR3 is downregulated in tumor hepatocytes and hepatocarcinogenesis is accelerated in the absence of TLR3. Consistently, a TLR3 downregulation is accompanied by a reduction in hepatocyte apoptosis without altering liver infiltration of T cells and NK cells.

Discussion

The poor prognosis associated with TLR3 downregulation in patients with HCC has mainly been attributed to the defective recruitment of immune cells and subsequent lack of killing of transformed hepatocytes. The present work demonstrates that TLR3 can also exert an anti-tumoral role in HCC by inducing apoptosis directly in transformed hepatocytes without

significantly modulating the tumor microenvironment. We therefore propose that during HCC development, the progressive decrease in TLR3 expression in transformed hepatocytes is an escape mechanism for cancer cells from apoptosis and drives tumor formation. In agreement with previous publications^{17,19}, we found that low TLR3 mRNA expression in human HCC is a predictive factor of HCC recurrence.

The direct apoptotic effect of TLR3 on HCC cells was first established when comparing normal vs. transformed hepatocytes. We found that TLR3 is expressed and triggers an inflammatory response^{31,32} but not apoptosis in normal primary hepatocytes. We further confirmed these results with immortalized, non-transformed HepaRG cells. Published data regarding transformed hepatocytes are more controversial, since human HCC cell lines have been reported either to express TLR3 and respond to Poly(I:C) or not, and except for HuH7³² and HepG2³³, the basis for this lack of response was not identified. In the present study, we show that TLR3 mRNA and protein expression are downregulated in 6/6 human HCC cell lines analyzed, which are not responsive to the TLR3 ligand. Importantly, the forced expression of TLR3 sensitized TLR3-deficient HCC cell lines not only to Poly(I:C)-induced inflammation, but also to apoptosis. Consistently, we have previously shown that TLR3 has a death-receptor function in cancerous bronchial epithelial cells³⁵. It is therefore highly pertinent that downregulation of TLR3 mRNA in patients was not observed during cirrhosis but became detectable once HCC had been diagnosed. Moreover, we demonstrated that downregulation of the TLR3 protein, which we observed in 50% of HCC, exclusively affects transformed hepatocytes but not the normal hepatocytes of the matched peritumoral parenchyma. Altogether, our results suggest that the progressive decrease in TLR3 expression by hepatocytes during their transformation may be a protective mechanism from TLR3-triggered apoptosis, thereby enhancing tumor formation.

Chew et al. previously reported that downregulation of TLR3 is included in a 14 immune-gene signature predicting poor patient survival in early HCC. They also proposed a model in which TLR3-dependent apoptosis is mostly indirect, resulting from the killing of HCC cells by cytotoxic immune cells^{15,36}. However, after examination of patient HCC samples, we could not confirm by IHC the presence of TLR3-positive tumor-infiltrating immune cells. Furthermore, the expression of TLR3 by human NK cells remains controversial³⁷. In addition, we found that the direct TLR3-triggered apoptosis of hepatocytes impaired HCC progression in transgenic AsVB mice independently of NK and T cell infiltration. The apparent discrepancy between our data and the results obtained by Chew et al. may be due to the different *in vivo* models used³⁸. While Chew et al. treated Tg(Alb-IHBV) mice with the

synthetic TLR3 ligand Poly(I:C) to activate TLR3 (and MDA5), we did not artificially induce the activation of TLR3 but rather compared spontaneous hepatocarcinogenesis in TLR3^{+/+} with that in TLR3^{-/-} AsVB mice. However, the ligand that triggers TLR3-dependent apoptosis in hepatocytes during their transformation remains unknown. In humans, potential candidates include HCV-derived dsRNA⁶, endogenous retroviruses³⁹, LINE's transcripts⁴⁰ and/or extracellular dsRNA from intestinal commensal bacteria⁴¹ or released by dying hepatocytes^{42,43}, particularly in cells with HBV or HCV. Similar sources of ligands may drive the downregulation of TLR3 in HCC in AsVB mice.

In conclusion, our work provides a set of evidence pointing to a restrictive role for hepatocyte-specific TLR3-triggered apoptosis during hepatocarcinogenesis. It also suggests that the expression of TLR3 in HCC cells may be a useful biomarker to guide clinicians during patient management, in particular to intensify treatments when TLR3 mRNA is downregulated. Moreover, TLR3 might become a promising target for treating 50% of HCC that do express the receptor.

ACKNOWLEDGMENT

Dr. Damien Sanlaville (Hospices Civils de Lyon, Lyon, France) for CGH analysis; the French “Centre de Ressources Biologiques” (CRB) for providing tumor samples for qPCR analysis. The authors thank Pr. M. Rivoire (Centre Léon Bérard, Lyon) for providing human liver tissues and Nicolas Gadot and Armelle Paquet from the Anipath platform (Université de Lyon) for their assistance in histologic and morphometric analyses. The editing assistance of Dr. Brigitte Manship (Centre de Recherche en Cancérologie de Lyon) is acknowledged.

REFERENCES

1. Levy, D. E., Marié, I. J. & Durbin, J. E. Induction and function of type I and III interferon in response to viral infection. *Curr Opin Virol* **1**, 476–486 (2011).
2. Takeuchi, O. & Akira, S. Innate immunity to virus infection. *Immunol. Rev* **227**, 75–86 (2009).
3. Alexopoulou, L., Holt, A. C., Medzhitov, R. & Flavell, R. A. Recognition of double-stranded RNA and activation of NF-kappaB by Toll-like receptor 3. *Nature* **413**, 732–8 (2001).
4. Tatematsu, M., Nishikawa, F., Seya, T. & Matsumoto, M. Toll-like receptor 3 recognizes incomplete stem structures in single-stranded viral RNA. *Nat Commun* **4**, 1833 (2013).
5. Karikó, K., Ni, H., Capodici, J., Lamphier, M. & Weissman, D. mRNA is an endogenous ligand for Toll-like receptor 3. *J Biol Chem* **279**, 12542–50 (2004).
6. Li, K. *et al.* Activation of chemokine and inflammatory cytokine response in hepatitis C virus-infected hepatocytes depends on Toll-like receptor 3 sensing of hepatitis C virus double-stranded RNA intermediates. *Hepatology* **55**, 666–675 (2012).
7. Eksioglu, E. A. *et al.* Characterization of HCV interactions with Toll-like receptors and RIG-I in liver cells. *PLoS ONE* **6**, e21186 (2011).
8. Salaun, B., Coste, I., Rissoan, M.-C., Lebecque, S. J. & Renno, T. TLR3 can directly trigger apoptosis in human cancer cells. *J Immunol* **176**, 4894–901 (2006).
9. Salaun, B., Lebecque, S., Matikainen, S., Rimoldi, D. & Romero, P. Toll-like receptor 3 expressed by melanoma cells as a target for therapy? *Clin. Cancer Res* **13**, 4565–4574 (2007).
10. Friboulet, L. *et al.* Poly(I:C) induces intense expression of c-IAP2 and cooperates with an IAP inhibitor in induction of apoptosis in cancer cells. *BMC Cancer* **10**, 327 (2010).
11. Nomi, N., Kodama, S. & Suzuki, M. Toll-like receptor 3 signaling induces apoptosis in human head and neck cancer via survivin associated pathway. *Oncol. Rep* **24**, 225–231 (2010).
12. Paone, A. *et al.* Toll-like receptor 3 triggers apoptosis of human prostate cancer cells through a PKC-alpha-dependent mechanism. *Carcinogenesis* **29**, 1334–42 (2008).
13. Estornes, Y. *et al.* dsRNA induces apoptosis through an atypical death complex associating TLR3 to caspase-8. *Cell death and differentiation* (2012). doi:10.1038/cdd.2012.22
14. Khvalevsky, E., Rivkin, L., Rachmilewitz, J., Galun, E. & Giladi, H. TLR3 signaling in a hepatoma cell line is skewed towards apoptosis. *J. Cell. Biochem* **100**, 1301–1312 (2007).
15. Chew, V. *et al.* Chemokine-driven lymphocyte infiltration: an early intratumoural event determining long-term survival in resectable hepatocellular carcinoma. *Gut* **61**, 427–438 (2012).
16. Combaz-Lair, C. *et al.* Immune biomarkers PD-1/PD-L1 and TLR3 in malignant pleural mesotheliomas. *Hum. Pathol.* **52**, 9–18 (2016).
17. Chen, X.-L., Xu, Y.-Y., Chen, L., Wang, G.-L. & Shen, Y. TLR3 Plays Significant Roles against HBV-Associated HCC. *Gastroenterol Res Pract* **2015**, 572171 (2015).
18. Yoneda, K. *et al.* Dual topology of functional Toll-like receptor 3 expression in human hepatocellular carcinoma: differential signaling mechanisms of TLR3-induced NF-kappaB activation and apoptosis. *Int. J. Oncol* **33**, 929–936 (2008).
19. Yuan, M.-M. *et al.* TLR3 expression correlates with apoptosis, proliferation and angiogenesis in hepatocellular carcinoma and predicts prognosis. *BMC Cancer* **15**, 245 (2015).
20. Wurbach, E. *et al.* Genome-wide molecular profiles of HCV-induced dysplasia and hepatocellular carcinoma. *Hepatology* **45**, 938–947 (2007).

21. Taura, M. *et al.* p53 regulates Toll-like receptor 3 expression and function in human epithelial cell lines. *Mol. Cell. Biol.* **28**, 6557–6567 (2008).
22. Lee, J.-S. The mutational landscape of hepatocellular carcinoma. *Clin Mol Hepatol* **21**, 220–229 (2015).
23. Woo, H. G. *et al.* Association of TP53 mutations with stem cell-like gene expression and survival of patients with hepatocellular carcinoma. *Gastroenterology* **140**, 1063–1070 (2011).
24. Parent, R., Marion, M.-J., Furio, L., Trépo, C. & Petit, M.-A. Origin and characterization of a human bipotent liver progenitor cell line. *Gastroenterology* **126**, 1147–1156 (2004).
25. Luangsay, S. *et al.* Expression and functionality of Toll- and RIG-like receptors in HepaRG cells. *J. Hepatol.* **63**, 1077–1085 (2015).
26. Toscano, F. *et al.* Cleaved/Associated TLR3 Represents the Primary Form of the Signaling Receptor. *J. Immunol.* (2012). doi:10.4049/jimmunol.1202173
27. http://p53.free.fr/Database/Cancer_cell_lines/HCC.html.
28. <http://p53.iarc.fr/CellLines.aspx>.
29. Lee, J. *et al.* Activation of innate immunity is required for efficient nuclear reprogramming. *Cell* **151**, 547–558 (2012).
30. Dubois, N. *et al.* Time-course development of differentiated hepatocarcinoma and lung metastasis in transgenic mice. *J. Hepatol.* **13**, 227–239 (1991).
31. Li, K., Chen, Z., Kato, N., Gale, M. & Lemon, S. M. Distinct poly(I-C) and virus-activated signaling pathways leading to interferon-beta production in hepatocytes. *J. Biol. Chem* **280**, 16739–16747 (2005).
32. Takii, Y. *et al.* Enhanced expression of type I interferon and toll-like receptor-3 in primary biliary cirrhosis. *Lab. Invest.* **85**, 908–920 (2005).
33. Li, K., Chen, Z., Kato, N., Gale, M. & Lemon, S. M. Distinct poly(I-C) and virus-activated signaling pathways leading to interferon-beta production in hepatocytes. *J. Biol. Chem.* **280**, 16739–16747 (2005).
34. Khvalevsky, E., Rivkin, L., Rachmilewitz, J., Galun, E. & Giladi, H. TLR3 signaling in a hepatoma cell line is skewed towards apoptosis. *J. Cell. Biochem.* **100**, 1301–1312 (2007).
35. Alkurdi, L. *et al.* Release of c-FLIP brake selectively sensitizes human cancer cells to TLR3-mediated apoptosis. *Cell Death & Disease* **9**, (2018).
36. Chew, V. *et al.* Toll-Like Receptor 3 Expressing Tumor Parenchyma and Infiltrating Natural Killer Cells in Hepatocellular Carcinoma Patients. *JNCI J Natl Cancer Inst* **104**, 1796–1807 (2012).
37. Adib-Conquy, M., Scott-Algara, D., Cavaillon, J.-M. & Souza-Fonseca-Guimaraes, F. TLR-mediated activation of NK cells and their role in bacterial/viral immune responses in mammals. *Immunol. Cell Biol.* **92**, 256–262 (2014).
38. Chew, V. TLR3-expressing Tumor Parenchyma and Infiltrating NK Cells Promote Tumor Control in Hepatocellular Carcinoma Patients.
39. Yu, P. *et al.* Nucleic acid-sensing Toll-like receptors are essential for the control of endogenous retrovirus viremia and ERV-induced tumors. *Immunity* **37**, 867–879 (2012).
40. Guler, G. D. *et al.* Repression of Stress-Induced LINE-1 Expression Protects Cancer Cell Subpopulations from Lethal Drug Exposure. *Cancer Cell* **32**, 221–237.e13 (2017).
41. Kawashima, T. *et al.* Double-Stranded RNA of Intestinal Commensal but Not Pathogenic Bacteria Triggers Production of Protective Interferon- β . *Immunity* (2013). doi:10.1016/j.immuni.2013.02.024
42. Zordevic-Khvalevsky, E. *et al.* Toll-like receptor 3 signaling attenuates liver regeneration. *Hepatology* **50**, 198–206 (2009).
43. Cavassani, K. A. *et al.* Toll like receptor 3 plays a critical role in the progression and

- severity of acetaminophen-induced hepatotoxicity. *PLoS ONE* **8**, e65899 (2013).
44. Gripon, P. *et al.* Infection of a human hepatoma cell line by hepatitis B virus. *Proc. Natl. Acad. Sci. U.S.A.* **99**, 15655–15660 (2002).
45. Merle, P. *et al.* Oncogenic role of the frizzled-7/ β -catenin pathway in hepatocellular carcinoma. *Journal of Hepatology* **43**, 854–862 (2005).

Materials and Methods

Origin and analysis of cell lines and patient samples

Human cell lines were purchased from the ATCC except HepaRG⁴⁴, FOCUS and HuH-7 that were maintained in our laboratory. We also purchased the mouse Hepa1-6 (ATCC[®] CRL-1830[™], C57L) and MH-22A (ECACC 96121721, C3HA) cells. We had previously generated the LL-11 HCC cell line (C57Bl/6 X DBA/2)⁴⁵. Primary human hepatocytes (PHH) and normal liver tissues (n = 11) were isolated from surgical resections of liver metastases developing from primary colorectal adenocarcinoma at the Centre Léon Bérard (France). Tumor samples (n = 126) for TLR3 mRNA and clinical data analyses were collected and performed at the Centre Français de Ressources Biologiques (ministerial agreements #AC-2013-1871 and DC-2013-1870). All of the patients had given their written informed consent in compliance with requirements of the local ethical committee of each institution.

HCC mouse models

All animal experiments, within this study, were performed according to animal protocol (A 69383 1201) which was approved by the Lyon SFR animal ethic committee (C2EA-15: CECCAPP). Mice (5 in 1 cage) were provided free access to food and water and kept in a colony room on a 12:12-hour light and dark cycle. The SV40 mouse model of HCC (AsVB, B6D2F1 hybrids) has been previously described³⁰. SV40 (male) and TLR3 WT or TLR3^{-/-} (female) mice (both C57BL/6) were crossed. F1 Male mice were systematically sacrificed at ages of 8 (4 TLR3 WT and 5 TLR3^{-/-}), 10 (8 TLR3 WT and 7 TLR3^{-/-}) and 12 weeks of life (7 TLR3 WT and 6 TLR3^{-/-}). Liver were examined macroscopically under 2X magnifying glasses. The right lobe of the liver was fixed in 10% buffered formalin and included in paraffin for histological analysis. Diethylnitrosamine (DEN) HCC mouse model: a single dose, 25 mg/kg of DEN was injected intra-peritoneally at day 15 in C57BL/6 TLR3 WT or TLR3^{-/-} male mice (n = 10) that were sacrificed at 9 months. Livers were harvested and fixed in 10% buffered formalin and included in paraffin.

Immunohistochemistry and quantitative analysis of tissue sections

Formaldehyde-fixed paraffin-embedded (FFPE) tissue sections from human HCC (obtained from the Pathology Department of the Groupement Hospitalier Lyon Nord, Hospices Civils de Lyon) and from livers of 4 month-old AsVB transgenic mice³⁰ expressing the SV40-TAg under the anti-thrombin III promoter were stained with antibodies specific for TLR3, cleaved caspase 3, Ki67, mCD3 and mCD8 as detailed in the Supplemental Methods. Expression of

TLR3 in tumor vs. non-tumor tissues was scored semi-quantitatively as follows: the percentage of hepatoma cells expressing no, less, equal or more TLR3 than peritumoral hepatocytes was counted in 10 random fields at 100X magnification and multiplied by a factor of 0, 1, 2 or 3, respectively. A sample with the same TLR3 expression in tumoral and peritumoral tissue was scored 200. Values ranging < 160 and > 240 were considered to be “low” and “high” values, respectively. For mouse HCC, 4 µm thick FFPE tissue sections were obtained from 3 random cutting planes from the right liver lobe. All staining was performed with the automated immunostainer (Ventana Discovery XT, Roche, Meylan, France) using the DABmap Kit according to the manufacturer’s instructions. All quantifications were performed blinded to the status of TLR3. Image analyses were performed under light microscope (Eclipse E400, Nikon France, Champigny, France) equipped with a tri-CDD video camera (Sony, Japan). The total surface of the liver section was determined by morphometric analysis (Histolab, Microvision Instruments, Evry, France). The median number of nodules was determined on H&E stained sections and reported to the total surface analyzed. Median size of nodules was automatically quantified by morphometric analysis and expressed in mm². Cleaved caspase 3 quantification was performed on at least 50 fields (magnification X100) randomly chosen on non-tumoral liver sections and reported to the total surface analyzed. Percentage of Ki67⁺ cells was determined automatically by morphometric analysis on 50 randomly chosen non-tumoral fields (median of 40000 nuclei analyzed). CD3⁺ and CD8⁺ lymphocytes quantification was measured on 50 fields randomly chosen on liver sections and reported to the total surface of the fields determined by morphometric analysis and presented as CD3 and CD8 density.

TLR3 mRNA quantification

TLR3 mRNA purified from tissue biopsies was quantified by qPCR with the Lightcycler 480 II apparatus (Roche Applied Science, Penzberg, GmbH) as detailed in the Supplemental Methods, and compared to the 50S ribosomal protein L15 (RPLO) for the study of the cell lines and to the 18S ribosomal RNA (18S) for the clinical samples, using the $\Delta\Delta C_t$ method as described by the manufacturer. The TLR3/18S value for normal livers (n = 8) was 4093 +/- 863 (mean +/- SEM). TLR3 mRNA expression < mean level in normal liver - 2 SEM (TLR3/18S < 2366) was recorded as low.

Statistics

Kaplan-Meier univariate survival analysis was performed and hazard ratios (HR), 95% confidence intervals (CIs), and *P*-values were calculated using the log-rank (Mantel–Cox) test. Paired *t*-tests, Mann Whitney test, Fisher’s exact, Wilcoxon matched-pairs signed rank test and two-ways analysis of variance tests were also used in experiments as indicated. All statistical tests were two-tailed.

Others methods

Additional information regarding the cell culture conditions, online data from normal and HCC livers, reagents, inducible lentiviral vectors, TLR3 mRNA quantification, Western blots, comparative genomic hybridization, flow cytometry analysis, the cytokines/chemokines dosage, and immunohistochemistry are described in the Supplemental Methods (available online).

FIGURES LEGENDS

FIGURE 1. Expression and prognostic value of TLR3 in human HCC. (A) Relative expression of TLR3 vs the actin protein determined by Western blot (WB) in 30 pairs of peritumoral (PT) vs. corresponding tumoral tissues (T) from the French Cohort of HCC. (B) Representative images of TLR3 expression detected by IHC in human HCC: upper pictures show the interface between peritumoral (PT) and tumoral (T) tissues in a TLR3⁺ HCC. Lower pictures show TLR3^{low} HCC. Bars correspond to 100 μ m. Original magnification: 100X (left) and 200X (right). (C) Disease-free survival (DFS) of 126 HCC patients of the French cohort according to the level of TLR3 mRNA in the tumor relative to normal liver. (D) Comparison of mRNA expression of TLR3 in normal, cirrhotic, preneoplastic and cancerous livers according to the EBI HCC cohort data available online (see results; n = 75). Statistics: A = Wilcoxon matched-pairs signed rank test; C = Log-rank test; D = two-tailed unpaired t-test.

FIGURE 2. Expression of TLR3 and response to Poly(I:C) in normal human hepatocytes. (A) TLR3 mRNA expression in unstimulated PHH and HepaRG cells. (B) WB analysis of TLR3 expression by unstimulated PHH and HepaRG cells. (C) FACS analysis of intracellular TLR3 expression by unstimulated PHH and HepaRG cells. Grey = control isotype Ab. Numbers indicate the percentage of TLR3⁺ cells. (D, E) PHH and HepaRG cells were treated for 24 h with the indicated concentrations of Poly(I:C) and the concentration of RANTES and IP-10 in the supernatant was measured by ELISA. (F) At the end of the experimental culture periods described above, the percentage of Annexin V⁺ apoptotic cells was measured by flow cytometry. (A-C) Data are representative of 3 independent experiments. Means \pm SEM of 3 independent experiments are shown (D-F). Statistics = two-tailed unpaired t-test (D, E).

FIGURE 3. TLR3 is suppressed in human hepatocarcinoma cell lines. (A) TLR3 mRNA expression in 6 unstimulated human HCC cell lines compared with HepaRG cells. (B) WB analysis of TLR3 expression by unstimulated PHH and HepaRG cells and in 6 human HCC cell lines. (C) FACS analysis of intracellular TLR3 expression by 6 human HCC cell lines. Grey = control isotype Ab. Numbers indicate the percentage of TLR3⁺ cells. Hep3B and HepG2 cells were treated for 24 h with 10 μ g/ml of Poly(I:C) and the concentration of RANTES (D) and IP-10 (E) in the supernatants was measured by ELISA, and the percentage of Annexin V⁺ apoptotic Hep3B and HepG2 cells was measured by flow cytometry (F). (A-C) Data are representative of 3 independent experiments. Means \pm SEM of 3 independent experiments are shown (D-F). Statistics = two-tailed unpaired t-test (D-F).

FIGURE 4. Re-expression of TLR3 restores inflammatory response and triggers apoptosis in human HCC cell lines. (A) Hep3B and HepG2 cells expressing either empty (E-LV) or TLR3-HA-encoding doxycycline-inducible lentiviral vectors (TLR3-LV) were treated with doxycycline for 24 h and the expression of TLR3 was analyzed by FACS using the anti-TLR3 mAb. Grey = control isotype Ab. Numbers indicate the percentage of TLR3⁺ cells. (B) The same cells were treated with increasing doses of doxycycline for 24 h and the expression of TLR3 was analyzed by WB with anti-TLR3 or anti-HA mAb, as indicated. The same cells were treated with doxycycline for 24 h followed by 24 h with increasing doses of Poly(I:C). Secretion of RANTES (C) and IP-10 (D) was measured by ELISA, and the percentage of Annexin V⁺ apoptotic Hep3B and HepG2 cells was measured by flow cytometry (E). (A, B) Data are representative of 3 independent experiments. Means \pm SEM of 3 independent experiments are shown (C-E). Statistics = two-tailed unpaired t-test (C-D).

FIGURE 5. HCC growth is accelerated in TLR3^{-/-} mice. Liver tissue sections from (A) 4 month-old transgenic female and male AsVB mice and from (B) 9 month-old male mice treated with DEN stained with mTLR3-specific antibody. Right images are enlarged views of the white rectangle area in corresponding images on the left (bar = 50 μ m). Original magnification = 200X. These images are each representative of >3 livers analyzed. (C) Macroscopic view and (D) number of macroscopic nodules detected at the surface of the livers of 10 weeks-old TLR3 WT and TLR3^{-/-} AsVB mice. (E) Density of microscopic nodules in livers from 8-, 10- and 12-week-old TLR3 WT and TLR3^{-/-} AsVB mice. (F) Percentage of proliferating Ki67⁺ cells detected by IHC in the liver of 8-, 10- and 12 weeks-old TLR3 WT and TLR3^{-/-} male AsVB mice. Means \pm SEM are shown. Statistics = two-tailed Mann-Whitney test (D-F).

FIGURE 6. Absence of TLR3 is associated with decreased apoptosis without alteration of T lymphocyte infiltration in mouse HCC. Density of (A) CD3⁺ T lymphocytes and (B) CD8⁺ T lymphocytes measured by IHC in the livers of 8-, 10- and 12-week-old TLR3 WT and TLR3^{-/-} male AsVB mice. Percentage of CD3⁺ (C), CD4⁺ (D) and NK cells (E) among CD45⁺ cells isolated from peritumoral (PT) and tumoral (T) liver tissues of 12-week-old TLR3 WT and TLR3^{-/-} male AsVB mice (n = 3) measured by flow cytometry. (F) Density of cleaved caspase-3⁺ apoptotic cells measured by IHC in the livers of 8-, 10- and 12-week-old TLR3 WT and TLR3^{-/-} male AsVB mice. (E) Liver tissue section from 12-month-old WT male AsVB mice stained with anti-Ki67 (upper panel) and anti-cleaved caspase 3 antibody (lower panel).

Bar = 50 μm . Original magnification = 100X. Means \pm SEM are shown. Statistics = two-tailed unpaired t-test (A-F).

Table 1
Univariate and multivariate analysis of clinical, pathological, and molecular variables for disease-free survival for patients treated by surgical resection as first line therapy (n = 126).

Variables	Univariate analysis		Multivariate analysis	
	HR (95% CI)	Wald test P value	HR (95% CI)	Wald test P value
Sex (male)	1.08 (0.62 - 1.88)	.77		
Age (>60 years)	0.99 (0.97 - 1.01)	.23		
Etiology (HBV)	1.31 (0.77 - 2.24)	.30		
Etiology (HCV)	0.64 (0.37 - 1.10)	.10		
Etiology (alcohol)	1.66 (0.66 - 1.92)	.54		
Etiology (NASH)	1.31 (0.77 - 2.22)	.32		
Cirrhosis	1.09 (0.69 - 1.71)	.72		
Tumor size > 50 mm	1.84 (1.15 – 2.95)	.01	1.80 (1.12-2.90)	.02
Microvascular invasion	1.12 (0.69 - 1.79)	.64		
AFP > 400 ng/mL	1.12 (0.63 - 2.02)	.69		
Presence of satellite nodules	1.45 (0.88 - 2.39)	.10		
Poor differentiation (OMS classification)	1.24 (0.91 – 1.68)	.16		
Low TLR3 mRNA	1.79 (1.04 - 3.06)	.03	1.73 (1.01-2.97)	.04

AFP: α-fetoprotein; HBV: hepatitis B virus, HCV: hepatitis C virus; HR, hazard ratio; NASH: Non-alcoholic steato-hepatitis

Patients with early perioperative mortality after resection (< 2 month), as well as patients treated by liver transplantation were removed from the initial cohort. Cox proportional hazards regression test was used for univariate and multivariate analysis. Parameters with p value ≤0.1 in univariate analysis were included in the multivariate model.

Figure 1






Figure 1




Supplemental Figure 1


A




B



C



D



E




Figure 2




Figure 3






Figure 4

A




B




C

RANTES




D

IP10






E

Annexin V⁺



HepG2



Supplemental Figure 2






Figure 5



Supplemental Figure 3



C






Figure 6



Supplemental Figure 4



normal hepatocyte
TLR3⁺

cancer hepatocyte
TLR3⁺

