

Impact of a thin plantar orthopaedic insert on posture and locomotion

Carole Puil, Armel Crétual, Anne-Hélène Olivier

▶ To cite this version:

Carole Puil, Armel Crétual, Anne-Hélène Olivier. Impact of a thin plantar orthopaedic insert on posture and locomotion. SOFPEL 2019 - 26ème congrès de la Société Francophone Posture, Équilibre et Locomotion, Dec 2019, Montréal, Canada. hal-02373802

HAL Id: hal-02373802 https://inria.hal.science/hal-02373802

Submitted on 25 Feb 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





Impact of a thin plantar orthopaedic insert on posture and locomotion

Carole PUIL^{1, 2}, Armel CRETUAL¹, Anne-Hélène OLIVIER¹
Univ Rennes, Inria, M2S - EA 7470, F-35000 Rennes, France¹
IFPEK²





Contact mail: carole.puil@gmail.com


INTRODUCTION

EMI® (Medio-Intern Element):

- Podiatrist tool included in insoles
- Correction of postural deficiency: Contralateral effect of a 3 mm high EMI® on ML

displacement of the CoP of healthy participants¹

- Effect on Eye convergence²
- => Lack of evaluation
- => Only on static tasks



AIM AND HYPOTHESIS

Aim: Evaluate the effect of the EMI® on posture and locomotion Hypothesis: effect of EMI® on ML and AP axis and on Ellipse surface


DESIGN AND ANALYSIS

- Participants: 19 healthy volunteers (9 \circlearrowleft , 10 \updownarrow , 27.05 yrs (±4.24))
- EMI® conditions: control (without), under left foot, under right foot

Static (Force plate ATMI®)

Conditions:

- Feet: parallel and 30° angle
- Eyes: open and close





Analysis:

CoP displacement on ML and AP axis:

- Average position: \overline{ML} \overline{AP}
- ⇒ Between trials within subject variability: BTV_{AP}, BTV_{ML}
- Within trial within subject variability


Ellipse Area


Locomotion (Vicon®)


Design based on Podokinetic after rotation (PKAR) protocol^{3,4} (30' of walking stimulation)

Conditions:

- Eyes: open and close
- Trajectory: straight, turn left 90°, turn right 90°








Analysis:

Comparison between the difference in final orientation of locomotion with/without vision, with a EMI® and without a EMI®




RESULTS

STATIC:

Only significant effects on feet position, eye condition and interaction between eye and feet position.

No significant effect of EMI®




Results of the 4 ways repeted mesures ANOVA (EYExFOOTxEMIxEMIside)

Significant effects are represented as well as their size effect (ŋ²p)

LOCOMOTION:

No significant effect of EMI® on the final orientation of the walking



CONCLUSION AND FUTURE WORKS

Our protocol, involving **healthy participants**, was not able to show significant effect of EMI® on posture and locomotion

Feet positions has significant effects on posture: clinical activities have to be standardized to be able to compare CoP movement.

Future works:

- Evaluate population with foot postural deficiency:
 More effect of EMI® on eye vergence on a population with foot postural deficiency²
- Normalize the evaluation of foot postural deficiency:
 No standardized procedure (compare foam effect² on CoP with Depron®)

BIBLIOGRAPHY