



Data-driven mechanistic modeling of metastasis: cancer at the organism scale

Sébastien Benzekry

► To cite this version:

Sébastien Benzekry. Data-driven mechanistic modeling of metastasis: cancer at the organism scale. RITS conference (Recherche en Imagerie et Technologies pour la Santé) of the French Society of Biomedical Engineering (SFGBM), May 2019, Tours, France. hal-02424452v2

HAL Id: hal-02424452

<https://inria.hal.science/hal-02424452v2>

Submitted on 27 Dec 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Data-driven mechanistic modeling of metastasis: cancer at the organism scale

S. Benzekry


Inria Bordeaux Sud-Ouest

21 mai 2019



Metastasis (μετά = beyond, στάσις = place)

- Metastases are the **main cause of death** (>90%) from solid cancers *Lambert and Weinberg, Cell, 2017*




Breast

- 94% of cases are local or regional at diagnosis but 30% will relapse
Pollard, N Eng J Med, 2016
- Estimate the amount of **residual distant disease** at diagnosis in order to **personalize** the adjuvant (chemo)-therapy
- Avoid unnecessary, heavy **toxicities**


Lung

- 57% of cases are metastatic
- Decide whether to use **whole brain radiation therapy** or just (stereotactic) surgery
- Avoid cognitive impairment of the patient



Institut Bergonié, Bordeaux

Mechanistic model of metastatic dissemination and growth



Growth rates of primary and secondary tumors g_p and g

$$\frac{dV_p}{dt} = (\alpha_p - \beta_p \ln(V_p)) V_p \quad (\text{Gompertz})$$

$$g(v) = (\alpha - \beta \ln(v)) v$$

Dissemination rate

$$d(V_p) = \mu V_p$$

Size distribution of the metastases $\rho(t, v)$



$$\begin{cases} \partial_t \rho(t, v) + \partial_v (g(v) \rho(t, v)) = 0 \\ g(V_0) \rho(t, V_0) = d(V_p(t)) \\ \rho(0, v) = \rho^0 \end{cases}$$

Metastatic burden (total number of metastatic cells)

$$M(t) = \int_{V_0}^{+\infty} v \rho(t, v) dv = \int_0^t d(V_p(t-s)) V(s) ds$$




Validation on animal data



Nonlinear **mixed-effects**
statistical model for inter-
animal variability

$$\theta^i = \theta_{pop} + \eta_i, \quad \eta_i \sim \mathcal{N}(0, \omega^2)$$



⇒ same growth for PT and mets: $\alpha_p = \alpha$, $\beta_p = \beta$

Current work: implementing the mechanistic model into a statistical learning model

- Use the model for predictions of **time-to-relapse (TTR)**


$$TTR^i = \inf \left\{ t > 0; N_{vis} \left(t, \theta^i \right) > 1 \right\}$$

- Implement clinical variables and biomarkers as **biologically meaningful** covariates

Features selection based on Cox regression

	P
tumor_size_clinical	0.00165
tumor_size_histological	0.4
grade	0.358
histology	0.0041
TNM_T	0.146
TNM_N	0.599
nb_invaded_ganglions	0.619
menopausal_status	0.000471
ER	0.543
PR	0.34
Ki67	3.09e-05
HER2	0.17
HER2.intensity	0.482
CK56	0.793
EGFR	0.016
VIM	0.226
ALDH1	0.674
CD24	0.894
CD44	0.397
E_cadherin	0.207
TRIO	0.0646
BCL2	0.727
age_at_diagnosis	0.0599
diagnosis_year	0.101
radio	0.276

x^i




$$\ln(\mu^i) = \ln(\mu_{pop}) + \beta_\mu^T x^i + \eta_\mu^i, \quad \eta_\mu^i \sim \mathcal{N}(0, \omega_\mu^2)$$

Parameter	Estimate	r.s.e. (%)	p-value
$\log \alpha_{pop}$	-8.883	10.151	
$\beta_{Ki67,\alpha}$	0.086	27.376	$2.59 \cdot 10^{-4}$
$\beta_{HER2,\alpha}$	0.029	42.883	0.820
$\beta_{CD44,\alpha}$	0.011	60.816	0.1
$\beta_{TRIO,\alpha}$	0.016	58.119	0.085
$\log \mu_{pop}$	-26.342	3.696	
$\beta_{EGFR,\mu}$	0.039	47.527	0.035
σ	0.606	23.104	
ω_α	2.062	22.715	
ω_μ	3.563	16.759	


	AUROC	Accuracy	NPV	PPV
Random forest	0.727	87	91.7	22.5
Logistic regression	0.772	90.1	91	25
Cox regression	0.728	87.4	91.1	16.7
Bio-based	0.641	89.7	91.3	31.2

AUROC = Area Under the ROC curve, NPV = Negative Predictive Value
PPV = Positive Predictive Value




Data of a NSCLC patient with brain mets


Primary tumor size



Metastases size




Number of visible mets




Lung CT





Brain CT scan





The model with dormancy could describe best the data




Objective function


Model	Patient 1	Patient 2
Base	5.51	2.53
Secondary	5.43	2.3
Delay	5.23	1.53
Dormancy	4.93	1.71
Diff. growth	4.95	1.79

Dormancy estimated to 133 days \pm 4.2%

Model



Data



$t = -65$ months



— 10 mm

*

Primary tumor (lung)

Metastases (brain)

$t = -65.1$ months




2 Combination bevacizumab - chemotherapy

Vascular normalization: a time window for improved pharmacokinetics?

- Bevacizumab = anti-VEGF monoclonal antibody ⇒ **anti-angiogenic** action (first approved in 2004)
- Only proved clinical efficacy when **combined (concomitantly) with cytotoxics**
- Possible explanation: transient **normalization** of the otherwise abnormal (leaky, tortuous) vascular architecture



Vakoc et al., Jain, 2009, Nat Med



Jain, Nat Med, 2001

Therapeutic question

What is the **optimal time gap** between administration of bevacizumab and cytotoxic chemotherapy? How to capture **inter-individual variability** for designing **personalized therapies**?

Hypothesis: sequential use of bevacizumab associated with chemotherapy would achieve better efficacy and modeling support could help to define the optimal time-window

Current beva-chemo
regimen are
underpowered



Experimental therapeutics

Breast cancer model

Mollard et al. (Benzekry), Oncotarget 2017

Lung cancer model


Experiment 1

Calibration



Experiment 2



Prediction



Modeling and Simulation

Imbs et al. (Benzekry), CPT: Pharmacometrics Syst Pharmacol, 2017

Semi-mechanistic mathematical model



Simeoni et al., Rocchetti, Cancer Res, 2004


Benzekry et al., CRAS, 2012

Millard et al., Benzekry, Oncotarget, 2017

Imbs et al., Benzekry, CPT: Pharmacometrics Syst Pharmacol, 2018



+ PK models for beva $A(t)$ and CT $C(t)$ concentrations

Control




Model fits: individual + population level (NLME)


Simultaneous





Median growth curves






Sequential C/B



Sequential B/C



Prediction of the optimal delay



⇒ to be **tested experimentally**

⇒ **personalized** scheduling

Acknowledgements

Biology

- Preclinical data of ortho-surgical animal models of metastases

*J. Ebos

*A. Tracz

*M. Mastri



Roswell Park Cancer Institute, Buffalo, NY, USA

- Beva + cytotoxics study



Dr. J. Ciccolini

Clinic

- Brain metastasis from lung tumors

*F. Chomy

Bergonié Institute, Bordeaux, FR



*F. Barlesi

*X. Muracciole

AP-HM, Marseille, FR

Modeling

*C. Nicolò



*D. Barbolosi

Thanks for listening!

That's all Folks!

We are looking for postdocs!

sebastien.benzekry@inria.fr