



HAL
open science

Regular factors in nearly regular graphs

Jean-Claude Bermond, Michel Las Vergnas

► **To cite this version:**

Jean-Claude Bermond, Michel Las Vergnas. Regular factors in nearly regular graphs. *Discrete Mathematics*, 1984, 50, pp.9-13. 10.1016/0012-365X(84)90030-X . hal-02426790

HAL Id: hal-02426790

<https://inria.hal.science/hal-02426790>

Submitted on 2 Jan 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

REGULAR FACTORS IN NEARLY REGULAR GRAPHS

Jean-Claude BERMOND*

Université Paris-Sud, Informatique, E.R.A. 452, Bâtiment 490, 91405 Orsay, France

Michel LAS VERGNAS*

Université Pierre et Marie Curie, U.E.R. 48—Mathématiques, 4 place Jussieu, 75005 Paris, France

We strengthen classical theorems of Petersen, Ore, Båbler, Gallai on the existence of regular factors in regular graphs to nearly regular graphs.

1. Introduction

An h -factor of a graph G , h an integer, is a spanning subgraph which is regular of degree h . More generally, given an integer-valued function f defined on $V(G)$, an f -factor is a subgraph H of G such that $d_H(x) = f(x)$ for all $x \in V(G)$. Several important theorems of the theory of graph factors give simple sufficient conditions for the existence of h -factors in regular graphs [1, 2, 5].

Theorem (Petersen [5]). *A regular graph of even degree k can be decomposed into $\frac{1}{2}k$ 2-factors.*

Theorem (Ore [4, Theorem 3.2.5]). *Let G be a 2-edge-connected regular graph of odd degree k and h be an even integer $2 \leq h \leq 2k/3$. Then G contains an h -factor.*

Theorem (k odd: Båbler [1]; k even: Gallai [2]). *Let G be a connected regular graph of degree k , with an even number of vertices. Let h be an odd integer, $h \leq \frac{1}{2}k$. If for all $X \subseteq V$, $|X|$ odd we have $e(X, V - X) \geq k/h$, then G has an h -factor.*

In the present note we give short proofs of these theorems by deriving them from Tutte's f -Factor Theorem [7]. We obtain slight generalizations: Due to integrality conditions these existence theorems remain valid for graphs which are almost regular (in a sense to be made precise below).

For completeness we recall Tutte's f -Factor Theorem [7].

* C.N.R.S.

Theorem. A graph has an f -factor if and only if

$$f(X) + d_G(Y) - f(Y) \geq e_G(X, Y) + \left(\begin{array}{l} \text{number of connected components} \\ \text{C of } G[V \setminus (X \cup Y)] \text{ such that} \\ e_G(C, Y) \not\equiv f(C) \pmod{2} \end{array} \right)$$

for all $X, Y \subseteq V(G)$ with $X \cap Y = \emptyset$.

The reader is referred to [3] for a short proof of the 1-Factor theorem [6], and to [8] for a short proof of the f -Factor Theorem from the 1-Factor theorem. Another necessary and sufficient condition for the existence of f -factors is given by Ore [4].

2. Notations

Graphs considered in the present note are finite, undirected, and may have loops and multiple edges (a loop incident to a vertex x contributes 2 to the degree of x).

Throughout the paper the following notations are used: Given a graph G the vertex-set of G is denoted by $V(G)$. The degree in G of a vertex x is $d_G(x)$. Given $X \subseteq V(G)$, we set $d_G(X) = \sum_{x \in X} d_G(x)$. The subgraph of G induced by X is denoted by $G[X]$, the number of edges of $G[X]$ by $e_G(X)$. Given $X, Y \subseteq V(G)$ with $X \cap Y = \emptyset$ we denote by $e_G(X, Y)$ the number of edges of G joining X and Y .

If no confusion results subscripts G will usually be omitted.

3. The theorems

Theorem 1 (even factors). Let G be a graph and h, k be two integers such that h is even and $2 \leq h \leq \frac{2}{3}k$. Suppose that $\sum_{x \in V(G)} |d_G(x) - k| < k/h$ and G is 2-edge connected. Then G contains an h -factor.

Proof. Tutte's condition for the existence of an h -factor, h even, is:

$$h |X| + d(Y) - f |Y| \geq e(X, Y) + \left(\begin{array}{l} \text{number of connected components } C \\ \text{of } G[V \setminus (X \cup Y)] \text{ such that} \\ e(C, Y) \text{ is odd} \end{array} \right)$$

for all $X, Y \subseteq V$ with $X \cap Y = \emptyset$.

The condition is clearly satisfied for $X = Y = \emptyset$. Let $X, Y \subseteq V$ with $X \cap Y = \emptyset$ and $X \cup Y \neq \emptyset$. We distinguish two types of connected components C of

$G[V \setminus (X \cup Y)]$ such that $e(C, Y)$ is odd:

(1) $e(C, X) = 0$. Since G is 2-edge-connected $e(C, Y) = e(C, V \setminus C) \geq 2$. Hence, $e(C, Y)$ being odd, we have $e(C, Y) \geq 3$.

(2) $e(C, X) \geq 1$. Since $e(C, Y)$ is odd, we have $e(C, Y) \geq 1$.

Let p_1, p_2 be the numbers of components of each type. We have clearly

$$p_2 + e(X, Y) \leq e(X, V \setminus X) \leq d(X),$$

$$3p_1 + p_2 + e(X, Y) \leq e(Y, V \setminus Y) \leq d(Y).$$

Hence the inequalities

$$p_2 + e(X, Y) \leq d(X),$$

$$3p_1 + p_2 + e(X, Y) \leq d(Y).$$

We multiply the first inequality by h/k , the second one by $(k-h)/k$ and add. We obtain

$$3 \frac{k-h}{k} p_1 + p_2 + e(X, Y) \leq \frac{h}{k} d(X) + \frac{k-h}{k} d(Y).$$

Since $3(k-h)/k \geq 1$ if $h \leq \frac{2}{3}k$ it follows, with $p = p_1 + p_2$, that

$$p + e(X, Y) \leq \frac{h}{k} d(X) + \frac{k-h}{k} d(Y).$$

Suppose

$$h |X| + d(Y) - h |Y| < p + e(X, Y).$$

Since both sides are integers we have

$$h |X| + d(Y) - h |Y| + 1 \leq p + e(X, Y) \leq \frac{h}{k} d(X) + \frac{k-h}{k} d(Y).$$

It follows that

$$d(X) - k |X| - (d(Y) - k |Y|) \geq \frac{k}{h}.$$

Since $X \cap Y = \emptyset$ we have

$$d(X) - k |X| - (d(Y) - k |Y|) \leq \sum_{x \in X \cup Y} |d(x) - k| < \frac{k}{h},$$

a contradiction. Hence Tutte's condition is satisfied. Theorem 1 follows. \square

The theorems of Petersen and Ore quoted in the introduction correspond to the particular cases of Theorem 1 when G is regular. To obtain Petersen's theorem it suffices to observe that the connected components of a regular graph of even degree are necessarily 2-edge-connected. (Proof: for $X \subseteq V$ we have $k |X| = d(X) = 2e(X) + e(X, V \setminus X)$, hence if k is even $e(X, V \setminus X)$ is also even.)

Theorem 2 (odd factors). Let G be a connected graph with an even number of vertices, and h, k be two integers such that h is odd and $1 \leq h \leq \frac{1}{2}k$. Suppose that $\sum_{x \in V(G)} |d(x) - k| < k/h$ and for all $X \subseteq V$ with $e(X, V - X)$ odd or $|X|$ odd we have $e(X, V - X) \geq k/h$. Then G contains an h -factor.

Note. The condition on $e(X, V - X)$ is obviously satisfied if the edge connectivity of G is at least k/h .

Proof. Tutte's condition for the existence of an h -factor, h odd, is:

$$h |X| + d(Y) - h |Y| \geq e(X, Y) + \left(\begin{array}{l} \text{number of connected components } C \\ \text{of } G[V \setminus (X \cup Y)] \text{ such that} \\ e(C, Y) \not\equiv |C| \pmod{2} \end{array} \right)$$

for all $X, Y \subseteq V$ with $X \cap Y = \emptyset$.

The condition is clearly satisfied for $X = Y = \emptyset$. Let $X, Y \subseteq V$ with $X \cap Y = \emptyset$ and $X \cup Y \neq \emptyset$. We distinguish four types of connected components C of $G[V \setminus (X \cup Y)]$ such that $e(C, Y) \not\equiv |C| \pmod{2}$.

(1) $|C|$ even, $e(C, X) = 0$. Since $e(C, Y)$ is odd we have $e(C, Y) = e(C, V \setminus C) \geq k/h$.

(2) $|C|$ even, $e(C, X) \geq 1$. Since $e(C, Y)$ is odd we have $e(C, Y) \geq 1$.

(3) $|C|$ odd, $e(C, Y) = 0$. We have $e(C, X) = e(C, V \setminus C) \geq k/h$ by hypothesis.

(4) $|C|$ odd, $e(C, Y) \geq 1$. Since $e(C, Y)$ is even we have $e(C, Y) \geq 2$.

Let p_1, p_2, p_3, p_4 be the number of components of each type. We have clearly

$$p_2 + \frac{k}{h} p_3 + e(X, Y) \leq e(X, V \setminus X) \leq d(X),$$

$$\frac{k}{h} p_1 + p_2 + 2p_4 + e(X, Y) \leq e(Y, V \setminus Y) \leq d(Y).$$

Hence the inequalities

$$p_2 + \frac{k}{h} p_3 + e(X, Y) \leq d(X),$$

$$\frac{k}{h} p_1 + p_2 + 2p_4 + e(X, Y) \leq d(Y).$$

We multiply the first inequality by h/k , the second one by $(k-h)/k$, and add. We obtain

$$\frac{k-h}{h} p_1 + p_2 + p_3 + 2 \frac{k-h}{k} p_4 + e(X, Y) \leq \frac{h}{k} d(X) + \frac{k-h}{k} d(Y).$$

Since $(k-h)/h \geq 1$ and $2(k-h)/k \geq 1$ if $h \leq \frac{1}{2}k$, it follows with $p = p_1 + p_2 + p_3 + p_4$, that

$$p + e(X, Y) \leq \frac{h}{k} d(X) + \frac{k-h}{k} d(Y).$$

As in the proof of Theorem 1 this inequality implies that Tutte's condition is satisfied. \square

The theorem of Bähler and Gallai corresponds to the particular cases of Theorem 2 when G is regular. If G is regular of degree k we have $k|X| = d(X) = 2e(X) + e(X, V - X)$. Hence if k is odd, $e(X, V - X)$ is odd if and only if $|X|$ is odd and if k is even, $e(X, V - X)$ is always even. Therefore the hypothesis of Theorem 2 reduces to: for all $X \subseteq V$, $|X|$ odd implies $e(X, V - X) \geq k/h$.

Finally, we note that the bounds on h and $\sum_{x \in V} |d(x) - k|$ in Theorems 1 and 2 are probably tight. (We leave as an exercise to the reader the construction of effective examples).

References

- [1] F. Bähler, Über die Zerlegung regulärer Streckenkomplexe ungerader Ordnung, *Comment. Math. Helvet.* 10 (1938) 275–287.
- [2] T. Gallai, On factorisation of graphs, *Acta Math. Acad. Sci. Hungar.* 1 (1950) 133–153.
- [3] L. Lovász, Three short proofs in graph theory, *J. Combinatorial Theory (B)* 19 (1975) 269–271.
- [4] O. Ore, Graphs and subgraphs, *Trans. Amer. Math. Soc.* 84 (1957) 109–136.
- [5] J. Petersen, Die Theorie der regulären Graphen, *Acta Math.*, 15 (1891) 193–220.
- [6] W.T. Tutte, The factorization of linear graphs, *J. London Math. Soc.* 22 (1947) 107–111.
- [7] W.T. Tutte, The factors of graphs, *Canadian J. Math.* 1 (1950) 44–49.
- [8] W.T. Tutte, A short proof of the factor theorem for finite graphs, *Canadian J. Math.* 6 (1954) 347–352.