



**HAL**  
open science

# Indetermination-free cytoarchitecture measurements in brain gray matter via a forward diffusion MRI signal separation method

Maëliss Jallais, Demian Wassermann

► **To cite this version:**

Maëliss Jallais, Demian Wassermann. Indetermination-free cytoarchitecture measurements in brain gray matter via a forward diffusion MRI signal separation method. ISMRM 2020 - 28th Annual Meeting & Exhibition, Aug 2020, Sydney / Virtual, Australia. hal-02477461

**HAL Id: hal-02477461**

**<https://inria.hal.science/hal-02477461v1>**

Submitted on 13 Feb 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Indetermination-free cytoarchitecture measurements in brain gray matter via a forward diffusion MRI signal separation method

Maëlliss Jallais<sup>1</sup> and Demian Wassermann<sup>1</sup>

<sup>1</sup>Université Paris-Saclay, Inria, CEA, Palaiseau, France

## Synopsis

Non-invasive imaging at the cellular level could lead us to quantify grey matter tissue cytoarchitecture, which has so far been accessible only through histology, or by indeterminate-prone approaches.

We propose a new dMRI-based index to study modulated by the size of the somas. This index can be extracted without indeterminations from any acquisition including three b-values superior to 3 ms/μm<sup>2</sup>. Simulations were experimentally confirmed by tests on the HCP MGH data set.

## Introduction

In this work we solve the problem of estimating soma sizes in grey matter with no indetermination. To extract a soma size-modulated mesurement, we combine approximations to a biophysical model at low b-values and its associated return-to-the-origin probability (RTOP)<sup>1</sup> for high b-values.

Such non-invasive measurement of cellular characteristics has the potential to quantify tissue cytoarchitecture, in a unique-solution system, which has so far been accessible only through histology.

Our contribution focuses on the human brain gray matter, which can be decomposed into three compartments, somas and processes that are modeled by spheres and 0-radius tubes, and extra-cellular water. This work shows the feasibility to extract factor that reflects the averaged diameter of the somas in the voxel with a unique solution. Our three-compartment model, under the hypothesis that there is no exchange between the three compartments<sup>2</sup>, is as follow :

$$\frac{S(q)}{S(0)} = f_t S_{tubes}(q, D_a) + f_s S_{spheres}(q, D_s, r_s) + (1 - f_t - f_s) S_{ecs}(q, D_{ecs})$$

$f_t$  and  $f_s$  represent the MR dendrite and soma signal fraction respectively,  $D_{ecs}$  and  $D_s$  the extra-cellular and soma diffusivities,  $D_a$  axial diffusivity and  $r_s$  the soma radius.

The tube signal is modeled by a power-law scaling<sup>3</sup>  $\frac{1}{4\sqrt{\pi\tau D_a}} \cdot q^{-1} + \gamma$ , and the sphere signal by Balinov's approximation<sup>4</sup>

$-\log S_{spheres}(q) = C(r_s, D_s) \cdot q^2$ . Here we exploit this relation to extract  $C = C(r_s, D_s)$  that, at fixed  $D_s$ , **modulated by soma size** with a method that can be applied to acquisitions with just 5 b-values. Three of them must be superior to a minimum b-value defined by cell soma size, see Fig2. The spacing of these points is determined by the SNR as the signal difference between points must not be noise-dominated.

Others<sup>5</sup> have already propose solutions to dMRI soma measurement for high b-values, and more b-values in all, but their method leads to indeterminate systems. Our direct method is indetermination free.

## Methods

By combining the expression to our biophysical model (1) for RTOP at large q (respectively b) and a Lemonade variant at low b (respectively q), we produce a method to extract the parameters of our model.

**RTOP: high b.** We compute a q-bounded RTOP, a direct measure of the restrictions of the diffusing fluid molecule motion that gives us information about the structure of the media<sup>6,7</sup>:

$$RTOP(q_{max}) = \frac{1}{(2\pi)^3} \int_0^{q_{max}} \frac{S(q)}{S(0)} dq_{max}$$

For  $q_{max}$  large enough, RTOP of the sphere signal converges towards a value that depends on C. We therefore obtain from eqs (1) and (2) assuming a large enough  $q_{max}$  :

$$RTOP(q_{max}) = \frac{(1 - f_t)}{(2\pi)^2 \cdot 8\sqrt{\pi} \cdot 4C^{3/2}} + \frac{f_t}{4(2\pi)^3 \sqrt{\pi\tau D_a}} \cdot q_{max}^2 + \frac{\gamma}{3(2\pi)^3} \cdot q_{max}^3$$

By solving an ordinary least square regression we can find the coefficients  $a_{fit}$ ,  $b_{fit}$  and  $c_{fit}$  of the previous polynomial and we end up with 2 equations with 3 unknowns.

**Spiked LEMONADE: low b.** We adapted the method proposed in <sup>8</sup> in adding the sphere contribution ( $f_s$  and C). Estimating the rotational invariant moment tensors  $M^{(2),0}$ ,  $M^{(2),2}$ , and  $M^{(4),2}$  for a b-value up to 2.5ms/μm<sup>2</sup> we can find 4 more equations with 6 unknowns.

**Total system.** We end up with a system of 6 equations for 6 unknowns that has no degeneracy.

Spiked LEMONADE, low b :

$$\begin{cases} M^{(2),0} = f_t D_a + 3f_s C + 3(1 - f_t - f_s) D_e \\ \frac{M^{(2),2}}{p_2} = f_t D_a \\ M^{(4),0} = f_t D_a^2 + 5f_s C^2 + 5(1 - f_t - f_s) D_e^2 \\ \frac{M^{(4),2}}{p_2} = f_t D_a^2 \end{cases}$$

RTOP, high b:

$$\begin{cases} a_{fit} = \frac{(1-f_t)\sqrt{\pi}}{(2\pi)^3 \cdot 4C^{3/2}} \\ b_{fit} = \frac{f_t\sqrt{\pi}}{2(2\pi)^4 \sqrt{\tau D_a^4}} \end{cases}$$

**Numerical simulations.** To mimic neurons, we modelled the sphere and tube signals using the DMIPY simulator<sup>7</sup> with soma radius equal to approximately 20 and 15 $\mu$ m, reproducing Von Economo and pyramidal neurons.

**Experimental Data.** We used the HCP MGH Adult Diffusion data set to study the variation of C on the insula, precentral and postcentral parts of the Destrieux atlas. It is composed of 35 subjects, with  $\delta/\Delta=10.6/43.1$ ms and  $b=1, 3, 5, 10$ ms/ $\mu$ m<sup>2</sup>.

## Results

**Simulations.** RTOP computation on soma simulations shows the convergence values specific to soma diameters of different neurons in the brain, obtained from Neuromorpho.org (**Fig.2a**). This value is proportional to C, and **Fig2b** shows that C is modulated by the size of the somas, and inversely proportional to it. The relative error of C with respect to the minimum b-value used for the RTOP computation is presented in **Fig2c**.

**Experiments.** **Fig3**, C value for three cortical regions on subject 1010 (right handed), preliminary evidence shows: larger soma size spread and asymmetry on the insula, agreeing with<sup>9</sup>. and larger somas on the left postcentral, possibly hemispheric dominance<sup>10</sup>. **Fig 4**. Soma-to-dendrite volume ratio showing soma predominance agreeing with literature<sup>10</sup>. Although further work is needed, this showcases our method's potential.

## Discussion and Conclusion

The minimal b-value needed to compute the large approximation of the signal RTOP must be at least 3ms/ $\mu$ m<sup>2</sup> to get a good C approximation in the case of VEN and pyramidal neurons. The proposed method does not need a large range of b-values as in<sup>5</sup>.

## Acknowledgements

This work was supported by the ERC-StG NeuroLang and the ANR/NSF NeuroRef grants.

## References


1. Partha P. Mitra and Pabitra N. Sen. Effects of microgeometry and surface relaxation on NMR pulsed-field-gradient experiments: Simple pore geometries. *Physical Review B*, 453(1):143–156, January 1992. ISSN 0163-1829, 1095-3795. doi: 10.1103/PhysRevB.45.143. URL <https://link.aps.org/doi/10.1103/PhysRevB.45.143>.
2. Palombo M, Shemesh N, Ianus A, et al., Abundance of cell bodies can explain the stickmodel's failure to describe high b-value diffusion signal in grey matter. *Proc. Int. Soc. Magn. Reson. Med.* 2018, 1096
3. Jelle Veraart, Els Fieremans, and Dmitry S. Novikov. On the scaling behavior of water diffusion in human brain white matter. *NeuroImage*, 185:379–387, January 2019. ISSN 10538119. doi: 10.1016/j.neuroimage.2018.09.075. URL <https://linkinghub.elsevier.com/retrieve/pii/S1053811918319475>
4. Balinov et al. The NMR Self-Diffusion Method Applied to Restricted Diffusion. Simulation of Echo Attenuation from Molecules in Spheres and between Planes. 1993
5. Marco Palombo, Clemence Ligneul, and Julien Valette. Modeling diffusion of intracellular metabolites in the mouse brain up to very high diffusion-weighting: Diffusion in long fibers (almost) accounts for non-monoexponential attenuation: Modeling Diffusion of Brain Metabolites in Vivo up to Very High Diffusion Weighting. *Magnetic Resonance in Medicine*, 77(1):343–350, January 2017. ISSN 07403194. doi: 10.1002/mrm.26548. URL <http://doi.wiley.com/10.1002/mrm.26548>
6. Rutger H.J. Fick, Demian Wassermann, Emmanuel Caruyer, and Rachid De-riche. MAPL: Tissue microstructure estimation using Laplacian-regularized MAP-MRI and its application to HCP data. *NeuroImage*, 134:365–385, July 2016. ISSN 10538119. doi: 10.1016/j.neuroimage.2016.03.046. URL <https://linkinghub.elsevier.com/retrieve/pii/S1053811916002512>
7. Mitra et al. Pulsed-field-gradient NMR measurements of restricted diffusion and the return-to-origin probability. *Journal of Magnetic Resonance*, 114:47–58, 1995
8. Dmitry S. Novikov, Jelle Veraart, Ileana O. Jelescu, and Els Fieremans. Rotationally-invariant mapping of scalar and orientational metrics

of neuronal microstructure with diffusion MRI. *NeuroImage*, 174:518–538, July 2018. ISSN 10538119. doi: 10.1016/j.neuroimage.2018.03.006. URL <https://linkinghub.elsevier.com/retrieve/pii/S1053811918301915>


9. Evrard, H. C., Forro, T. & Logothetis, N. K. Von Economo Neurons in the Anterior Insula of the Macaque Monkey. *Neuron* 74, 482–489 (2012)

10. Katrin Amunts and Karl Zilles. Architectonic Mapping of the Human Brain beyond Brodmann. *Neuron*, 88(6):1086–1107, December 2015. ISSN 08966273. doi: 10.1016/j.neuron.2015.12.001. URL <https://linkinghub.elsevier.com/retrieve/pii/S0896627315010727>


## Figures



**Figure 1:** Visual abstract. On the top left we illustrate a multi-shell dMRI. On the one hand we are using gradient directions at low b-values to estimate rotational invariants moment tensors. On the other hand we are computing  $RTOP(q_{max})$  for high q-values. We end up with a determinate system and get  $C$ , a measurement that is modulated by soma size.




**Figure 2:** (a)  $RTOP$  of the soma signal depends on the sphere diameter. The minimum b-value used for the large b approximation is related to the diameter of the somas acquired. (b)  $C$  is modulated by the sphere diameter. (c) Relative error of  $C$  estimation. The bigger the soma, the lower b-values for a better relative error. The minimum b-value used for the large b approximation is lower for VEN than pyramidal neurons.



**Figure 3:**  $C$  value for three cortical regions on subject 1010 (right handed), Preliminary evidence shows: larger soma size spread and asymmetry on the insula, agreeing with <sup>9</sup>. ; and larger somas on the left postcentral, possibly hemispheric dominance <sup>10</sup>.





**Figure 4:** Soma-to-dendrite volume ratio showing soma predominance agreeing with literature<sup>9</sup>. Although further work is needed, this showcases our method's potential.