HAL open science

Optimization of chemical batch reactors using temperature control

Jérémy Rouot, Bernard Bonnard

To cite this version:

Jérémy Rouot, Bernard Bonnard. Optimization of chemical batch reactors using temperature control. 9th International Congress on Industrial and Applied Mathematics, Jul 2019, Valencia, Spain. hal-02483219

HAL Id: hal-02483219

https://inria.hal.science/hal-02483219v1

Submitted on 18 Feb 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Optimization of chemical batch reactors using temperature control

Jérémy Rouot

EPF, France

joint work with B. Bonnard (INRIA & UBFC)

Chemical Networks with mass action kinetics

Graph Model :

Species $\{X_1, \ldots, X_m\}$.

Notations : $\mathscr{R}$ is the set of reactions of the form :

$$\sum_{i=1}^{m} \alpha_i X_i \longrightarrow \sum_{i=1}^{m} \beta_i X_i$$

α_i, β_i are stochiometric coefficients.

Feinberg-Horn-Jackson graph

- Vertices : $\mathbf{y} = (\alpha_1, \ldots, \alpha_m)^\top$, $\mathbf{y}' = (\beta_1, \ldots, \beta_m)^\top$
- Orientation : $\mathbf{y} \to \mathbf{y}'$

$$\begin{tikzcd} A \arrow[r, shift left] & B \arrow[l, shift left] \arrow[r] & C \arrow[ll, bend left] \end{tikzcd} \tag{1}$$

Rate dynamics $\mathbf{y} \to \mathbf{y}'$ (Mass kinetics)

$$K(\mathbf{y} \to \mathbf{y}') = k(T)\, c^{\mathbf{y}}$$

- $k(T) = A\exp(-\frac{E}{RT})$: Arrhenius law
  E, A parameters, T temperature and R is the gas constant
- $c = (\mathbf{c_1}, \ldots, c_m)^\top$
  c_i : concentrations of the species X_i with

$$c^{\mathbf{y}} = \mathbf{c_1}^{\alpha_1} \ldots c_m^{\alpha_m}$$

$\Rightarrow K(\mathbf{y} \to \mathbf{y}')$ **depends only on $\mathbf{y}$.**

Dynamics for the network

$$\dot{\mathbf{c}}(\mathbf{t}) = F(\mathbf{c}(\mathbf{t}), T) = \sum_{\mathbf{y}\to\mathbf{y}'\in\mathscr{R}} K(\mathbf{y}\to\mathbf{y}')\,(\mathbf{y}'-\mathbf{y})$$

The dynamics is defined by the graph.
More explicit representation of the dynamics.

- **Stochiometric subspace**

$$S = \textbf{span}\left\{\mathbf{y}'-\mathbf{y},\ \mathbf{y}\to\mathbf{y}' \in \mathscr{R}\right\}$$

- **Positive class** (strict if > 0)

$$(\mathbf{c}(\mathbf{0}) + S) \cap \mathbb{R}^m_{\geq 0}$$

Lemma

The class $(\mathbf{c}(\mathbf{0}) + S) \cap \mathbb{R}^m_{>0}$ *is* **invariant** *for the dynamics.*

Notations

- **Complex matrix :** $Y = (y_1, \ldots, y_n)$ (n : number of complexes).
- **Incidence connectivity matrix :** $\boldsymbol{A = (a_{ij})_{ij}}$
  with for instance $a_{21} = k_1$ indicating a reaction with constant k_1 from the first node of the graph to the second.
- **Laplacian matrix :**

$$\tilde{A} = A - \text{diag}\left(\sum_{i=1}^{n} a_{i1}, \ldots, \ldots, \sum_{i=1}^{n} a_{in}\right)$$

One has

$$\dot{\mathbf{c}}(\mathbf{t}) = f(\mathbf{c}(\mathbf{t}), T) = Y\,\tilde{A}\,c^{Y}$$

where $c^{Y} = (c^{y_1}, \ldots, c^{y_n})^{\top}$.

Zero deficiency theorem

Definition (Deficiency)

Feinberg and Horn-Jackson : articles in Archive Rational Mechanics

Graph concept : *deficiency :* $\boldsymbol{\delta = n - l - s}$ *where*

- n *: number of vertices*
- l *: number of connecting components*
- s *: dimension of the stochiometric subspace*

Definition

The network is **weakly reversible** *if* $\forall$ *vertices* (i, j) *such that* $\exists$ *oriented path joining* i *to* j*, there exists an oriented path joining* j *to* i*.*

Assumption $\boldsymbol{\delta = 0}$ *(Zero deficiency assumption)*

Theorem

1. *If the network is* **not weakly reversible** *then for arbitrary kinetics, the differential equation* **cannot have a positive equilibrium nor a positive periodic trajectory.**
2. *If the network is* **weakly reversible**, *there exists within each strictly positive compatibility class precisely* **one equilibrium** c^*, *this equilibrium is locally asymptotically stable with (pseudo-Helmholtz) Lyapunov function* $V(c, c^*) = \sum_i \left[c_i(\ln(c_i) - \ln(c_i^*) - 1) + c_i^* \right]$.
   Moreover there is non trivial periodic orbit.

Application : Test bed cases :

case 1 : $A \xrightarrow{k_1} B \xrightarrow{k_2} C$ $\delta = 3 - 1 - 2 = 0$: not weakly reversible

case 2 : (**McKeithan network)**

$$\begin{tikzcd} A \arrow[r, bend left, "k_1"] \arrow[rrr, bend right, "k_4"'] & B \arrow[l, bend left, "k_3"] \arrow[r, "k_2"] & C \end{tikzcd}$$

$\delta = 3 - 1 - 2 = 0$: one single equilibirum globally asymptotically stable

Equilibrium for the McKeithan network



FIGURE – Phase portrait for the McKeithan model. (left) Focus ; (right) Node.

Geometric Optimal Control

Optimal Control Problem

$$\frac{\mathrm{d}\mathbf{c}}{\mathrm{d}t} = f(\mathbf{c}, T), \quad \frac{\mathrm{d}T}{\mathrm{d}t} = u, \quad u \in [u_-, u_+]$$

$u(\cdot)$ tracked the derivative of the temperature (related to the Goh Transformation).

Single input C^ω-control system :

$$\begin{cases} \dot{\mathbf{q}} = F(\mathbf{q}) + u\,G(\mathbf{q}), \quad |u| \leq 1, \\ \mathbf{q} = (\mathbf{c}, T) \in \mathbb{R}^n \end{cases}$$

Formulation :

$$\max \mathbf{c_1}(t_f) \quad t_f : \text{ time batch duration}$$

Formulated as

$$\begin{cases} \min\ t_f, & |u| \leq 1 \\ \mathbf{c_1}(t_f) = d \text{ is a desired quantity} \end{cases}$$

$\mathbf{N}$: terminal manifold of codimension 1.



Necessary optimality conditions Pontryagin Maximum Principle (1956)

Statement :

$$\begin{cases} \dot{\mathbf{q}} = F(\mathbf{q}) + u\,G(\mathbf{q}), \quad |u| \leq 1, \\ \min\ t_f, \quad \mathbf{q}(t_f) \in \mathbf{N} \end{cases}$$

- $H(\mathbf{q}, p, u) = p \cdot (F(\mathbf{q}) + u\,G(\mathbf{q}))$, $p \in \mathbb{R}^n \setminus \{0\}$: adjoint vector
- H : pseudo-Hamiltonian and the maximized Hamiltonian is

$$M(\mathbf{q}, p) = \max_{|u| \leq 1} H(\mathbf{q}, p, u), \qquad \mathbf{q}, p \text{ are fixed}$$

Theorem

Assume $(q^*(\cdot), p^*(\cdot))$ *is a time minimal solution on* $[0, t_f^*]$ *then there exists* $p^*(\cdot)$ *such that a.e. on* $[0, t_f^*]$ *:*

$$\dot{q}^*(\cdot) = \frac{\partial H}{\partial p}(q^*(t), p^*(t), u^*(t)), \quad \dot{p}^*(\cdot) = -\frac{\partial H}{\partial q}(q^*(t), p^*(t), u^*(t)) \tag{2}$$

the maximization condition is satisfied

$$H(q^*(t), p^*(t), u^*(t)) = M(q^*(t), p^*(t)).$$

Moreover

- $t \mapsto M(q^*(t), p^*(t))$ is constant and ≥ 0 and at the final time one has the *transversality condition* :

$$p^*(t_f) \perp T_{q^*(t_f)}\mathbf{N} \tag{3}$$

Extremals are solutions of (2). BC-extremal : transversality condition (4) satisfied.

Maximization condition

- *regular :*

$$u(t) = \text{sign}\, p(t) \cdot G(\mathbf{q}(\mathbf{t})) \text{ a.e.}$$

  Finite number of switches : **Bang-Bang**

- *singular :*

$$p(t) \cdot G(\mathbf{q}(\mathbf{t})) = 0 \quad \forall\, t$$

Exceptional extremals : $M = 0$.

Moreover

- $t \mapsto M(q^*(t), p^*(t))$ is **constant** and $\geq \mathbf{0}$.
- *Transversality condition* :

$$p^*(t_f) \perp T_{q^*(t_f)}\mathbf{N} \tag{4}$$

Extremals are solutions of (2). BC-extremal : transversality condition (4) satisfied.

Maximization condition

- *regular :*

$$u(t) = \text{sign}\, p(t) \cdot G(\mathbf{q}(\mathbf{t})) \text{ a.e.}$$

  Finite number of switches : **Bang-Bang**
- *singular :*

$$p(t) \cdot G(\mathbf{q}(\mathbf{t})) = 0 \quad \forall\, t$$

Exceptional extremals : $M = 0$.

Computations of singular extremals and properties

Notation : X, Y : two vector fields on $\mathbb{R}^n$

Lie bracket :

$$[X, Y](q) = \frac{\partial X}{\partial q} Y(q) - \frac{\partial Y}{\partial q} X(q)$$

$z = (q, p)$ $H_X(z) = p \cdot X(q)$

Poisson bracket :

$$\{H_X, H_Y\}(z) = p \cdot [X, Y](q)$$

Computations $H_G(z) = p \cdot G(q) = 0$

Differentiating twice w.r.t. time gives the two equations

$$\frac{\mathrm{d}}{\mathrm{d}t} H_G(z) = \mathrm{d}H_G \cdot \dot{z} = \{H_G, H_F + u\, H_G\} = \mathbf{\{H_G, H_F\} = 0}$$

$$\mathbf{\{\{H_G, H_F\}, H_F\}(z) + u\,\{\{H_G, H_F\}, H_G\}(z) = 0}$$

Then if $\{\{H_G, H_F\}, H_G\}(z) \neq 0$ then we compute $\hat{u}$ and plug it in H to obtain the *true Hamiltonian*.

Generalized Legendre-Clebsch condition

$$\{\{H_G, H_F\}, H_F\}(z) \geq 0$$

$\Rightarrow$ necessary optimality condition (High Order Maximum Principle, Krener).

Strict Legendre-Clebsch condition

$$\{\{H_G, H_F\}, H_F\}(z) > 0$$

Classification of singular extremals

$M = H_F$: constant value

- $M = 0$: **Exceptional case**
- $M > 0$: $\{\{H_G, H_F\}, H_G\}(z) > 0$: **Hyperbolic case** (**fast**)
- $M > 0$: $\{\{H_G, H_F\}, H_G\}(z) < 0$: **Elliptic case** (**slow**)

Classification of regular extremals (Ekeland - IHES, Kupka - TAMS)
Denote :

- σ_+ : bang arc with $u = +1$
- σ_- : bang arc with $u = -1$
- σ_s : singular arc $u = u_s \in]-1, 1[$

$\sigma_1\sigma_2$ is the arc σ_1 followed by σ_2.

Switching surface :

- $\Sigma : \{p \cdot G(q) = 0\}$

$\Phi(t) = p(t) \cdot G(\mathbf{q(t)})$ is the **switching function**.
$\dot{\Phi}(t) = p(t) \cdot [G, F](\mathbf{q(t)})$
$\ddot{\Phi}(t) = p(t) \cdot \big([[G, F], F](\mathbf{q(t)}) + u(t)\,[[G, F], G](\mathbf{q(t)})\big)$

Ordinary Switching time : $t \in]0, t_f[$ such that $\Phi(t) = 0$ and $\dot{\Phi}(t) \neq 0$

Lemma

Near $z(t)$ every extremal solution projects onto $\sigma_+\sigma_-$ if $\dot{\Phi}(t) < 0$ and $\sigma_-\sigma_+$ if $\dot{\Phi}(t) > 0$

Fold case : If $\Phi(t) = \dot{\Phi}(t) = 0$ then $z(t) \in \Sigma'$

$\ddot{\Phi}_\varepsilon(z(t)) = p(t) \cdot \left([[G,F],F](\mathbf{q(t)}) + \varepsilon\,[[G,F],G](\mathbf{q(t)})\right), \quad \varepsilon = \pm 1$

Assumption : Σ' : surface of codimension two, $\ddot{\Phi}_\varepsilon(z(t)) \neq 0$ for $\varepsilon = \pm 1$.

$z(t)$: fold point

- Case 1 : **parabolic case** $\quad \ddot{\Phi}_+(t)\ddot{\Phi}_-(t) > 0$
- Case 2 : **hyperbolic case** $\quad \ddot{\Phi}_+(t) > 0$ and $\ddot{\Phi}_-(t) < 0$
- Case 3 : **elliptic case** $\quad \ddot{\Phi}_+(t) < 0$ and $\ddot{\Phi}_-(t) > 0$

u_s is the singular control defined by

$$p(t) \cdot \left([[G,F],F](\mathbf{q(t)}) + u_s(t)\,[[G,F],G](\mathbf{q(t)})\right) = 0$$



Fold case

In the parabolic case $|u_0| > 1$ and the singular arc is not admissible.

Theorem

Kupka TAMS In the neighborhood of $z(t)$ every extremals projects onto :

- *Parabolic case : $\sigma_+\sigma_-\sigma_+$ or $\sigma_-\sigma_+\sigma_-$*
- *Hyperbolic case : $\sigma_\pm\sigma_s\sigma_\pm$*
- *Elliptic case : every extremal is of the form $\sigma_+\sigma_-\sigma_+\sigma_-\ldots$ (Bang-Bang) but the number of switches is not uniformly bounded.*

Application to Chemical Networks

Time minimal synthesis for chemical systems

$$\begin{cases} \min\ t_f \qquad |u| \leq 1 \\ \dot{\mathbf{q}} = F(\mathbf{q}) + u\,G(\mathbf{q}) \\ \mathbf{c_1}(t_f) \in \mathbf{N} = \{\mathbf{c_1} = d\} \end{cases}$$

Methods : Two steps :

1. Calculation of the time minimal syntheses near the terminal manifold
2. Bounds on the number of switches



`Step 1`: Take $q_0 \in \mathbf{N}$, $z_0 = (q_0, n(q_0))$ where $n(q_0)$ is the normal vector of $\mathbf{N}$ at q_0.
Find, in a small neighborhood U of q_0, the time minimal closed loop control $u^*(q)$ to reach $\mathbf{N}$ starting from $\mathbf{q}$ in minimal time.

Computations : $\dot{\mathbf{q}} = F(\mathbf{q}) + u\,G(\mathbf{q})$, $q(t_f) \in \mathbf{N}$

Synthesis : $u^*(\mathbf{q})$ means

- determine the **switching locus**

  Ex. :



- determine the **splitting locus** or the **cut locus** C where two distinct optimal trajectories occur.

  Ex. :



Tools : Singularity theory $\mathbf{N} = \{f^{-1}(0)\}$

- *expand* at q_0 with Taylor series : jet spaces.
- *compute :* Normal form to estimate W, C near q_0. The tools are simple but the classification is complicated.

Tools : Singularity theory $\mathbf{N} = \{f^{-1}(0)\}$

- *expand* at q_0 with Taylor series : jet spaces.
- *compute :* Normal form to estimate W, C near q_0. The tools are simple but the classification is complicated.

Ex. : *Two reactions only*. $(C, T) \in \mathbb{R}^3$ $\dot{\mathbf{q}} = F + uG$ and $\mathbf{N} = f^{-1}(0)$.

Generic case $z_0 = (q_0, n(q_0))$.

***G* is tangent to** $\mathbf{N}$: Then $p \cdot G = 0$ so p is normal to $\mathbf{N}$.

Using classification of extremals at a point such that $p \cdot G(\mathbf{q}) = 0$, $p \cdot [G, F](\mathbf{q}) \neq 0$:



depending on the sign of $p \cdot [G, F](q_0)$.

... but there are more complicated situations

Define :

$\mathcal{S}$ the singular locus : $\{\mathbf{q} \in \mathbf{N};\ n \cdot [G, F](\mathbf{q}) = 0\}$

$\mathcal{E}$ the exceptional locus : $\{\mathbf{q} \in \mathbf{N};\ n \cdot F(\mathbf{q}) = 0\}$

For $\mathcal{S}$: from the classification near a fold point one has :

- *Hyperbolic* case
- **Elliptic** case
- *Parabolic* case

To make the analysis we construct a semi-normal form : $\mathbf{q} = (x, y, z)$ near 0

$$\begin{cases} \dot{x} = 1 + a(x)\, z^2 + 2b(x)\, yz + c(x)\, y^2 + \dots \\ \dot{y} = d(x)\, y + e(0) + \dots \\ \dot{z} = (u - \hat{u}(x)) + f(x)\, y + g(0)\, z + \dots \end{cases}$$

with

- $\mathbf{N}$ is identified to $x = 0$
- the singular arc is identified to $\sigma_s : t \to (t, 0, 0)$ with singular control $\hat{u}$.
- $a(0) < 0$ hyperbolic if $|\hat{u}| < 1$.
- $a(0) > 0$: elliptic if $|\hat{u}| < 1$.
- parabolic if $|\hat{u}| > 1$.

Synthesis : There exists a C^0-foliation by planes such that in each plane the synthesis is :

Case : Hyperbolic.



Note that the synthesis is $\sigma_+\sigma_s\sigma_-$ hence the temperature is not constant.

Case : Parabolic. For instance, a synthesis is



The McKeithan network



Stratification of the terminal manifold :



Bridge phenomenon

Local (planar) simplified model (inspired from the saturation problem in Magnetic Resonance Imaging) :

$$\min_{u(\cdot)} \; t_f \qquad \dot{q}(t) = F(q(t)) + u\,G(q(t)), \quad t \in [0, t_f]$$

where

$$q = (x, y), \quad F = (1 - x^2 y)\frac{\partial}{\partial y}, \quad G = -(y-1)\frac{\partial}{\partial x} + x\frac{\partial}{\partial y}.$$

Singular lines $\mathscr{L}$: $\det(G, [G, F])(q) = 0$

- **fast** if $D(q) \cdot D''(q) > 0$
- **slow** if $D(q) \cdot D''(q) < 0$

where $D(q) = \det(G(q), [G, F](q), [[G, F], G](q))$
and $D'(q) = \det(G(q), [G, F](q), [[G, F], F](q))$

Singular sets



The **Singular control** along $\mathscr{L}$ is

$$u_s(q) = -\frac{D'(q)}{D(q)}$$

and is not bounded.

Trajectories : bridge



Bridge connecting two switching points of the singular set.

Conclusion

General techniques to handle complicated networks :
Geometric approach : Find coordinates to analyze the syntheses
→ applicable to general networks

Details :
T. Bakir, B. Bonnard, J. Rouot *Geometric Optimal Control Techniques to Optimize the Production of Chemical Reactors using Temperature Control* (submitted 2019)
B. Bonnard, G. Launay, M. Pelletier,
Classification générique de synthèses temps minimales avec cible de codimension un et applications,
Annales de l'I.H.P. Analyse non linéaire **14** no.1 (1997) 55–102.

Even a simple network $A \to B \to C$ can give complex optimal solution : work in progress on the *McKeithan network*.