



HAL
open science

From point to field scale: How consistent are agro-ecosystem models in terms of changes in soil texture?

Kurt-Christian Kersebaum, Evelyn Wallor, Elsa Coucheney, Munir Hoffmann, Anuschka Neira, Elisabet Lewan

► To cite this version:

Kurt-Christian Kersebaum, Evelyn Wallor, Elsa Coucheney, Munir Hoffmann, Anuschka Neira, et al.. From point to field scale: How consistent are agro-ecosystem models in terms of changes in soil texture?. ICROP2020: Second International Crop Modelling Symposium , Feb 2020, Montpellier, France. hal-02950300

HAL Id: hal-02950300

<https://inria.hal.science/hal-02950300>

Submitted on 27 Sep 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

From point to field scale: How consistent are agro-ecosystem models in terms of changes in soil texture?

Kersebaum Kurt-Christian¹ (ckersebaum@zalf.de), Wallor Evelyn¹, Coucheney Elsa², Hoffmann Munir³, Neira Anuschka¹, Lewan Elisabet²

¹ *Research Platform Models and Simulation, Leibniz Centre for Agric Landsc Research, Muencheberg, Germany;* ² *Department of Soil and Environment, Swedish University of Agric Science, Uppsala, Sweden;* ³ *Research Area Landscape Synthesis, Leibniz Centre for Agric Landsc Research, Muencheberg, Germany*

Introduction

Field scale variability of soil texture contributes to the site-specific performance of water dynamics, nitrogen turnover and crop development in the soil-plant-atmosphere system. In agricultural practice, management recommendations that operate on the basis of spatially and temporally variable soil states are rarely considered, even though, having knowledge about it could help to avoid unwanted environmental impact and waste of resources (Fowler et al., 2013). Agro-ecosystem models are able to derive spatially and temporally high resolution state variables considering both, soil and crop processes. To assess selected models for future application in agricultural practice (e.g. precision farming), this study examines the consistency of yield simulation in relation to soil texture changes at the field scale.

Materials and Methods

The data set used in the present study results from intensive mapping and sampling activities at a 20 ha large agricultural field in north western Germany. Data is extensively described in Wallor et al. (2018) and includes three years of on-the-go measured yields and results of soil analysis at 60 sampling points arranged as a regular grid. The soil inventory of the field consists of seven homogeneous soil zones that show increasing clay contents from 5% to 45%. Winter wheat yields of the years 2000 and 2001 show a clear response to these changing soil conditions. We exemplarily tested the models APSIM (AS; Keating et al., 2003), COUP (CO; Jansson, 2012), and HERMES (HE; Kersebaum, 2007) regarding their consistency in yield simulation with respect to changes in clay content and hence, soil water contents. A four-year period is simulated including one spin-up year and one year for model calibration.

Results and Discussion

When comparing the mean deviation of simulated and observed yields with the mean deviation of simulated and observed soil water contents for each model, AS showed highly variable deviations in yield predictions although water deviations were only slightly biased (cf. Figure 1). Especially for sampling points that have less than 25% clay the response of yield formation to soil water is over-predicted. HE produced the most consistent results showing a relative small band of deviations for water and yields, independently from soil texture. CO's simulation was similarly consistent in this sense, but showed larger deviations for yield predictions. The overestimation of yields was accompanied by over-predicted water contents without showing a clear relation to soil texture.

Conclusions

In terms of the highly sufficient depiction of heterogeneous soil water contents, which mainly affect yield performance in this field, the assessed models performed mostly consistent. This suggests that the cause for the deviations between simulation and observation can be found in the model-specific structure and equations, as the influence of model inherent estimation of soil hydrological parameters was excluded before.

Acknowledgements

This work was conducted within the Project I4S which is part of the BonaRes programme funded by the German Federal Ministry of Education and Research (BMBF), Grant Number 031B0513I. It is an in-depth evaluation of a modeling exercise that was carried out within the framework of FACCE JPI MACSUR2. The contributing authors thank the National Ministries and the European Union for funding the projects (grant 0339740 and 031B0039C).

.../...




Figure 1. Model-based relation between the mean deviation of simulated and observed yields and the mean deviation of simulated and observed soil water contents (WC) grouped by soil texture class (SI2, SI3, SI4 = 5 to 17% clay; Ls3, Ls4 = 17 to 25% clay; Lt2, Lt3 = 25 to 45% clay); mean deviation was calculated based on all observations per sampling point of the entire simulation period

References:

1. Fowler D, Coyle M, Skiba U, Sutton MA, Cape JN, Reis S, et al. (2013) The global nitrogen cycle in the twenty-first century. *Philos T R Soc B*, 368:1-3.
2. Jansson PE, (2012) CoupModel: Model Use, Calibration, and Validation. *Trans. ASABE*, 55(4):1335-1344.
3. Keating BA, Carberry PS, Hammer GL, Probert ME, Robertson MJ, Holzworth D, et al. (2003) An overview of APSIM, a model designed for farming systems simulation. *Eur J Agron*, 18(3-4):267-288.
4. Kersebaum KC (2007) Modelling nitrogen dynamics in soil-crop systems with HERMES. *Nutr Cycl Agroecosys*, 77(1):39-52.
5. Wallor E, Kersebaum KC, Lorenz K, Gebbers R (2018) A comprehensive dataset demonstrating the within-field variability of soil properties and crop growth conditions in north western Germany. *ODJAR*, 5:1-10.