



HAL
open science

Left Atrial Appendage Electrical Isolation Reduces Atrial Fibrillation Recurrences

Ali Gharaviri, Simone Pezzuto, Mark Potse, Sander Verheule, Giulio Conte,
Rolf Krause, Ulrich Schotten, Angelo Auricchio

► **To cite this version:**

Ali Gharaviri, Simone Pezzuto, Mark Potse, Sander Verheule, Giulio Conte, et al.. Left Atrial Appendage Electrical Isolation Reduces Atrial Fibrillation Recurrences: a simulation study. *Circulation. Arrhythmia and electrophysiology*, 2020, 10.1161/CIRCEP.120.009230 . hal-03088377

HAL Id: hal-03088377

<https://inria.hal.science/hal-03088377>

Submitted on 26 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Left atrial appendage electrical isolation reduces atrial fibrillation recurrences: a**
2 **simulation study**

3
4 **Gharaviri:** Atrial fibrillation recurrences after catheter ablation

5 Ali Gharaviri, PhD^a, Simone Pezzuto^a, PhD, Mark Potse, PhD^{c,d}, , Sander Verheule, PhD^e,
6 Giulio Conte, MD, PhD^{a,f}, Rolf Krause, PhD^a, Ulrich Schotten, MD, PhD^e, and
7 Angelo Auricchio, MD, PhD^{a,f}.

8
9 a: Center for Computational Medicine in Cardiology, Institute of Computational Science,
10 Università della Svizzera italiana, Lugano, Switzerland.

11 b: Carmen team, Inria Bordeaux Sud-Ouest, Talence, France.

12 c: Université de Bordeaux, IMB, UMR 5251, F-33400 Talence, France.

13 d: IHU Liryc, Electrophysiology and Heart Modeling Institute, foundation Bordeaux
14 Université, Bordeaux, France.

15 e: Department of Physiology, Maastricht University, Maastricht, The Netherlands.

16 f: Fondazione Cardiocentro Ticino, Lugano, Switzerland.
17

18 **Corresponding author:**

19 Ulrich Schotten, MD, PhD
20 Department of Physiology, Maastricht University
21 PO Box 616, 6200 MD Maastricht
22 The Netherlands
23 Phone: +31 43 3881077
24 Fax: +31 43 3884166
25 Email: Schotten@maastrichtuniversity.nl
26

27 **Journal Subject Terms:** Arrhythmia, Atrial fibrillation, Catheter ablation

28 **Keywords:** Fibrosis, Pulmonary vein isolation, Atrial Fibrillation recurrence, Left atrial
29 appendage electrical isolation, in-silico study.

30 **Non-standard Abbreviations and Acronyms:**

31 PV = Pulmonary vein

32 PVI = Pulmonary vein isolation

33 AF = atrial fibrillation

34 (persAF) = persistent atrial fibrillation patients

35 LAA = left atrial appendage

36 LAAI = left atrial appendage isolation

37 CA = catheter ablation

38

39 Pulmonary vein (PV) isolation (PVI) improves freedom from atrial tachyarrhythmia
40 recurrences in patients with paroxysmal and persistent atrial fibrillation (AF).¹ However, PVI
41 has a limited success rate in persistent AF patients (persAF) and ablation techniques
42 adjunctive to PVI isolating non-PV triggers remains an area of debate in these patients.¹
43 Recently, there has been a growing interest in the electrical and anatomical role of the left
44 atrial appendage (LAA) in triggering and sustaining AF, particularly in persAF or after
45 repeated AF ablation procedures.² Whether isolation of triggers originating from the LAA or
46 substrate modification caused by LAA isolation (LAAI) is the underlying mechanism for AF
47 recurrence prevention is still unknown.

48 We investigated the effect of LAAI in a structurally detailed model of the human atria,
49 extensively described previously³ (supplemental material). The model entails wall thickness
50 heterogeneities, endocardial trabeculated network, and a subepicardial layer with realistic
51 fiber orientations. **The material properties for the atria were set to produce an approximately**
52 **normal P-wave during sinus rhythm (Table S1).**

53 In total, 20 different pacing locations were selected in both atria based on reported possible
54 sources of extra-PV ectopic focal activity in AF patients⁴ (Figure).

55 In each simulation, AF was initiated by applying incremental pacing with the duration of 2
56 seconds at a selected pacing site. The pacing cycle length started with 280ms and gradually
57 reduced to 124ms. AF initiation likelihood was compared between control (no ablation) and
58 catheter ablation (CA) simulations with different degrees of fibrosis. CA simulations
59 included four groups: PV isolation (PVI), PVI plus linear lesions encircling the left atrial
60 posterior wall (BOX), PVI plus LAA circumferential isolation (PVI + LAAI), and BOX +
61 LAAI. Virtual ablation lesions consisted of tissue volumes modeled by non-conductive
62 elements. The outcome of the stimulation protocol was analyzed in terms of the type of self-
63 sustained rhythm after 2 seconds of stimulations. If no activity was observed, the initiation
64 was considered unsuccessful. Otherwise, a distinction between AF and atrial flutter (AFL)
65 was made (supplemental material), with the latter being considered unsuccessful.

66 The spatial distribution of fibrosis was uneven, with patches or islands presenting higher
67 degree of fibrosis (Figure S1).³ Simulations were performed without fibrosis, with moderate
68 fibrosis, and severe fibrosis, in which 0%, 50%, and 70% of elements were fibrotic.³

69 In total, 300 simulations were performed. Each simulation group entails 60 simulations (20
70 simulations for each fibrosis degree).

71 In control simulations, an increase in the fibrosis degree led to a significant increase in AF
72 initiation likelihood. In simulations without fibrosis or with moderate fibrosis (which may
73 represent paroxysmal AF patients), PVI and BOX caused a significant reduction in AF
74 initiation likelihood, either by preventing initiation of fibrillation or by converting AF into
75 AFL (Figure). In contrast, in PVI and BOX simulations with severe fibrosis (which may
76 represent persAF) we observed no significant reduction in AF initiation likelihood. In
77 simulations with LAAI on top of PVI or BOX, we observed a comparable reduction in AF

78 initiation likelihood as in PVI only or BOX simulations in the absence of fibrosis or in the
79 presence of moderate fibrosis. In severe fibrosis, adding LAAI to PVI or BOX further
80 reduced AF initiation likelihood by at least one third.

81 An increase in fibrosis degree in control simulations led to a significant increase fibrillation
82 wave generation rate and therefore a significant increase in AF conduction pattern
83 complexity, quantified as number of waves per cycle (Figure). In simulations without fibrosis
84 or with moderate fibrosis, CAs caused a significant reduction in fibrillation wave generation
85 rate and AF conduction pattern complexity. In contrast, in severe fibrosis simulations, a
86 significant reduction in wave generation rate and AF pattern complexity was only observed in
87 simulations with LAAI plus PVI or BOX.

88 In our control group, the averaged AF cycle length (AFCL) was 149ms, value previously
89 reported in AF patients. In simulations without fibrosis or with moderate fibrosis, CAs caused
90 a significant AFCL prolongation (Table S2). In severe fibrosis simulations only the presence
91 of LAAI concomitant with CAs could significantly increase AFCL.

92 The simulations were performed on a single atrial geometry and inter-subject variability
93 could have effects on simulations results. However, the in-depth study of the atrial anatomy
94 effect on AF initiation is beyond the scope of this study.

95 We modelled LAAI and investigated its effect on AF recurrences in the presence of different
96 degrees of fibrosis. As presented, adding LAAI to either PVI or BOX significantly lowered
97 AF recurrence rate in severe fibrosis simulations, which is consistent with clinical studies
98 mentioning the significant effect of LAAI in lowering AF recurrence rates in persAF.²

99 Finally, we could demonstrate that the mechanisms underlying AF initiation prevention in the
100 presence of LAAI is not due to isolating triggers in LAA, but rather related to the reduction
101 of AF conduction pattern complexity as a result of substrate modification due to LAAI.

102

103 **Funding:** This work was supported by the Swiss National Supercomputing Centre (CSCS
104 project ID s778), by grants to US from the Netherlands Heart Foundation (CVON2014-09,
105 RACE V), and the EU (ERACoSysMED H2020 ERA-NET). Dr Conte is supported by the
106 Swiss National Foundation (SNF) (Ambizione grant no PZ00P3_180055 / 1).

107 **Disclosure:** AA is a consultant to Boston Scientific, Backbeat, Biosense Webster, Cairdac,
108 Corvia, Microport CRM, Philips, Radcliffe Publisher. He received speaker fee from Boston
109 Scientific, Medtronic, and Microport. He participates in clinical trials sponsored by Boston
110 Scientific, Medtronic, Philips. He has intellectual properties with Boston Scientific, Biosense
111 Webster, and Microport CRM. US received consultancy fees or honoraria from Johnson &
112 Johnson, Roche Diagnostics, and Bayer Healthcare. US is co-founder and shareholder of
113 YourRhythmics BV. He holds intellectual property with Roche and YourRhythmics BV. The
114 other authors have nothing to declare.

117 **Reference:**

- 118 1. Verma A, Jiang CY, Betts TR, Chen J, Deisenhofer I, Mantovan R, Macle L, Morillo
119 CA, Haverkamp W, Weerasooriya R, Albenque JP, Nardi S, Menardi E, Novak P, Sanders P
120 and Investigators SAI. Approaches to catheter ablation for persistent atrial fibrillation. *N Engl*
121 *J Med.* 2015;372:1812-22.
- 122 2. Di Biase L, Burkhardt JD, Mohanty P, Mohanty S, Sanchez JE, Trivedi C, Gunes M,
123 Gokoglan Y, Gianni C, Horton RP, Themistoclakis S, Gallingshouse GJ, Bailey S, Zagrodzky
124 JD, Hongo RH, Beheiry S, Santangeli P, Casella M, Dello Russo A, Al-Ahmad A, Hranitzky
125 P, Lakkireddy D, Tondo C and Natale A. Left Atrial Appendage Isolation in Patients With
126 Longstanding Persistent AF Undergoing Catheter Ablation: BELIEF Trial. *J Am Coll*
127 *Cardiol.* 2016;68:1929-1940.
- 128 3. Gharaviri A, Bidar E, Potse M, Zeemering S, Verheule S, Pezzuto S, Krause R,
129 Maessen JG, Auricchio A and Schotten U. Epicardial Fibrosis Explains Increased Endo-
130 Epicardial Dissociation and Epicardial Breakthroughs in Human Atrial Fibrillation. *Front*
131 *Physiol.* 2020;11:68.
- 132 4. Chang HY, Lo LW, Lin YJ, Chang SL, Hu YF, Li CH, Chao TF, Chung FP, Ha TL,
133 Singhal R, Chong E, Yin WH, Tsao HM, Hsieh MH and Chen SA. Long-term outcome of
134 catheter ablation in patients with atrial fibrillation originating from nonpulmonary vein
135 ectopy. *J Cardiovasc Electrophysiol.* 2013;24:250-8.

140 **Figure 1.** A) Catheter ablation patterns and pacing points (pacing locations located outside of
141 ablated area were indicated by black stars and the pacing location within the ablated area was
142 indicated by yellow). B) AF and AFL initiation likelihood in control and catheter ablation
143 simulations with different degrees of fibrosis. C) Number of waves per cycle. Wave life span
144 and wave generation rate (the slope of fitted line) in D) control E) PVI F) PVI + LAAI
145 simulations with sever fibrosis.

146

