

On the logical structure of choice and bar induction principles

Nuria Brede, Hugo Herbelin

▶ To cite this version:

Nuria Brede, Hugo Herbelin. On the logical structure of choice and bar induction principles. 2021. hal-03144849v1

HAL Id: hal-03144849 https://inria.hal.science/hal-03144849v1

Preprint submitted on 17 Feb 2021 (v1), last revised 5 Jul 2022 (v5)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On the logical structure of choice and bar induction principles

Nuria Brede¹ and Hugo Herbelin²

¹University of Potsdam, Germany ²Inria Paris, Université de Paris, CNRS, IRIF, France

Abstract

We develop an approach to choice principles and their contra-positive bar-induction principles as extensionality schemes connecting an "effective" or "intensional" view of respectively ill- and well-foundedness properties to an "extensional" or "ideal" view of these properties. After classifying and analysing the relations between different intensional definitions of ill-foundedness and well-foundedness, we introduce, for a domain A, a codomain B and a "filter" T on finite approximations of functions from A to B, a generalised form GDC_{ABT} of the axiom of dependent choice and dually a generalised bar induction principle GBI_{ABT} such that:

 GDC_{ABT} intuitionistically captures

- the strength of the general axiom of choice expressed as $\forall a \exists b R(a, b) \Rightarrow \exists \alpha \forall a R(a, \alpha(a)))$ when T is a filter that derives point-wise from a relation R on $A \times B$ without introducing further constraints,
- the strength of the Boolean Prime Filter Theorem / Ultrafilter Theorem if B is the two-element set \mathbb{B} ,
- the strength of the axiom of dependent choice if $A = \mathbb{N}$,

and up to weak classical reasoning

• the (choice) strength of Weak König's Lemma if $A = \mathbb{N}$ and $B = \mathbb{B}$.

 GBI_{ABT} intuitionistically captures

- the strength of Gödel's completeness theorem in the form validity implies provability when $B = \mathbb{B}$,
- the strength of the Bar Induction when $A = \mathbb{N}$,
- the (choice) strength of the Weak Fan Theorem when $A = \mathbb{N}$ and $B = \mathbb{B}$.

Contrastingly, even though GDC_{ABT} and GBI_{ABT} smoothly capture several variants of choice and bar induction, some instances are inconsistent, e.g. when A is $\mathbb{B}^{\mathbb{N}}$ and B is \mathbb{N} .

1 Introduction

1.1 Bar induction, dependent choice and their variants as extensionality principles

For a domain A, there are different ways to define a well-founded tree branching over A. A first possibility is to define it as an inductive object built from leaves and from nodes associating a subtree to each element in A. We will call this definition *intensional*. Using a syntax familiar to functional programming languages or Martin-Löf-style type theory, such intensional trees correspond to inhabitants of an inductive type:

type tree = Leaf | Node of (A ightarrow tree)

A second possibility is a definition which we shall call *extensional* and which is probably more standard in the context of non type-theoretic mathematics. Let A^* denote the set of finite sequences of elements of A, with $\langle \rangle$ denoting the empty sequence and $u \star a$ the extension of the sequence u with a from A. Then an extensional tree T is a downwards-closed predicate over A^* . Finite sequences $u \in A^*$ are interpreted as finite paths from the root of a tree and the predicate determines which paths are contained in T. We say that T is *extensionally well-founded* if for all infinite paths α in $A^{\mathbb{N}}$, the path eventually "leaves" the tree, i.e. there is an initial finite prefix u of α such that u (as path from the root) is not contained in T.

The intensional definition is stronger: to any inductively-defined tree t, we can associate an extensionally well-founded tree T(t) by recursion on t as follows:

$$\begin{array}{lll} u \in T(\texttt{Leaf}) & \triangleq & \bot \\ u \in T(\texttt{Node}(f)) & \triangleq & \top \\ u \star a \in T(\texttt{Node}(f)) & \triangleq & u \in T(f(a)) \end{array}$$

and we can prove by induction on t that $\forall \alpha \exists n T(t)(\alpha_{|n})$, where $\alpha_{|n}$ is the restriction of α to its first n values.

Bar induction, introduced by Brouwer and further analysed by Kleene and Vesley [25] is the converse property, namely that any extensionally well-founded T can be turned into an inductively-defined tree. I.e. as a scheme it asserts the equivalence of the extensional and intensional definitions of well-foundedness¹. This makes explicit the view developed in particular by Howard and Kreisel [15] that bar induction transforms extensional well-foundedness into well-founded induction (where the latter is seen as the ability to exhibit an explicit inductively-defined tree-like proof of being *inductively barred*).

To clarify what this means, and in particular to clarify in which case an inductively-defined t denotes the same tree as an extensionally well-founded T tree, we can define a realisability relation between t and T as follows:

- Leaf realises T at u if $\langle \rangle \notin T$
- Node(f) realises T at u if for all a, f(a) realises T at $u \star a$

Then, we can prove by induction on t that the subtree of t rooted in u realises T(t) at u.

The notion *inductively well-founded* combines the data of a tree t and the property that this tree t realises a predicate: We define T is inductively well-founded at u as the smallest predicate which satisfies

- if $u \notin T$ then T is inductively well-founded at u
- if, for all a, T is inductively well-founded at $u \star a$, then T is inductively well-founded at u

In particular, the existence of a tree t such that t realises T at u is equivalent to T inductively well-founded at u: the inductive structure of the subtree of t rooted in u is then reflected in the structure of the proof of T inductively well-founded at u.

This allows to rephrase the correspondence between the existence of an inductively-defined tree t and the existence of an extensionally well-founded tree T into a correspondence talking in both cases of an arbitrary (a priori not well-founded) tree:

- if T is inductively well-founded, it is extensionally well-founded (this is provable by induction on the proof of T inductively well-founded)
- if T is extensionally well-founded, it is inductively well-founded (this is a reformulation of bar induction using the more informative property of inductively well-founded called *inductively barred* when it applies to the complement of the tree, as in the original formulation of bar induction rather than just the existence of an inductively-defined tree)

Now, if bar induction is considered as an extensionality principle, the same should be the case for its contrapositive which is logically equivalent to the *axiom of dependent choice*: The axiom of dependent choice can be rephrased as a principle asserting that if a tree is *coinductively ill-founded*, then it is *extensionally ill-founded* (i.e. an infinite branch can be found). We will detail this in Section 2, together with precise relations between these principles and their restriction on finitely-branching trees, namely *König's lemma* and the *Fan Theorem*, introducing a systematic terminology to characterise and compare these different variants.

Note that the approach to consider bar induction and choice principles as extensional principles is also consistent with the methodology developed e.g. by Coquand and Lombardi: to avoid the necessity of choice or bar-induction axioms, mathematical theorems are restated using the (co-)inductively-defined notions of well- and ill-foundedness rather than the extensional notions [9, 10].

1.2 Weak König's Lemma at the intersection of Boolean Prime Filter Theorem and Dependent Choice

We know from reverse mathematics of the subsystems of second order arithmetic [24] that the binary form of König's lemma, namely *Weak König's Lemma* (WKL) has the strength of *Gödel's completeness theorem* (for a countable language). Reverse mathematics of the axiom of choice and its variants in set theory [13, 14, 18, 11] also tells that Gödel's completeness theorem has the strength of the *Boolean Prime Filter Theorem* (for a language of arbitrary cardinal). This suggests that the Boolean Prime Filter Theorem is the "natural" generalisation of WKL from countable to arbitrary cardinals.

On the other side, Weak König's Lemma is a consequence² of the axiom of Dependent Choice, the same way as its dual, the Weak Fan Theorem, is an instance of Bar Induction, itself classically equivalent to the axiom of Dependent

¹Kleene and Vesley [25] used respectively the terms "explicit" and "inductive".

²Note that König's Lemma is a theorem of set theory and that we need to place ourselves in a sufficiently weak metatheory, e.g. RCA_0 , to state this result.

ill-founded-style	well-founded-style	
functions from A to B		
$AC_{ABR} = GDC_{ABR_{\top}}$		
branching from \mathbb{N} over arbitrary B		
$DC^{productive}_B = GDC_{\mathbb{N}B}$	$Bl_B^{ind}=GBl_{\mathbb{N}B}$	
branching from \mathbb{N} over non-empty finite B		
$KL = DC^{staged}_{\mathrm{fin}}$	$FT=Bl^{uniform}_{\mathrm{fin}}$	
binary branching from N		
$WKL=DC^{staged}_{\mathbb{B}}$	$WFT=Bl^{uniform}_{\mathbb{B}}$	
$WKL =_{co-intuit.} GDC_{\mathbb{NB}}$	$WFT =_{\mathit{intuitionistic}} GBI_{\mathbb{NB}}$	
binary branching from arbitrary A		
$Compl^A(\mathcal{T}) = GDC_{A\mathbb{B}\mathcal{T}^C}$	$Compl^+_A(\mathcal{T}) = GBl_{A\mathbb{B}\mathcal{T}}$	
$BPF_{Free(A)}(F_{\mathcal{T}}) = GDC_{A\mathbb{B}\mathcal{T}^C}$		

Table 1: Summary

Choice. This suggests that there is common principle which subsumes both the Axiom of Dependent Choices and the Boolean Prime Filter Theorem with Weak König's Lemma at their intersection.

Such a principle is stated in Section 3 where it is shown that the ill-founded version indeed generalises the axiom of Dependent Choice and the well-founded version generalises Bar Induction. In the same section, we also show that one of the instance of the ill-founded version captures the general Axiom of Choice, but that, in its full generality, the new principle is actually inconsistent.

Section 4 is devoted to show that the Boolean Prime Filter Theorem is an instance of the generalised axiom of Dependent Choice. In particular, this highlights that the notions of *ideal* and *filter* generalise the notion of a binary tree where the prefix order between paths of the tree is replaced by an inclusion order between non-sequentially ordered paths now seen as finite approximations of a function from \mathbb{N} to the two-element set \mathbb{B} .

1.3 Methodology and summary

For our investigations to apply both to classical and to intuitionistic mathematics, we carefully distinguish between the choice axioms (seen as ill-foundedness extensionality schemes) and bar induction schemes (seen as well-foundedness extensionality schemes).

All in all, the classification we obtain is summarised in table 1 where the definitions of the different notions can be found in the respective sections of the paper.

2 The logical structure of dependent choice and bar induction and principles

2.1 Metatheory

We place ourselves in a metatheory capable to express arithmetic statements. In addition to the type \mathbb{N} of natural numbers together with induction and recursion, we assume the following constructions to be available:

- The type \mathbb{B} of Boolean values 0 and 1 together with a mechanism of definition by case analysis. It shall be convenient to allow the definition of propositions by case analysis as in if b then P else Q, whose logical meaning shall be equivalent to $(b = 0 \land P) \lor (b = 1 \land Q)$.
- For any type A, the type A^* of finite sequences over A whose elements shall generally be ranged over by the letters $u, v \dots$ We write $\langle \rangle$ for the empty sequence and $u \star a$ for the extension of sequence u with element a. We write |u| for the length of u and u(n) for the nth element of u when n < |u|. We write v@u for the concatenation of v and u.

We write $u \leq_s v$ to mean that u is an initial prefix of v. This is inductively defined by:

$$\frac{u \leq_s v}{u \leq_s u} \qquad \frac{u \leq_s v}{u \leq_s v \star a}$$

We shall also support case analysis over finite sequences under the form of a case operator.

- For any two types A and B, the type $A \to B$ of functions from A to B. Functions can be built by λ -abstraction as in $\lambda x.t$ for x in A and t in B and used by application as in t(u) for t in $A \to B$ and u in A. To get closer to the traditional notations, we shall also abbreviate $t(u_1) \dots (u_n)$ into $t(u_1, \dots, u_n)$.
- A type Prop reifying the propositions as a type. The type $A \to \text{Prop}$ shall then represent the type of predicates over A. We shall allow predicates to be defined inductively (smallest fixpoint) or coinductively (greatest fixpoint), using respectively the μ and ν notations.
- For any type A and predicate P over A, the subset $\{a:A \mid P(a)\}$ of elements of A satisfying P.

This is the language of higher-order arithmetic but in practice, we shall need quantification just over functions and predicates superficially of rank 1 (i.e. of the form $A_1 \rightarrow \ldots \rightarrow A_n \rightarrow A$ or $A_1 \rightarrow \ldots \rightarrow A_n \rightarrow Prop$ with no arrow types in A and the A_i). We however also allow arbitrary type constants to occur, so we can think of our effective metatheory as a second-order arithmetic generic over arbitrary more complex types. In practice, our metatheory could typically be the image of arithmetic in set theory or in an impredicative type theory. We shall in any case use the notation $a \in A$ to mean that a has type A when A is a type, which, if in set theory, shall mean a set satisfying some specific property.

We shall leave open the meaning of the connectives. Using linear logic as a reference for the semantics of connectives, we shall interpret implication either linearly (i.e. as linear logic's $P \multimap Q$ where P can be used only once in a proof of Q), intuitionistically (i.e. as $!P \multimap Q$ where ! is the of-course modality of linear logic expressing that P can be used an arbitrary number of times in the proof), or classically (i.e. as $!P \multimap ?Q$ where ? is the why-not modality of linear logic, expressing that classical reasoning can be used to prove Q). Similarly, $\exists a^A Q$ shall have an intuitionistic/linear meaning (i.e. as $\bigoplus_{a \in A} Q$ using notations of linear logic) or a classical meaning (i.e. as $? \bigoplus_{a \in A} Q$).

2.2 Infinite sequences

We write $A^{\mathbb{N}}$ for the infinite (countable) sequences of elements of A. There are different ways to represent such an infinite sequence:

- We can represent it as a function, i.e. as a functional object of type $\mathbb{N} \to A$.
- We can represent it as a functional relation, i.e. as a relation R of type $\mathbb{N} \to A \to \mathsf{Prop}$ such that $\forall n \exists ! a R(n, a)$.
- Additionally, when A is \mathbb{B} , an extra possible representation shall be as predicate P over N with intended meaning 0 if P(n) holds and 1 if $\neg P(n)$ holds (and unknown meaning otherwise).

The representation as a functional relation is weaker in the sense that a function α mechanically induces a functional relation $\lambda n. \lambda a. \alpha(n) = a$ but the converse requires the axiom of unique choice. In the sequel, we shall use the notation $\alpha(n) = a$ and $\alpha(n) \triangleq a$ to mean different things depending on the representation chosen for $\mathbb{N} \to B$.

In the first case, $\alpha(n) = a$ shall mean $\alpha(n) = a$ where the equality sign of the notation is interpreted as the equality on A. Similarly, $\alpha(n) \triangleq a$ shall define the function $\alpha \triangleq \lambda n. a$.

In the second case, $\alpha(n) = a$ shall however mean $\alpha(n, a)$ and $\alpha(n) \triangleq a$ shall define the functional relation $\alpha \triangleq \lambda(n, a')$. a' = a where n can occur in a.

When A is \mathbb{B} , the representation as a predicate P is even weaker in the sense that a functional relation R induces a predicate $\lambda n. R(n, 0)$ but the converse requires classical reasoning. We can easily turn a predicate P into a relation $\lambda n. \lambda b.$ if b then P(n) else $\neg P(n)$ but proving $\forall n \exists ! b$ if b then P(n) else $\neg P(n)$ requires a call to excluded-middle on P(n).

When A is \mathbb{B} and α is a predicate, we shall define $\alpha(n) = 0$ as $\alpha(n)$ and $\alpha(n) = 1$ as $\neg \alpha(n)$. Technically, this means seeing $\alpha(n) = b$ as a notation for if b then $\alpha(n)$ else $\neg \alpha(n)$. Similarly, $\alpha(n) \triangleq b$ shall define $\alpha \triangleq \lambda n$ if b then \top else \bot .

In particular, this means that all choice and bar induction statements of this paper have two readings of a different logical strength (depending on the validity of the axiom of unique choice in the metatheory), or even three readings (depending on the validity of the axiom of unique choice and of classical reasoning) when the codomain of the function mentioned in the theorems is \mathbb{B} .

T is a tree	T is monotone
(closure under restriction)	(closure under extension)
$\forall u \forall a (u \star a \in T \Rightarrow u \in T)$	$\forall u \forall a (u \in T \Rightarrow u \star a \in T)$
downwards arborification of T	upwards monotonisation of T
$(\downarrow^{-}T)$	$(\uparrow^+ T)$
$\lambda u. \forall u' (u' \leq_s u \Rightarrow u' \in T)$	$\lambda u. \exists u' (u' \leq_s u \land u' \in T)$
upwards arborification of T	downwards monotonisation of T
$(\uparrow T)$	$(\downarrow^+ T)$
$\lambda u. \exists u' (u \leq_s u' \land u' \in T)$	$\lambda u. \forall u' (u \leq_s u' \Rightarrow u' \in T)$

Table 2: Logically equivalent dual concepts on dual predicates (I)

If $\alpha \in A^{\mathbb{N}}$, we write $u \prec_s \alpha$ to mean that u is an initial prefix of f. This shall be defined inductively by the following clauses:

$$\frac{u \prec_s \alpha \qquad \alpha(|u|) =}{u \star a \prec_s f}$$

If $a \in A$ and $\alpha \in A^{\mathbb{N}}$, we write $a@\alpha$ for the sequence β defined by $\beta(0) \triangleq a$ and $\beta(n+1) \triangleq \alpha(n)$. We have the following easy property:

Proposition 1 If $u \prec_s \alpha$ then $a@u \prec_s a@\alpha$.

2.3 Trees and monotone predicates

Let B be a type and T be a predicate on B^* . We overload the notation $u \in T$ to mean that T holds on $u \in B^*$. We say that T is finitely-branching if B is in bijection with a non-empty bounded subset of N (i.e. to $\{n : \mathbb{N} \mid n \leq p\}$ for some p).

We say that T is a *tree* if it is closed under restriction, and, dually, that T is *monotone* if it is closed under extension (the formal definitions are given in table 2). Classically, we have T monotone iff $\neg T$ is a tree, and, dually, $\neg T$ monotone iff T is a tree. In particular, another way to describe a tree is as an antimonotone predicate³. It is convenient for the underlying intuition to restrict oneself to predicates which are trees, or which are monotone, even if it does not always matter in practice. When it matters, a predicate is turned into a tree either by pruning the predicate from sequences not connected to the root or by completing it with missing sequences from the root: these are respectively the *downwards arborification* $\uparrow^- T$ of a predicate. We dually write $\uparrow^+ T$ and $\downarrow^+ T$ for the *upwards monotonisation* and *downwards monotonisation* of T. Arborification and monotonisation are idempotent. We shall in general look for minimal definitions of the concept involved in the paper, and thus consider arbitrary predicates as much as possible, turning them into trees or monotone predicates only when needed to give sense to the definitions.

2.4 Well-foundedness and ill-foundedness properties

We list properties on predicates which are relevant for stating ill-foundedness axioms (i.e. choice axioms), and their dual well-foundedness axioms (i.e. bar induction axioms). Duality can be understood both under a classical or linear interpretation of the connectives, where the predicate T in one column is supposed to be dual of the predicate T occurring in the other column (dual predicates if in linear logic, negated predicates if in classical logic). Table 3 details properties which differ by contraposition and are thus logically equivalent (in classical and linear logic). On the other side, tables 4 and 5 detail properties which are logically opposite.

We indicated with (*) concepts for which we did not find an existing terminology in the literature. Thus, the terminology is ours. Also, what we called *staged infinite* is often simply called *infinite*. We used *staged infinite* to make explicit the difference from a definition based on the presence of an infinite number of nodes. Thereby we also obtain a symmetry with the notion of *staged barred*. What we call *having an infinite branch* could alternatively be called *ill-founded*, or *having a*

³From a categorical perspective, a tree is a contravariantly functorial predicate over the preorder generated by $u \leq_s v$, while a monotone predicate is covariantly functorial.

T is progressing at u (*)	T is hereditary at u
$u \in T \Rightarrow (\exists a u \star a \in T)$	$(\forall a u \star a \in T) \Rightarrow u \in T$
T is progressing (*)	T is hereditary
$\forall u (T \text{ is progressing at } u)$	$\forall u (T \text{ is hereditary at } u)$

Table 3: Basic logically equivalent dual properties on dual predicates

Table 4: Logically opposite dual properties on dual predicates

ill-foundedness properties	well-foundedness properties	
Intensione	al concepts	
T is a spread	T is resisting (*)	
$\langle \rangle \in T \wedge T$ progressing	$T \text{ hereditary} \Rightarrow \langle \rangle \in T$	
pruning of T	adherence $(*)$ of T	
$\nu X.\lambda u. \ (u \in T \land \exists a \ u \star a \in X)$	$\mu X.\lambda u. (u \in T \lor \forall a u \star a \in X)$	
T is productive	T is inductively barred	
$\langle \rangle \in \text{pruning of } T$	$\langle \rangle \in adherence of T$	
Intensional concepts relevant for the finite case		
T has unbounded paths	T is uniformly barred	
$\forall n \exists u (u = n \wedge u \in \downarrow^{-} T)$	$\exists n \forall u (u = n \Rightarrow u \in \uparrow^{\!\!+} T)$	
T is staged infinite	T is staged barred (*)	
$\forall n \exists u (u = n \land u \in T)$	$\exists n \forall u (u = n \Rightarrow u \in T)$	
Extensional concepts		
T has an infinite branch	T is barred	
$\exists \alpha \forall u (u \prec_s \alpha \Rightarrow u \in T)$	$\forall \alpha \exists u (u \prec_s \alpha \wedge u \in T)$	

choice function. In particular, the terminology having an infinite branch applies here to any predicate and is not restricted to trees. Note that *well-founded* in the standard meaning is the same as *barred* for the dual predicate. In particular, when opposing ill-foundedness and well-foundedness, we adopt a bias towards the tree view, i.e. towards the left column. We have the following:

Proposition 2 If T is a tree, then having unbounded paths is equivalent to being staged infinite. Dually, if T is monotone, being a uniform bar is equivalent to being staged barred.

PROOF: Because trees and monotone predicates are invariant under arborification and monotonisation.

As a consequence, it is common to use the notion of staged infinite, which is simpler to formulate, when we know that T is a tree. Otherwise, if T is an arbitrary predicate which is not necessarily a tree, there is no particular interest in using the notion of staged infinite. Similarly, staged barred is a simpler way to state uniformly barred when T is monotone, i.e., conversely, uniform bar is the expected refinement of staged barred when T is not known to be monotone.

By a direct coinduction, any spread is productive, and further, any spread is productive at any $u \in T$. Thus, we have:

Proposition 3 T spread implies T productive, and, dually, T inductively barred implies T resisting.

Conversely, a productive T may be productive at $\langle \rangle$ without being productive at all $u \in T$, so we may need to prune T to extract from it a spread. Dually, not all *resisting* predicates are inductive bars but we can saturate them into inductive bars, using the notion of *adherence*. We make this formal in the following proposition:

$\it ill$ -foundedness-style	well-founded ness-style	
Relativised intensional concepts		
T is productive from u	T is inductively barred from u	
$u \in \text{pruning of } T$	$u \in adherence of T$	
Relativised intensional concepts relevant for the finite case		
T has unbounded paths from u	T is uniformly barred from u n	
$\forall n \exists u' (u' = n \wedge u @u' \in {\downarrow^{-}} T)$	$\exists n \forall u' (u' = n \Rightarrow u @u' \in \uparrow^+ T)$	
T is staged infinite from u	T is staged barred from u'	
$\forall n \exists u' (u' = n \wedge u @u' \in T)$	$\exists n \forall u' (u' = n \Rightarrow u @u' \in T)$	
Extensional concepts		
T has an infinite branch from u	T is barred from u	
$\exists \alpha \forall u' (u' \prec_s \alpha \Rightarrow u @ u' \in T)$	$\forall \alpha \exists u' (u' \prec_s \alpha \wedge u @ u' \in T)$	

Table 5: Useful relativisation of some of the concepts of Table 4

Proposition 4 If T is productive then its pruning is a spread. Dually, if T resists then its adherence is an inductive bar.

PROOF: That $\langle \rangle$ is in the pruning of T is direct from T productive. That the pruning of T is progressing on all u is also direct by construction of the pruning. The other part of the statement is by duality.

Also, we can relate productive and spread, as well as inductive bar and resisting as follows:

Proposition 5 T is productive iff there exists $U \subset T$ which is a spread. Dually, T is an inductive bar iff all $U \supset T$ resists.

PROOF: By duality, it is enough to prove the first equivalence. From left to right, we use Proposition 4, observing that the pruning of T is included in T. From right to left, a spread is productive and a coinduction suffices to prove that inclusion preserves productivity.

On the other side, having unbounded paths is equivalent to being a spread or to productive only when T is finitelybranching. Similarly for uniformly barred compared to inductive bar and resisting. Moreover, none of the equivalence holds linearly. The second one requires intuitionistic logic, i.e. requires the ability to use an hypothesis several times while the first one, dually, requires a bit of classical reasoning⁴.

For S being a class of formulae and P and Q ranging over S, let D_S be the principle $\forall xy (P(x) \lor Q(y)) \Rightarrow (\forall x P(x)) \lor (\forall y Q(y))$. Dually, let C_S be $(\exists x P(x)) \land (\exists y Q(y)) \Rightarrow \exists x \exists y (P(x) \land Q(y))$.

Proposition 6 If B is non-empty finite, then productive is equivalent to having unbounded paths and being an inductive bar is equivalent to uniformly barred. The first statement holds in a logic where D_S holds and the second in a logic where C_S holds, for S a class of formulae containing arithmetical existential quantification over T.

PROOF: Relying on duality, we only prove the first statement. Without loss of generality, we also assume that B is \mathbb{B} . Our proof relies on an argument found in [3, 16] and proceeds by proving more generally that T is productive from u iff T has unbounded paths from u.

From left to right, we reason by induction on n. If n is 0 this is direct from T productive by defining $u' \triangleq \langle \rangle$. Otherwise, by T productive from u, we get a such that T is productive from $u \star a$, obtaining by induction u' of length n-1 such that $(u \star a)@u' = u@(a@u') \in \downarrow^{-} T$, showing that a@u' is the expected sequence of length n.

From right to left, we reason coinductively. To prove that $u \in T$, we take a path of length 0. Then, in order to apply the coinduction hypothesis and prove the coinductive part, we prove that there is b such that T has unbounded paths from $u \star b$. By D_S , it is enough to prove that for all n_0 and n_1 , there is a path u_0 of length n_0 and a path u_1 of length n_1 such that either $(u \star 0)@u_0$ or $(u \star 1)@u_1$ is in $\downarrow^- T$. So, let n_0 and n_1 be given lengths. By unbounded paths from u,

⁴or cointuitionistic reasoning to be more precise

ill-foundedness-style	well-foundedness-style
Tree-based Dependent Choice (DC ^{spread})	Dual to Tree-based Dep. Choice $(B ^{\mathrm{resists}})$
T spread $\Rightarrow T$ has an infinite branch	T barred $\Rightarrow T$ resists
Dual to Generic Bar Induction (DC ^{productive})	Generic Bar Induction (Bl ^{ind})
T productive $\Rightarrow T$ has an infinite branch	T barred $\Rightarrow T$ inductively barred
Dual to Fan Theorem $(DC_{fin}^{unbounded})$	Fan Theorem $(BI_{\mathrm{fin}}^{\mathrm{uniform}} = FT)$
T finitely-branch. w. unbounded paths $\Rightarrow T$ has an infinite branch	T barred and finitely-branching $\Rightarrow T$ uniform bar
König's Lemma $(DC^{\mathrm{staged}}_{\mathrm{fin}} = KL)$	Dual to König's Lemma $(Bl_{\mathrm{fin}}^{\mathrm{staged}})$
T finitely-branching staged-infinite tree $\Rightarrow T$ has an infinite branch	T barred, finitely-branching and monotone $\Rightarrow T$ staged barred

Table 6: Tree-based choice principles and bar induction principles

we get a sequence u'' of length $max(n_0, n_1) + 1$ such that $u@u'' \in \downarrow^- T$. This is a non-empty sequence, hence a sequence of the form b@u' so that we have either $(u \star 0)@u' \in \downarrow^- T$ or $(u \star 1)@u' \in \downarrow^- T$ for u' of length $max(n_0, n_1)$. By closure of $\downarrow^- T$, prefixes u_0 of length n_0 and u_1 of length n_1 of u' can be extracted which both are in $\downarrow^- T$.

Remark: Based on the decomposition of WKL for decidable trees into a choice principle and the Least Principle of Omniscience (LPO), we suspect that we actually have the stronger result that the equivalence of unbounded paths and productivity implies D_S for the corresponding underlying class of formula S, and similarly with C_S and the dual statement.

2.5 Standard bar induction and dependent choice principles

We now formulate standard statements of bar induction and dependent choice axioms from the literature using our definitions. What we call Bar Induction is a rather standard form [25], apart from the fact that we do not fix in advance the logical complexity of B – such as being countable or not – or the arithmetic strength of T – i.e. whether it is decidable, or recursively enumerable, etc. For Dependent Choice⁵, we consider here a pruned-tree-based definition, which we shall later compare to other logically equivalent definitions in Section 2.7.

The Fan Theorem is sometimes stated over finitely-branching trees, where the definition of finite itself may vary [25, 16], but it is also sometimes considered by default to be on a binary tree [4, 2, 3, 7, 9, 17] in which case the finite version is sometimes called extended. What we call here Fan Theorem is a definition where the finite branching is uniform at all levels and we shall later call the binary version Weak Fan Theorem in Section 4. The statement of the Fan Theorem sometimes relies on the notion of inductive bar (e.g. [9]), or on the definition of staged barred for monotone predicates (as a variant in [17]), or on the dual notions of finite tree (i.e., technically of staged barred for the negation of a tree) and well-founded tree (i.e., technically of inductively barred for the negation of a tree) in e.g. [5]. But it also often relies on the definition of uniform bar [2, 3, 4, 7, 16, 17, 25] over an arbitrary predicate. It is this last variant which here we will explicitly call *Fan Theorem*. Note that as in the case of bar induction, we omit the usual restriction of the statement of the Fan Theorem to decidable predicates.

König's Lemma is generally stated as T infinite implies T has an infinite branch, but the definition of T infinite may differ from author to author. We here use the name König's Lemma for the definition from [5, 16] which expresses explicitly that the infinity can only be in depth. It does so by requiring arbitrary long branches rather than an infinite number of nodes. The exact definition of arbitrarily long branches also depends on authors. For instance, [26] relies (up to classical reasoning) on having unbounded paths for arbitrary predicates, but most of the time it is about what we call staged infinite tree [3, 16, 17].

Note that there is a huge body of literature on bar induction, dependent choice, König's Lemma and the Fan Theorem, especially in the context of constructive mathematics. We apologise for not being exhaustive.

Otherwise said, the Fan Theorem and König's Lemma are dual up to the (small and somewhat insignificant) difference that in the literature we found König's Lemma mostly formulated on trees while the Fan Theorem was mostly formulated

⁵ or Dependent Choices for some authors, e.g. [18]

for arbitrary predicates. In particular, there is a standard way to go from arbitrary predicates to trees or monotone predicates by associating to each predicate its tree or monotone closure. Using the closure under restriction and closure under extension of a predicate, we can connect König's Lemma to the dual of the Fan Theorem and the Fan Theorem to the dual of König's Lemma.

Proposition 7 As schemes, it is equivalent to state König's Lemma on trees using staged-infinity $(DC_{\text{fin}}^{\text{staged}})$ or on arbitrary predicates using has unbounded paths $(DC_{\text{fin}}^{\text{unbounded}})$. Similarly, it is equivalent to state the Fan Theorem on monotone predicates using staged barred $(Bf_{\text{fin}}^{\text{staged}})$ or on arbitrary predicates using uniformly barred $(Bf_{\text{fin}}^{\text{uniform}})$.

PROOF: We treat the first equivalence. From left to right, if T is a predicate, we apply $\mathsf{DC}_{fin}^{staged}$ to $\downarrow^- T$. The resulting infinite branch is an infinite branch in T because $\downarrow^- T \subset T$. From right to left, the statement holds by Proposition 2. The second equivalence is by duality.

Similarly, Tree-based Dependent Choice and Bar Induction are dual except that Tree-based Dependent Choice is mostly stated for spreads while Bar Induction is stated for (the dual of) productive. Again, there are standard ways to pass from spread or co-spread to productive trees or inductive bars, and from a statement on serial or monotone relations to a statement on productive or inductively barred trees.

Proposition 8 As schemes, DC^{spread} and DC^{productive} are equivalent, and so are Bl^{spread} and Bl^{productive}.

PROOF: By Proposition 5.

2.6 Choice and bar induction as relating intensional and extensional concepts

The intensional definitions are stronger than the extensional ones, which implies that the choice and bar induction axioms can alternatively be seen as stating the logical equivalence of the intensional and extensional versions of ill-foundedness and well-foundedness properties (of various strengths).

Theorem 1 T inductively barred implies T barred. Dually, T has an infinite branch implies T is productive.

PROOF: We prove by induction on the definition of T inductively barred that T inductively barred at u implies T barred from u where the latter requires that for all α , there is $v \prec_s \alpha$ such that $uv \in T$.

If $u \in T$, then it is enough to take $\langle \rangle$ for v to get $u@\langle \rangle \in T$ for any α . If T is barred from $u \star b \in T$ for all $b \in B$, this means that there is $v \prec_s \beta$ such that $(u \star b)@v \in T$ for any β . For a given α , set $b \triangleq \alpha(0)$ and $\beta(n) \triangleq \alpha(n+1)$ so that we can find $v \prec_s \beta$, hence $b@v \prec_s b@\beta$, i.e. $b@v \prec_s \alpha$ (by Prop. 1) together with $u@(b@v) \in T$.

The dual proof builds T productive at u from T has an infinite branch from u by coinduction. From the infinite branch α from u and $\langle \rangle \prec_s \alpha$ we get $u@\langle \rangle \in T$, i.e. $u \in T$. It remains to find b such that T is productive from $u \star b$ and it suffices to take $\alpha(0)$ since T has an infinite branch $\beta(n) \triangleq \alpha(n+1)$ from $u \star \alpha(0)$ simply because $v \prec_s \alpha$ implies $\alpha(0)@v \prec_s \alpha(0)\star\beta$ (by Prop. 1) and $(u \star \alpha(0))@v \in T$ from $u@(\alpha(0)@v) \in T$.

2.7 Relation to other formulations of Dependent Choice and to countable Zorn's Lemma

For R a relation on B, it is common to formulate dependent choice as $\forall b^B \exists b'^B R(b,b') \Rightarrow \forall b_0^B \exists f^{\mathbb{N} \to B} (f(0) = b_0 \land \forall n R(f(n), f(n+1)))$. Let us call *serial* a (homogeneous) relation such that $\forall b^B \exists b'^B R(b,b')$ holds. In this section, we formally compare the resulting statement of dependent choice to $\mathsf{DC}_{BT}^{\mathsf{productive}}$, examining also dual statements.

Let R be a serial relation, i.e. a relation such that $\forall b^B \exists b'^B R(b,b')$. Using a seed b_0 , each such relation R can be turned into a predicate on B^* under the two following ways:

- The chaining $R^*_{\top}(b_0)$ from b_0 is probably the most natural one: it says that $u \in R^*_{\top}(b_0)$ if all steps in u from b_0 are in R.
- The alignment $R^{\triangleright}_{\top}(b_0)$ from b_0 artificially uses non-empty sequences to represent pairs of elements. We have $u \in R^{\triangleright}_{\top}(b_0)$ if the last two elements are related by R, and, in case one element is in the sequence, compare it by R to b_0 , and, in case the sequence is empty, make the predicate hold by default.

Reasoning by induction on $v \leq_s u$ in one direction and on u in the other direction, we can show that both are related:

Proposition 9 $u \in R^*_{\top}(b_0)$ iff $u \in \downarrow^{-} R^{\triangleright}_{\top}(b_0)$

ill-foundedness-style	well-foundedness-style
Intensional concepts	
R serial (or left-total)	${\cal R}$ has a "least" element
$\forall b\exists b'R(b,b')$	$\exists b\forall b'R(b,b')$
R left-not-full (*)	R has a "maximal" element
$\forall b \exists b' \neg R(b,b')$	$\exists b\forall b'\neg R(b,b')$
R choosable	R barred
$\exists \alpha \forall a (\alpha(a) = b \Rightarrow R(a, b))$	$\forall \alpha \exists a (\alpha(a) = b \wedge R(a, b))$
chaining of R from $b_0 (R^*_{\top}(b_0))$	antichaining of R from b_0 $(R^*_{\perp}(b_0))$
$\mu X.\lambda b. \lambda u.$ case u of	$\nu X.\lambda b. \lambda u.$ case u of
$ \left[\begin{array}{cc} \langle \rangle & \mapsto \top \\ b' \star u \mapsto R(b, b') \wedge X(b', u) \end{array} \right] $	$\left[\begin{array}{cc} \langle \rangle & \mapsto \bot \\ b' \star u \mapsto R(b, b') \lor X(b', u) \end{array} \right]$
	$\begin{bmatrix} \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\ \\$
angument of n from b_0 $(n_{\pm}(b_0))$	blockings of R from $\theta_0 (R_{\perp}(\theta_0))$
$\begin{bmatrix} \langle \rangle & \mapsto \top \\ b & \mapsto R(b_0, b) \\ u' \star b \star b' \mapsto R(b, b') \end{bmatrix}$	$\begin{bmatrix} \langle \rangle & \mapsto \bot \\ b & \mapsto R(b_0, b) \\ u' \star b \star b' \mapsto R(b, b') \end{bmatrix}$

Table 7: Logically opposite dual concepts on dual homogeneous relations

Table 8: Dependent choice and bar induction principles

ill-foundedness-style	well-foundedness-style
Dependent Choice $(DC_{BRb_0}^{\text{serial}})$	Dual to Dependent Choice $(Bl_{BRb_0}^{\text{least}})$
$R \text{ serial} \Rightarrow R^{\triangleright}_{\top}(b_0)$ has an infinite branch	$R^{\triangleright}_{\perp}(b_0)$ barred $\Rightarrow R$ has a least element

Dually, we can define *antichaining* and *blockings* such that:

Proposition 10 $u \in R^*_{\perp}(b_0)$ iff $u \in \uparrow^+ R^{\triangleright}_{\perp}(b_0)$

The formal definitions are given below, where we can notice that the use of μ vs. ν does not matter in practice since the structure of the relation is a function of |u|.

We are now in position to state a relatively standard form of Dependent Choice which we call $\mathsf{DC}_{BRb_0}^{\mathrm{serial}}$ for R relation on B and b_0 seed in B. Though to our knowledge uncommon in the literature, we also mention its dual which we call $\mathsf{Bl}_{BRb_0}^{\mathrm{least}}$.

We state a few results that allow to show the equivalence of $\mathsf{DC}_{BRb_0}^{\text{serial}}$ and $\mathsf{DC}_{BT}^{\text{productive}}$ as schemes. We have the following properties.

Proposition 11 R serial implies $R^{\triangleright}_{\top}(b_0)$ productive for any b_0 . Dually, if $R^{\triangleright}_{\perp}(b_0)$ is inductively barred then R has a least element.

PROOF: We prove by coinduction that $u \in R^{\triangleright}_{\top}(b_0)$ implies $R^{\triangleright}_{\top}(b_0)$ productive from u. If u is empty, $R^{\triangleright}_{\top}(b_0)(\langle\rangle)$ holds by definition and there is by seriality a b_1 such that $R^{\triangleright}_{\top}(b_0)(b_1)$. This allows to conclude by coinduction hypothesis. If u has the form $u' \star b$, there is also by seriality a b' such that $R^{\triangleright}_{\top}(b_0)(u' \star b \star b')$ and we can again conclude by coinduction hypothesis. The productivity of $R^{\triangleright}_{\top}(b_0)$ finally follows because $R^{\triangleright}_{\top}(b_0)(\langle\rangle)$ holds by definition. The dual statement is by dual (inductive) reasoning.

Table 9: Logically opposite dual concepts on dual relations

ill-foundedness-style	well-foundedness-style
R choosable	R barred
$\exists \alpha \forall a (\alpha(a) = b \Rightarrow R(a,b))$	$\forall \alpha \exists a (\alpha(a) = b \wedge R(a, b))$

Table 10: Countable choice and weak bar induction principles

$\it ill$ -foundedness-style	$well\-foundedness\-style$
Countable Choice (CC_{BR})	Dual to Countable Choice (WBI_{BR})
R left-total $\Rightarrow R$ choosable	R barred $\Rightarrow R$ has a least element

Conversely, for T a predicate, let B_T be defined by $B_T \triangleq \{u \in B^* \mid T \text{ is productive from } u\}$ and let R_T be the relation on B_T defined by $R_T(u, u') \triangleq \exists b (u \star b = u')$. The relation R_T is serial by construction: for u such that T is productive from u, there is a such that T is productive from $u \star a$ and $u \star a \in T$. Also, $\langle \rangle \in B_T$ as soon as T is productive.

We can now formally state the correspondence in our language:

Theorem 2 As schemes, $DC_{BT}^{\text{productive}}$ and $DC_{BRb_0}^{\text{serial}}$ are logically equivalent.

PROOF: From left to right, we take $R^{\triangleright}_{\top}(b_0)$ and use Proposition 11. From right to left, we take B_T and R_T , obtaining $\langle \rangle \in B_T$ from T productive. We get an infinite branch β in $(B_T)^{\mathbb{N}}$ from which we get an infinite branch α in $B^{\mathbb{N}}$ by setting $\alpha(n) \triangleq b$ knowing that $\beta(n+1)$ has necessarily the form $\beta(n) \star b$. Any $u \prec_s \alpha$ happens to coincide with $\beta(|u|)$ which is in T because $u \in B_T$ implies T being productive from u.

As a final remark, let us mention countable Zorn's lemma [27]: If a partial order S on a countable set has no countable chain, it has a maximal element. It corresponds to the instantiation on $\neg S$ of the generalisation of the scheme $R^*_{\perp}(b_0)$ barred implies R has a least element over all b_0 , using our definitions up to classical reasoning, and dropping the partial order requirement. This is the case because a least element is a maximal one in the complement of a relation and because, classically, the barring of all antichainings of $\neg S$ is the same as the absence of countable chains in a partial order S.

2.8 Relation to countable choice

In Table 9, we introduce definitions allowing to state in Table 10 the axiom of countable choice, CC, and its dual, which we call *weak bar induction*.

We shall prove that CC is derivable from $DC^{productive}$ and introduce for that the *alignment* of a sequential relation over $\mathbb{N} \times A$ as a predicate over A^* (see Table 11). We have:

Theorem 3 For B and R given, CC_{BR} is equivalent to $DC_{BR_{T}}^{\text{productive}}$. Dually, WBI_{BR} is equivalent to $BI_{BR_{\perp}}^{\text{ind}}$.

ill-foundedness-style	well-foundedness-style	
Intensional concepts		
sequential positive alignment of R $(R_{\top}^{\mathbb{N}})$	sequential negative alignment of R $(R_{\perp}^{\mathbb{N}})$	
λu . case u of	λu . case u of	
$\left[\begin{array}{cc} \langle \rangle & \mapsto \top \\ u \star b \mapsto R(u ,b) \end{array} \right]$	$egin{bmatrix} \langle angle & \mapsto oldsymbol{ar{L}} \ u \star b \mapsto R(u ,b) \end{bmatrix}$	

Table 11: Logically opposite dual concepts on dual relations

PROOF: The correspondence between R left-total and $R_{\perp}^{\mathbb{N}}$ productive is obtained by coinduction from left to right and, from right to left, by extracting the n^{th} element of the proof of $R_{\perp}^{\mathbb{N}}$ to get the image of n by R. The correspondence between R choosable (as a relation) and $R_{\perp}^{\mathbb{N}}$ choosable (as a predicate on B^*) is obtained as follows. If α is the witness of R choosable, we define an infinite branch by $\alpha'(n) \triangleq (n, \alpha(n))$. Conversely, if α' is an infinite branch, we define $\alpha(n) \triangleq b$ whenever $\alpha'(n)$ is necessarily of the form (n, b). Similarly for the dual case.

We do not conversely expect to be able in general to express $DC^{productive}$ in term of CC since countable choice is strictly weaker than dependent choice, and similarly for Bl^{ind} in terms of WBI. However, if *B* is countable, the statements of DC and CC become (notoriously) mutually expressible using classical-reasoning-based minimisation: their common strength as choice principle then is not greater than the axiom of unique choice. The latter itself is a tautology if functions are represented as functional relations. It has however the logical effect of reifying functional relations as proper functions if functions are represented as proper objects in a functional type. We conjecture however that the equivalence of BI and WBI with countable codomain is provable intuitionistically.

3 Non sequential generalisation of dependent choice and bar induction

In the previous section, we considered predicates branching countably many times over a domain B. In this section, we investigate how to generalise countable sequences of branchings to branching in an arbitrary order over a non-necessarily countable domain A.

When B is \mathbb{B} , we shall obtain principles equivalent to the Boolean Prime Ideal/Filter Theorem (ill-founded case), or to the Completeness Theorem but we shall recover the strength of dependent choice (ill-founded case) and bar induction (well-founded case) when A is countable. In particular we will obtain the strength of the Weak Fan Theorem (well-founded case) and Weak König's Lemma (ill-founded case), up to classical reasoning, when A is countable and B is \mathbb{B} .

For a certain instance, we will get the strength of the full axiom of choice. However, the new principle is limited. For instance, for $A \triangleq \mathbb{B}^{\mathbb{N}}$ and $B \triangleq \mathbb{N}$, we end up with an inconsistent axiom.

3.1 Finite approximations of functions

Let A be a domain whose elements are ranged over by the letters a, a', ... and B a codomain whose elements are ranged over by the letters b, b', ... Let T be a predicate over $(A \times B)^*$ i.e. over sequences of pairs in A and B, thought as a set of possible finite approximations of a function from A to B. We order $(A \times B)^*$ by set inclusion, which we write \subset . We overload the notations $\downarrow^- T$ and $\uparrow^+ T$ to now respectively mean the downwards closure under restriction and upwards closure under extension with respect to \subset . In particular, since $v \subset v'$ for any v' obtained from v by permutation or duplication, $\downarrow^- T$ and $\uparrow^+ T$ are stable by permutation. We write $v \sim v'$ for $v \leq v'$ and $v' \leq v$, i.e. for the equivalence of vand v' as finite sets.

Note that we do not prevent that a sequence may contain several occurrences of the same pair (a, b). However, such a sequence shall be equivalent to a sequence without redundancies (this design choice is somewhat arbitrary, we just found it more convenient not to enforce the absence of redundancies).

We use v to range over approximations. We write $(a, b) \in v$ to mean that (a, b) is one of the elements of the sequence. For $v \in (A \times B)^*$, we write dom(v) for the set of a such that there is some b such that $(a, b) \in v$. For $\alpha \in A \to B$ and $v \in (A \times B)^*$, we define $v \prec \alpha$ to mean $\alpha(a) = b$ for all $(a, b) \in v$, or more formally for the predicate defined by the following clauses:

$$\frac{v \prec \alpha \qquad \alpha(a) = a}{v \star (a, b) \prec \alpha}$$

We think of $(A \times B)^*$ as finite approximations of functions from A to B and of predicates over finite approximations as constraints generating an ideal or a filter.

In Table 12, we generalise the notion of productive over (morally) trees into a coinductive notion of A-B-approximable relative to a valid finite set of approximations, and dually, we generalise the notion of inductively barred from holding on a sequence to holding relative to a finite set of approximations.

3.2 Generalised Dependent Choice and Generalised Bar Induction

We state the generalisation of dependent choice and bar induction to non-sequential choices over a non-necessarily countable domain in Table 13. Called GDC_{ABT} (shortly GDC_{AB} or GDC as schemes) and GBI_{ABT} (shortly GBI_{AB} or GBI as schemes), they are generalisations in the sense that they respectively capture $DC^{productive}$ and BI^{ind} for countable A.

ill-foundedness-style	$well\-foundedness\-style$	
Intensional concepts		
T A-B-approximable from v	T inductively A - B -barred from v	
$\nu X.\lambda v. \begin{pmatrix} v \in \downarrow^{-} T \land \\ \forall a \notin dom(v) \\ \exists b (v \star (a, b) \in X) \end{pmatrix}$	$\mu X.\lambda v. \begin{pmatrix} v \in \uparrow^+ T \lor \\ \exists a \notin dom(v) \\ \forall b (v \star (a, b) \in X) \end{pmatrix}$	
T A-B-approximable	T inductively A - B -barred	
T A-B-approximable from $\langle\rangle$	T inductively $A\text{-}B\text{-}\mathrm{barred}$ from $\langle\rangle$	
Extensional concepts		
T has an A - B -choice function	T is A - B -barred	
$\exists \alpha \forall u (u \prec \alpha \Rightarrow u \in T)$	$\forall \alpha \exists u (u \prec \alpha \land u \in T)$	

Table 12: Logically opposite dual concepts on dual predicates

Table 13: Dual axioms on dual predicat
--

ill-founded ness-style	well-foundedness-style	
Generalised Dependent Choice (GDC_{ABT})	Generalised Bar Induction (GBI_{ABT})	
T A-B-approximable \Rightarrow T has an A-B-choice function	$T A$ - B -barred $\Rightarrow T$ inductively A - B -barred	

To prove it, let us assume without loss of generality that A is \mathbb{N} itself. We say that $v \in (\mathbb{N} \times B)^*$ is sequential whenever either v is empty or v has the form $v' \star (|v'|, b)$ with v' itself sequential. To each $u \in B^*$ we can associate a sequential element ord(u) by $ord(\emptyset) \triangleq \emptyset$ and $ord(u \star b) \triangleq ord(u) \star (|u|, b)$. We then extend \subset from a relation on $(\mathbb{N} \times B)^*$ to an heterogeneous relation on $(\mathbb{N} \times B)^*$ and B^* by writing $v \subset u$ to mean $v \subset ord(u)$.

To each T over $(\mathbb{N} \times B)^*$, we can associate ||T|| on B^* by $u \in ||T|| \triangleq ord(u) \in T$. Conversely, to each T over B^* , we can associate \widehat{T} on $(\mathbb{N} \times B)^*$ defined by $v \in \widehat{T} \triangleq \exists u \in T (v = ord(u))$. We have:

Proposition 12 $u \in T$ iff $u \in ||\widehat{T}||$

PROOF: Direct by definition.

Proposition 13 T is \mathbb{N} -B-approximable iff ||T|| is productive, and, \hat{T} is \mathbb{N} -B-approximable iff T is productive. Dually, T is inductively \mathbb{N} -B-barred iff ||T|| is inductively barred, and, \hat{T} is inductively \mathbb{N} -B-barred iff T is inductively barred.

PROOF: By duality and Proposition 12, it is enough to prove the first item. The proof is by coinduction in both directions. From left to right, we prove $T \mathbb{N}$ -B-approximable from ord(u) implies ||T|| productive from u. We take |u| for a in the

definition of \mathbb{N} -*B*-approximable from ord(u), get some *b* and pass it to the definition of ||T|| productive from *u*. From right to left, we prove more generally that if ||T|| is productive from *u* then *T* is \mathbb{N} -*B*-approximable from *v* for all $v \subset u$. By definition of $u \in ||T||$, we have $ord(u) \in T$ and thus $v \in \downarrow^{-} T$ by closure of *T*. Now, take $n \notin dom(v)$. If n < |u|, we set *b* to be u(n) and apply the coinduction hypothesis with *v* extended with *b*, which still satisfies $v \star b \subset u$ by a combinatorial argument. If $n \ge |u|$, we explore the proof of productivity of ||T|| one step further, getting some *b* such that $u \star b \in ||T||$ and ||T|| is productive from $u \star b$. The property $v \subset (u \star b)$ continues to hold and we reason by induction on |u| - n until falling into the first case.

Similarly, we have:

Proposition 14 $\downarrow^- T$ has an \mathbb{N} -B-choice function iff ||T|| has an infinite branch, and, $\downarrow^- \widehat{T}$ has a \mathbb{N} -B-choice function iff T has an infinite branch. Dually, $\uparrow^+ T$ is barred iff ||T|| is B-barred, and, $\uparrow^+ \widehat{T}$ is \mathbb{N} -B-barred iff T is barred.

ill-foundedness style	well-foundedness-style		
Intensional concepts			
positive alignment of $R(R_{\top})$	negative alignment of R (R_{\perp})		
$\lambda u.$ case u of	$\lambda u.$ case u of		
$\left[\begin{array}{c} \langle \rangle & \mapsto \top \\ u \star (a,b) \mapsto R(a,b) \end{array} \right]$	$\left[\begin{array}{c} \langle \rangle & \mapsto \bot \\ u \star (a,b) \mapsto R(a,b) \end{array} \right]$		

Table 14: Logically opposite dual concepts on dual homogeneous relations

Table 15: The axiom of the Choice and its dual

ill-foundedness-style	well-foundedness-style	
Standard Axiom of Choice (AC_{ABR})	Dual to Standard Axiom of Choice $(co-AC_{ABR})$	
R left-total $\Rightarrow R$ has an A - B -choice function	$R A$ - B -barred $\Rightarrow R$ has a least element	

PROOF: By duality and Proposition 12, it is enough to prove the first item. From left to right, if $u \prec_s \alpha$, it is enough to consider $ord(u) \prec \alpha$. From right to left, if $v \prec \alpha$, we consider $u \triangleq \alpha_n$ where n is |v|. We have $u \prec_s \alpha$ thus $u \in T$ and $ord(u) \in T$. Since $v \subset ord(u)$, we get $v \in \downarrow^- T$.

Consequently, we have:

Proposition 15 $DC_B^{\text{productive}}$ iff $GDC_{\mathbb{N}B}$ and Bl_B^{ind} iff $GBl_{\mathbb{N}B}$.

PROOF: We mediate by the property that $GDC_{\mathbb{N}BT}$ is equivalent as a scheme to "T A-B-approximable implies $\downarrow^{-} T$ has an A-B-choice function". In one direction this is the case because $\downarrow^{-} T \subset T$, in the other direction, because \downarrow^{-} is idempotent. The other equivalence holds by duality.

3.3 Inconsistency of the unconstrained form of Generalised Dependent Choice and Generalised Bar Induction

In its full generality, the generalisation of GDC and GBI obtained by allowing non-countable branchings over an arbitrary codomain B is inconsistent: for large enough A and B, it may happen that some T is coinductively A-B-approximable without T having a (full) A-B-choice function. Indeed, take $A \triangleq \mathbb{B}^{\mathbb{N}}$ and $B \triangleq \mathbb{N}$ and filter the choice function so that it is injective. That is, we define $u \in T$ as follows: if u is $\langle \rangle$ or u is a singleton then $u \in T$ holds; if u is $u' \star (f, n) \star (f', n')$ with f extensionally different from f' then $n \neq n'$.

Then, T is coinductively $\mathbb{B}^{\mathbb{N}}$ - \mathbb{N} -approximable by successively extending u with (f, |u|) for any f not already in dom(u). But there is no total choice function α from $\mathbb{B}^{\mathbb{N}}$ to \mathbb{N} , since, by Cantor's theorem, such a function is necessarily noninjective. Thus, taking f and f' distinct such that $n \triangleq \alpha(f) = \alpha(f')$, we get that the sequence $(f, n), (f', n) \prec_n$ is not in T.

Therefore, we have:

Proposition 16 $GDC_{\mathbb{B}^N\mathbb{N}}$ and $GBl_{\mathbb{B}^N\mathbb{N}}$ are inconsistent.

3.4 Relation to the general axiom of choice

We state the standard axiom of choice in Table 15 and prove that it is equivalent to an instance of the generalised dependent choice GDC. To do so, we generalise in Table 14 the notion of sequential alignment introduced in Section 2.8 to the notion of alignment of a relation on $A \times B$ as a predicate over $(A \times B)^*$.

Theorem 4 AC_{ABR} is logically equivalent to $GDC_{ABR_{\top}}$

Table 16: Dual axioms on dual predicates

ill-foundedness-style	well-foundedness-style	
Generalised Weak König's Lemma $(GDC_{A\mathbb{B}T})$	Generalised Weak Fan Theorem $(GBI_{A\mathbb{B}T})$	
$T A$ - \mathbb{B} -approximable $\Rightarrow T$ has an A - \mathbb{B} -choice function	$T A$ -B-barred $\Rightarrow T$ inductively A -B-barred	

PROOF: The proof is similar to the one of Theorem 3. It consists in first showing that any partial choice function α is actually total for a left-total R. Indeed, for $a \in A$, there is a b such that R(a,b) by left-totality. Thus, from any u such that $u \in R_{\top}$ and $u \star (a, b) \notin R_{\top}$, we get a contradiction by definition of R_{\top} .

4 The Boolean instances of generalised dependent choice and bar induction: relation to the Boolean Prime Ideal/Filter Theorem and completeness theorems

4.1 Generalised Weak König Lemma and Generalised Weak Fan Theorem

By instantiating the codomain B to \mathbb{B} in GDC and GBI, we obtain extensions $\text{GBI}_{A\mathbb{B}}$ of the Weak Fan Theorem (more precisely of $\text{BI}_{\mathbb{B}}^{\text{ind}}$) and $\text{GDC}_{A\mathbb{B}}$ of the Weak König Lemma (more precisely of $\text{DC}_{\mathbb{B}}^{\text{productive}}$) which replace the countable sequence of branching made on a "tree" (in practise predicates) by a countable sequence of choices in arbitrary order over a non-necessarily countable domain. This shall be proved classically equivalent to the Boolean Prime Ideal/Filter Theorem. This is consistent with the standard reverse mathematics results which show that the completeness theorem is equivalent to the Weak König's Lemma on countable theories [24] but equivalent to the Boolean Prime Filter Theorem on theories of arbitrary cardinality [13, 14, 18, 11].

4.2 Logical reading: relation to completeness theorem

We can give a logical reading to $(A \times \mathbb{B})^*$ as follows. We call *atom* any element of A. We interpret pairs in $A \times \mathbb{B}$ as *literals*, i.e. as atoms together with a polarity indicating whether the atom is negated or not (we adopt the convention that 0 stands for non-negated and 1 for negated). We call *clause* any unordered sequence of elements in $A \times \mathbb{B}$. We call *context* any unordered sequence of elements of A. We range over clauses by the letters C, D and over contexts by the letters Γ , Δ , ...

Any clause C can canonically be represented as a pair of two contexts Γ and Δ with Γ the subset of non-negated elements of A in C and Δ the subset of negated elements. We write $\Gamma \triangleright \Delta$ for such a pair. We call a set of clauses a *theory* and use the letter \mathcal{T} to range over theories. We write $(\Gamma \triangleright \Delta) \in \mathcal{T}$ to mean that there is a clause of \mathcal{T} associated to the pair $\Gamma \triangleright \Delta$. We write $\Gamma \triangleright \Delta$ to mean that Γ and Δ have an atom in common.

We consider (a variant of) Scott's notion of entailment relation [23], i.e. of a preorder relation up to "side contexts". Let \mathcal{T} be a theory on A. We define the *entailment relation* generated by \mathcal{T} to be the relation on sequents, written $\Gamma \vdash_{\mathcal{T}} \Delta$, such that the following holds:

$$\frac{\Gamma \, \emptyset \, \Delta}{\Gamma \vdash_{\mathcal{T}} \Delta} \mathsf{A} \mathsf{x} \quad \frac{(\Gamma \triangleright \Delta) \in \uparrow^{+} \mathcal{T}}{\Gamma \vdash_{\mathcal{T}} \Delta} \mathsf{A} \mathsf{x}_{T} \quad \frac{\Gamma \vdash_{\mathcal{T}} \Delta, F \quad \Gamma, F \vdash_{\mathcal{T}} \Delta}{\Gamma \vdash_{\mathcal{T}} \Delta} \mathsf{Cut}$$

It is usual to add an explicit weakening rule to the definition of entailment relation but here we shall consider it as an admissible rule. Formally, the existence of a derivation of $\Gamma \vdash_{\mathcal{T}} \Delta$ using the inferences rules above is the same as

$$\vdash_{\mathcal{T}} \triangleq \mu X.\lambda(\Gamma \triangleright \Delta). \left(\begin{array}{c} (\Gamma \triangleright \Delta) \in \uparrow^+ \mathcal{T} \\ \forall \exists F \notin (\Gamma \cup \Delta) \left(\begin{array}{c} (\Gamma, F \triangleright \Delta) \in X \\ \land (\Gamma \triangleright \Delta, F) \in X \end{array} \right) \end{array} \right)$$

Thus, $\Gamma \vdash_{\mathcal{T}} \Delta$ exactly says that \mathcal{T} is inductively A-B-barred from $\Gamma \triangleright \Delta$.

Conversely, let us consider $\Gamma \not\vdash_{\mathcal{T}} \Delta$. We could define it by negation of $\Gamma \vdash_{\mathcal{T}} \Delta$ but we instead give a direct explicit definition which we call positive disprovability. Let \mathcal{T}^C denotes the complement of \mathcal{T} . The positive disprovability $\Gamma \not\vdash_{\mathcal{C}} \Delta$

$\it ill-foundedness-style$	well-foundedness-style		
Intensional concepts			
\mathcal{T} is (positively) consistent	\mathcal{T} is inconsistent		
$ \forall au$	$\vdash_{\mathcal{T}}$		
Extensional concepts			
\mathcal{T} has a model	${\cal T}$ is unsatisfiable		
$\exists \alpha \alpha \vDash \mathcal{T} \qquad \qquad \forall \alpha \alpha \not\vDash \mathcal{T}$			

Table 17: Logically opposite dual concepts of logic on the same predicate

Table 18: Reformulation of Table 16 as statements about a given logical theory

$\it ill$ -foundedness-style	well-foundedness-style	
Existence-style Completeness Theorem $(Compl^-(A\mathcal{T}))$	Provability-style Completeness Theorem $(Compl^+(A\mathcal{T}))$	
\mathcal{T} consistent $\Rightarrow \mathcal{T}$ has a model	\mathcal{T} unsatisfiable $\Rightarrow \mathcal{T}$ inconsistent	

can be characterised as the \mathcal{T}^C A-B-approximability from $\Gamma \triangleright \Delta$, that is, formally:

$$(\Gamma \triangleright \Delta) \in \nu X. \lambda(\Gamma \triangleright \Delta). \left(\begin{array}{c} (\Gamma \triangleright \Delta) \in \downarrow^{-} \mathcal{T}^{C} \\ \land \forall F \notin (\Gamma \cup \Delta) \begin{pmatrix} (\Gamma, F \triangleright \Delta) \in X \\ \lor (\Gamma \triangleright \Delta, F) \in X \end{pmatrix} \right) \right)$$

Let α be a function from A to \mathbb{B} . It can be interpreted as a model over A with 1 to indicate that the atom is true in the model and 0 to indicate that the atom is false in the model.

Truth $\alpha \models \mathcal{T}$ of a theory \mathcal{T} in a model α can be defined by

$$\alpha \vDash \mathcal{T} \triangleq \forall (\Gamma \triangleright \Delta) \in \mathcal{T} (\Gamma \subset \alpha \Rightarrow \Delta \not) \alpha)$$

where we use the notation $\Gamma \subset \alpha$ to mean that $\forall a \in \Gamma \alpha(a) = 1$ and the notation $\Delta \ \alpha$ to mean $\neg \forall a \in \Delta \alpha(a) = 0$. Then, \mathcal{T} has a model if there exists α such that $\alpha \models \mathcal{T}$.

Falsity $\alpha \not\models \mathcal{T}$ of a theory \mathcal{T} in a model α can be defined explicitly by

$$\alpha \not\models \mathcal{T} \triangleq \exists (\Gamma \triangleright \Delta) \in \mathcal{T} \Gamma \subset \alpha \land \Delta \subset \overline{\alpha}$$

where $\Delta \subset \overline{\alpha}$ stands for $\forall a \in \Delta \alpha(a) = 0$. The theory \mathcal{T} is *unsatisfiable* if for all $\alpha, \alpha \not\models \mathcal{T}$.

Then, we get that \mathcal{T} A-B-barred corresponds exactly to the unsatisfiability of \mathcal{T} . Also, noticing that, $\exists \alpha \forall u (u \prec \alpha \Rightarrow u \in \mathcal{T}^C)$ is linearly equivalent to $\exists \alpha \forall u (u \in \mathcal{T} \Rightarrow \neg u \prec \alpha)$ and that $\neg u \prec \alpha$ is linearly equivalent to $\Gamma \subset \alpha \Rightarrow \Delta \land \alpha$, we get that \mathcal{T}^C has an A-B-choice function if and only if there exists a model for \mathcal{T} (see Table 17 where $\vdash_{\mathcal{T}}$ and $\not\vdash_{\mathcal{T}}$ refer to the provability and positive disprovability of the empty clause).

The completeness theorem of logic is conventionally expressed either as the existence of a model for any consistent theory, or contrapositively, that if a theory is unsatisfied in all theories, then it is inconsistent, as shown on Table 18.

For instance, see Rinaldi, Schuster and Wessel [22] for the statement of a completeness theorem such as $\text{Compl}^+(\mathcal{T})$, up to the identification of some \exists with $\neg \neg \exists$. See also e.g. [21] for an algebraic reading.

Summing up, we have:

Generalised Weak König's Lemma over \mathcal{T}^C , i.e. $\mathsf{GDC}_{A\mathbb{B}\mathcal{T}^C}$, logically reads as $\mathsf{Compl}^-(A\mathcal{T})$

Generalised Weak Fan Theorem over \mathcal{T} , i.e. $\mathsf{GBl}_{A\mathbb{B}\mathcal{T}}$, logically reads as $\mathsf{Compl}^+(A\mathcal{T})$

where it should be reminded that, to preserve the duality, we used in $\mathsf{Compl}^-(\mathcal{T})$ an explicit definition of $\Gamma \not\vdash_{\mathcal{T}} \Delta$ which is classically equivalent to but intuitionistically stronger than the negation of $\Gamma \vdash_{\mathcal{T}} \Delta$.

ill-foundedness-style	well-foundedness-style
Boolean Prime Filter Theorem ($BPF_{\mathcal{B}}(F)$ for F a filter)	"Boolean Full Filter Theorem" (co- $BPF_{\mathcal{B}}(F)$ for F a filter)
F proper $\Rightarrow F$ extensible into a prime filter	F not extensible into a prime filter $\Rightarrow F$ full
Boolean Prime Ideal Theorem $(BPI_{\mathcal{B}}(I) \text{ for } I \text{ an ideal})$	"Boolean Full Ideal Theorem" (co- $BPI_{\mathcal{B}}(I)$ for I an ideal)
I proper \Rightarrow I extensible into a prime ideal	I not extensible into a prime ideal $\Rightarrow I$ full

Table 19: Reformulation of Table 16 as statements about a given Boolean algebra

4.3 Algebraic reading: relation to the Boolean Prime Ideal/Filter Theorem

The previous reasoning based on entailment relations can also be expressed in terms of Boolean algebras, connecting Generalised Weak König's Lemma to the Boolean Prime Ideal/Filter Theorem. There is however a caveat: the standard definition of proper filter and proper ideal relies on an intrinsically classical definition, so that the relation holds only with a classical reading of the connectives involved in the definition of approximable.

Let $(\mathcal{B}, \lor, \land, \bot, \neg, \neg)$ be a Boolean algebra and \vdash the canonical order relation associated to it $(b \vdash b' \triangleq b \land b' = b)$. We call *filter* over \mathcal{B} any non-empty subset F of \mathcal{B} which is closed under \land and closed under \vdash on the right. A filter is *proper* if it does not contain \bot . Otherwise, it coincides with \mathcal{B} and we call it *full*. We call *ultrafilter* a maximal proper filter. A maximal filter in a Boolean algebra can be described as a map U from \mathcal{B} to \mathbb{B} such that $b_1 \land b_2 \in U$ iff $b_1 \in U \land b_2 \in U$, $b_1 \lor b_2 \in U$ iff $b_1 \in U \lor b_2 \in U$, $\neg b \in U$ iff $\neg(b \in U)$, $\neg \in U$, and $\bot \notin U$. In a Boolean algebra, the notion of maximal filter coincides with the notion of prime filter where a filter F is *prime* if $(b_1 \lor b_2) \in F$ implies $b_1 \in F$ or $b_2 \in F$.

Dually, we call *ideal* over \mathcal{B} any non-empty subset I of \mathcal{B} which is closed under \lor and closed under \vdash on the left. An ideal is *proper* if it does not contain \top , and *full* otherwise. A *prime* ideal I is such that $(b_1 \land b_2) \in I$ implies $b_1 \in I$ and $b_2 \in I$ and this coincides with the notion of maximal proper ideal. A prime/maximal proper ideal can be characterised the same way as a prime/maximal proper filter, i.e. as a map U from \mathcal{B} to \mathbb{B} such that $b_1 \land b_2 \in U$ iff $b_1 \in U \land b_2 \in U$, $\neg b \in U$ iff $\neg (b \in U)$, except that this time $\bot \in U$ and $\top \notin U$.

There is a family of provably equivalent theorems about the existence of maximal/prime ideals/filters in Boolean algebras (see e.g. Jech [18, 2.3]) called Boolean Prime Ideal Theorem in arbitrary Boolean algebras, or Ultrafilter Theorem in the Boolean algebras of subsets of a set. We consider in Table 19 the case of a general Boolean algebra and state the Boolean Prime Ideal Theorem in its two "ideal" and "filter" flavours. We also consider their contrapositive.

We now show that the Boolean Prime Ideal and Prime Filter Theorems and Generalised Weak König's Lemma, i.e. $GDC_{A\mathbb{B}}$, are mutual instances of each other. We first show that the Generalised Weak König's Lemma is an instance of the Boolean Prime Ideal and Boolean Prime Filter Theorems.

To any domain A we can associate a freely generated Boolean algebra ($\mathsf{Free}(A), \dot{\lor}, \dot{\land}, \dot{\perp}, \dot{\top}, \dot{\neg}$) by considering the set of algebraic expressions built from $\dot{\lor}, \dot{\land}, \dot{\perp}, \dot{\top}$ and $\dot{\neg}$, all quotiented by the axioms of a Boolean algebra, namely by:

$a \lor (b \lor c)$	$= (a \lor b) \lor c$	$a \land (b \land c)$	$= (a \land b) \land c$
$a \stackrel{.}{\lor} b$	$=b \stackrel{.}{\lor} a$	$a \stackrel{.}{\wedge} b$	$=b \stackrel{.}{\wedge} a$
$a\stackrel{.}{\vee}(a\stackrel{.}{\wedge}b)$	= a	$a\mathrel{\dot{\wedge}} (a\mathrel{\dot{\vee}} b)$	= a
$a \lor \perp$	= a	$a \land \dot{\top}$	= a
$a\stackrel{,}{\vee}(b\stackrel{,}{\wedge}c)$	$= (a \stackrel{.}{\vee} b) \stackrel{.}{\wedge} (a \stackrel{.}{\vee} c)$	$a\mathrel{\dot{\wedge}} (b\mathrel{\dot{\vee}} c)$	$= (a \stackrel{.}{\wedge} b) \stackrel{.}{\vee} (a \stackrel{.}{\wedge} c)$
$a \lor \dot{\neg} a$	$=\dot{\top}$	$a\mathrel{\dot{\wedge}}\dot{\neg} a$	$=$ \perp

As in the previous section, any v in $(A \times \mathbb{B})^*$, can be written under the form $\Gamma \triangleright \Delta$ and two possible interpretations in $\mathsf{Free}(A)$ are now possible. The first one is to interpret a sequent $\Gamma \triangleright \Delta$ as the formula $(\bigvee \neg \Gamma) \lor (\bigvee \Delta)$ and to interpret any predicate over $(A \times \mathbb{B})^*$ seen as a logical theory \mathcal{T} as the corresponding subset \mathcal{T}^{\vdash} of formulas of this form in $\mathsf{Free}(A)$. It is then standard that the set $F_{\mathcal{T}}$ of logical consequences of \mathcal{T}^{\vdash} defined by $b \in F_{\mathcal{T}} \triangleq \mathcal{T}^{\vdash} \vdash b$ is a filter.

There is also a dual interpretation of $\Gamma \triangleright \Delta$ as the formula $(\bigwedge \Gamma) \land (\bigwedge \neg \Delta)$ leading to interpret \mathcal{T} as a "theory" of consequences, say $\vdash \mathcal{T}$. Then, the set $I_{\mathcal{T}}$ of logically stronger assumptions of $\vdash \mathcal{T}$ defined by $b \in I_{\mathcal{T}} \triangleq b \vdash \vdash \mathcal{T}$ is an ideal.

This can be rephrased in the following statement where $\overline{\mathcal{T}}$ swaps the role of 0 and 1 in \mathcal{T} , i.e. $v \in \overline{\mathcal{T}} \triangleq \overline{v} \in \mathcal{T}$, where \overline{v} itself turns each (a, 0) in v into (a, 1) and vice-versa.

For \mathcal{T} a theory and $\vdash_{\mathcal{T}}$ the entailment relation canonically built from it, it is standard that the set of logical consequences of \mathcal{T}^{\vdash} in Free(A) forms a filter and the set of logically stronger assumptions of $\vdash_{\mathcal{T}}$ in Free(A) forms an ideal. In particular,

the filter is proper if $\mathcal{T}^{\vdash} \not\vdash \bot$, that is if \mathcal{T} is consistent, that is, by Section 4.2, if \mathcal{T}^{C} is A-B-approximable where the connectives involved in the definition of A-B-approximable are interpreted classically. Similarly, the ideal is proper if $\top \not\vdash^{\vdash} \mathcal{T}$, that is if $\overline{\mathcal{T}}$ is consistent, that is if $\overline{\mathcal{T}}^{C}$ is A-B-approximable where, again, the connectives are interpreted classically.

Similarly, we have the general result that prime filters and prime ideals on a free Boolean algebra, here Free(A), are characterised by their intersection with generators, here A. Whether other elements of Free(A) belong or not to a prime filter or prime ideal is canonically determined⁶ by:

$\alpha(a \lor a') = b''$	≜	$\alpha(a) = b \land \alpha(a') = b$	$b' \wedge b'' = b + b'$
$\alpha(a \stackrel{.}{\wedge} a') = b''$	\triangleq	$\alpha(a) = b \land \alpha(a') = b$	$b' \wedge b'' = b \cdot b'$
$\alpha(\perp) = b$	\triangleq	b = 0	for ideals
$\alpha(\dot{\top}) = b$	\triangleq	b = 1	for ideals
$\alpha(\dot{\perp}) = b$	\triangleq	b = 1	for filters
$\alpha(\dot{\top}) = b$		b = 0	for filters
$\alpha(\dot{\neg} a) = b'$	\triangleq	$\alpha(a) == b \wedge b' = -b$	

where $+, \cdot, -$ are the corresponding operations on \mathbb{B} .

In particular, the existence of a function from A to \mathbb{B} characterising a prime filter that extends the filter $F_{\mathcal{T}}$ on $\mathsf{Free}(A)$ is the same, by Section 4.2, as a model of \mathcal{T} and as an A- \mathbb{B} -choice function for \mathcal{T} . Similarly, the existence of a function from A to \mathbb{B} characterising a prime ideal that extends the ideal $I_{\mathcal{T}}$ on $\mathsf{Free}(A)$ is the same as a model of $\overline{\mathcal{T}}$ and as an A- \mathbb{B} -choice function for $\overline{\mathcal{T}}$.

Consequently, we get the following:

Theorem 5 $GDC_{A\mathbb{B}\mathcal{T}^C}$, where the connectives are interpreted classically, is equivalent to $BPF_{Free(A)}(F_{\mathcal{T}})$ and $BPI_{Free(A)}(I_{\overline{\mathcal{T}}})$. On its side, $GBI_{A\mathbb{B}\mathcal{T}}$ is equivalent to co- $BPF_{Free(A)}(F_{\mathcal{T}})$ and co- $BPI_{Free(A)}(F_{\overline{\mathcal{T}}})$, even with a linear or intuitionistic interpretation of the connectives.

5 Further questions

While we did not explicitly state and check it, we suspect that most of our proofs are essentially expressible in a contractionfree logic such as relevant logic or a linear logic without contraction [12], to the notable exception of Proposition 6. Might such a careful check be the purpose of future work.

There is a rich literature on choice axioms and on principles equivalent to choice axioms. Not all of them can be classifyed as either ill- or barred/well-foundedness-style, though. For instance, open induction and update induction [20, 8, 6], are classically equivalent to bar induction and dependent choice but are formulated as well-foundedness of some order on functions.

The study could also be extended to stronger choice principles such as Zorn's lemma, the ordinal variants of the axiom of dependent choices by Lévy [19] and the ordinal variants of Zorn's lemma [27] by Wolk.

Acknowledgments

We thank the communities of researchers who contributed to develop the material we built on, and in particular Camille Noûs, from the Cogitamus Lab, who embodies the collective and collaborative nature of scientific research.

The ideas in Section 4.3 derived from investigations led by Charlotte Barot [1]. The second author thanks Valentin Blot and Étienne Miquey for numerous fruitful discussions on the axiom of dependent choice and bar induction.

⁶We define the value of α as equations to remain agnostic on the representation of a function to \mathbb{B} , see 2.2.

References

- [1] Charlotte Barot. Two [choice] principles involved in the intuitionistic completeness theorem. Master thesis, University Paris-Diderot, 2017.
- [2] Josef Berger. The fan theorem and uniform continuity. In Proceedings of the First International Conference on Computability in Europe: New Computational Paradigms, CiE'05, page 18–22, Berlin, Heidelberg, 2005. Springer-Verlag.
- [3] Josef Berger. A decomposition of Brouwer's fan theorem. J. Logic & Analysis, 1, 2009.
- [4] Josef Berger and Hajime Ishihara. Brouwer's fan theorem and unique existence in constructive analysis. Mathematical Logic Quarterly, 51(4):360–364, 2005.
- [6] Ulrich Berger. A computational interpretation of open induction. In 19th IEEE Symposium on Logic in Computer Science (LICS 2004), 14-17 July 2004, Turku, Finland, Proceedings, page 326. IEEE Computer Society, 2004.
- [7] Crystal Cheung. Brouwer's Fan Theorem: An Overview. Master thesis, Cornell University, 2015.
- [8] Thierry Coquand. A note on the open induction principle. Chalmers University, 1997.
- [9] Thierry Coquand. About brouwer's fan theorem. Revue Internationale de Philosophie, 230:483–489, 2004.
- [10] Thierry Coquand and Henri Lombardi. A logical approach to abstract algebra. Mathematical Structures in Computer Science, 16(5):885-900, 2006. See also http://www.cse.chalmers.se/~coquand/formal.html.
- [11] Christian Espíndola. Semantic completeness of first-order theories in constructive reverse mathematics. Notre Dame J. Formal Logic, 57(2):281–286, 2016.
- [12] Jean-Yves Girard. Linear logic. Theor. Comput. Sci., 50:1–102, 1987.
- [13] Leon Henkin. The completeness of the first-order functional calculus. J. Symb. Log., 14(3):159–166, 1949.
- [14] Jean E. Rubin Herman Rubin. Equivalents of the Axiom of Choice. Elsevier, 1963.
- [15] William A. Howard and Georg Kreisel. Transfinite induction and bar induction of types zero and one, and the role of continuity in intuitionistic analysis. J. Symb. Log., 31(3):325–358, 1966.
- [16] Hajime Ishihara. Constructive reverse mathematics: Compactness properties. From Sets and Types to Topology and Analysis: Towards Practicable Foundations for Constructive Mathematics, 48, 10 2005.
- [17] Hajime Ishihara. Weak König's Lemma Implies Brouwer's Fan Theorem: A Direct Proof. Notre Dame J. Formal Logic, 47(2):249–252, 04 2006.
- [18] Thomas J. Jech. The Axiom of Choice. Dover Books on Mathematics Series. Courier corporation, 1973.
- [19] Azriel Lévy. The interdependence of certain consequences of the axiom of choice. Fundamenta Mathematicae, 54(2):135–157, 1964.
- [20] Jean-Claude Raoult. Proving open properties by induction. Information Processing Letters, 29(1):19–23, 1988.
- [21] Davide Rinaldi and Peter Schuster. A universal Krull-Lindenbaum theorem. Journal of Pure and Applied Algebra, 220(9):3207 - 3232, 2016.
- [22] Davide Rinaldi, Peter Schuster, and Daniel Wessel. Eliminating disjunctions by disjunction elimination. The Bulletin of Symbolic Logic, 23(2):181–200, 2017.
- [23] Dana Scott. Completeness and axiomatizability. In Proceedings of the Tarski symposium, pages 411–435, 1974.
- [24] Stephen G. Simpson. Subsystems of Second Order Arithmetic. Perspectives in Logic. Cambridge University Press, 2nd edition, 2009.

- [25] Richard E. Vesley Stephen Kleene. The foundations of intuitionistic mathematics. North-Holland, 1965.
- [26] Wim Veldman. Brouwer's fan theorem as an axiom and as a contrast to kleene's alternative. Arch. Math. Log., 53(5–6):621–693, August 2014.
- [27] Elliot S. Wolk. On the principle of dependent choices and some forms of Zorn's lemma. Canad. Math. Bull., 26(3), 1983.